

The vascular flora of Tarsia (Calabria, southern Italy)

Lorenzo Peruzzi¹

¹ PLANTSEED Lab, Department of Biology, University of Pisa, Pisa, Italy

Corresponding author: Lorenzo Peruzzi (lorenzo.peruzzi@unipi.it)

Academic editor: Fabrizio Bartolucci | Received 7 January 2026 | Accepted 3 February 2026 | Published 12 February 2026

Citation: Peruzzi L (2026) The vascular flora of Tarsia (Calabria, southern Italy). Italian Botanist 21: 35–46. <https://doi.org/10.3897/italianbotanist.21.184523>

Abstract

A list of the vascular flora occurring in the municipality of Tarsia (province of Cosenza, Calabria) is provided. The list is based on a bibliographic analysis and on field studies carried out in the years 2018–2025. A total of 618 specific and subspecific taxa are recorded in the study area (including 64 aliens), plus 35 cultivated taxa. *Chenopodium ficifolium* Sm. and *Orobanche cernua* Loebl. are new records for Calabria.

Keywords

Alien species, biodiversity, endemics, floristic data, Italy, phytogeography

Introduction

A flora is a useful source of information for biogeographical, ecological and evolutionary studies (Peruzzi 2018; D’Antraccoli et al. 2022b; Bartolucci et al. 2024), so that floristic inventories are crucial to provide suitable data for decision-making processes in biodiversity conservation and landscape planning, either in urban (Peruzzi et al. 2025) or more natural (Carta et al. 2018; Gestri et al. 2023; Pierini and Peruzzi 2025) areas.

The territory in the municipality of Tarsia (Calabria, southern Italy) has received little attention from botanists in the past, except for the southern portion where the northern shores of Lake Tarsia are located, an artificial basin created in 1959. For this area, considered of low floristic knowledge by Bernardo et al. (2005), only 41 floristic records have been published over time (Crisafulli et al. 2010; Maiorca et al. 2013) before this study.

The aim of this study is to present a complete and updated floristic inventory of the vascular flora of Tarsia, in the framework of a series of contributions whose goal is to improve the floristic knowledge of Calabria (La Valva and Ricciardi 1978; La Valva 1984; Ballelli and Venanzoni 1993; Maiorca and Spampinato 1994; Bernardo and Maiorca 1996; Maiorca et al. 2002, 2007; Musarella and Tripodi 2004; Maiorca and Puntillo 2009, 2015; Bernardo et al. 2011, 2012; Di Marco et al. 2013; Roma-Marzio et al. 2015; Spampinato et al. 2019; Caruso et al. 2022; Spampinato et al. 2022; Capuano and Caruso 2023; Capuano et al. 2024; Laface et al. 2024; Mauro et al. 2024; Musarella et al. 2024).

Material and methods

Study area

The territory of the municipality of Tarsia (province of Cosenza) covers an area of 48.28 km² in northern Calabria, lower Crati Valley, at elevations between 40 and 370 m a.s.l. (Fig. 1). The vegetation of the area reflects a Mediterranean bioclimatic context; a small portion in the northeast belongs to the southern edaphically indifferent series of downy oak (*Quercus virgiliana* (Ten.) Ten., a current synonym of *Q. pubescens* Willd. subsp. *pubescens*); the southern portion bordering the Crati River belongs to the central-southern peninsular hygrophilous geosigmetum of lowland and riparian vegetation; the remaining territory is included in the southern Apennine Tyrrhenian acidophilous series of downy oak (*Q. virgiliana*) (Bernardo et al. 2010a, 2010b). Most of the territory is devoted to agriculture, mainly olive groves and vineyards, with *Citrus* orchards to a much lesser extent. Spontaneous vegetation is therefore confined to a few relict patches, resulting in low environmental variability and high degree of human disturbance.

Methods

Besides the analysis of the few literature sources available (see Introduction), field investigations were carried out in the last 20 years, more intensively in the years 2018–2025. The most significant species were reported during this multi-year research (Puntillo and Peruzzi 2008; Gangale et al. 2010; Peruzzi et al. 2010; Astuti et al. 2015, 2021; Peruzzi et al. 2016; Bartolucci et al. 2017; Roma-Marzio et al. 2018; Galasso et al. 2020, 2021; Laface et al. 2020; Sáez et al. 2022). For the most significant taxa, herbarium specimens (PI, herbarium acronyms follow Thiers 2026) were prepared and/or photographic documentation was produced. In some cases, seeds were collected and deposited in the seed bank of the Botanic Garden and Museum of the University of Pisa. To outline the entire vascular flora, extensive use was made of field observations, all archived and freely accessible in the online database Wikiplantbase #Italia (Peruzzi et al. 2019 onwards). These data were supplemented by information derived from herbarium specimens preserved in CLU, FI, and NAP.




Figure 1. Localisation and delimitation of the municipality of Tarsia (province of Cosenza, Calabria, southern Italy); the dominance of cultivated and urbanized areas is evident.

Nomenclature and circumscription of the taxa follows Bartolucci et al. (2024), Galasso et al. (2024) and their updates periodically appearing in the Portal to the Flora of Italy (<https://dryades.units.it/floritaly/index.php>; Martellos et al. 2020). Angiosperm families are arranged according to APG IV (2016). Within families, genera, species and subspecies are listed in alphabetical order. Life forms and chorotypes were attributed according to Pignatti et al. (2017a, 2017b, 2018), not considering casual aliens and cultivated species. We also highlighted those taxa included in the National Red List (Rossi et al. 2013, 2021; Orsenigo et al. 2021). Chorotypes were simplified as follows: Eurosiberian, Eurosiberian-Mediterranean, Mediterranean, wide distribution, Italian endemic, alien, as in Peruzzi (2023). The endemism status was based on Peruzzi et al. (2014, 2015). The complete dataset assembled for the present study is available in Suppl. material 1.

The expected number (S) of entire and alien taxa occurring in the study area (A) were calculated using Arrhenius' species-area relationship $S = cA^z$, and using the c (241.2 and 10.1 for the entire and alien taxa, respectively) and z (0.281 and 0.404 for entire and alien taxa, respectively) coefficients empirically calculated for the floristic richness of Italy by D'Antraccoli et al. (2024).

At the end of the study, the method proposed by D'Antraccoli et al. (2022a) was applied to obtain a final Map of Relative Floristic Ignorance. This allows a proper planning of possible further floristic investigations of the area. All the floristic records (10,726) available in Peruzzi et al. (2019 onwards) for the province of Cosenza were used.

Results

The complete floristic inventory is reported in the Suppl. material 1. The expected number of species/subspecies was 803, of which 38 aliens. A total of 618 specific and subspecific taxa were documented for the study area, including 64 aliens (21 casual, 25 naturalized, 18 invasive).

The available records for the municipality of Tarsia amount to 6,821, more than 95% of which are based on field observations, the remaining part being constituted by herbarium (148) and photographic (135) documentation. The recorded flora consists of 618 spontaneous species/subspecies (64 of which are alien) and 35 cultivated taxa, belonging to 401 genera and 99 families. It should be noted that some species reported in the literature for the Tarsia area are, strictly speaking, actually present in adjacent areas (among the most relevant examples: *Fritillaria messanensis* Raf. subsp. *messanensis*).

The most represented families are Asteraceae (90 taxa), Poaceae (66 taxa) and Fabaceae (61 taxa), while the richest genera are *Trifolium* (18 taxa), *Euphorbia* (10 taxa), *Silene* (8 taxa), *Ophrys* and *Plantago* (7 taxa each).

From a chorological perspective, the Mediterranean geographic element is predominant (43%), followed by widely distributed (18.5%) and Eurosiberian taxa (15.5%), with 10% of taxa having a distribution spanning both the Mediterranean and Eurosiberian regions. Italian endemic species account for 2.8% of the flora, while alien species represent 10.2%. Regarding life forms, therophytes prevail (39.2%), followed by hemicryptophytes (28.6%), geophytes (14.7%), phanerophytes (12.8%), chamaephytes (3.1%), and hydrophytes (1.6%).

Among the taxa of particular phytogeographical and conservation interest are the Italian endemics *Artemisia campestris* L. subsp. *variabilis* (Ten.) Greuter, *Crocus biflorus* Mill., *Daucus broteroi* Ten., *Euphorbia corallioidea* L., *E. meuslii* Geltman, *Gelasia villosa* (Scop.) Cass. subsp. *columnae* (Guss.) Bartolucci, Galasso & F.Conti, *Gypsophila arrostoi* Guss. subsp. *arrostoi*, *Micromeria consentina* (Ten.) N.Terracc., *Onobrychis alba* (Waldst. & Kit.) Desv. subsp. *echinata* (G.Don) P.W.Ball, *Ophrys incubacea* Bianca subsp. *brutia* (P.Delforge) Kreutz, *Ornithogalum exscapum* Ten., *Picris scaberrima* Guss. (described from the Tarsia area; see Astuti et al. 2015; Fig. 1), *Salvia haematodes* L., *Silene echinata* Otth, *Stipa austroitalica* Martinovský, *Thalictrum calabricum* Spreng., and *Visnaga crinita* (Guss.) Giardina & Raimondo. Unfortunately, the latter species is locally extinct and never found again after 1827 (NAP-GUSS!).

The taxa with the highest frequency of observation in the area are: *Oloptum thomasii* (Duby) Banfi & Galasso (162), *Artemisia campestris* subsp. *variabilis* (155 records), *Daucus carota* L. subsp. *carota* (140), *Avena barbata* Pott ex Link (139), *Dittrichia viscosa* (L.) Greuter subsp. *viscosa* (127), *Anethum foeniculum* L. (117), *Cynodon dactylon* (L.) Pers. (108), *Picris hieracioides* L. subsp. *hieracioides* (99), *Quercus pubescens* Willd. subsp. *pubescens* (95), and *Ferula communis* L. subsp. *communis* (86) among native species; *Erigeron canadensis* L. (81), *Arundo donax* L. (77), *Opuntia ficus-indica* (L.) Mill. (73), *Sorghum halepense* (L.) Pers. (70), and *Robinia pseudoacacia* L. (44) among alien species; *Olea europaea* L. (111),

Prunus amygdalus Batsch (54), *Eucalyptus camaldulensis* Dehnh. (34), *Vitis vinifera* L. (28), and *Nerium oleander* L. subsp. *oleander* (8) among cultivated taxa.

Four species documented by herbarium specimens have not been confirmed, and could be possibly extinct at the local level: besides the above-mentioned *Visnaga crinita*, *Euphorbia aleppica* L. (CLU!) *Lotus angustissimus* L. (NAP-GUSS!), and *Trifolium ligusticum* Balb. ex Loisel. (NAP-GUSS!).

Discussion

The species richness of the established flora (native plus naturalized and invasive aliens) is 25% below expectations, while aliens are 11% below the expected. Both these data could be explained by the low environmental variability and high degree of human disturbance of the study area. The low number of species could, in part, also be due to undersampling in some areas. The Map of Relative Floristic Ignorance highlights the cells that are best known on floristic grounds (Fig. 2). The cells near the northern and south-western border of the municipality show the highest relative floristic ignorance, and more intensive studies in these areas may change the picture.

On phytogeographical grounds, the area clearly shows a Mediterranean affinity, considering both chorological and biological spectra, with 43.3% of Mediterranean taxa and a H/T ratio of 0.73.

Among the taxa recorded for the study area, 34 are included in the Red List of the Italian Flora (Rossi et al. 2013, 2021; Orsenigo et al. 2021): 21 were assessed as Least Concern, 8 as Data Deficient (among which the Italian endemics *Euphorbia meuselii*, *Gypsophila arrostoi* subsp. *arrostoi*, and *Picris scaberrima*), 4 as Near Threatened (*Artemisia campestris* subsp. *variabilis*, *Onobrychis alba* subsp. *echinata*, *Silene echinata*, *Squilla pancration* Steinh.), 1 as Vulnerable (*Dracunculus vulgaris* Schott), 1 as Endangered (*Fritillaria messanensis* subsp. *messanensis*), and 1 as Critically Endangered (Possibly Extinct) (*Visnaga crinita*).

There are also species that are rare in Calabria, such as *Lophiolepis scabra* (Poir.) Del Guacchio, Bureš, Iamónico & P.Caputo, *Isoetes duriei* Bory, *Myosurus minimus* L., *Notobasis syriaca* (L.) Cass., *Ornithogalum refractum* Kit. ex Willd., *Ranunculus sceleratus* L., and *Spirodela polyrhiza* (L.) Schleid., as well as species otherwise typically associated with coastal environments, such as *Artemisia arborescens* (Vaill.) L. and *Atriplex halimus* L., both occurring on the cliffs below the town of Tarsia. *Chenopodium ficifolium* Sm. and *Orobanche cernua* Loeffl. are new records for the region.

Finally, concerning aliens, noteworthy is the presence of 18 invasive alien taxa, some of which are particularly widespread. Among alien taxa, two are included in the list of invasive alien species of Union concern under the European regulation UE 1143/2014 and the application regulations UE 2016/1141, 2017/1263, 2019/1262, 2022/1203, and 2025/1422: *Ailanthus altissima* (Mill.) Swingle and *Reynoutria japonica* Houtt., the latter so far only cultivated.




Figure 2. Map of Relative Floristic Ignorance of the municipality of Tarsia (province of Cosenza, Calabria, southern Italy), computed with a cellsize = 1000 m and a temporal coefficient tau = 20. The colour scale represents values of Index of Relative Floristic Ignorance (IRFI), ranging from 0 (blue) to a maximum value (red) representing the cells showing lowest and highest relative ignorance, respectively. Coordinates are projected in EPSG 3035: ETRS89/ETRS-LAEA.

Acknowledgements

The author wishes to thank: Liliana Bernardo (Università della Calabria) for kindly providing data on the available bibliography for the area and on the herbarium specimens preserved in CLU; Annalisa Santangelo (Università di Napoli “Federico II”) for kindly providing data on herbarium specimens preserved in NAP-GUSS; Giannantonio Domina (Università di Palermo) for the identification of *Orobanchae*; Giuseppe Antonelli and Marco D’Antraccoli (Università di Pisa) for the preparation of the Map of Relative Floristic Ignorance and of Tarsia, respectively. The financial support by Fondazione per la Flora Italiana for the publication fee is also gratefully acknowledged.

References

Angiosperm Phylogeny Group (2016) An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG IV. Botanical Journal of the Linnean Society 181(1): 1–20. <https://doi.org/10.1111/boj.12385>

- Astuti G, Barone G, Di Gristina E, Domina G, Giacò A, Orsenigo S, Peruzzi L (2021) Chromosome numbers for the Italian flora: 11. *Italian Botanist* 11: 145–153. <https://doi.org/10.3897/italianbotanist.11.70173>
- Astuti G, Roma-Marzio F, Peruzzi L (2015) The genus *Picris* (Asteraceae) in southern Italy: contribution to its systematic knowledge. *Phytotaxa* 207(1): 106–114. <https://doi.org/10.11646/phytotaxa.207.1.5>
- Ballelli S, Venanzoni R (1993) Contributo alla conoscenza della flora e della vegetazione di un settore del Monte Gariglione (Parco Nazionale della Calabria “Sila Piccola”). *Atti 5° Workshop Progetto Strategico Clima, Ambiente e Territorio nel Mezzogiorno (I)*. Amalfi, 1993, 28–30.
- Bartolucci F, Domina G, Adorni M, Alessandrini A, Ardenghi NMG, Banfi E, Baragliu GA, Bernardo L, Bertolli A, Biondi E, Carotenuto L, Casavecchia S, Cauzzi P, Conti F, Crisanti MA, D’Amico FS, Di Cecco V, Di Martino L, Faggi G, Falcinelli F, Forte L, Galasso G, Gasparri R, Ghillani L, Gottschlich G, Guzzon F, Harpke D, Lastrucci L, Lattanzi E, Maiorca G, Marchetti D, Medagli P, Olivieri N, Pascale M, Passalacqua NG, Peruzzi L, Picollo S, Prosser F, Ricciardi M, Salerno G, Stinca A, Terzi M, Viciani D, Wagensommer RP, Nepi C (2017) Notulae to the Italian native vascular flora: 3. *Italian Botanist* 3: 29–48. <https://doi.org/10.3897/italianbotanist.3.13200>
- Bartolucci F, Peruzzi L, Galasso G, Alessandrini A, Ardenghi NMG, Bacchetta G, Banfi E, Barberis G, Bernardo L, Bouvet D, Bovio M, Calvia G, Castello M, Cecchi L, Del Guacchio E, Domina G, Fascetti S, Gallo L, Gottschlich G, Guarino R, Gubellini L, Hofmann N, Iberite M, Jiménez-Mejías P, Longo D, Marchetti D, Martini F, Masin RR, Medagli P, Peccenini S, Prosser F, Roma-Marzio F, Rosati L, Santangelo A, Scoppola A, Selvaggi A, Selvi F, Soldano A, Stinca A, Wagensommer RP, Wilhalm T, Conti F (2024) A second update to the checklist of the vascular flora native to Italy. *Plant Biosystems* 158(2): 219–296. <https://doi.org/10.1080/11263504.2024.2320126>
- Bernardo L, Bartolucci F, Cancellieri L, Costalonga S, Galasso G, Galesi R, Gargano D, Iberite M, Iocchi M, Lattanzi E, Lavezzo P, Magrini S, Peccenini S, Sciandrello S, Scoppola A, Signorino G, Tilia A, Spampinato G (2012) Contributo alla conoscenza floristica della Calabria: resoconto dell’escursione del Gruppo di Floristica (S.B.I) nel 2008 nella Presila Catanzarese. *Informatore Botanico Italiano* 44(1): 125–151.
- Bernardo L, Gargano D, Passalacqua NG, Peruzzi L, Spampinato G (2005) Carta dello stato delle conoscenze floristiche della Calabria. In: Scoppola A, Blasi C (Eds) *Stato delle conoscenze sulla flora vascolare d’Italia*. Palombi editori, Roma, 197–201.
- Bernardo L, Maiorca G (1996) The vascular flora of the Cassano Structural High (NE Calabria, Italy). *Giornale Botanico Italiano* 130(2–3): 531–556. <https://doi.org/10.1080/11263509609430323>
- Bernardo L, Passalacqua NG, Spampinato G (2010a) Le serie di vegetazione della regione Calabria. In: Blasi C (Ed.) *La vegetazione d’Italia*. Palombi & Partner S.r.l., Roma, 411–428.
- Bernardo L, Passalacqua NG, Peruzzi L (Eds) (2011) *Flora vascolare della Calabria – Prodrómo*. *Informatore Botanico Italiano* 43(2): 185–332.
- Bernardo L, Passalacqua NG, Spampinato G (2010b) Carta delle serie di vegetazione della regione Calabria. In: Blasi C (Ed) *La vegetazione d’Italia, Carta delle Serie di Vegetazione, scala 1:500.000*. Palombi & Partner S.r.l., Roma.

- Capuano A, Montepaone G, Lazzeri V, Caruso G (2024) *Salsola tragus* L. (Amaranthaceae) and *Thymelaea gussonei* Boreau (Thymelaeaceae): new records for the native vascular flora of Calabria (S-Italy) with some remarks about their taxonomy and distribution. Research Journal of Ecology and Environmental Sciences 4(1): 61–79. <https://doi.org/10.31586/rjees.2024.883>
- Capuano A, Caruso G (2023) New records for the alien vascular flora of Calabria (S-Italy). Research Journal of Ecology and Environmental Sciences 3(2): 1–35. <https://doi.org/10.31586/rjees.2023.723>
- Carta A, Forbicioni L, Frangini G, Pierini B, Peruzzi L (2018) An updated inventory of the vascular flora of Elba island (Tuscan Archipelago, Italy). Italian Botanist 6: 1–22. <https://doi.org/10.3897/italianbotanist.6.26568>
- Caruso G, Casavecchia S, Biondi E (2022) The vascular flora of Uria basin (Catanzaro, S-Italy) and its conservation relevance. Flora Mediterranea 32: 99–116. <https://doi.org/10.7320/FlMedit32.099>
- Crisafulli A, Cannavò S, Maiorca G, Musarella CM, Signorino G, Spampinato G (2010) Aggiornamenti floristici per la Calabria. Informatore Botanico Italiano 42(2): 431–442.
- D’Antraccoli M, Bedini G, Peruzzi L (2022a) Maps of relative floristic ignorance and virtual floristic lists: an R package to incorporate uncertainty in mapping and analysing biodiversity data. Ecological Informatics 67: 101512. <https://doi.org/10.1016/j.ecoinf.2021.101512>
- D’Antraccoli M, Bedini G, Peruzzi L (2022b) Next Generation Floristics: a workflow to integrate novel methods in traditional floristic research. Plant Biosystems 156(2): 594–597. <https://doi.org/10.1080/11263504.2022.2056650>
- D’Antraccoli M, Peruzzi L, Conti F, Galasso G, Roma-Marzio F, Bartolucci F (2024) Floristic Richness in a Mediterranean Hotspot: A Journey across Italy. Plants 13(1): 12. <https://doi.org/10.3390/plants13010012>
- Di Marco F, Bernardo L, Peruzzi L (2013) Contribution to the vascular flora of Papasidero (north-western Calabria, Italy). Atti della Società Toscana di Scienze Naturali, Memorie, serie B 119(2012): 33–50.
- Galasso G, Conti F, Peruzzi L, Alessandrini A, Ardenghi NMG, Bacchetta G, Banfi E, Barberis G, Bernardo L, Bouvet D, Bovio M, Calvia G, Castello M, Cecchi L, Del Guacchio E, Domina G, Fascetti S, Gallo L, Guarino R, Gubellini L, Guiggi A, Hofmann N, Iberite M, Jiménez-Mejías P, Longo D, Marchetti D, Martini F, Masin RR, Medagli P, Musarella CM, Peccenini S, Podda L, Prosser F, Roma-Marzio F, Rosati L, Santangelo A, Scoppola A, Selvaggi A, Selvi F, Soldano A, Stinca A, Wagensommer RP, Wilhalm T, Bartolucci F (2024) A second update to the checklist of the vascular flora alien to Italy. Plant Biosystems 158: 297–340. <https://doi.org/10.1080/11263504.2024.2320129>
- Galasso G, Domina G, Adorni M, Angiolini C, Apruzzese M, Ardenghi NMG, Assini S, Aversa M, Bacchetta G, Banfi E, Barberis G, Bartolucci F, Bernardo L, Bertolli A, Bonali F, Bonari G, Bonini I, Bracco F, Brundu G, Buccomino G, Buono S, Calvia G, Cambria S, Castagnini P, Ceschin S, Dagnino D, Di Gristina E, Di Turi A, Fascetti S, Ferretti G, Fois M, Gentili R, Gheza G, Gubellini L, Hofmann N, Iamónico D, Ilari A, Király A, Király G, Laface VLA, Lallai A, Lazzaro L, Lonati M, Longo D, Lozano V, Lupoletti J, Magrini S, Mainetti A, Manca M, Marchetti D, Mariani F, Mariotti MG, Masin RR, Mei G, Menini

- F, Merli M, Milani A, Minuto L, Mugnai M, Musarella CM, Olivieri N, Onnis L, Pas-salacqua NG, Peccenini S, Peruzzi L, Pica A, Pinzani L, Pittarello M, Podda L, Prosser F, Ravetto Enri S, Roma-Marzio F, Rosati L, Sarigu M, Scafidi F, Sciandrello S, Selvaggi A, Spampinato G, Stinca A, Tavilla G, Toffolo C, Tomasi G, Turcato C, Villano C, Nepi C (2020) Notulae to the Italian alien vascular flora: 9. *Italian Botanist* 9: 47–70. <https://doi.org/10.3897/italianbotanist.9.53401>
- Galasso G, Domina G, Andreatta S, Argenti E, Bacchetta G, Bagella S, Banfi E, Barberis D, Bardi S, Barone G, Bartolucci F, Bertolli A, Biscotti N, Bonali F, Bonini F, Bonsanto D, Brundu G, Buono S, Caldarella O, Calvia G, Cambria S, Campus G, Caria MC, Conti F, Coppi A, Dagnino D, Del Guacchio E, Di Gristina E, Farris E, Ferretti G, Festi F, Fois M, Furlani F, Gigante D, Guarino R, Gubellini L, Hofmann N, Iamónico D, Jiménez-Mejías P, La Rosa A, Laface VLA, Lallai A, Lazzaro L, Lonati M, Lozano V, Luchino F, Lupoletti J, Magrini S, Mainetti A, Marchetti D, Marenzi P, Marignani M, Martignoni M, Mei G, Menini F, Merli M, Mugnai M, Musarella CM, Nicoletta G, Noor Hussain A, Olivieri N, Orlandini S, Peccenini S, Peruzzi L, Pica A, Pilon N, Pinzani L, Pittarello M, Podda L, Probo M, Prosser F, Raffaelli C, Ravetto Enri S, Riviaccio G, Rosati L, Sarmati S, Scafidi F, Selvi F, Sennikov AN, Sotgiu Cocco G, Spampinato G, Stinca A, Tavilla G, Tomaselli V, Tomasi D, Tomasi G, Trenchi M, Turcato C, Verloove F, Viciani D, Villa M, Wagensommer RP, Lastrucci L (2021) Notulae to the Italian alien vascular flora: 11. *Italian Botanist* 11: 93–119. <https://doi.org/10.3897/italianbotanist.11.68063>
- Gangale C, Uzunov D, Peruzzi L (2010) Notula: 39. *Veronica peregrina* L. subsp. *peregrina* (Plantaginaceae). In: Nepi C, Peccenini S, Peruzzi L (Eds) *Notulae alla flora esotica d'Italia* 2. *Informatore Botanico Italiano* 42(2): 533.
- Gestri G, Pierini B, D'Antraccoli M, Bernardini A, Peruzzi L (2023) An updated inventory of the vascular flora of the Cerbaie hills (Tuscany, Italy). *Italian Botanist* 15: 165–175. <https://doi.org/10.3897/italianbotanist.15.105302>
- Laface VLA, Musarella CM, Noto D, Siclari A, Tralongo S, Spampinato G (2024) The Peat Bogs of Aspromonte National Park (Southern Italy): Flora and Vegetation. In: Calabrò F, Madureira L, Morabito FC, Piñeira Mantiñán MJ (Eds) *Networks, Markets & People. NMP 2024. Lecture Notes in Networks and Systems*, vol 1185. Springer, Cham, 405–414. https://doi.org/10.1007/978-3-031-74672-7_40
- Laface VLA, Musarella CM, Spampinato G, Iamónico D, Noor Hussain A, Fortini P, Peruzzi L, Giardini M, Angeloni D, Meneguzzo E, Meneguzzo A, Banfi E, Busnardo G, Galasso G, Kleih M, Wallnöfer B, Bernardo L, Roma-Marzio F, Brentan M, Marcucci R (2020) Nuove segnalazioni floristiche italiane 9. *Flora vascolare* (67–78). *Notiziario della Società Botanica Italiana* 4(2): 229–233.
- La Valva V (1984) Flora e vegetazione dell'isola di Cirella. *Delpinoa* 23–24(1981–1982): 27–51.
- La Valva V, Ricciardi M (1978) Flora e vegetazione dell'isola di Dino. *Delpinoa* 18–19(1976–1977): 127–176.
- Maiorca G, Puntillo D (2009) The vascular flora of the “Bosco di Mavigliano”, a Site of Community Importance (SIC) in Northern Calabria (South-Italy). *Webbia* 64(1): 75–100. <https://doi.org/10.1080/00837792.2009.10670853>

- Maiorca G, Puntillo D (2015) Contributi alla conoscenza floristica della Catena Costiera (Calabria). 1. La forra della Fiumara dei Bagni. *Informatore Botanico Italiano* 47(2): 187–204.
- Maiorca G, Spampinato G (1994) The vascular flora of Argentino River valley, a nature reserve in NW Calabria (Italy). *Flora Mediterranea* 4: 49–100.
- Maiorca G, Spampinato G, Caprio A (2002) Flora e vegetazione dei laghi costieri La Vota (Calabria centro-occidentale). *Fitosociologia* 39(1): 81–108.
- Maiorca G, Spampinato G, Crisafulli A, Puntillo D, Caridi D, Cameriere P (2013) Carta della Biodiversità vegetale del Lago di Tarsia (Calabria settentrionale). Progetto PHYTOS.I.S., Monografia n. 4, ARSSA, Luigi Pellegrini ed. Cosenza.
- Maiorca G, Spampinato G, Crisafulli A, Cameriere P (2007) Flora vascolare e vegetazione della riserva regionale “Foce del Fiume Crati” (Calabria, Italia meridionale). *Webbia* 62(2): 121–174. <https://doi.org/10.1080/00837792.2007.10670821>
- Martellos S, Bartolucci F, Conti F, Galasso G, Moro A, Pennesi R, Peruzzi L, Pittao E, Nimis PL (2020) FlorItaly – the portal to the Flora of Italy. *Phytokeys* 156: 55–71. <https://doi.org/10.3897/phytokeys.156.54023>
- Mauro LR, Musarella CM, Laface VLA, Morabito A, Patti M, Cannavò S, Spampinato G (2024) Floristic Update of the SAC “Pentidattilo” (Calabria, Southern Italy). In: Calabrò F, Madureira L, Morabito FC, Piñeira Mantiñán MJ (Eds) *Networks, Markets & People. NMP 2024. Lecture Notes in Networks and Systems*, vol 1185. Springer, Cham, 415–424. https://doi.org/10.1007/978-3-031-74672-7_41
- Musarella CM, Laface VLA, Angiolini C, Bacchetta G, Bajona E, Banfi E, Barone G, Biscotti N, Bonsanto D, Calvia G, Cambria S, Capuano A, Caruso G, Crisafulli A, Del Guacchio E, Di Gristina E, Domina G, Fanfarillo E, Fascetti S, Fiaschi T, Galasso G, Mascia F, Mazzacuva G, Mei G, Minissale P, Motti R, Perrino EV, Picone RM, Pinzani L, Podda L, Potenza G, Rosati L, Stinca A, Tavilla G, Villano C, Wagensommer RP, Spampinato G (2024) New Alien Plant Taxa for Italy and Europe: An Update. *Plants* 13(5): 620. <https://doi.org/10.3390/plants13050620>
- Musarella CM, Tripodi G (2004) La flora della rupe e dei ruderi di Pentidattilo (Reggio Calabria). *Informatore Botanico Italiano* 36(1): 3–12.
- Orsenigo S, Fenu G, Gargano D, Montagnani C, Abeli T, Alessandrini A, Bacchetta G, Bartolucci F, Carta A, Castello M, Cogoni D, Conti F, Domina G, Foggi B, Gennai M, Gigante D, Iberite M, Peruzzi L, Pinna MS, Prosser F, Santangelo A, Selvaggi A, Stinca A, Villani M, Wagensommer RP, Tartaglini N, Duprè E, Blasi C, Rossi G (2021) Red list of threatened vascular plant species in Italy. *Plant Biosystems* 155(2): 310–335. <https://doi.org/10.1080/11263504.2020.1739165>
- Peruzzi L (2018) Floristic inventories and collaborative approaches: a new era for checklists and floras? *Plant Biosystems* 152(2): 177–178. <https://doi.org/10.1080/11263504.2017.1419997>
- Peruzzi L (2023) The vascular flora of Empoli (Tuscany, central Italy). *Italian Botanist* 15: 21–33. <https://doi.org/10.3897/italianbotanist.15.101748>
- Peruzzi L, Astuti G, Bartolucci F, Conti F, Roma-Marzio F (2016) Chromosome numbers for the Italian flora: 2. *Italian Botanist* 2: 29–42. <https://doi.org/10.3897/italianbotanist.2.10900>

- Peruzzi L, Bernardo L, Domina G (2010) Notula: 1698. *Orobanche schultzei* (Orobanchaceae). In: Nepi C, Peccenini S, Peruzzi L (Eds) Notulae alla Checklist della flora italiana: 10. Informatore Botanico Italiano 42(2): 515.
- Peruzzi L, Conti F, Bartolucci F (2014) An inventory of vascular plants endemic to Italy. Phytotaxa 168(1): 1–75. <https://doi.org/10.11646/phytotaxa.168.1.1>
- Peruzzi L, Domina G, Bartolucci F, Galasso G, Peccenini S, Raimondo FM, Albano A, Alessandrini A, Banfi E, Barberis G, Bernardo L, Bovio M, Brullo S, Brundu G, Brunu A, Camarda I, Carta L, Conti F, Croce A, Iamónico D, Iberite M, Iiriti G, Longo D, Marsili S, Medagli P, Pistarino A, Salmeri C, Santangelo A, Scassellati E, Selvi F, Soldano A, Stinca A, Villani M, Wagensommer RP, Passalacqua NG (2015) An inventory of the names of vascular plants endemic to Italy, their loci classici and types. Phytotaxa 196(1): 1–217. <https://doi.org/10.11646/phytotaxa.196.1.1>
- Peruzzi L, Pierini B, Arduini I, Bedini G, Franzoni J (2025) The vascular flora of Pisa (Tuscany, Central Italy). Plants 14(3): 307. <https://doi.org/10.3390/plants14030307>
- Peruzzi L, Roma-Marzio F, Pinzani L, Bedini G [Eds] (2019) [onwards] Wikiplantbase #Italia. <https://bot.biologia.unipi.it/wpb/index>
- Pierini B, Peruzzi L (2025) The vascular flora of the Apuan Alps (Tuscany, Central Italy). Italian Botanist 19: 45–63. <https://doi.org/10.3897/italianbotanist.19.145947>
- Pignatti S, Guarino R, La Rosa M (2017a) Flora d'Italia 1, Ed. 2. Edagricole di New Business Media, Milano, 1064 pp.
- Pignatti S, Guarino R, La Rosa M (2017b) Flora d'Italia 2, Ed. 2. Edagricole di New Business Media, Milano, 1178 pp.
- Pignatti S, Guarino R, La Rosa M (2018) Flora d'Italia 3, Ed. 2. Edagricole di New Business Media, Milano, 1286 pp.
- Puntillo D, Peruzzi L (2008) Notula: 1491. *Myosurus minimus* L. (Ranunculaceae). In: Nepi C, Peruzzi L, Scoppola A (Eds) Notulae alla checklist della flora italiana: 6. Informatore Botanico Italiano 40(2): 254–255.
- Roma-Marzio F, Bernardo L, Liguori P, Peruzzi L (2015) Vascular Flora of Monte Sparviere (Southern Italy, Pollino Massif). Atti della Società Toscana di Scienze Naturali, Memorie, serie B, 122: 73–88.
- Roma-Marzio F, Lastrucci L, Guzzon F, Ardenghi NMG, Peruzzi L, Mossini S (2018) Nuove segnalazioni floristiche italiane 5. Flora vascolare (28–46). Notiziario della Società Botanica Italiana 2(2): 205–210.
- Rossi G, Montagnani C, Gargano D, Peruzzi L, Abeli T, Ravera S, Cogoni A, Fenu G, Magrini S, Gennai M, Foggi B, Wagensommer RP, Venturella G, Blasi C, Raimondo FM, Orsenigo S [Eds] (2013) Lista Rossa della Flora Italiana. 1. Policy Species e altre specie minacciate. Comitato Italiano IUCN e Ministero dell'Ambiente e della Tutela del Territorio e del Mare, 54 pp.
- Rossi G, Orsenigo S, Gargano D, Montagnani C, Peruzzi L, Fenu G, Abeli T, Alessandrini A, Astuti G, Bacchetta G, Bartolucci F, Bernardo L, Bovio M, Brullo S, Carta A, Castello M, Cogoni D, Conti F, Domina G, Foggi B, Gennai M, Gigante D, Iberite M, Lasen C, Magrini S, Nicoletta G, Pinna MS, Poggio L, Prosser F, Santangelo A, Selvaggi A, Stinca A, Tartaglini N, Troia A, Villani MC, Wagensommer RP, Wilhelm T, Blasi C (2021) Lista

- Rossa della Flora Italiana. 2. Endemiti e altre specie minacciate. Ministero dell’Ambiente e della Tutela del Territorio e del Mare.
- Sáez L, Mesbah M, López-Alvarado J, Bacchetta G, El Mokni R, Peruzzi L, Oxelman B (2022) Re-establishment of *Silene neglecta* Ten. (Caryophyllaceae) with taxonomic notes on some related taxa. *PhytoKeys* 195: 143–160. <https://doi.org/10.3897/phytokeys.195.81370>
- Spampinato G, Laface VLA, Posillipo G, Cano Ortiz A, Quinto Canas R, Musarella CM (2022) Alien flora in Calabria (Southern Italy): an updated checklist. *Biological Invasions* 24: 2323–2334. <https://doi.org/10.1007/s10530-022-02800-y>
- Spampinato G, Sciandrello S, Giusso del Galdo G, Puglisi M, Tomaselli V, Cannavò S, Musarella CM (2019) Contribution to the knowledge of Mediterranean wetland biodiversity: Plant communities of the Aquila Lake (Calabria, Southern Italy). *Plant Sociology* 56(2): 53–68. <https://doi.org/10.7338/pls2019562/04>
- Thiers B (2026) Index Herbariorum: a global directory of public herbaria and associated staff. New York, Botanical Garden’s Virtual Herbarium. <https://sweetgum.nybg.org/science/ih/> [accessed on 7 January 2026]

Supplementary material I

Additional information

Authors: Lorenzo Peruzzi

Data type: pdf

Explanation note: 1. Floristic list and records. 2. References cited in the supplementary material.

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/italianbotanist.21.184523.suppl1>