

The alkaloids of *Corydalis hendersonii* Hemsl. contribute to the cardioprotective effect against ischemic injury in mice by attenuating cardiomyocyte apoptosis via p38 MAPK signaling pathways

fuxing ge

Beijing University of Chinese Medicine

xiaoli gao

Beijing University of Chinese Medicine

xioachun zhou

Beijing University of Chinese Medicine

junjun li

Beijing University of Chinese Medicine

xiaojing ma

Beijing University of Chinese Medicine

meiwen huang

Beijing University of Chinese Medicine

shana wuken

Beijing University of Chinese Medicine

pengfei tu

Beijing University of Chinese Medicine

chao an

Beijing University of Chinese Medicine

Xingyun Chai (✉ xingyunchai@yeah.net)

Beijing University of Chinese Medicine <https://orcid.org/0000-0003-1321-7543>

Research Article

Keywords: *Corydalis hendersonii* Hemsl., Tibetan folk medicine, Myocardial ischemia, Apoptosis, p38 MAPK signaling pathway, Alkaloids

Posted Date: October 7th, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-2110884/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Chinese Medicine on March 17th, 2023. See the published version at <https://doi.org/10.1186/s13020-023-00726-8>.

Abstract

Background: There is a characteristic Tibetan folk medicine in China named *Corydalis hendersonii* Hemsl. (CH) has been used for treatment of cardiovascular related diseases, called “plethora” in Tibetan medicine. Previous studies demonstrated that ethanol extract of CH shows anti-acute myocardial infarction (AMI) effect through inhibiting fibrosis and inflammation. Rich alkaloids fraction (RAF) is isolated from CH, but whether RAF possessing an equivalent effect with CH ethanol extract and by which mechanism it protects against AMI has not yet reported. The paper aimed to study whether RAF protects myocardial injury in mice and its underlying mechanism.

Material and methods: The cation exchange resin was used to obtain RAF and poor alkaloidal fraction (PAF). Liquid chromatography mass spectrometry-ion trap-time of flight (LCMS-IT-TOF) was used to analyze the chemical profile and isolate pure compounds. The ligation of left anterior descending (LAD) of coronary artery in mice was used to evaluate the in vivo anti-AMI effect, by dividing into eight groups: Sham, Model, Fosinopril (10 mg/kg, i.g.), total extract (TE, 400 mg/kg, i.g.), PAF (300 mg/kg, i.g.), and RAF (25, 50, and 100 mg/kg, respectively, i.g.) groups. Echocardiography was used to evaluate mice heart function through the index of left ventricular internal diameter (LVEDs), left ventricular internal diameter (LVEDd), fractional shortening (FS) and ejection fraction (EF). We detected the lactate dehydrogenase (LDH) and creatine kinase-MB (CK-MB) in the serum and the plasma level of angiotensin II (AngII). The apoptosis of mice myocardial tissue was verified by TUNEL assay. The expression of p38 mitogen-activated protein kinases (p38 MAPK), Bcl-2 and X Protein associated Bcl-2 (Bax) were detected through immunofluorescence staining, qRT-PCR and western blot in heart tissue and H9c2 cells.

Results: Echocardiography data indicated that the values of LVEDd and LVEDs were reduced and the values of FS and EF were improved by TE and RAF significantly. RAF also decreased the levels of LDH, CK-MB and AngII and significantly inhibited inflammatory cells in the marginal zone of myocardial infarction. The TUNEL assay results showed that RAF significantly attenuated cell apoptosis. Immunofluorescence and qRT-PCR assay showed that RAF inhibited p38 MAPK, Bax, and Bcl-2 proteins in mice myocardium. Western blot results validated that the expressions of key proteins were inhibited by RAF. Also, the apoptotic cells and apoptosis-related proteins were dramatically reduced by RAF in vivo and in vitro. Besides, RAF and PAF were analyzed by LCMS-IT-TOF to identify the main compounds and to demonstrate the difference between them. The results showed that a total of 14 alkaloids were identified, which indicated that the isoquinoline alkaloids were the main ingredients in RAF may contributing to the cardioprotective effect in mice.

Conclusions: RAF improves cardiac function by inhibiting apoptosis via p38 MAPK signaling pathways, and RAF contributed to the effect against myocardial ischemic injury of TE in mice, which provided a substantial reference for the clinical application against ischemia heart disease and quality control of CH.

Highlights

- The total extract of *Corydalis hendersonii* Hemsl. (CH) and the rich alkaloids fraction (RAF) of CH showed significant cardioprotective effect on mice following acute myocardial infarction.
- RAF improved cardiac function by inhibiting apoptosis via p38 mitogen-activated protein kinases (p38 MAPK) signaling pathways *in vitro* and *in vivo*.
- The results of liquid chromatography mass spectrometry-ion trap-time of flight (LCMS-IT-TOF) showed that the 14 isoquinoline alkaloids in RAF as the main ingredients contribute to the cardioprotective effect.

Introduction

Ischemic heart disease (IHD), caused by the imbalance between coronary blood flow and myocardial demand due to changes in the coronary circulation, has a high incidence and fatality rate [1, 2]. Although the drug development and treatment methods in this area have been developed to a certain extent, the mortality rate of IHD has been rising rapidly since 2005, and the mortality rate of IHD is higher than that in the rural areas, and correspondingly the research and development of drugs for cardiovascular diseases treatment has been vigorously developed at home and abroad and seeking drugs for the treatment of IHD including from traditional medicine resources increasingly became one of hotspots in the aspects [3–5].

Our group has been focusing on ethnomedicines in China used to cure acute myocardial infarction (AMI)-related IHD, for instance the *Syringa pinnatifolia* Hemsl. [6–8], and *Corydalis hendersonii* Hemsl. (CH) and *Meconopsis horridula* Hook. f. & Thomson [9, 10].

Corydalis hendersonii Hemsl. is one of the characteristic Tibetan folk medicine called “plethora” used to cure of high-altitude polycythemia (HAPC), hypertension and other diseases, which provides a new effective way for discovery of new therapeutic strategy against ischemic heart disease [11, 12]. Previously, the anti-AMI effect of ethanol total extract (TE) and underlying mechanism of anti-inflammation via JAK2-STAT3 and NF- κ B signaling pathway has been investigated in our group [10]. However, whether alkaloid fraction exerts the anti-AMI effect equivalently with TE and whether it is underlying another mechanism need to further explore. In the present study, the effect and mechanisms of rich alkaloids fraction (RAF) against ischemic injury were evaluated by ligating the left anterior descending (LAD) coronary artery in ICR mice and H9c2 cells under hypoxic conditions.

Material And Methods

Preparation of TE, RAF and PAF

The whole plant of *Corydalis hendersonii* Hemsl. (the plant name has been checked with <http://www.theplantlist.org>) was collected from Shannan Autonomous Prefecture of Tibet in July 2017. Prof. Yuan Zhang (Beijing University of Chinese Medicine, BUCM) authenticated CH. The voucher specimen of CH was No. CH201707 and was stored in BUCM. About 160 g of total extract (TE) was

obtained after extraction. Disperse the TE with hydrochloric acid aqueous solution and adjust the pH value to 2–3. Then the strong acid cation exchange resin is applied for separation and purification of alkaloids with the condition that passing through the hydrogen cation exchange resin column at the rate of 2 bed volumes per hour (BV/h). The resin column was flushed with water at the rate of 2 BV/h to neutral and the poor alkaloids fraction (PAF, 120 g) was obtained by collecting the combined eluent concentrated in vacuum. At last, the column was eluted at the rate of 2 BV/h to obtain rich alkaloids fraction (RAF, 30 mg) with 90% alkaline ethanol whose pH value at 10–11.

Lcms-it-tof

The research protocol was as the same with previous study [10]. The mobile phase (1.0 mL/min) consists of acetonitrile (A)–0.1% aqueous with formic acid (B) with a gradient program as follows: 0–15 min, 5–15% A; 16–50 min, 15–30% B; 51–80 min, 30–95% B; Roughly 20% portion of the effluent was introduced into the ESI (source by splitting the effluent via two polyetheretherketone tubes with length ratio of 1:4). The diode array detector (DAD) detector scans its absorption wavelength in the full wavelength range of 190–400 nm, so we set a 10 µL injection volume and 254 nm detection wavelength.

Experimental animal and ethics statement

The ICR mice (27–28 g) were obtained from Sabei Fu Biotechnology Co., Ltd. (Beijing, China) and reared in a specific pathogen-free (SPF) environment. The mice were housed with regular food and free water in a 12 h light-dark cycles, under conditions of $22 \pm 2^\circ\text{C}$. According to the previous reports, the AMI model of mice LAD ligation was used to evaluate the effect of TE, RAF and PAF [9]. The Ethics Committee of BUCM had approved the experiments (the ethics permit number: BUCM-3–2015090701-3003), and the research was implemented according to the US guidelines (NIH publication #85 – 23, revised in 1996).

Eight groups were randomly divided: sham, model, Fosinopril (10 mg/kg, i.g.), TE (400 mg/kg, i.g.), PAF (300 mg/kg, i.g.), RAF high dose (RAF-H, 100 mg/kg, i.g.), RAF medium dose (RAF-M, 50 mg/kg, i.g.), RAF low dose (RAF-L, 25 mg/kg, i.g.) groups. Fosinopril was used as the positive control. The drugs were dissolved in 0.5% CMC-Na and were orally administered once a day for 7 days. The control groups were only given solvent.

Echocardiography

The mice were administered drugs or solvent for 7 days and then isoflurane (Ward Life Science and Technology Co., Ltd., Shenzhen, China) was used to anesthetize for 2D M-mode and B-mode echocardiography ultrasonic. The levels of left ventricular internal diameter (LVEDd), ejection fraction (EF), left ventricular internal diameter (LVEDs) and fractional shortening (FS) were calculated for measurement of the cardiac function.

Sample collection

After the echocardiography, 0.5% pentobarbital sodium were used to anaesthetize the mice at 50 mg/kg by intraperitoneal injection, and then they were sacrificed for collecting blood samples and heart samples. The serum and plasma were obtained through centrifugation for 20 min at 3000 rpm and then stored at -80°C . The serum levels of lactate dehydrogenase (LDH) and creatine kinase-MB (CK-MB) were detected by Hitachi 17080 Automatic Biochemical Analyzer (Hitachi Co., Ltd., Japan). The plasma level of AngII was detected by ELISA. Heart was divided into two pieces along the ligature line and then the lower half was immediately put in liquid nitrogen for frozen, then it was stored at -80°C until testing. The left of the heart was prepared for paraffin section, which was placed in 4% paraformaldehyde for 72 h.

Histological examination

After 72 h fixing, cardiac tissues were cut into 4 μm thin slices and prepared for pathological examination. The paraffin sections were stained with haematoxylin and eosin (HE), masson stain and sirius red stain, and then were observed with microscope.

TUNEL assay

The cardiac tissue samples were embedded in paraffin and sectioned into 3 μm thick flakes to determine apoptotic cells. Paraffin sections were immersed two times with xylene for 5 min each and then putted in 100%, 95%, 90%, 80%, 70% ethanol solutions for 3 min, respectively. Then treated tissue with Proteinase K working solution for 15–30 min, at $21-37^{\circ}\text{C}$. After incubation with 2'-deoxyuridine 5'-triphosphate for 1 h at 37°C , the nucleus was counterstained with 4',6-Diamidino-2-phenylindole dihydrochloride (DAPI).

qRT-PCR

The TRIzol reagent (Gibco-BRL, Paisley, UK) was used to extract total RNA according to the manufacturer's protocols. The concentration of the samples was determined by Nano Drop 2000 (Thermo Scientific, USA). Subsequently, the PCR Mix kit (Transgen Biotech Co.) was used to amplify the mice tissue cDNA. Afterwards, the expression of mRNA was measured by qRT-PCR: 1 cycle at 94°C for 30 s and 40 cycles at 94°C for 5 s and 50°C for 30 s and then determined by melting curve. The primer sequences were showed as follows: p38 mitogen-activated protein kinases (p38 MAPK) F, CCTATCCTGGAAGAGCCATACT; p38 MAPK R, ACTTTGTCACGCTGACCAGAT; Bax F, AGACAGGGGCCTTTTTGCTAC, Bax R, AATTCGCCGGAGACACTCG; Bcl-2 F, GAGCCTGTGAGAGACGTGG, Bcl-2 R, CGAGTCTGTGTATAGCAATCCCA; GAPDH F, TATGACTCTACCCACGGCAAG, GAPDH R, TACTCAGCACCAGCATCACC.

Immunofluorescence detection of tissue

After deparaffinized and immersed in xylene, the paraffin slices were eluted with an alcohol gradient. The tissue slices were placed in EDTA (pH 8.0) for antigen retrieval. The interval of power-off after 8 min at medium heat to boiling is 16 min and then low heat for 7 minutes to boiling. The sections were incubated with anti-p38 MAPK (8690T, Cell Signaling Technology), anti-Bcl-2 (26593, Proteintech) and anti-Bax (50599, Proteintech) at 4°C overnight. And then incubated with goat anti-rabbit IgG (LP1001B, ABGENT) at room temperature for 50 min. Counter-stain the cell nucleus with DAPI and incubate for 10 min in the

dark at room temperature. After washing with PBS, the images were captured with the laser scanning confocal microscope (FV1000 Olympus, Japan).

Cell viability assay

H9c2 cells (China Infrastructure of Cell Line Resources, Beijing, China) was cultured under the condition of 37°C with 5% CO₂ and the medium was DMEM (Corning, Manassas, VA, USA) mixed with 10% FBS (Corning, Manassas, VA, USA). DMSO (Sigma, USA) was used to dissolve RAF and the initial concentration was 40 mg/mL. Firstly, the H9c2 cells were seeded in 96-well plates with the density of 8×10^3 cells/well for 24 h before incubating with RAF (1.25, 2.5, 5, 10, 20, 40, 80, 160, and 320 µg/mL) for 24 h. Secondly, we put the 96-well plates into the hypoxia box and added RAF at the concentrations of 0.63, 1.25, 2.5, 5, and 10 µg/mL for 8 h. Control group was cultured without the hypoxia box and other conditions are the same. Finally, each well was added 100 µL CCK-8 and incubated for 2 h, then a microplate reader was used to measure the OD value of the plates at 450 nm.

Immunofluorescence detection of cells

The H9c2 cells were seeded into 6-well plates and the density was 2×10^5 cells/mL under the condition of 37°C with 5% CO₂ for 24 h. After removed the supernatant, each well was added 500 µL of PBS and discarded. Then the cells were fixed in 4% paraformaldehyde and Hoechst 33258 was (1 µg/mL) was stained for 10 min in dark. Afterwards, the cells were washed with PBS and dried while protected from light. The fluorescence microscope (Leica Micro systems GmbH) was used to observe the apoptotic cells at 350 nm and 460 nm.

Western blot

Myocardial tissue (50 mg) and the H9c2 cells treated with hypoxia were harvested and fragmented in RIPA lysis buffer (Applygen Technologies Inc., Beijing, China) supplemented with 1% protease inhibitor and phosphatase inhibitor (Pierce™, Thermo Fisher Scientific, USA). After the sample content was measured by BCA method (Applygen Technologies Inc., Beijing, China), the protein was boiled at 99°C for 10 min. The protein samples of tissue and cells were separated by 10% SDS-PAGE gels and then they were transferred onto PVDF membranes (Millipore, Boston, MA, USA). After that, the primary antibodies (1:1000 or 1:2000, Table 1) and secondary antibodies (1:2000) were incubated with the membranes at 4°C overnight and at room temperature for 2 h, respectively. Then the bands were reacted with ECL (ECL Plus, GE Healthcare, United States) in the dark for 1 min, and were analyzed through the Image lab software.

Table 1
Primary antibodies used in western blot experiment

Protein	Primary antibody	Concentration
p38 MAPK	Anti-p38 MAPK 8690T CST	1 :1000
p-p38 MAPK	Anti-p-p38 MAPK 4511T CST	1 :1000
MKK3	Anti-MKK3 ab195037 Abcam	1 :1000
MKK6	Anti-MKK6 ab33866 Abcam	1 :1000
p-MKK3/6	Anit-MKK3/6 12280 CST	1 :1000
Bax	Anti-Bax 50599 Proteintech	1 :2000
Bcl-2	Anti-Bcl2 26593 Proteintech	1 :2000
GAPDH	GAPDH 51332S CST	1 :1000

Statistical analysis

In the study, all data were showed as the mean $\pm$ SEM and analyzed by GraphPad Prism 5.0 (GraphPad Software, Inc., San Diego, CA, United States). The differences among groups were calculated by one-way analysis of variance (ANOVA) with post hoc Dunnett's test. $P < 0.05$ represented significant.

Results

LCMS-IT-TOF profiles of RAF and PAF

HPLC chromatograms showed that RAF contains ingredients during 0–40 min retention time with moderate and high polarity, whereas PAF mainly consists of hydrophilic constituents in the first 20 min (Fig. 1). Based on data analysis of LC-MS and those previously isolated compounds [13], 14 major peaks of alkaloids were assigned as *N*-trans-p-coumaroyl noradrenaline (**1**); *N*-trans-p-coumaroyloctopamine (**2**); magnoflorine (**3**); *N*-trans-feruloyloctopamine (**4**); berberine (**5**); dehydrocheilanthifoline (**6**); isomer-dehydrocheilanthifoline (**7**); tetrahydropalmatine (**8**); bicuculine (**9**); 6,7-methylenedioxy-2-(6-acetyl-2,3-methylenedioxybenzyl)-1(2H)-isoquinolinone (**10**); protopine (**11**); allocryptopine (**12**); hendersonine B (**13**); stylophine (**14**).

Effect on echocardiography

Compared with the sham group, the values of LVEDd and LVEDs were increased significantly and the values of EF and FS were decreased significantly ($P < 0.001$, $P < 0.001$) in the model group. The TE and RAF group could decrease the levels of LVEDd and LVEDs and increase the levels of EF and FS, as shown in Fig. 2A. Moreover, after treatment for seven days, RAF existed the same effect compared with TE in ameliorating mice cardiac function, but PAF showed negative result, as shown in Fig. 2B, C. As described above, it suggested that RAF contributes to the cardioprotective effect whereas PAF didn't.

Effect on the levels of CK-MB, LDH and AngII

The levels of LDH, CK-MB, and AngII were markedly high ($P < 0.001$) in the model group. Compared with the model group, RAF and TE could reduce the rising levels of LDH and CK-MB in serum ($P < 0.01$, $P < 0.05$), whereas PAF had no significant influence (Fig. 3A). Our further research found that RAF also improved the myocardial enzymes and mitigated the increase in plasma AngII level (Fig. 3B).

Effect on myocardial tissue pathology

HE staining displayed that the cells were normal and arranged regularly in the sham group, while a great quantity of inflammatory cells infiltrate in the interstitium related to the model group (Fig. 4A). MASSON and Sirius red staining in the sham group showed that the interstitium contained a small amount of collagen fibers and no fibroblast proliferation, but the destruction of the collagen network and the myocardial fibrosis were serious after modeling. However, treatment RAF for 7 days, the inflammatory cell infiltration and collagen deposition were decreased, demonstrating the ischemic injury of mice myocardium were improved (Fig. 4B, C).

The TUNEL assay result illustrated that the myocardial cells after modeling were more apoptotic since the blue-green fluorescence was increased. RAF treatment markedly reduced myocardial apoptosis, protecting the ischemic myocardial damage (Fig. 4D).

Effect on p38 MAPK, Bax, Bcl-2 mRNA expression

The expression of p38 MAPK and Bax mRNA were detected through qRT-PCR. In model group, the mRNA expression of p38 MAPK and Bax was significantly up-regulated, suggesting that myocardial ischemia could increase p38 MAPK expression. RAF down-regulated the mRNA levels of p38 MAPK and Bax, and up-regulated the mRNA level of the anti-apoptotic gene Bcl-2 (Fig. 5A).

Effect on p38 MAPK, Bax, Bcl-2 immunofluorescence staining

The green fluorescence intensity was increased in the model group, indicating that the content of p38 MAPK and Bax protein increased. Those signals of the treatment group were weaker than the model group, suggesting that RAF may inhibit the protein expression of p38 MAPK and Bax (Fig. 5B, C). While the signals of Bcl-2 immunofluorescence were weak in the model group, after administered with RAF, the signals were enhanced (Fig. 5D).

Effects on p38 MAPK-mediated apoptotic proteins

Compared with the sham group, the expression of p-p38 MAPK, p-MKK3/6 and Bax proteins were obviously increased in mice after myocardial infarction, while RAF had a significant inhibitory effect against apoptosis by down-regulating the expression of these proteins. The RAF administration group also up-regulated the anti-apoptotic protein expression of Bcl-2 to varying degrees. The protein results revealed that the anti-AMI effect of RAF was related to the p38 MAPK signaling pathway (Fig. 6).

Effects on H9c2 cells viability

After 24 hours of administration, cell survival rates of RAF (1.25–10 µg/mL) groups were not lower than the control group ($P > 0.05$), indicating that RAF had no apparently cytotoxic to H9c2 cells. After 8 hours of culture under hypoxia, serum-free and low glucose conditions, RAF (10 µg/mL) improved cell viability, but significantly improved cell viability at a concentration of 5 µg/mL, indicating that RAF had protective effect against hypoxia-induced injury (Fig. 7A).

Effects on apoptosis of H9c2 cells

The fluorescence intensity was remarkably decreased by RAF (1.25, 2.5, and 5 µg/mL), indicating that RAF could suppress apoptosis in varying degrees. Through the microscope we found that all along the dose increased in the RAF administration group, live cells increased and dead cells decreased, further indicating that RAF could improve the survival rate of cells (Fig. 7B).

Effects on protein expression of H9c2 cells

When myocardial ischemia occurred, the expression of p-p38 MAPK and p-MKK3 were up-regulated and promoted Bax expression, which suggested that hypoxia injury could activated the p38 MAPK-mediated apoptotic signaling pathway. However, RAF reversed the tendency of p-p38 MAPK, p-MKK3 and Bax/Bcl-2 expression compared to the model group (Fig. 7C). The above results showed that RAF exhibited anti-apoptotic effect via the p38 MAPK signaling pathway.

Discussion

The present results showed that TE (400 mg/kg) and RAF (100 mg/kg) treatments could ameliorate mice cardiac function, suggesting RAF and TE, to some extent, had the equivalent effects against AMI and alkaloids rather than non-alkaloids of CH contributes the cardioprotective effect. Furthermore, RAF (25–100 mg/kg) reduced the activity of serum myocardial enzymes CK-MB and LDH, decreased the apoptosis and collagen synthesis of myocardial pathological changes, reduced the increased expression of p38 MAPK, MKK3, and MKK6 both in mice and H9c2 cells. These results indicated that RAF exerts the cardioprotective effect against ischemic injury through inhibiting apoptosis underlying p38 MAPK signaling pathway.

Coronary artery ligation leads to interruption of blood supply. After MI, oxygen supply decreases and energy metabolism are abnormal, and the heart fails to maintain normal work [14]. Continuous ischemia causes a wide range of myocardial infarction, further amplifies the inflammatory response, and aggravates myocardial damage. Apoptosis is one of the main ways of myocardial injury caused by coronary artery occlusion, and there are many apoptotic cells in ischemic myocardial injury [15]. More and more researches gradually verified that apoptosis is one of the most importantly mode of cardiomyocyte death when AMI happened [16–18]. Therefore, inhibiting myocardial cell apoptosis after myocardial infarction is one of the important treatment measures to improve cardiac function [19, 20]. As a key factor of myocardial apoptosis signal transduction, p38 MAPK is involved in a series of processes

after AMI [21, 22]. This signaling pathway phosphorylates p38 MAPK through a cascade of three key kinases, MEK1/2, TAK1, MKK3/6, and p38 MAPK. The upstream activators of p38 MAPK are MKK3, MKK4, and MKK6. Unlike MKK4, MKK3 and MKK6 are only specific. Under stress, MAPK kinase 3/6 is activated, which in turn activates p38 MAPK and activates cardiomyocyte apoptosis. Moreover, the proteins of Bcl-2 family are critical in the mitochondrial transfer pathway and the "master switch" of apoptosis [23–25]. Our findings depicted evident that RAF restored of MKK3, MKK6, and p38 phosphorylation and increased the ratio of Bcl-2 / Bax to attenuate cardiomyocyte apoptosis.

By analysis of LC-MS data, 14 alkaloids were characterized in RAF, including the major ones (peaks **3, 6–12**) and minor in its contents (peaks **1, 2, 4, 5, 13, and 14**). Among them, compounds **3, 6** and **11–14** were identified from TE in our previous study. The difference between the previous TE and the present RAF was ascribed to the different collection time of plant and extract procedure in this study. The CH in our previous study was collected in June 2013 and extracted with 80% ethanol and the material of CH in the current study was purchased in July 2017 and RAF was obtained by column chromatography of ion exchange resin after ethanol extraction.

Some alkaloids had the effects of anti-inflammatory and inhibiting platelet aggregation, which benefits to the anti-AMI activity of TE [26, 27]. Some alkaloids had also been reported in anti-oxidation and anti-apoptosis [28], which further explained the contribution of RAF to anti AMI. At the same time, it suggested that TCM played an overall role with the multi-component and multi-pathway paradigm.

PAF accounting for 75% of TE in mass ratio didn't show the effect against AMI and at the same time, PAF had little signal in LC-MS. Whether PAF could contribute to other aspects of traditional efficacy and which substances could exert efficacy need follow-up study.

Conclusion

In summary, both the rich-alkaloid fraction and total extract of *Corydalis hendersonii*, one of Tibetan folk medicines, have the cardioprotective effect against AMI mice. The rich-alkaloid fraction limits myocardial ischemia through attenuating the cardiomyocyte apoptosis underlying the mechanism of p38 MAPK signaling pathway in H9c2 cells and mice. The major isoquinoline and phenylpropionamide alkaloids contributed to the efficacy and could be considered as the biologically active substances against ischemic heart disease. Our research provides the substantial foundation for the development of potential drugs for the treatment of ischemic heart diseases, and provides a reference for utilization of natural CH resource as traditional medicines.

Abbreviations

CH

Corydalis hendersonii Hemsl

TE

total extract
RAF
rich alkaloids fraction
PAF
poor alkaloidal fraction
HAPC
high-altitude polycythemia
AMI
acute myocardial infarction
LVEDd
left ventricular internal diameter
LDH
lactate dehydrogenase
EF
ejection fraction
LVEDs
left ventricular internal diameter
FS
fractional shortening
AngII
angiotensin II
CK-MB
creatine kinase-MB
Bax
X protein associated Bcl-2
IHD
ischemic heart disease
BV/h
bed volumes per hour
DAD
diode array detector
MKK
MAP kinase kinase
LCMS-IT-TOF
liquid chromatography mass spectrometry-ion trap-time of flight
p38 MAPK
p38 mitogen-activated protein kinases.

Declarations

Ethics approval and consent to participate

The Ethics Committee of BUCM had approved the experiments (the ethics permit number: BUCM-3–2015090701-3003), and the research was implemented according to the US guidelines (NIH publication #85-23, revised in 1996).

Consent for publication

Not applicable.

Availability of data and material

Data can be shared with the author's consent.

Competing interests

There is no conflict of interest in this paper.

Funding

National Natural Science Foundation of China (Nos. 81774001, 81473426).

Authors' contributions

Fuxing Ge, Xiaoli Gao, Chao An and Xingyun Chai prepared and revised the manuscript. Fuxing Ge, Xiaoli Gao, Junjun Li, Xiaojing Ma and Meiwen Huang contributed to the animal and molecular biology experiments. Xiaochun Zhou, Shana Wuken and Xingyun Chai performed the LCMS-IT-TOF experiments and data analysis. Xingyun Chai and Pengfei Tu are responsible for the research design and approved this article.

Acknowledgments

We thank National Natural Science Foundation of China (Nos. 81774001, 81473426) supported the study.

References

1. Ostadal B, Kolar F. Cardiac adaptation to chronic high-altitude hypoxia: beneficial and adverse effects. *Resp Physiol Neurobi.* 2007;158:224–36.
2. Rustamova Y, Lombardi M. Ischemic heart disease. In: *Cardiac Magnetic Resonance Atlas*. Cham: Springer; 2020. pp. 63–91.
3. Wang X, Ji Y, Yang B. Chinese innovation in cardiovascular drug discovery. *Br J Pharmacol.* 2015;172:5425–29.

4. Li X, Zhang J, Huang J, Ma A, Yang J, Li W, et al. A multicenter, randomized, double-blind, parallel-group, placebo-controlled study of the effects of qili qiangxin capsules in patients with chronic heart failure. *J Am Coll Cardiol*. 2013;62:1065–72.
5. Li XL, Liu JX, Guo YJ. Effect of shuangshen ningxin formula on energy metabolism of myocardial ischemia/reperfusion rats. *Chin J Chin Mater Med*. 2013;38:2874–7.
6. Li J, Ge F, Wuken S, Jiao S, Huang M, Tu P, et al. Zerumbone, a humulane sesquiterpene from *Syringa pinnatifolia*, attenuates cardiac fibrosis through inhibition of the TGF- β /Smads signaling pathway after MI in mice. *Phytomedicine*. 2022;100:154078.
7. Feng X, Zhang R, Li J, Cao Y, Zhao F, Du X, et al. *Syringa pinnatifolia* Hemsl. fraction protects against myocardial ischemic injury by targeting the p53-mediated apoptosis pathway. *Phytomedicine*. 2019;52:136–46.
8. Cao Y, Wang J, Su G, Wu Y, Bai R, Zhang Q, et al. Anti-myocardial ischemia effect of *Syringa pinnatifolia* Hemsl. by inhibiting expression of cyclooxygenase-1 and-2 in myocardial tissues of mice. *J Ethnopharmacol*. 2016;187:259–68.
9. Zhao F, Bai R, Li J, Fen X, Jiao S, WuKen S, et al. *Meconopsis horridula* Hook. f. & Thomson extract and its alkaloid oleracein E exert cardioprotective effects against acute myocardial ischaemic injury in mice. *J Ethnopharmacol*. 2020;258:112893.
10. Bai R, Yin X, Feng X, Cao Y, Wu Y, Zhu Z, et al. *Corydalis hendersonii* Hemsl. protects against myocardial injury by attenuating inflammation and fibrosis via NF- κ B and JAK2-STAT3 signaling pathways. *J Ethnopharmacol*. 2017;207:174–83.
11. Danzhengpengcuo DME. *Jingzhu Materia Medica (Tibetan)*. National Publishing House. 1986; 374.
12. Editorial committee of the administration bureau of traditional Chinese medicine. Volume for Tibetan medicine of Chinese Materia Medica. Vol. 12. Shanghai: Shanghai Science & Technology Press; 2002. p. 321.
13. Yin X, Bai R, Guo Q, Su G, Wang J, Yang X, et al. Hendersine a, a novel isoquinoline alkaloid from *corydalis hendersonii*. *Tetrahedron Lett*. 2016; 4858–62.
14. Frangogiannis GN. Regulation of the inflammatory response in cardiac repair. *Circ Res*. 2012;110:159–73.
15. Sui H, Chen WW, Wang W. Interpretation of report on cardiovascular diseases in China (2015). *Chin J Cardiovasc Med*. 2016;21:259–61.
16. Chen YF, Yang YJ. Ventricular remodeling after acute myocardial infarction: significance of cardiomyocyte apoptosis. *Chin J Cardiovasc Med*. 2003;31:629–31.
17. Krijnen PAJ, Nijmeijer R, Meijer CJLM, Visser CA, Hack CE, Niessen HWM. Apoptosis in myocardial ischemia and infarction. *J Clin Pathol*. 2002;55:801–11.
18. Bayat H, Swaney JS, Ander AN, Dalton N, Kennedy BP, Hammond HK, et al. Progressive heart failure after myocardial infarction in mice. *Basic Res Cardiol*. 2002;97:206–13.

19. Baldi A, Abbate A, Bussani R, Patti G, Melfi R, Angelini A, et al. Apoptosis and post-infarction left ventricular remodeling. *J Mol Cell Cardiol.* 2002;34:165–74.
20. Song XJ. The mechanism of atorvastatin inhibiting cardiomyocyte apoptosis in rats with heart failure after myocardial infarction. Jilin University. 2011.
21. Wang M, Chen SL. Advances in the research of correlations between P38 MAPK and cardiovascular diseases. *Prog Mod Biomed.* 2012;12:5968–70.
22. Lee HY, Lee JS, Kim HG, Kim WY, Lee SB, Choi YH, et al. The ethanol extract of *Aquilariae Lignum* ameliorates hippocampal oxidative stress in a repeated restraint stress mouse model. *BMC Complem Altern M.* 2017;17:397.
23. Hu Y, Lin Y, Zhao Y. Progress on foreign research of treating on apoptosis induced myocardial ischemia/reperfusion. *Chin Med Her.* 2013;10:32–4.
24. Ni YL, Lin SD. Regulation of Bcl-2 protein family on the growth and development of follicles and early embryos. *Carcinog Teratogenesis Mutagen.* 2008;20:166–8.
25. Lalier L, Cartron PF, Juin P, Nedelkina S, Manon S, Bechinger B, et al. Bax activation and mitochondrial insertion during apoptosis. *Apoptosis.* 2007;12:887–96.
26. Shang WQ, Chen YM, Gao XL, Pu C, Tu PF, Chai XY. Phytochemical and pharmacological advance on Tibetan medicinal plants of *Corydalis*. *Chin J Chin Mater Med.* 2014;39:1190–8.
27. Gao YP, Wu Q, Liang J, Zhong GY. Chemical constituents of *Corydalis hendersonii*. *Chin J Exp Tradit Med Form.* 2016;22:60–3.
28. Jantamat P, Weerapreeyakul N, Puthongking P. Cytotoxicity and apoptosis induction of coumarins and carbazole alkaloids from *Clausena Harmandiana*. *Molecules.* 2019;24:3385.

Figures




Figure 1

HPLC-DAD (A) and positive-mode (B) LCMS-IT-TOF chromatograms of RAF, and HPLC-DAD (C) and positive-mode (D) LCMS-IT-TOF chromatograms of PAF. (1) *N*-trans-p-coumaroyl noradrenaline; (2) *N*-trans-p-coumaroyloctopamine; (3) magnoflorine; (4) *N*-trans-feruloyloctopamine; (5) berberine; (6) dehydrocheilanthifoline; (7) isomer-dehydrocheilanthifoline; (8) tetrahydropalmatine; (9) bicuculine; (10) 6,7-methylenedioxy-2-(6-acetyl-2,3-methylenedioxybenzyl)-1(2H)-isoquinolinone; (11) protopine; (12) allocryptopine; (13) hendersonine B; (14) stylopine.




Figure 2

Effect on cardiac function in AMI mice. **(A)** Mice were administered with TE (400 mg/kg, i.g.), RAF (100 mg/kg, i.g.), PAF (300 mg/kg, i.g.), 0.5% CMC-Na (i.g.) or fosinopril (10 mg/kg, i.g.) for 7 days. **(B)** Mice were administered with RAF-L (25 mg/kg, i.g.), RAF-M (50 mg/kg, i.g.), RAF-H (100 mg/kg, i.g.), 0.5% CMC-Na (i.g.) or fosinopril (10 mg/kg, i.g.) for 7 days. The LVEDd, LVEDs, EF and FS of AMI mice were measured by echocardiography. Data are showed as mean $\pm$ SEM of three independent experiments ($n = 8$). * $P < 0.05$, ** $P < 0.01$, and *** $P < 0.001$ compared with the Model group. **(C)** The representative echocardiography images.




Figure 3

Effect on CK-MB, LDH and AngII in AMI mice. **(A)** Mice were administered with TE (400 mg/kg, i.g.), RAF (100 mg/kg, i.g.), PAF (300 mg/kg, i.g.), 0.5% CMC-Na (i.g.) or fosinopril (10 mg/kg, i.g.) for 7 days. **(B)** Mice were administered with RAF-L (25 mg/kg, i.g.), RAF-M (50 mg/kg, i.g.), RAF-H (100 mg/kg, i.g.), 0.5% CMC-Na (i.g.) or fosinopril (10 mg/kg, i.g.) for 7 days. The serum levels of CK-MB, LDH were analyzed by Automatic Biochemical Analyzer and the plasma level of AngII was detected by ELISA. Data are showed as mean $\pm$ SEM of three independent experiments ($n = 8$). * $P < 0.05$, ** $P < 0.01$, and *** $P < 0.001$ compared with the Model group.




Figure 4

Effect on myocardial histopathology and apoptosis after AMI modeling. Mice were administered with RAF-L (25 mg/kg, i.g.), RAF-M (50 mg/kg, i.g.), RAF-H (100 mg/kg, i.g.), 0.5% CMC-Na(i.g.) or fosinopril (10 mg/kg, i.g.) for 7 days. **(A)** HE staining (images magnification at 400 ×); **(B)** Masson staining (collagen is stained blue, myocardial cells are stained red) (images magnification at 400 ×); **(C)** Sirius red

staining (collagen is stained red, myocardial cells are stained yellow) (images magnification at 400 ×);
(D) TUNEL staining (images magnification at 40 ×).




Figure 5

Effect on p38 MAPK, Bax and Bcl-2 in heart tissue. Mice were administered with RAF-L (25 mg/kg, i.g.), RAF-M (50 mg/kg, i.g.), RAF-H (100 mg/kg, i.g.) or 0.5% CMC-Na (i.g.) for 7 days. The expressions of p38

MAPK, Bax and Bcl-2 were analyzed by qRT-PCR (A) and immunofluorescence staining (B, C, D). Data are showed as mean $\pm$ SEM of three independent experiments (n = 3). * P < 0.05, ** P < 0.01, and *** P < 0.001 compared with the Model group.





Figure 6

Effect on protein expression of p-p38 MAPK, p-MKK3/6 and Bax/Bcl-2 in AMI mice. Mice were administered with RAF-L (25 mg/kg, i.g.), RAF-M (50 mg/kg, i.g.), RAF-H (100 mg/kg, i.g.) or 0.5% CMC-Na (i.g.) for 7 days. The protein expression was detected by western blot. Data are showed as mean $\pm$ SEM of three independent experiments (n = 3). * P < 0.05, ** P < 0.01, and *** P < 0.001 compared with the Model group.



control for 8 h, the protein expression of p-p38 MAPK, p-MKK3 and Bax/Bcl-2 was analyzed by western blot. Data are showed as mean $\pm$ SEM of three independent experiments (n = 3). * P < 0.05, ** P < 0.01 compared with the Mod group.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [floatimage1.jpeg](#)
- [SFig1.jpg](#)