Journal of Soil Science and Plant Nutrition (2026) 26:5724–5736
https://doi.org/10.1007/s42729-026-03225-5

ORIGINAL PAPER

Green Synthesis of Ag and Pd Nanoparticles using *Melia azedarach* Leaf Extract: Effects on Carbon Mineralization in Clay and Sandy Textured Soils

Nacide Kizildag Ozdal[1] · Bahar Meryemoglu[1] · Husniye Aka Sagliker[2]

Received: 8 October 2025 / Accepted: 17 March 2026 / Published online: 10 April 2026
© The Author(s) 2026

Abstract
This study investigated the effects of green-synthesized silver (Ag) and palladium (Pd) nanoparticles (NPs) on soil carbon mineralization under controlled incubation conditions (28 °C and costant moisture for 42 days). AgNPs and PdNPs were synthesized using extracts of *Melia azedarach* L. leaves and characterized by UV-Vis spectroscopy, SEM-EDS, FTIR, and XRD analyses. The nanoparticles were applied to two different soil textures (clay and sandy) at three concentrations (10, 50 and 100 mg kg^{-1}) to evaluate their impacts on microbial activity through carbon mineralization measurements over 42 days. The results indicated that carbon mineralization was generally higher in clay soils than in sandy soils, with the highest mineralization rate reaching 15.43% in clay soils compared to 10.21% in sandy soils. AgNPs exhibited stronger inhibitory effects on microbial activity than PdNPs, with the highest AgNP dose (100 mg kg^{-1}) reducing carbon mineralization rates to 7.79% in clay soil and 7.16% in sandy soil. Conversely, the addition of *Melia azedarach* leaves and extracts enhanced mineralization, likely due to the presence of easily degradable organic compounds. This study revealed a distinct soil structure dependent pattern, in which clay fractions provide a significant buffering effect against the toxicity of biogenic nanoparticles. These results indicated that the surface coating derived from *Melia azedarach* extract interacted with clay surfaces, potentially reducing the bioavailability of nanoparticles compared with sandy soils.

Keywords Silver nanoparticles · Palladium nanoparticles · *Melia azedarach* · Soil microbial activity · Carbon mineralization

1 Introduction

The use of nanoparticles (NPs), which play a significant role in nanotechnology and are regarded as a key technology for the future, has increased significantly in recent years due to rapid advancements in various sectors (Kumari et al. 2023; Sedefoğlu 2024). NPs smaller than 100 nm exhibit a large surface area area to volume ratio and high surface energy (Khan et al. 2019; Saleh 2020; Shrestha et al. 2020; Taha et al. 2022; Nassem and Durrani 2023). NPs have been synthesized using physical methods like plasma arc, ball milling, pyrolysis, and laser desorption and chemical methods including sol-gel, chemical vapor deposition, sono-chemical precipitation, and solvothermal techniques, among others for many years (Gleiter 2000; Kumari et al. 2023; Aigbe and Osibote 2024). Physical and chemical synthesis methods have many disadvantages such as being complex and very expensive, requiring the use of toxic substances, and unsuitable for pharmacological and biomedical applications (Abid et al. 2022). More recently, with advancements in technology, biological synthesis of NPs has become feasible, utilizing biological agents in addition to these traditional methods. Biomaterials like plants, fungus, and microorganisms are used in the green synthesis of NPs; these materials serve as capping and reducing agents (Gohil et al. 2024). Green synthesis method has also become the preferred method for the researchers because it is easier to apply, relatively economical, and environmentally friendly (Hong et al. 2016; Singh et al. 2016; Annu and

✉ Nacide Kizildag Ozdal
nkizildag@cu.edu.tr

[1] Present address: Central Research Laboratory, Cukurova University, Adana 01330, Türkiye

[2] Faculty of Engineering and Nature Science, Department of Biology, Osmaniye Korkut Ata University, Osmaniye 80000, Türkiye

Ahmed 2018; Kizildag et al. 2019; Sedefoğlu et al. 2022; Osman et al. 2024; Shah et al. 2025). Though green-synthesized is considered an environmentally friendly approach to producing nanoparticles, it has several disadvantages (Dikshit et al. 2021). Because the chemical composition of biological extracts can vary widely, it becomes challenging to standardize the process and obtain nanoparticles with consistent properties. In addition, controlling factors such as particle size, shape, and distribution is often more difficult than in traditional chemical methods. Green synthesis can also lead to low yields, obstacles in scaling up production, and the need for extra purification steps due to remaining biomolecules. Lastly, since the reaction mechanisms are not yet fully understood, optimizing the process and ensuring reproducibility can be problematic (Fahim et al. 2024).

The increasing use of metal NPs in multiple industrial commodities has elevated the risk of environmental exposure, particularly to air, water, and soil (Collins et al. 2012a; Simonin and Richaume 2015). Among these environmental components, soil is the primary medium where NPs tend to accumulate (Xu et al. 2015; Ahmed et al. 2021). NPs can enter the soil through several pathways, including the application of sewage sludge, landfill disposal, or accidents during industrial production processes (Keller and Lazareva 2013; Keller et al. 2013).

The process by which microorganisms decompose organic matter and release carbon as CO_2 is known as soil carbon (C) mineralization. This process is a trustworthy measure of the activity of soil microbial communities since it is directly caused by microbial activity (Pascual et al. 2000; Rodríguez-Albarracín et al. 2025). In order to grow and obtain energy, microorganisms eat organic substrates, and the CO_2 they release is a reflection of both their metabolic activity and the functional dynamics of the soil ecosystem (Liang et al. 2024). According to Zhang et al. (2023), measuring carbon mineralization is a commonly used method of assessing microbial activity because it provides insight into the efficiency and rate at which bacteria utilize organic materials. Since it might be challenging to measure microbial numbers or metabolism directly, examining carbon mineralization offers a reliable but indirect approach (Bruni et al. 2025). In summary, measuring soil carbon mineralization provides important information about microbial activity and general ecosystem processes (Rieke et al. 2022). In general, the residual NPs in the soil mainly contaminate the food chain by entering the plant system (Melo et al. 2020). In addition, NPs can be integrated into plant and microbial systems, thus leading to absorption of NPs in the food chain, that subsequently threatens the quality and safety of human health (Ali et al. 2020). Many studies have reported toxic effects of NPs on microbial activity in some soils (Schlich and Hund-Rinke 2015; Zhu et al. 2019; Ali et al. 2020; Saleh 2020; Egbuna et al. 2021; Kumar et al. 2025). However, depending on the soils physicochemical properties, low concentrations of certain nanoparticles have been reported to exert negligible or stimulatory effects on soil enzymes and microbial biomass (Schlich and Hund-Rinke 2015; Shah and Belozerova 2009). These conflicting findings highlighted the need for further extensive and detailed research to elucidate the specific mechanisms and environmental factors, such as soil texture and biogenic coatings, that modulate the magnitude of NPs toxicity in diverse natural ecosystems (Szőllősi et al. 2020; Muzammil et al. 2023). Even though NPs are increasingly being researched in various fields, there is still a lack of scientific data on how NPs impact soil microbial activity (You et al. 2018). Research on soil microbial responses to NPs is increasing, but studies are often focused on specific microbial groups, particular NPs types, or controlled laboratory conditions. Consequently, the broader ecological effects particularly on microbial functions such as decomposition remain poorly understood, highlighting the need for the impacts of Ag and Pd nanoparticles to be systematically investigated in natural soil ecosystems. The response or tolerance of microorganisms in the soil to NPs will provide meaningful and valuable information on the effects of these compounds on the ecological balance. These abilities are important because the decomposition activities of microorganisms ensure the continuity of ecosystems and, therefore, the sustainability of life. In this context, understanding how soil microorganisms respond to green-synthesized Ag and Pd nanoparticles particularly in different soil textures fills an important knowledge gap and provides novel insight into how these NPs may influence key microbial processes such as carbon mineralization.

The accumulation of NPs in soil ecosystems, depending on exposure time, poses a threat to microorganisms living in the soil and may disrupt the ecological balance (Rajput et al. 2018; Kizildag et al. 2019). Texture types and organic matter contents of soils are significant parameters for soil microbial activities and also the toxicity of NPs (Simonin and Richaume 2015; Shao et al. 2022). Although the literature on nanoparticle interactions within different soil textures is limited, recent studies have highlighted the importance of soil properties in nanotoxicity (You et al. 2018; Kizildag et al. 2019; Sharma and Kumar 2024). For instance, Colman et al. (2013) reported that even low doses of AgNPs (0.14 mg kg^{-1}) negatively impacted bacterial community composition in sandy loam soils. Regarding metal oxide nanoparticles, You et al. (2018) demonstrated that CeO_2, Fe_3O_4, TiO_2, and ZnO NPs, applied at 100 and 500 mg kg^{-1}, significantly altered both bacterial communities and the activities of key enzymes (catalase, invertase, phosphatase, and urease) in saline-alkali and black soils. Similarly, Chai et al. (2015) observed that ZnO and CeO_2 NPs tested at 500 mg kg^{-1} reduced thermogenic metabolism, the abundance of Azotobacter and phosphate/potassium-solubilizing bacteria, as well as overall enzymatic activities.

While the toxicity of chemically synthesized nanoparticles is well-documented, there is a significant gap in understanding how green-synthesized NPs capped with phytochemicals behave across different soil textures. Understanding whether soil components (like clay or sand) can buffer the potential toxicity of these biogenic. NPs is crucial for their safe application in agriculture. Green synthesis was selected not only for its environmental sustainability but also for the specific phytochemical coating that governs the NPs surface properties. While the impacts of conventionally synthesized nanoparticles are well documented, the interactions between biogenic surface coatings and complex soil structures remain poorly understood. This study aimed to address this gap by investigating how the interplay between NPs surface properties and soil texture influences microbial responses. Based on this gap, we hypothesized that the physicochemical properties of soil texture significantly modulate the impact of green-synthesized AgNPs and PdNPs on microbial activity. Specifically, we predicted that clay-rich soils would buffer the inhibitory effects on carbon mineralization more effectively than sandy soils, possibly due to the interaction between clay surfaces and the organic capping layer of the NPs. To test this hypothesis, the primary objective of this study was to compare the mineralization rates of C in two contrasting soil textures (clay, sandy) amended with green-synthesized AgNPs and PdNPs, thereby determining the role of soil texture in regulating nanoparticle ecotoxicity.

In the light of all these information, the aim of this study was to determine the carbon mineralization of clay and sandy textured soils at increasing doses (10, 50 and 100 mg kg^{-1}) of silver (Ag) and palladium (Pd) nanoparticles (NPs) green-synthesized from extract of *Melia azedarach* L. leaves, and to reveal their potential effects on the ecosystem.

There are almost no studies on the effects of NPs on carbon mineralization in Turkish soils. Since soil characteristics are known to influence the toxicity of NPs, this exploratory work is relevant due to the use of clay and sandy soils commonly found in Türkiye. This study results is significant in terms of adding contributed new data to the existing knowledge about the inhibitory effects of green synthesized AgNPs and PdNPs by monitoring carbon mineralization of microorganisms in two different soil types (clay and sandy), enabling better environmental risk assessment.

2 Materials and Methods

This study was carried out with the clay and sandy textured soils collected from agricultural lands of Çukurova University Balcali campus area in Adana, Türkiye. Both soils were taken from 0 to 20 cm depth after upper layer cleaning, and they were air dried and sieved with a 2-mm sieve in the laboratory conditions.

2.1 Green Synthesis of AgNPs and PdNPs

For the green synthesis of AgNPs and PdNPs, leaves of *Melia azedarach* L. (Meliaceae) plant naturally growing on the Çukurova University Balcalı campus was used in this study. Plant samples were washed with distilled water and dried in drying oven (60 °C and over night). 5 g of dried leaves were held in 50 ml of distilled water at about 90 °C for 60 min. The extracted materials were then centrifuged for 20 min at 4000 rpm. These extracts were used to produce NPs. Ag and PdNPs were prepared using water-soluble salts of these metals. 1 mM solutions of $AgNO_3$ and $PdCl_2$ were prepared. The NPs were produced by heating 90 mL of metal solution and 10 mL of leaves extract under reflux for 30 min at 80 °C. After then, this solution was left in the dark for 24 h. Several spectroscopic techniques were used to characterize the produced metal NPs after dried.

2.2 Characterization of AgNPs and PdNPs

The characterization studies of AgNPs and PdNPs carried out at Çukurova University Central Research Laboratory. The UV-Visible Spectroscopy (UV-Vis) (Agilent Carry 100 UV-Vis) were used to get the absorption spectrum of NPs. The X-ray diffraction was used to determine the crystal structure of the NPs (PANalytical, XRD). Scanning electron microscopy was used to analyze the surface morphology (FEI-SEM). FTIR examinations were used to identify the material surface functional groups. A Jasco FTIR spectrometer was used to record spectra in ATR (Attenuated Total Reflectance) mode within the 4000–400 cm^{-1} range.

2.3 Soil Analysis

The soil texture was determined by the Bouyoucos hydrometer (Bouyoucos 1951), field capacity (%) by a vacuum pump at 1/3 atmospheric pressure (Demiralay 1993) and pH with a WTW Inolab 720 pH-meter in 1:2.5 soil-water suspension (Jackson 1958). Total organic carbon (TOC) and total nitrogen (TN) contents of soils (%) were determined by modified Walkley and Black method and Kjeldahl method, respectively (Duchaufour 1970).

2.4 Soil Carbon Mineralization

For carbon mineralization; three different doses (10, 50 and 100 mg kg^{-1}) of AgNPs and PdNPs green-synthesized from *Melia azedarach* L. leaves were mixed into the clay and sandy soils by 80% of field capacity and incubated at 28 °C for 42 days (Schaefer 1967). The leaves were air-dried, ground into powder, and mixed into the soil. To measure normal soil microbial activity in both soils, the controls

included soil with no addition of AgNPs or PdNPs, soil amended with leaves at a level equivalent to the carbon content of each soil, and soil treated with *Melia* extract.

CO_2 produced by microbial activity was captured in 40 ml of saturated $Ba(OH)_2$ solution placed in beakers at the center of the soil samples. The beakers were subsequently transferred to an incubator set at 28 °C for 42 days. The amount of CO_2 produced was measured by titration with oxalic acid (Benlot 1977). Empty vessels were used as blank. The carbon mineralization rate (%) was calculated by dividing the cumulative amount of $C(CO_2)$ produced in 42 days into total organic carbon.

2.5 Statistical Analysis

A Repeated Measures (General Linear Model) analysis was performed to determine the differences in carbon mineralization between the two different textured soils (clay and sandy), two NPs (AgNPs and PdNPs), and their different doses (10, 50 and 100 mg kg^{-1}) (Kleinbaum et al. 1998). Three replicates were used for each combined soil for statistical comparisons. Data were analyzed by a series of analyses of variance. Differences between data were assumed significant at $p<0.05$. All statistical analyses were carried out using SPSS 21.0.

3 Results

3.1 Characterization of NPs

Spectroscopic methods were used in the characterization of nanoparticles. The comparative UV-VIS spectra of AgNPs and PdNPs green-synthesized using extract of *Melia azedarach* leaves were presented in Fig. 1a. AgNPs were expected to exhibit a peak within the wavelength range of 400–450 nm according to UV-VIS, with the AgNPs showing this peak around 400 nm. The absorption band for Pd ions was generally observed at 415 nm. The absence of this peak for PdNPs provides as proof of the decrease of Pd ions (Fig. 1a).




Fig. 1 Characterization of *Melia azedarach* leaf extract and green-synthesized nanoparticles. **a** UV-Vis absorption spectra of the extract, AgNPs, and PdNPs; **b** Scanning Electron Microscopy (SEM) images with Energy Dispersive X-ray Spectroscopy (EDS) profiles; **c** Fourier Transform Infrared (FT-IR) spectra; **d** X-ray Diffraction (XRD) patterns. Abbreviations: AgNPs: Silver Nanoparticles; PdNPs: Palladium Nanoparticles; UV-Vis: Ultraviolet-Visible; SEM-EDS: Scanning Electron Microscopy-Energy Dispersive X-ray Spectroscopy; FT-IR: Fourier Transform Infrared; XRD: X-ray Diffraction

The morphological characterization of the synthesized AgNPs and PdNPs via SEM analysis confirmed their spherical structure (Hasanzadeh et al. 2021). The presence of AgNPs and PdNPs was confirmed by the presence of intense peaks corresponding to Ag and Pd, respectively, in the elemental composition analysis carried out using EDX, as illustrated in Fig. 1b. Additionally, the detection of minor peaks for elements such as carbon and oxygen was attributed to the presence of phytochemical residues associated with the particles (Patra and Baek 2016; Ramkumar et al. 2017).

The obtained FT-IR spectra exhibited fundamental vibration bands corresponding to organic functional groups (Fig. 1c). These bands include O–H stretching vibrations around 3300 cm^{-1}, C–H stretching vibrations associated with alkanes observed at 2917 cm^{-1}, and vibration bands encompassing phenolic O–H and C–O bonds around 1300 cm^{-1}, as well as stretching bands in the range of 1050–1025 cm^{-1} (Singh et al. 2016).

The XRD pattern of the AgNPs and PdNPs were presented in Fig. 1d. As shown, diffraction peaks corresponding to AgNPs were observed at 38°, 44°, 64°, and 77°. The XRD pattern of the PdNPs exhibited five distinct reflections, typically corresponding to the (111), (200), (220), (311), and (222) planes at 40°, 46°, 68°, 81°, and 78°, respectively. These diffraction planes were clearly evident in the synthesized palladium NPs (Fig. 1d). The sharp peak observed at the Pd (111) plane indicates the high crystallinity of the PdNPs (Narasaiah et al. 2017).

3.2 Soil Carbon Mineralization

To investigate the potential use of green-synthesized NPs in soil applications, carbon mineralization was evaluated across two distinct soil textures (clay and sandy). The physicochemical properties of the soils are detailed in Table 1. The clay soil was characterized as a clay loam with a slightly alkaline pH of 7.85 ± 0.07. The carbon and nitrogen contents for this soil were determined as 1.83 ± 0.03 and 0.21 ± 0.01, respectively, which align with typical levels reported in the literature (Loveland and Webb 2003; Poeplau et al. 2018). In contrast, the sandy soil, classified as sandy loam with a pH of 7.55 ± 0.06, exhibited lower carbon (1.32 ± 0.02) and nitrogen (0.18 ± 0.01) contents. Consequently, the C/N ratios were calculated as 8.71 ± 0.22 and 7.33 ± 0.14, respectively.

Table 1 Physical and chemical properties of the clay and sandy textured soils used in the experiments. Values represent mean ± standard error (SE, $n = 3$)

	Clay Soil	Sandy Soil
Sand	38.55 ± 1.12	65.93 ± 1.20
Silt	30.90 ± 0.82	12.70 ± 1.63
Clay	30.55 ± 1.84	22.17 ± 0.28
Texture	**Clay loam**	**Sandy loam**
Field Capacity	32.20 ± 0.71	27.20 ± 0.82
pH	7.85 ± 0.07	7.55 ± 0.06
C (%)	1.83 ± 0.03	1.32 ± 0.02
N (%)	0.21 ± 0.01	0.18 ± 0.01
C/N	8.71 ± 0.22	7.33 ± 0.14

AgNPs and PdNPs were green-synthesized using extract from leaves of *Melia azedarach* L. naturally grown on the Çukurova University campus in Adana city of Türkiye. The characterization of these NPs was determined, and they were added to clay and sandy soils with different organic matter contents in varying doses to assess their effects on carbon mineralization. Additionally, the leaves and extract of *Melia azedarach* were mixed into both clay and sandy soils based on their carbon-equivalent content to investigate their potential impacts on soil microorganisms.

After a 42-day incubation period, cumulative carbon mineralization was generally higher in clay soils compared to sandy soils. PdNPs application resulted in increased mineralization in both clay and sandy soils. Furthermore, the addition of the leaves and their extracts to the soils as a source of carbon led to increased mineralization in both soils after application of the extract. Higher mineralization was observed in clay control soils and soils mixed with leaves and extract compared to AgNPs treatments (Fig. 2a). Among the soils treated with AgNPs, the highest mineralization occurred in the Ag(50) mixed application (13.29 mg $C(CO_2)$/100 g soil), while the lowest mineralization was observed in the Ag(100) mixed soils (10.47 mg $C(CO_2)$/100 g soil).

It was observed that easily decomposable organic matter was rapidly utilized by microorganisms during the first 7 days, mineralization slowed between days 7–21, and then resumed a faster pace after day 21 (Fig. 2a). A significant difference was found between the control soil and all treatments at the $p < 0.05$ level, with no significant difference observed between the Ag(10) and Ag(50) treatments.

With the exception of the soil mixed with Pd(50), higher cumulative carbon mineralization was observed in the clay control, leaves and extract-amended soils compared to the Pd treatments. Statistically significant differences were found between all treatments, except for the control with Pd(10) and the leaves with Pd(50) applications ($p < 0.001$) (Fig. 2b).

As depicted in Fig. 3, the highest mineralization was observed in soils mixed with leaves (17.87 mg $C(CO_2)$). Upon examining the carbon mineralization in soils mixed with leaves and NPs (Leaves+Ag100, Leaves+Pd100), the 42-day cumulative $C(CO_2)$ values were similar to those of the control soil, with no significant differences observed. However, the differences between Ag100 and Leaves+Ag100, as well as between Pd100 and Leaves+Pd100, were statistically highly significant ($p < 0.001$).




Fig. 2 Cumulative carbon mineralization [mg $C(CO_2)$/100 g] in clay textured soils treated with *Melia azedarach* leaves, extract, and green-synthesized nanoparticles. a Effects of Silver Nanoparticles (AgNPs); b Effects of Palladium Nanoparticles (PdNPs) applied at 10, 50, and 100 mg kg^{-1}. Values represent mean ± standard error (SE, n = 3). Different lowercase letters above bars indicate significant differences between treatments according to Tukey's HSD test at $p < 0.05$

The highest mineralization in sandy soils (14.30 mg $C(CO_2)$) was observed in soils amended with extract of Melia leaves, while the lowest mineralization (10.04 mg $C(CO_2)$) occurred in soils with the addition of Ag(100). Additionally, higher mineralization was observed in soils mixed with extract and leaves compared to the control soils. The addition of different doses of AgNPs to the soils resulted in a reduction in mineralization, indicating the effect of NPs (Fig. 4a).

The effects of Pd nanoparticle applications on cumulative carbon mineralization in sandy soils were found to be similar, with no statistically significant differences observed between the treatments (Fig. 4b).




Fig. 3 Cumulative carbon mineralization [mg $C(CO_2)$/100 g] in clay textured soils. The figure displays the comparative effects of Control, *Melia azedarach* leaves alone, nanoparticles alone (100 mg kg^{-1} AgNPs and PdNPs), and their combined applications (Leaves + AgNPs; Leaves + PdNPs). Values represent mean ± standard error (SE, $n = 3$). Different lowercase letters above bars indicate significant differences between treatments according to Tukey's HSD test at $p < 0.05$

As shown in Fig. 5, higher mineralization was observed in sandy soils mixed with leaves compared to the other treatments, with a significant difference only observed between the Ag(100) application ($p < 0.01$). Additionally, no statistically significant difference was observed between the Melia leaves plus Ag(100) and Melia leaves plus Pd(100) treatments.


The carbon mineralization ratios of clay soils were highest in the soil amended with extract of Melia leaves (15.43%) and lowest in the Ag(100) application (7.79%) (Fig. 6a). Statistically significant differences were observed between the control soil and all other treatments, except for Pd(10), Leaves + Ag(100), and Leaves + Pd(100) ($P < 0.005$). A highly significant difference ($p < 0.001$) was found between the soil mixed with leaves and the other soils amended with extract and NPs.

When evaluating the carbon mineralization rates of Ag nanoparticle applications, a significant difference was observed between Ag(10) and Ag(100) ($P < 0.05$), whereas no statistically significant difference was found between Ag(10) and Ag(50) nanoparticle treatments. In contrast, a different pattern was observed for Pd nanoparticle applications. A significant difference was detected between soils treated with Pd(10) and Pd(50) ($P < 0.001$), while no significant difference was found in soils treated with Pd(100).

The Leaves+Ag100 application exhibited highly significant differences when compared to both the leaves and Ag(100) treatments ($p < 0.001$). Similarly, the Leaves+Pd100 application showed significant differences when compared to both the leaves and Pd(100) treatments ($p < 0.005$) (Fig. 6a). When the ratios of carbon mineralization in sandy soils were examined (Fig. 6b), the soil modified with extract showed the highest mineralization rate (10.21%), whereas the Ag(100) treatment showed the lowest rate (7.16%). No significant differences were detected among the various AgNPs concentrations (Ag(10), Ag(50), and Ag(100)). In contrast, when evaluating the PdNPs applications, a statistically significant difference was found between Pd(10) and both Pd(50) and Pd(100) treatments ($p < 0.05$) (Fig. 6b).

4 Discussion

The AgNPs and PdNPs used in this study were obtained by green synthesis method using *Melia azedarach* leaf extract. The investigation of the analysis performed in this context revealed that the UV results were in accordance with those of NPs produced from plant extracts that reported in the literature (Sasikala et al. 2014; Sinha et al. 2014). *Melia azedarach* leaves are rich in phenolic compounds and contain a significant number of hydroxyl, carbonyl, and carboxyl groups in their structure. According to FT-IR results, the formation of NPs led to changes in the absorption bands at 1375 cm^{-1} and 3300 cm^{-1}, depending on the concentration






Fig. 4 Cumulative carbon mineralization [mg $C(CO_2)$/100 g] in sandy textured soils treated with *Melia azedarach* leaves, extract, and green-synthesized nanoparticles. **a** Effects of Silver Nanoparticles (AgNPs); **b** Effects of Palladium Nanoparticles (PdNPs) applied at 10, 50, and 100 mg kg^{-1}. Values represent mean ± standard error (SE, $n = 3$). Different lowercase letters above bars indicate significant differences between treatments according to Tukey's HSD test at $p < 0.05$




Fig. 5 Cumulative carbon mineralization [mg $C(CO_2)$/100 g] in sandy textured soils. The figure displays the comparative effects of Control, *Melia azedarach* leaves alone, nanoparticles alone (100 mg kg^{-1} AgNPs and PdNPs), and their combined applications (Leaves + AgNPs; Leaves + PdNPs). Values represent mean ± standard error (SE, $n = 3$). Different lowercase letters above bars indicate significant differences between treatments according to Tukey's HSD test at $p < 0.05$




Fig. 6 Carbon mineralization ratios in (**a**) clay and (**b**) sandy textured soils treated with *Melia azedarach* leaves, extract, and green-synthesized nanoparticles. Treatments include Control, Leaves, Extract, and varying doses (10, 50, and 100 mg kg^{-1}) of AgNPs and PdNPs. Values represent mean ± standard error (SE, $n = 3$). Different lowercase letters indicate significant differences between treatments according to Tukey's HSD test at $p < 0.05$

of Ag and Pd. The decrease in peak intensity at 3300 cm^{-1}, corresponding to O–H stretching vibrations, indicated that the hydroxyl groups of the plant extract contribute to metal ion reduction and stabilization (Akturk et al. 2019; Thakur and Gupta 2025). The high similarity between the spectra of the synthesized NPs and the plant extract suggests that the NPs were stabilized by biomolecules present in the plant, such as phenolics and carbohydrate derivatives with high hydroxyl group content (Eker et al. 2025).

The XRD results were compared with previous studies in the literature, and the findings were found to be consistent. In a previous study, it was reported that AgNPs exhibit a face-centered cubic (FCC) structure, with the diffraction peaks at 38°, 44°, 64°, and 78° corresponding to the (111), (200), (220), and (311) crystal planes, respectively (Veisi et al. 2018; Raj et al. 2018).

Numerous studies have demonstrated that NPs exhibit different physicochemical and, consequently, toxicological behaviors depending on the soil type (Cornelis et al. 2014; Schlich and Hund Rinke 2015). Factors such as pH, soil moisture, soil temperature, organic matter, and microorganism count were some of the parameters influenced by NPs. Similarly, in a study by Rahmatpour et al. (2017), NPs applied to clay soils exhibited lower toxicity than those applied to sandy soils, which was attributed to the immobilization of NPs through heteroaggregation with clay particles (Rahmatpour et al. 2017).

The PdNPs and AgNPs were added to clay and sandy soils. The application of PdNPs caused both clay and sandy soils to become more mineralized. Additionally, after the extract was applied to the soil, both soils showed increased mineralization as a result of the addition of leaves and their extracts as a carbon source. This is because the extract was in a more easily decomposable state, which makes it easier for soil microorganisms to utilize it as a source of carbon. Results indicate that copiotrophic microorganisms most likely caused a rapid carbon mineralization phase to occur during the first week of incubation, which was followed by a slower phase that was mostly carried out by an oligotrophic microbial community. While oligotrophic (autochthonous) microorganisms are largely in charge of the breakdown of more resistant organic matter when C depletion occurs due to competition among microorganisms, copiotrophic (zygomorphic) microorganisms have a tendency to multiply when readily accessible organic matter is added to the soil (Dilly 2002; Yarwood 2021; Medina et al. 2025). Unlike conventional chemically synthesized NPs which often show direct toxicity, the green-synthesized NPs used in this study exhibited a texture-dependent impact pattern. This suggests that the organic corona derived from the plant extract may interact with clay surfaces differently than bare chemical NPs, a finding that adds a new dimension to the current understanding of nanoparticle-soil interactions. Our results suggested that the natural organic coating of NPs were a key determinant of their interactions with soils of differing structure. In clay-rich soils, this layer likely enhanced interactions with clay minerals, thereby regulating nanoparticle bioavailability and potentially mitigating their toxicity. These findings underscored that surface modifications from green synthesis are a critical determinant of NPs behavior in soils and should be carefully considered in environmental risk assessments. Some studies have shown that the microbial communities responsible for decomposition were significantly susceptible to NPs (Collins et al. 2012b; Batista et al. 2017; Grün et al. 2019). Pradhan et al. (2011) studied the effects of AgNPs on leaf microbial decomposition and demonstrated that exposure to these NPs resulted in a reduction in the rate of leaf decomposition (Pradhan et al. 2011). It was known that the interaction of NPs with organic compounds, such as leaves, can limit the bioavailability of essential nutrients to microorganisms, which in turn can reduce microbial activity (Simonet and Valcárcel 2009). In this study, both NPs added to the soil with leaves were found to restrict soil microbial activity. Moreover, AgNPs toxicity in soil microorganisms was reduced by increasing the soil pH, clay content and types; however, it was not affected significantly by organic content of soil (Schlich and Hund-Rinke 2015; Zou et al. 2016; Wang et al. 2020).

5 Conclusions

This study confirms the hypothesis that soil texture acts as a decisive factor in modulating the toxicity of green-synthesized nanomaterials. The findings demonstrate that both silver nanoparticles (AgNPs) and palladium nanoparticles (PdNPs) exert a negative impact on soil microbial activity; however, this effect is significantly more pronounced in sandy soils compared to clay soils. The differences in carbon mineralization between soils with distinct textures may be associated with the potential of clay minerals to mitigate nanoparticle toxicity through physicochemical interactions, such as reduced bioavailability, as opposed to the sandy soil where such interactions are limited. Additionally, the toxicity of AgNPs was found to be higher than that of PdNPs, likely due to their potent antibacterial characteristics. Furthermore, soils enriched with the plant leaf extract maintained higher rates of mineralization, indicating that the biochemical composition of the synthesis medium can support microbial recovery. This study contributes to the state of the art by providing comparative evidence that soil texture is a decisive factor in modulating the toxicity of green-synthesized nanoparticles. Our findings highlight that clay fractions provide a buffering effect against biogenic NPs, a critical parameter that must be considered in future environmental risk assessments of green nanomaterials.

Authors Contribution Nacide Kizildag Ozdal: Conceptualization, Data curation, Formal analysis, Funding acquisition, Methodology, Writing – original draft, Writing – review & editing. Bahar Meryemoglu: Conceptualization, Methodology, Formal analysis, Writing – original draft, Writing – review & editing. Husniye Aka Sagliker: Writing – original draft, Writing – review & editing.

Funding Open access funding provided by the Scientific and Technological Research Council of Türkiye (TÜBİTAK). This work was funded by Cukurova University Research Grant (Project number: FBA-2020-13343).

Declarations

Competing interests The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/.

References

Abid N, Khan AM, Shujait S, Chaudhary K, Ikram M, Imran M, Haider J, Khan M, Khan Q, Maqbool M (2022) Synthesis of nanomaterials using various top-down and bottom-up approaches, influencing factors, advantages, and disadvantages: A review. Adv Colloid Interface Sci 300:102597. https://doi.org/10.1016/j.cis.2021.102597

Ahmed B, Rizvi A, Ali K, Lee J, Zaidi A, Khan MS, Musarrat J (2021) Nanoparticles in the soil–plant system: a review. Environ Chem Lett 19:1545–1609. https://doi.org/10.1007/s10311-020-01138-y

Aigbe UO, Osibote OA (2024) Green synthesis of metal oxide nanoparticles, and their various applications. J Hazard Mater Adv 13:100401. https://doi.org/10.1016/j.hazadv.2024.100401

Akturk O, Gun Gok Z, Erdemli O, Yigitoglu M (2019) One-pot facile synthesis of silk sericin-capped gold nanoparticles by UVC radiation: Investigation of stability, biocompatibility, and antibacterial activity. J Biomed Mater Res 107:2667–2679. https://doi.org/10.1002/jbm.a.36771

Ali MA, Ahmed T, Wu W, Hossain A, Hafeez R, Islam Masum MM, Wang Y, An Q, Sun G, Li B (2020) Advancements in plant and microbe-based synthesis of metallic nanoparticles and their antimicrobial activity against plant pathogens. Nanomaterials 10:1146. https://doi.org/10.3390/nano10061146

Annu A, Ahmed S (2018) Green synthesis of metal, metal oxide nanoparticles, and their various applications. Handb Ecomaterials, 1–45

Batista D, Pascoal C, Cassio F (2017) Temperature modulates AgNP impacts on microbial decomposer activity. Sci Total Environ 601–602:1324–1332. https://doi.org/10.1016/j.scitotenv.2017.05.229

Benlot C (1977) Recherches sur les activites biochimiques dans les successions de sols derives de cendres volcaniques sous climat tropical humide (Zaire- Indonesie), ENS Lab. De Zoologie, Paris, pp 73–76. (In French)

Bouyoucos GS (1951) A recalibration of the hydrometer for making mechanical analysis of soil. Agron J 43:434–438

Bruni E, Yuste JC, Menichetti L, Flores O, Guasconi D, Guenet B, Manzoni S (2025) Microbial biomass–not diversity–drives soil carbon and nitrogen mineralization in Spanish holm oak ecosystems. Geoderma 460:117408

Chai H, Yao J, Sun J, Zhang C, Liu W, Zhu M, Ceccanti B (2015) The effect of metal oxide nanoparticles on functional bacteria and metabolic profiles in agricultural soil. Bull Environ Contam Toxicol 94:490–495. https://doi.org/10.1007/s00128-015-1485-9

Collins D, Luxton T, Kumar N, Shah S, Walker VK, Shah V (2012a) Assessing the Impact of Copper and Zinc Oxide Nanoparticles on Soil: A Field Study. PLoS ONE 7:e42663. https://doi.org/10.1371/journal.pone.0042663

Collins D, Luxton T, Kumar N, Shah S, Walker V (2012b) Influence of silver nanoparticle toxicity on freshwater microbial communities. J Hazard Mater 213:323–332

Colman BP, Arnaout CL, Anciaux S, Gunsch CK, Hochella MF (2013) Low concentrations of silver nanoparticles in biosolids cause adverse ecosystem responses under realistic field scenario. PLoS ONE 8:e57189. https://doi.org/10.1371/journal.pone.0057189

Cornelis G, Hund-Rinke K, Kuhlbusch T, Van den Brink N, Nickel C (2014) Fate and bioavailability of engineered nanoparticles in soils: A review. Crit Rev Environ Sci Technol 44:2720–2764. https://doi.org/10.1080/10643389.2013.829767

Demiralay I (1993) Toprak Fiziksel Analizleri, Atatürk Üniversitesi Ziraat Fakültesi Yayınları, Erzurum (In Turkish)

Dikshit PK, Kumar J, Das AK, Sadhu S, Sharma S, Singh S, Gupta PK, Kim BS (2021) Green Synthesis of Metallic Nanoparticles: Applications and Limitations. Catalysts 11(8):902. https://doi.org/10.3390/catal11080902

Dilly O (2002) Microbial energetics in soils. Biol Fertil Soils 35:367–376

Duchaufour P (1970) Precis de Pedologie, Masson et C^{ie} (In French)

Egbuna C, Parmar VK, Jeevanandam J, Ezzat SM, Patrick-Iwuanyanwu KC, Adetunji CO, Khan J, Onyeike EN, Uche CZ, Akram M, Ibrahim MS, El Mahdy NM, Awuchi CG, Saravanan K, TijjaniH, Odoh UE, Messaoudi M, Ifemeje JC, Olisah MC, Ezeofor NJ, Chikwendu CJ, Ibeabuchi CG (2021) Toxicity of nanoparticles in biomedical application: nanotoxicology. Hindawi J Toxic 21:9954443. https://doi.org/10.1155/2021/9954443

Eker F, Akdaşçi E, Duman H, Bechelany M, Karav S (2025) Green Synthesis of Silver Nanoparticles Using Plant Extracts: A Comprehensive Review of Physicochemical Properties and Multifunctional Applications. Int J Mol Sci 26:6222. https://doi.org/10.3390/ijms26136222

Fahim M, Shahzaib A, Nishat N, Jahan A, Bhat TA, Inam A (2024) Green synthesis of silver nanoparticles: a comprehensive review of methods, influencing factors, and applications. JCIS Open 16:100125. https://doi.org/10.1016/j.jciso.2024.100125

Gleiter H (2000) Nanostructured Materials: Basic Concepts and Microstructure. Acta Mater 48:1–29. https://doi.org/10.1016/S1359-6454(99)00285-2

Gohil K, Dhakhda S, Patel V, Rathod A, Singh PK (2024) Green Synthesis Of Silver Nanoparticles Using Various Plantextracts And Evaluation Of Their Antimicrobial Activities. Biosci Biotechnol Res Asia 21:765–777. https://doi.org/10.13005/Bbra/3263

Grün AL, Manz W, Kohl YL, Meier F, Straskraba S, Jost C, Drexel R, Emmerling C (2019) Impact of silver nanoparticles (AgNP) on soil microbial community depending on functionalization, concentration, exposure time, and soil texture. Environ Sci Eur 31:15. https://doi.org/10.1186/s12302-019-0196-y

Hasanzadeh A, Gholipour B, Rostamnia S, Eftekhari A, Tanomand A, Valizadeh KA, Khaksar S, Khalilov R (2021) Biosynthesis of AgNPs onto the urea-based periodic mesoporous organosilica

(AgxNPs/Ur-PMO) for antibacterial and cell viability assay. J Colloid Interface Sci 585:676–683. https://doi.org/10.1016/j.jcis.2020.10.047

Hong J, Wang L, Sun Y, Zhao L, Niu G, Tan W, Rico CM, Peralta-Videa JR, Gardea-Torresdey JL (2016) Foliar applied nanoscale and microscale CeO_2 and CuO alter cucumber (*Cucumis sativus*) fruit quality. Sci Total Environ 563–564:904–911. https://doi.org/10.1016/j.scitotenv.2015.08.029

Jackson ML (1958) Soil Chemical Analysis. Prentice-Hall, Inc., Englewood Cliffs

Keller AA, Lazareva A (2013) Predicted releases of engineered nanomaterials: from global to regional to local. Environ Sci Technol Lett 1:65–70. https://doi.org/10.1021/ez400106t

Keller AA, McFerran S, Lazareva A, Suh S (2013) Global life cycle releases of engineered nanomaterials. J Nanopart Res 15:1692. https://doi.org/10.1007/s11051-013-1692-4

Khan I, Saeed K, Khan I (2019) Nanoparticles: Properties, applications and toxicities. Arab J Chem 12:908–931. https://doi.org/10.1016/j.arabjc.2017.05.011

Kizildag K, Cenkseven S, Koca FD, Aka Sagliker H, Darici C (2019) How titanium dioxide and zinc oxide nanoparticles do affect soil microorganism activity? Eur J Soil Biol 91:18–24. https://doi.org/10.1016/j.ejsobi.2019.01.001

Kleinbaum DG, Kupper LL, Muller KE, Nizam A (1998) Applied Regression Analysis and Other Multivariable Methods. Duxbury, Pacific Grove, CA, USA

Kumar R, Kumar A, Bhardwaj S, Sikarwar M, Sriwastaw S, Sharma G, Gupta M (2025) Nanotoxicity unveiled: Evaluating exposure risks and assessing the impact of nanoparticles on human health. J Trace Elem Med Biol 13:100252. https://doi.org/10.1016/j.jtemin.2025.100252

Kumari S, Raturi S, Kulshrestha S, Chauhan K, Dhingra S, András K, Thu K, Khargotra R, Singh T (2023) A comprehensive review on various techniques used for synthesizing nanoparticles. J Mater Res Technol 27:1739–1763. https://doi.org/10.1016/j.jmrt.2023.09.291

Liang G, Stefanski A, Eddy WC, Bermudez R, Montgomery RA, Hobbie SE, Reich PB (2024) Soil respiration response to decade-long warming modulated by soil moisture in a boreal forest. Nat Geosci 17(9):905–911

Loveland P, Webb J (2003) Is there a critical level of organic matter in the agricultural soils of temperate regions: a review. Soil Tillage Res 70:1–18. https://doi.org/10.1016/S0167-1987(02)00139-3

Medina J, Panettieri M, Calabi-Floody M, Plaza C, Knicker H (2025) Biochar and metallic nanoparticle additives in agricultural residues composting modulate the mineralization patterns of end-Products when added to tropical soils. J Soil Sci Plant Nutr 25:2261. https://doi.org/10.1007/s42729-025-02266-6

Melo APZ, Maciel M, Sganzerla WG, Almeida AR, Armas RD, Machado MH, Rosa CG, Nunes MR, Bertoldi FC, Barreto PLM, Barreto (2020) Antibacterial activity, morphology, and physicochemical stability of biosynthesized silver nanoparticles using thyme (*Thymus vulgaris*) essential oil. Mater Res Express 7:015087. https://doi.org/10.1088/2053-1591/ab6c63

Muzammil S, Ashraf A, Siddique MH, Aslam B, Rasul I, Abbas R, Afzal M, Faisal M, Hayat S (2023) A review on toxicity of nanomaterials in agriculture: Current scenario and future prospects. Sci Prog 106:4. https://doi.org/10.1177/00368504231221672

Narasaiah P, Mandal BK, Sarada NC (2017) Biosynthesis of Copper Oxide nanoparticles from Drypetes sepiaria Leaf extract and their catalytic activity to dye degradation. IOP Conf Ser Mater Sci Eng 263:022012. https://doi.org/10.1088/1757-899X/263/2/022012

Nassem T, Durrani T (2023) The role of some important metal oxide nanoparticles for wastewater and antibacterial applications: A review. Environ Chem Ecotoxic 3:59–75. https://doi.org/10.1016/j.enceco.2020.12.001

Osman AI, Zhang Y, Farghali M, Rashwan AK, Eltaweil AS, Abd El-Monaem EM, Mohamed IMA, Badr MM, Ihara I, Rooney DW, Yap PS (2024) Synthesis of green nanoparticles for energy, biomedical, environmental, agricultural, and food applications: A review. Environ Chem Lett 22:841–887. https://doi.org/10.1007/s10311-023-01682-3

Pascual JA, Garcia C, Hernandez T, Moreno JL, Ros M (2000) Soil microbial activity as a biomarker of degradation and remediation processes. Soil Biol Biochem 32(13):1877–1883

Patra JK, Baek KH (2016) Green nanobiotechnology: Factors affecting synthesis and characterization techniques. J Nanomater 3531725:1–2. https://doi.org/10.1155/2016/3531725

Poeplau C, Don A, Six J, Kaiser M, Benbi D, Chenu C, Cotrufo MF, Derrien D, Gioacchini P, Grand S, Gregorich E, Griepentrog M, Gunina A, Haddix M, Kuzyakov Y, Kühnel A, Macdonald LM, Soong J, Trigalet S, Vermeire ML, Rovira P, van Wesemael B, Wiesmeier M, Yeasmin S, Yevdokimov I, Nieder R (2018) Isolating organic carbon fractions with varying turnover rates in temperate agricultural soils – a comprehensive method comparison. Soil Biol Biochem 125:10–26. https://doi.org/10.1016/j.soilbio.2018.06.025

Pradhan A, Seena S, Dobson M, Pascoal C (2011) Interactive effects of copper and silver nanoparticles on leaf decomposition and associated fungi. Environ Sci Techno 45:9366–9372

Rahmatpour S, Shirvani M, Mosaddeghi MR (2017) Effects of silver nanoparticles on soil microbial biomass and enzyme activities in calcareous and acidic soils. Environ Monit Assess 189:1–12

Raj S, Singh H, Trivedi R, Pandey A (2018) Green synthesis and structural characterization of silver nanoparticles using plant extracts. Int J Nanosci 17:1850013

Rajput VD, Minkina T, Sushkova S, Behal A, Singh R, Mandzhieva S (2018) Effects of nanoparticles on crops and soil microbial communities. J Soils Sediments 18:2179–2187

Ramkumar V, Shanmugam S, Sundar S (2017) Plant-mediated synthesis of silver nanoparticles and their antimicrobial activity. Appl Nanosci 7:253–259

Rieke EL, Cappellazzi SB, Cope M, Liptzin D, Mac Bean G, Greub KL, Honeycutt CW (2022) Linking soil microbial community structure to potential carbon mineralization: a continental scale assessment of reduced tillage. Soil Biol Biochem 168:108618. https://doi.org/10.1016/j.soilbio.2022.108618

Rodríguez-Albarracín HS, Demattê JA, Rosin NA, Rosas JTF, Darghan AE, de Carvalho HWP (2025) A progressive knowledge strategy for monitoring soil microbiological activity by microscale spectroscopic detection. Soil Adv 4:100058

Saleh TA (2020) Nanomaterials: Classification, properties, and environmental toxicities. Environ Technol Innov 20:101067. https://doi.org/10.1016/j.eti.2020.101067

Sasikala A, Tamilselvi S, Arulmozhi R (2014) Green synthesis of silver nanoparticles using plant extracts and their antimicrobial activities: A review. Int J Curr Microbiol Appl Sci 3:235–243

Schaefer R (1967) Caracteres et evolution des activites microbiennes dans une chaine de sols hydromorphes mesotrophiques de la plaine d'alsace première partie: Cadre géographique et milieu édaphique. Rev Ecol Biol Sol 4:567–592 (In French)

Schlich K, Hund-Rinke K (2015) Influence of soil properties on the effects of silver nanoparticles on microbial activity in soils. Environ Pollut 196:321–330

Sedefoğlu N (2024) Green synthesis of ZnO nanoparticles by *Myrtus communis* plant extract with investigation of effect of precursor, calcination temperature and study of photocatalytic performance. Ceram Int 50:9884–9895. https://doi.org/10.1016/j.ceramint.2024.01.387

Sedefoğlu N, Zaloğlu N, Bozok F (2022) Green synthesized ZnO nanoparticles using *Ganoderma lucidum*: Characterization and *In*

Vitro Nanofertilizer effects. J Alloys Compd 918:165695. https://doi.org/10.1016/j.jallcom.2022.165695

Shah V, Belozerova I (2009) Influence of metal nanoparticles on the soil microbial community and germination of lettuce seeds. Water Air Soil Pollut 197(1):143–148

Shah ST, Sari IP, Yanto DHY, Chowdhury ZZ, Bashir MN, Badruddin IA, Lee JS (2025) Nature's nanofactories: biogenic synthesis of metal nanoparticles for sustainable technologies. Green Chem Lett Rev 18:1. https://doi.org/10.1080/17518253.2024.2448171

Shao M, Zhang S, Niu B, Pei Y, Song S, Lei T, Yun H (2022) Soil texture influences soil bacterial biomass in the permafrost-affected alpine desert of the Tibetan plateau. Front Microbiol 13:1–12

Sharma R, Kumar V (2024) Nano enabled agriculture for sustainable soil. Waste Manage Bull 2:152–161. https://doi.org/10.1016/j.wmb.2024.01.002

Shrestha S, Wang B, Dutta P (2020) Nanoparticle processing: Understanding and controlling aggregation. Adv Colloid Interface Sci 279:102162. https://doi.org/10.1016/j.cis.2020.102162

Simonet BM, Valcárcel M (2009) Monitoring nanoparticles in the environment. Anal Bioanal Chem 393:17–21

Simonin M, Richaume A (2015) Impact of engineered nanoparticles on the activity, abundance, and diversity of soil microbial communities: A review. Environ Pollut 202:119–129

Singh P, Kim YJ, Zhang D, Yang DC (2016) Biological synthesis of nanoparticles from plants and microorganisms. Trends Biotechnol 34:588–599

Sinha T, Ahmaruzzaman M, Sil AK (2014) Green synthesis of silver nanoparticles using plant extract and their application in dye degradation. J Environ Chem Eng 2:66–72

Szőllősi R, Molnár Á, Kondak S, Kolbert Z (2020) Dual Effect of Nanomaterials on Germination and Seedling Growth: Stimulation vs. Phytotoxicity Plants 9:1745. https://doi.org/10.3390/plants9121745

Taha T, Barzinjy A, Hussain F, Nurtayeva T (2022) Nanotechnology and computer science: trends and advances. Memories - Mater Devices Circuits Syst 2:100011. https://doi.org/10.1016/j.memori.2022.100011

Thakur S, Gupta R (2025) Medicinal plants derived green synthesis of nanoparticles: classification, methods, applications and environmental impacts. Nanotechnol Environ Eng 10:27. https://doi.org/10.1007/s41204-025-00421-z

Veisi H, Azizi S, Mohammadi P (2018) Green synthesis of silver nanoparticles using plant extracts and their applications: A review. J Clean Prod 170:231–248

Wang Y, Liu Y, Yang K, Lin D (2020) Reciprocal interference of clay minerals and nanoparticulate zero-valent iron on their interfacial interaction with dissolved organic matter. Sci Total Environ 739:140372. https://doi.org/10.1016/j.scitotenv.2020.140372

Xu C, Peng C, Sun L, Zhang S, Huang H, Chen Y, Shi J (2015) Distinctive effects of TiO_2 and CuO nanoparticles on soil microbes and their community structures in flooded paddy soil. Soil Biol Biochem 86:24–33. https://doi.org/10.1016/j.soilbio.2015.03.011

Yarwood SA (2021) Microbial ecology Gentry, T.J. Fuhrmann, J.J. Zuberer D.A. (Eds.), Principles and Applications of Soil Microbiology (3rd ed.), Elsevier, Amsterdam, 239–267

You T, Liu D, Chen J, Yang Z, Dou R, Gao X, Wang L (2018) Effects of metal oxide nanoparticles on soil enzyme activities and bacterial communities in two different soil types. J Soils Sediments 18:211–221. https://doi.org/10.1007/s11368-017-1716-2

Zhang F, Liu Y, Zhang Y (2023) Bibliometric analysis of research trends in agricultural soil organic carbon mineralization from 2000 to 2022. Agriculture 13(6):1248

Zhu Y, Wu J, Chen M, Liu X, Xiong Y, Wang Y, Feng T, Kang S, Wang X (2019) Recent advances in the biotoxicity of metal oxide nanoparticles: impacts on plants, animals and microorganisms. Chemosphere 237:124403. https://doi.org/10.1016/j.chemosphere.2019.124403

Zou X, Li P, Huang Q, Zhang H (2016) The different response mechanisms of Wolffia globosa: lightinduced silver nanoparticle toxicity. Aquat Toxicol 176:97–105. https://doi.org/10.1016/j.aquatox.2016.04.019

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.