

Content is available at: CRDEEP Journals
Journal homepage: <http://www.crdeepjournal.org/category/journals/ijls/>

International Journal of Life Sciences
(ISSN: 2277-193x) (Scientific Journal Impact Factor: 6.106)

UGC Approved-A Peer Reviewed Quarterly Journal



Research Paper

Leaf Dust Accumulation and Its Impact on Chlorophyll of Three Tree Species Growing in The Region of Mauganj District

Anshu Rani Patel^{1*} and Kajal Singh Patel¹

¹Department of Environmental Biology, A.P.S. University, Rewa (M.P.), 486003 – India.

ARTICLE DETAILS

Corresponding Author:

Anshu Rani Patel

Key words:

Rhizosphere, plant microbiome, root exudates, chemical signaling, plant growth-promoting rhizobacteria, symbiosis

ABSTRACT

This has been conducted in Mauganj M.P. which is recently announced as district. This study determined the dust deposition and its impact on foliar chlorophyll contents of three tree species (*Azadirachta indica*, *Bauhinia variegata* and *Bombax ceiba*) growing in the Mauganj (M.P.). Results indicated higher dust load and lower chlorophyll content in the leaves of observed plants at polluted sites, as compared to control site. A negative correlation has been calculated between dust load and chlorophyll contents of leaves sampled from polluted sites. There was marked seasonal variation in the foliar dust accumulation and chlorophyll contents. Leaf samples collected during winter months showed higher accumulation of dust and lower chlorophyll contents as compared to summer and monsoon months leaf samples.

1. Introduction

Mauganj is a district and administrative headquarters in the Indian state of Madhya Pradesh, located in the northeastern region near the Uttar Pradesh border. Carved out of the Rewa district, it operates as the 53rd district in the state and is particularly well known for its scenic waterfalls and rich historical roots. Growing urbanization and development make this district strong polluted site. Plants play significant role in removal of particulate matters from ambient air of industrial as well as urban regions by intercepting and retaining them on their leaf surface. The foliage of plants acts as persistent absorber and filter of air. Leaves of the plants can effectively absorb air particulates (Freer-Smith et al. 1997, Prusty et al. 2006, Nowak et al. 2006, Jamil et al. 2009, Qiu et al. 2009) and reduce air pollution (Nowak et al. 2006, Hofman et al. 2014, Lin et al. 2017). The leaf morphology and biochemical characteristics such as ascorbic acid, relative water and chlorophyll contents are influenced by accumulation of dust on plant leaves.

There is variation in ability of each plant species to absorb and adsorb pollutants by their foliar surface and depends on several biochemical, physiological and morphological characteristics (Seyyednejad et al. 2011). The leaf tissues are directly injured by cement dusts with PH values of 9 or higher (Vardaka et al. 1995) or indirectly injured through alteration of soil pH (Hope et al. 1991, Auerback et al. 1997). The present study was conducted to assess dust accumulation on leaves of three tree species growing in the J.P. cement plant region and to study the correlation between dust load and chlorophyll contents of leaves.

2. Materials and Methods:

This study was conducted in the Mauganj (M.P.) which is situated about 70 km away toward north-west direction from Rewa city. The study site is situated at latitude 24° -20' N and longitude 81° -20' E. Dust deposited fresh leaves of three tree species namely *Azadirachta indica*, *Bauhinia variegata* and *Bombax ceiba* growing in cement plant region were collected during winter, summer and monsoon months of 2025. Fifteen replicates of nearly equal size leaves from branches of almost similar height of each plant species growing in different sites of Mauganj region as well as in A.P.S. university campus (as control) was collected and packed in pre-weighted polythene and brought to the laboratory. The dust adhering

¹Corresponding Author can be contacted at: dr.anshuapsu@gmail.com

Received: 06-06-2026; Sent for Review on: 12-06-2026; Draft sent to Author for corrections: 16-06-2026; Accepted on: 18-06-2026; Online Available from 20-06-2026

DOI: [10.5281/zenodo.20768351](https://doi.org/10.5281/zenodo.20768351)

IJLS-8880/© 2026 CRDEEP Journals. All Rights Reserved.

on the dorsal and ventral surface of leaves was carefully cleaned with the help of fine brush in the same polythene. Again the weight of polythene was taken with the help of electronic balance in order to determine the amount of dust present on the leaf surface. After cleaning, the leaf area was calculated with the amount of graph paper. Each leaf sample outline was drawn on a graph paper and then the number of squares was counted in cm^2 to obtain the leaf area. The amount of dust deposited on leaf surface in mg/cm^2 was calculated by the following formula (Prajapati and Tripathi 2008).

$$W = (W_2 - W_1)/a$$

Where; W is dust content (mg/cm^2), w_1 is initial weight of polythene, W_2 is final weight of polythene with dust and ' a ' is total area of leaf (cm^2).

2.1 Estimation Of Chlorophyll:

Chlorophyll extraction was done as per method of Arnon (1994). One gram of fresh leaves was blended and then extracted with 20-40 ml of 80% acetone and left for 15 minutes. The liquid portion was taken in to another tube and was then centrifuged at 5000-10000 rpm for 5 minutes. The supernatant of the sample was transferred and the absorbance was taken at 645 nm and 663 nm with the help of spectrophotometer. The absorbance at 645 nm and 663nm against the solvent (acetone) blank was also taken.

The concentration of chlorophyll 'a', chlorophyll 'b' and total chlorophyll were calculated using the following equation:

Chlorophyll 'a' : $12.7(A_{663}) - 2.69(A_{645})$.

Chlorophyll 'b' : $22.9(A_{645}) - 4.68(A_{663})$.

Total chlorophyll : $20.2(A_{645}) + 8.02(A_{663})$.

3. Results and Discussion:

Foliar dust accumulation varied seasonally on the leaves of three plant species at both polluted and control sites (Table-1). Winter leaf samples of three species accumulated higher dust followed by summer and monsoon leaf samples. The higher dust accumulation in winter months may be due to cool and foggy condition and wet surface of leaves that help in dust capturing, preventing particulate dispersion as suggested by Rai and Panda (2014). Whereas in monsoon month, washing of leaves due to rains might have accounted lower dust accumulation. High wind speed may be the reason for comparatively lower dust deposition on leaves during summer months (Prajapati and Tripathi 2008).

Table 1: Seasonal dust accumulation (mg/cm^2) on the leaves of three tree species growing in Mauganj region (polluted) and control site

Species	Polluted Site			Control Site		
	Winter	Summer	Raining	Winter	Summer	Raining
<i>Azadirachta indica</i>	0.854	0.548	0.290	0.196	0.153	0.069
<i>Bauhinia variegata</i>	0.521	0.298	0.173	0.098	0.052	0.041
<i>Bombax ceiba</i>	0.598	0.365	0.231	0.079	0.063	0.050

Average dust deposition (mg/cm^2) has been determined for tree species growing at polluted and control sites (Fig.1). *Azadirachta indica* leaves accumulated the higher amount of dust as $0.582 \text{ mg}/\text{cm}^2$ in J.P. cement plant region followed by *Bombax cieba* ($0.398 \text{ mg}/\text{cm}^2$) and *Bauhinia variegata* ($0.330 \text{ mg}/\text{cm}^2$). The average foliar dust accumulation at control site was observed as $0.139 \text{ mg}/\text{cm}^2$, $0.064 \text{ mg}/\text{cm}^2$ and $0.063 \text{ mg}/\text{cm}^2$ for *Azadirachta indica*, *Bombax ceiba* and *Bauhinia variegata* respectively. Thus, *Azadirachta indica* leaves accumulated 4 times more dust at polluted site than control site, while *Bauhinia variegata* accumulated 5 times more dust than control site. The plant species *Bombax ceiba* growing at polluted site accumulated 6 times more dust than that of control site.




Fig. 1 : Average dust accumulation (mg/cm^2) on the leaves of tree species in polluted and control site.

Dust deposition on leaf is influenced by micro-morphology (epidermal and cuticular features, presence /absence of grooves and trichomes, wax coating), orientation (horizontal or vertical), sessile or semi- sessile nature, size and thickness, and folded margins of leaves and height and canopy of plant species (Davison and Blackmore 1976, Garg et al. 2000, Pal et

al. 2002, Prusty et al. 2005, Prajapati and Tripathi 2008, Keler et al. 2016, Sharma et al. 2017). Leaf micro-morphology such as groove area and trichomes have direct effect on particulate deposition (Chen et al. 2017). Higher dust accumulation on *Azadirachta indica* leaves may be due to rough foliar surface with depressions in the middle of the leaf and small petiole. Small petiole reduces movement of leaves leading to accumulation of dust. Comparatively lesser accumulation of dust on *Bombax ceiba* leaves may be attributed to smooth surface and thin lamina of leaf. The medium height of *Bauhinia variegata* might be responsible for least accumulation of dust on leaves. A negative correlation has been calculated between dust load and chlorophyll content of leaves of *Azadirachta indica* ($r = -0.990$), *Bauhinia variegata* ($r = -0.784$) and *Bombax ceiba* ($r = -0.999$) growing at polluted sites i.e. higher the dust deposition; lower the leaf chlorophyll content of the plant species.

Leaf samples of plant species growing at polluted and control site were analysed for chlorophyll 'a', chlorophyll 'b' and total chlorophyll contents for three seasons (Table-2). Results indicated marked seasonal variation in chlorophyll contents of tree species. Monsoon month's leaves showed higher content of chlorophyll followed by summer and winter month's samples at both the sites. The leaf samples of observed plants exhibited reduced contents of chlorophyll 'a', chlorophyll 'b' and total chlorophyll at polluted sites as compared to control ones. Several recent studies have demonstrated that dust accumulation causes degradation in photosynthetic pigments (Singh and Siddiqui 2003, Prajapati and Tripathi 2008, Rai and Panda 2014, Singh and Pal 2017, Singh et al. 2018, Walia et al. 2019). Prajapati and Tripathi (2008) have suggested that dust particles might be having various metals and polycyclic hydrocarbons, inhibiting the enzymes necessary for synthesizing chlorophyll. Gunamani et al. (1991) are of the opinion that alkaline conditions prevailing due to solubility of dust particulate in the cell sap are responsible for chlorophyll degradation.

Table 2: Seasonal content of chlorophyll 'a','b' and total chlorophyll (mg/g) in leaves of three tree species growing at polluted and controlled sites.

Species	Sites	Season								
		Winter			Summer			Monsoon		
		Chl-a	Chl-b	TC	Chl-a	Chl-b	TC	Chl-a	Chl-b	TC
<i>Azadirachta indica</i>	Polluted	2.110	1.002	3.112	3.063	1.102	4.165	4.887	1.815	6.702
	Controlled	2.740	1.240	3.980	3.334	1.281	4.615	5.412	1.837	7.249
<i>Bauhinia variegata</i>	Polluted	7.814	3.114	10.928	10.411	5.001	15.412	12.001	6.320	20.321
	Controlled	13.202	1.022	14.224	16.321	3.132	19.453	20.458	8.022	26.474
<i>Bombax ceiba</i>	Polluted	15.216	3.563	18.779	19.564	5.181	24.745	22.713	7.820	30.526
	Controlled	15.513	6.982	22.495	19.578	9.820	29.398	22.660	12.616	35.267

Chl 'a'-Chlorophyll 'a', Chl 'b'- Chlorophyll 'b', TC -Total Chlorophyll.
Polluted - Mauganj Sites, Controlled - University Campus.

4. Conclusion

This study concludes that plant species are reducing the particulate loads of ambient air in the Mauganj district by intercepting and accumulating them on their foliar surface. This study also indicates that particulate matters of ambient air in the region are adversely affecting the plant species. The accumulated dust on foliar surface appeared to be responsible for degradation of chlorophyll, a photosynthetic pigment.

Acknowledgement: Authors are thankful to M.P.C.S.T. Bhopal (M.P.) for financial assistance

References

- Arnon, D. I. 1949. Copper enzyme in isolated chloroplasts, polyphenoloxidase in *Beta vulgaris* plant physiology, 24:1-15.
- Auerbach, N.A., Walker, M.D. and Walker, D.A. 1997. Effects of roadside disturbance on substrate and vegetation properties in arctic tundra. Ecol. Appl., 7: 218-235.
- Chen, L., Lin, L., Zhang, L., Zou, R. and Zhang, Z. 2017. Variation in tree species ability to capture and retain airborne fine particulate matter (PM 2.5). Scientific Reports, 7:3206 (DOI): 10. 1038/5 41598-017-03360-1.
- Davison, A. and Blackmore, J. 1976. Effects of air pollution on plants. In : T.A. Mansfield (Ed.), Cambridge University Press, Cambridge, pp. 209.
- Freer- Smith, P.H., Sophy, H. and Goodman, A. 1997. The uptake of particulates by an urban woodland: site description and particulate composition. Environ. Pollut., 95: 27-35.
- Garg, S.S., Kumar, N. and Das, G 2000. Effect of Bansal Ramraj mill dust on vegetation and health at Jaitwara, Distt Satna. Indian Journal of Environmental Protection, 20 : 326-328.
- Gunamani, T., Gurusamy, R. and Swamynathan, K. 1991. Effect of dust pollution on the dermal appendages and anatomy of leaves in some herbaceous plants. J. Swamy Boli club, 8(3-4) : 79-85.
- Hofman, J., Bartholomeus, H., Calders, K., Wittenberghe, S.V., Wuyts, K. and Samson, R. 2014. On the relation between tree crown morphology and particulate matter deposition on urban tree leaves: A ground based LIDAR approach. Atmos. Environ., 99 : 130-139.

- Hope, A.S., Fleming, J.B., Stow, D.A. and Aguado, E. 1991. Tussock tundra albedos on the north slope of Alaska: Effect of illumination, vegetation composition and dust deposition. *Journal of Applied Meteorology*, 30 (8) : 1200-1206.
- Jamil, S., Abhilash, P.C., Singh, A., Singh, N. and Behl, H.M. 2009. Fly ash trapping and metal accumulation capacity of plants: Implication for green belt around thermal power plants. *Landscape urban plan.*, 92 : 136-147.
- Keller, N.S., Bhardwaj, S.K., Pant, K.S. and Rai, T.S. 2016. Determination of leaf dust accumulation on certain plant species grown alongside National Highway-22, India. *Current World Environment*, 11(1): 77-82.
- Liu, Y., yang, Z., Zhu, M and Yin, J. 2017. Role of plant leaves in removing airborne dust and associated metals on Beijing roadsides. *Aerosol and Air Quality Research*, 17: 2566-2584.
- Nowak, D.J., Crane, D.E. and Stevens, J.C. 2006. Air pollution removal by urban trees and shrubs in the United States. *Urban for. Urban greening*, 4:115-123.
- Pal, A., Kulshreshtha, K., Ahmad, K. J. and Behl, H.M. 2002. Do leaf surface characters play a role in plant resistance to auto exhaust pollution? 197 : 47-55.
- Prajapati, S. K. and Tripathi B.D. 2008. Seasonal variation of leaf dust accumulation and pigment content in plant species exposed to urban particulates pollution. *Journal of Environmental Quality*, 37: 865-870.
- Prusty, B.A.K., Mishra, P.C. and Azeez, P.A. 2005. Dust accumulation and leaf pigment concentration in vegetation near the national highway at Sambalpur, Orissa, India. *Ecotoxicol. Environ. Saf.*, 60 : 228-235.
- Qiu, Y., Guan, D., Song, W. and Huang, K. 2009. Capture of heavy metals and sulfur by foliar dust in urban Huizhou, Guangdong province, China. *Chemosphere*, 75:447- 452.
- Rai, P. K., and Panda, L.S.2014. Leaf dust deposition and its impact on biochemical aspects of some roadside plants of Aizawl, Mizoram, North East India. *International Research Journal of Environmental Science*, 3(ii) : 14-19.
- Seyyednejad, S.M., Majdian, K., Koochak, H. and Niknejad, M. 2011. Air pollution tolerance indices of some plants around industrial zone in south of Iran. *Asian Journal of Biological Sciences*, 4: 300-305.
- Sharma, B., Chanda, S. K. and Bhuyan, M. 2017. Impact of dust accumulation on three roadside plants and their adaptive responses at National Highway 37, Assam, India. *Tropical Plant Research*, 4(1) : 61-167.
- Singh, I. and Siddique, Z.A. 2003. Effect of *Alternaria tritici* and foliar flyash deposition on growth, yield, photosynthetic pigment, proteins and lysine contents of three cultivars of wheat. *Bioresour Technol.*, 26 (2) :189-192.
- Singh, P. and Pal, A. 2017. Response of dust accumulation on roadside plant species due to open cast mining at Jhansi-Allahabad NH-76, Uttar Pradesh, India. *Tropical Plant Research*, 4(3) : 461-467.
- Singh, P., Mishra, R.M. and Shrivastava, R. 2018. Leaf dust deposition and its impact on chlorophyll content of *Cassia fistula* and *Encalyptus globulus* growing in the vicinity of Jaypee cement plant, Mauganj (M.P.). *IJE SRT*, 7 (8) : 471-476.
- Vardaka, E., Cook, C.M., Lanaras, T., Sgardelis, S.P. and Pantis, J.D. 1995. Effect of dust from a limestone quarry on the photosynthesis of *Quercus Coccifera* and evergreen sclerophyllous shrub. *Bull. Environ. Contam. Toxicol.*, 54 : 414-419.
- Walia, K., Aggarwal, R.K. and Bhardwaj, S.K. 2019. Leaf dust accumulation and its relationship with biochemical parameters of different plant species growing along national highway- 22, India. *International Journal of Chemical Studies*, 7 (1) : 1386-1390.