

Nephroprotective Potential of Bauhinia variegata: Phytochemistry, Pharmacological Activities, and Therapeutic Perspectives

Sarika More^{*1}, Ansari MD Huzaifa Siraj Ahmad², Taarique Deshmukh³, Rashid Akhtar Nehal Ahmad⁴, Shripad Bairagi⁵

^{1,2,3,4,5} Royal College of Pharmaceutical Education and Research, Malegaon, Nashik, Maharashtra 423203.

ABSTRACT

Kidney diseases are a problem for people all around the world and they make a lot of people very sick. Sometimes the medicines we take can hurt our kidneys, which's why we need to find safe and good ways to protect them. Plants that are used in medicine are looking like they could be an option because they have lots of different things in them that can help us. Bauhinia variegata Linn. Or Kachnar, is a plant that has been used for a time to help with lots of health problems like when we are inflamed have problems with the way our bodies work get infections or hurt our tissues. This plant has lots of things in it like flavonoids, phenolic acids, tannins, glycosides, saponins and alkaloids that make it work well for our health. Many studies have shown that Bauhinia variegata is good at stopping cells from getting hurt reducing inflammation fighting off germs helping with diabetes protecting our livers and keeping our cells safe. It seems like these good things about Bauhinia variegata can help protect our kidneys from getting hurt by things that can cause stress and damage to our kidney cells. This review is going to talk about what Bauhinia variegata looks like what is in it how it has been used and what it can do for our health how it can help our kidneys. We will also talk about what researchers have found out if it is safe to use and what we might be able to do with Bauhinia variegata in the future to help people with kidney problems. What we know far says that Bauhinia variegata is a good natural way to prevent and treat kidney problems and we should do more studies to see if it really works. Bauhinia variegata is a kidney disease solution that is looking very promising. We need to do more research on it to see how it can help people, with kidney diseases.

Keywords: Bauhinia variegata, Nephroprotection, Kidney Disorders, Medicinal Plants, Phytochemicals, Oxidative Stress, Antioxidant Activity, Renal Protection, Drug-Induced Nephrotoxicity, Herbal Medicine.

INTRODUCTION

The kidneys are important organs that help keep our body in balance by controlling the amount of fluid, salt and other substances in our blood. They also help get rid of waste products that our body does not need. The kidneys do a lot more than clean our blood they also make hormones that help our body work properly. If the kidneys are not working well it can cause problems and even lead to kidney disease. Kidney disease is an issue that affects a lot of people around the world.

Some medicines can actually hurt our kidneys. Cause them to stop working properly. This is called kidney

damage. It can happen when we take certain medicines like antibiotics pain relievers and cancer treatments. One medicine that is known to hurt the kidneys is Gentamicin. It is an antibiotic that doctors use to treat infections. It can only be used for a short time because it can cause kidney damage.

Plants have been used for a time to help treat different health problems. They have a lot of chemicals that can help our body heal and stay healthy. One plant that has been studied a lot is called Bauhinia variegata or Kachnar. It is a plant that grows in places and has been used in traditional medicine for a long time to treat things like inflammation, infections and tissue damage.

Relevant conflicts of interest/financial disclosures: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Bauhinia variegata has a lot of chemicals that are good for our body. It has things like flavonoids, phenolic compounds and alkaloids that can help reduce inflammation and protect our kidneys. Scientists have studied this plant. Found that it can help prevent kidney damage.

This article is going to talk about the Bauhinia variegata plant and how it can help our kidneys. We will discuss what the plant looks like what kind of chemicals it has and how it has been used in traditional medicine. We will also talk about how it can help protect our kidneys and what scientists have learned from studying this plant. The goal is to see if Bauhinia variegata can be used as a treatment, for kidney disease.

2. Botanical Profile of Bauhinia variegata

Bauhinia variegata Linn. is a medium-sized deciduous tree belonging to the family Fabaceae (subfamily Caesalpinioideae). It is one of the most famous species of the genus Bauhinia, which includes many flowering plants that grow in tropical and subtropical regions of the world. The plant popularly known as Kachnar in India and has been extensively used in traditional medicine for its multiple therapeutic properties. This species is appreciated not only for its medicinal value but also for its ornamental value with attractive orchid-like flowers and unique bilobed leaves.

Taxonomical Classification

- **Kingdom:** Plantae
- **Phylum:** Tracheophyta
- **Class:** Magnoliopsida
- **Order:** Fabales
- **Family:** Fabaceae
- **Subfamily:** Caesalpinioideae
- **Genus:** Bauhinia
- **Species:** Bauhinia variegata Linn.

Vernacular Names

Bauhinia variegata is known by different names in various regions:

- **English:** Orchid Tree, Mountain Ebony
- **Hindi:** Kachnar
- **Marathi:** Kanchan
- **Sanskrit:** Kovidara
- **Tamil:** Mandarai
- **Telugu:** Devakanchanam
- **Bengali:** Rakta Kanchan
- **Gujarati:** Kanchanar

Geographical Distribution

The plant is native to the South and Southeast Asia and is widely distributed in India, Nepal, Pakistan, Bangladesh, Myanmar, China and Thailand. Bauhinia variegata is common in tropical and subtropical areas of India especially in forests, along roadsides, in gardens and cultivated landscapes. It can tolerate many different climate conditions and is used often as an ornamental tree for its beauty.

Morphological Characteristics

Tree:

Bauhinia variegata is a deciduous tree growing to 10–15 m tall. The trunk is with moderate branches, the bark grayish brown and rough in texture.

Leaves:

The leaves are simple, alternate and characteristically bilobed like a camel's hoof. They are broad, leathery leaves, normally 8 to 15 cm long. The odd shape of the leaves is an important identifying characteristic of the species.

Flowers:

The flowers are large and fragrant and showy, appearing in clusters in flower season. Their colours



range from white to pink, purple or lavender and they have five petals with irregular markings. They look like orchid flowers and this is the reason for the plant's common name "Orchid Tree".

Fruits:

The fruit is a flattened, elongated pod 15-30 cm long. Mature pods hold many seeds and when they become dry they split apart, scattering the seeds.

Seeds:

The seeds are hard, brown, and flattened. They have good germination potential and they contribute to the propagation of the species.

Parts Used for Medicinal Purposes

Various parts of *Bauhinia variegata* have been employed in traditional medicine, including:

- Leaves
- Bark
- Flowers
- Flower buds
- Roots
- Seeds

These plant parts contain a number of bioactive constituents responsible for the pharmacological activities of the species. They are used in herbal preparations and scientific investigations. The bark, leaves and flower buds are most often made use of.

Medicinal Significance

The extensive traditional use of *Bauhinia variegata* has led to many scientific studies to validate its therapeutic potential. Different parts of the plant showed antioxidant, anti-inflammatory, antimicrobial, antidiabetic, hepatoprotective, anticancer, wound healing and nephroprotective activities. These results confirm the importance of this plant as a valuable medicinal resource, and justify the continued study of its phytochemical composition and pharmacological applications.

3. Phytochemical Constituents of *Bauhinia variegata*

Truthfully, procrastination is the one habit I cannot seem to shake no matter how many times I try. I go to bed sure that morning-me will sit down and get it done, and then it is 11 p.m. and I have been glued to my phone for hours. The part that always surprises me is how manageable the work turns out to be — once I finally open the document, a decent opening paragraph usually appears in about ten minutes flat. The real toll is everything leading up to that: the anxiety humming in the background, the false starts, the crushing guilt about wasted hours. I have started to believe procrastination is not really about laziness. It is about wanting to escape the discomfort of getting started.

Being completely honest, procrastination is the one pattern I keep swearing I will fix and keep falling right back into. I make a plan to hit the assignment in the morning, and before I know it the clock reads 11 p.m. and I have done nothing but scroll. The strange thing is the task itself is almost never as bad as the buildup — once I force myself to open the document, the first paragraph usually lands in about ten minutes. Everything that precedes that moment is where the damage lives: the gnawing anxiety, the half-starts, the guilt about hours I threw away. I genuinely think procrastination is not about being lazy. It is about flinching away from the feeling of beginning.

Flavonoids form one of the most important classes of phytochemicals found in *Bauhinia variegata*. They have been known to possess anti-lipid peroxidation properties, antioxidative actions, and act on inflammatory pathways. Flavonoids play a significant role in contributing to the protective effect of the plant for the kidneys by preventing injury to the renal cells from free radicals. The same applies to phenolics, which possess antioxidant actions.

Another category of bioactive compounds discovered in *Bauhinia variegata* is tannins. The astringent, antimicrobial, and anti-inflammatory effects are some of the benefits exhibited by tannins, in addition to protecting biological membranes from oxidative stress. On the other hand, saponins found in *Bauhinia variegata* possess immunomodulating and



cytoprotective effects, whereas alkaloids play a role in antimicrobial and anti-inflammatory effects.

Terpenes and phytosterols found in different parts of the plant have also become subjects of scientific inquiry for their potential use in medicine. They can help in the stabilization of membranes, regulate inflammatory agents, and protect from damage. In addition, glycosides and other secondary metabolites contained within the plant also improve its medicinal value.

Phytoconstituents of *Bauhinia variegata* have been shown to offer scientific evidence for the use of this plant in multiple conditions. In addition, the possible synergy of all bioactive ingredients can further improve the efficiency of this medicinal herb and explain its wide pharmacological effects. The presence of antioxidants and anti-inflammatory ingredients in this plant is especially relevant for nephroprotection since these factors are responsible for kidney damage and dysfunction.

Table 1. Major Phytochemical Constituents Reported in *Bauhinia variegata*

Phytochemical Class	Major Constituents	Reported Biological Activities
Flavonoids	Kaempferol, Quercetin derivatives	Antioxidant, Anti-inflammatory
Phenolic Compounds	Phenolic acids	Free radical scavenging, Cytoprotection
Tannins	Hydrolysable and condensed tannins	Antimicrobial, Astringent, Antioxidant
Saponins	Triterpenoid saponins	Immunomodulatory, Cytoprotective
Alkaloids	Various alkaloidal compounds	Antimicrobial, Anti-inflammatory
Glycosides	Flavonoid glycosides	Therapeutic and protective effects
Terpenoids	Terpene derivatives	Anti-inflammatory, Tissue protection
Steroids	Phytosterols	Membrane stabilization, Anti-inflammatory

4. Traditional and Ethnomedicinal Uses of *Bauhinia variegata*

Traditional use of medicinal plants has been a part of the traditional health care system for many centuries. *Bauhinia variegata* has a special place in Ayurveda, Unani medicine, and several folk remedies because of its wide range of therapeutic uses. Various parts of the plant have been used to treat and control a variety of diseases such as the bark, leaf, flower, flower bud, root and seed. The *Bauhinia variegata* has been widely used in traditional medicine owing to the apparent efficacy and as such has been the subject of scientific research on its pharmacology.

Bauhinia variegata is also known as Kanchanar in Ayurvedic medicine and is used for its astringent, anti-inflammatory, and healing properties. Bark is often incorporated into traditional medicines for treating glandular swellings, skin conditions, ulcers and wounds. It also has some benefits in diseases related to abnormal proliferation and inflammation of tissue. The bark has been used to make a decoction, which is a tea, to ease digestive tract conditions and to improve health.

The flower buds of *Bauhinia variegata* are used as a vegetable throughout several parts of India, it is said that it has medicinal properties. The flower buds have been used traditionally for gastrointestinal disorders such as diarrhea and dysentery. The flowers are also considered as cooling agent and have been used to alleviate symptoms of inflammatory conditions.

The leaves of the plant have been used in folk medicine to treat skin infections, wounds and inflammatory disorders. The extracts of leaves are also used topically for the treatment of wounds in order to decrease the inflammation in the affected area. Leaves are used in some traditional medicine as poultices to treat pain and swelling from injuries.

The root and root bark has been used as a medicine for respiratory problems like cough, bronchitis and asthma. Root preparations are also recommended by traditional healers for some urinary diseases and gynaecological health issues. Occasionally seeds of the plant are used in indigenous medicine for their digestive and antimicrobial properties.



Bauhinia variegata has also been traditionally used as a health-promoting plant because of its multiple health benefits. Its preparations have been utilized to help the liver, digestion, metabolic disorders and to make resistance to infections better. This extensive use indicates that there are several bioactive compounds that could bring several therapeutic effects.

Modern pharmacological studies have confirmed a few of the traditional uses of *Bauhinia variegata*. Many of its ethnomedicinal applications have been supported by research which has shown antioxidant, anti-inflammatory, antimicrobial, antidiabetic, hepatoprotective, and nephroprotective activities of various plant extracts. The close correlation between the plant and the treatment of inflammatory and metabolic diseases is relevant, since these diseases are known to play a key role in the genesis and progression of kidney disease.

The persist use of *Bauhinia variegata* in traditional medicine indicates its therapeutic significance and calls for additional work to determine the active components, the mechanism of action and the clinical effect of this plant. The incorporation of traditional knowledge along with scientific findings can pave the way for the creation of new plant-based medicines for a range of ailments, including renal disorders.

Table 2. Traditional Uses of Different Parts of *Bauhinia variegata*

Plant Part	Traditional Uses
Bark	Wounds, ulcers, inflammation, glandular swellings, digestive disorders
Leaves	Skin diseases, wound healing, pain, swelling, infections
Flowers	Inflammatory conditions, digestive ailments, general health promotion
Flower Buds	Diarrhea, dysentery, nutritional supplement
Roots	Cough, bronchitis, asthma, urinary disorders
Seeds	Digestive disorders, antimicrobial applications

5. Pharmacological Activities of *Bauhinia variegata*

The therapeutic value of *Bauhinia variegata* has attracted considerable scientific attention over the past few decades. Numerous experimental studies have demonstrated that different parts of the plant possess a wide range of pharmacological activities. These biological effects are largely attributed to the presence of diverse phytochemicals, including flavonoids, phenolic compounds, tannins, alkaloids, glycosides, saponins, and terpenoids. The synergistic action of these bioactive constituents contributes to the plant's medicinal potential and supports its traditional use in various disease conditions.

5.1 Antioxidant Activity

Oxidative stress plays a central role in the pathogenesis of many chronic diseases, including kidney disorders, diabetes, cardiovascular diseases, and cancer. Reactive oxygen species generated during cellular metabolism can damage lipids, proteins, and nucleic acids, leading to tissue injury and functional impairment. Various extracts of *Bauhinia variegata* have demonstrated significant antioxidant activity in both in vitro and in vivo studies.

The antioxidant potential of the plant is primarily attributed to its high content of flavonoids and phenolic compounds, which possess strong free radical scavenging properties. These compounds help neutralize reactive oxygen species, inhibit lipid peroxidation, and enhance endogenous antioxidant defense mechanisms. Such antioxidant effects are particularly important in protecting renal tissues from oxidative damage associated with nephrotoxic agents and pathological conditions.

5.2 Anti-inflammatory Activity

Inflammation is a complex biological response that contributes to the development and progression of numerous diseases. Persistent inflammation can result in tissue damage and organ dysfunction, including renal impairment. Studies have reported that extracts of *Bauhinia variegata* exhibit significant anti-inflammatory activity by suppressing inflammatory mediators and reducing inflammatory responses.

The anti-inflammatory effects of the plant are believed to be associated with flavonoids, tannins, and other phenolic compounds that modulate cytokine production and inhibit inflammatory pathways. These properties may help prevent tissue injury and promote recovery in inflammatory disorders.

5.3 Antidiabetic Activity

Diabetes mellitus is a major metabolic disorder characterized by chronic hyperglycemia and associated complications affecting multiple organs, including the kidneys. Experimental studies have demonstrated that *Bauhinia variegata* possesses antidiabetic activity through mechanisms involving improved glucose utilization, enhanced insulin sensitivity, and reduction of oxidative stress.

Plant extracts have been shown to lower blood glucose levels and improve biochemical parameters in diabetic animal models. The presence of flavonoids and other bioactive constituents may contribute to the regulation of carbohydrate metabolism and protection against diabetic complications. These findings support the traditional use of the plant in managing metabolic disorders.

5.4 Antimicrobial Activity

Microbial infections remain a significant public health concern worldwide. Several investigations have demonstrated that extracts of *Bauhinia variegata* exhibit antimicrobial activity against a variety of bacterial and fungal pathogens. The antimicrobial properties are primarily attributed to phenolic compounds, tannins, and flavonoids that interfere with microbial growth and survival.

The ability of the plant to inhibit pathogenic microorganisms may contribute to its traditional use in the treatment of wounds, skin infections, and gastrointestinal disorders. Furthermore, antimicrobial activity can support tissue healing by reducing infection-related complications.

5.5 Hepatoprotective Activity

The liver is highly susceptible to damage caused by toxins, drugs, and oxidative stress. Research studies have reported that *Bauhinia variegata* possesses

hepatoprotective properties capable of protecting liver tissues from chemical-induced injury. Experimental evidence suggests that plant extracts can reduce liver enzyme levels, improve antioxidant status, and preserve hepatic architecture.

The hepatoprotective effects are believed to result from antioxidant and anti-inflammatory mechanisms that prevent cellular damage and enhance tissue repair. These findings indicate the potential utility of the plant in managing liver disorders and promoting hepatic health.

5.6 Wound-Healing Activity

Wound healing involves a coordinated process of inflammation, tissue regeneration, and remodeling. Traditional medicine has long utilized *Bauhinia variegata* for the treatment of wounds and skin injuries. Scientific studies have supported these applications by demonstrating enhanced wound contraction, accelerated epithelialization, and improved tissue regeneration following treatment with plant extracts.

The wound-healing activity may be attributed to antimicrobial, antioxidant, and anti-inflammatory properties that facilitate tissue repair and reduce the risk of infection.

5.7 Anticancer Activity

Recent investigations have explored the anticancer potential of *Bauhinia variegata*. Various extracts have shown cytotoxic effects against selected cancer cell lines and have demonstrated the ability to inhibit cellular proliferation. These effects are believed to be mediated through antioxidant activity, induction of apoptosis, and modulation of signaling pathways involved in cell growth.

Although the available evidence is promising, further studies are required to identify the specific bioactive compounds responsible for these effects and to evaluate their clinical significance.

5.8 Nephroprotective Activity

Among the various pharmacological properties of *Bauhinia variegata*, nephroprotective activity has




gained increasing attention. Kidney damage induced by drugs, toxins, and metabolic disorders is often associated with oxidative stress, inflammation, and cellular injury. Experimental studies suggest that extracts of *Bauhinia variegata* can protect renal tissues by reducing oxidative damage, enhancing antioxidant defenses, and suppressing inflammatory responses.

The antioxidant constituents present in the plant help preserve renal cellular integrity and prevent functional impairment. Furthermore, the anti-inflammatory effects contribute to the reduction of tissue injury and support renal recovery. These findings indicate that *Bauhinia variegata* may serve as a promising natural source for the development of nephroprotective agents.

Table 3. Major Pharmacological Activities of *Bauhinia variegata*

Pharmacological Activity	Proposed Mechanism
Antioxidant	Free radical scavenging, inhibition of lipid peroxidation
Anti-inflammatory	Suppression of inflammatory mediators and cytokines
Antidiabetic	Regulation of glucose metabolism and oxidative stress
Antimicrobial	Inhibition of bacterial and fungal growth
Hepatoprotective	Prevention of oxidative liver damage
Wound-Healing	Enhanced tissue regeneration and repair
Anticancer	Induction of apoptosis and inhibition of cell proliferation
Nephroprotective	Reduction of oxidative stress and renal inflammation



6. Nephroprotective Potential of *Bauhinia variegata*

The kidneys are very special organs that keep water and electrolytes in balance, remove metabolic wastes, control blood pressure and have several endocrine roles. The kidneys have a high blood flow and a very active role in the excretion and filtration of substances and are thus especially susceptible to damage from drugs, toxins, environmental chemicals and metabolic

diseases. Nephrotoxicity is a significant health problem globally and often a factor in the onset and development of acute kidney injury (AKI) and chronic kidney disease (CKD). Therefore, the search for naturally occurring compounds that have renoprotective activity and can help to maintain kidney function has become more and more important.

The medicinal plants are emerging as potential sources of nephroprotective agents due to their rich phytochemical composition and relatively good safety. Of these plants, the plant of the genus *Bauhinia variegata* has received scientific interest, because of the presence of many biologically active compounds and the extensive pharmacological activities. The extracts from various parts of the plant have been shown in several studies to have properties that could be used to prevent or reduce renal damage resulting from oxidative stress, inflammation and toxic insults.

Oxidative stress plays one of the main mechanisms for kidney damage. Excessive generation of ROS may be responsible for lipid peroxidation, protein oxidation, DNA damage and cellular dysfunction during nephrotoxicity, a phenomenon that can overwhelm the body's endogenous antioxidant defenses. These events lead to renal tissue structure and function deterioration. The antioxidant constituents of flavonoids, phenolic compounds, tannins, and so on in *B. variegata* suggest that this plant has nephroprotective effects. These phytochemicals can neutralize free radicals, lower oxidative load and help to prevent damage to the cell components.

Experimental studies have shown that antioxidants found in plants can be very effective for improving renal health by stimulating the activity of the endogenous antioxidant enzymes, including those of superoxide dismutase, catalase and glutathione peroxidase. The nephroprotective effect of *B. variegata* must be further investigated to define the mechanism of its protective action, but the available evidence suggests that the antioxidant activity of *B. variegata* may be a key mechanism in the prevention of nephrotoxic agent-induced kidney injury.

Another important factor which plays a role in renal damage is inflammation. After exposure to toxic agents or pathogenic agents, inflammatory mediators are released which attracts immune cells and further intensifies the pathological changes. Chronic inflammation can speed up the progression of kidney disease and lead to a decrease in the renal tissue's ability to regenerate. The bio-active components of *Bauhinia variegata* have been reported as anti-inflammatory which might modulate inflammatory response. The plant may be able to maintain renal

structure and function by decreasing the production of pro-inflammatory cytokines and inflammatory cell infiltration.

Renal injury is one of the most commonly used experimental models to assess nephroprotective drugs, with gentamicin being the most widely used agent to induce renal injury. Gentamicin is taken up into renal tubular cells where it produces excessive production of ROS, resulting in oxidative stress, mitochondrial dysfunction and cell necrosis. The pathological changes are accompanied by increased blood urea nitrogen and serum creatinine, both markers of decreased kidney function. Natural antioxidants have demonstrated significant nephroprotective properties in the renoprotection from gentamicin damage and the phytochemical profile of the plant *Bauhinia variegata* indicates that it might also have similar properties.

The nephroprotective effects of *Bauhinia variegata* seems to be of a multifactorial mechanism. The plant could also show antioxidant and anti-inflammatory effects and may also have cytoprotective and membrane stabilizing properties. Cellular membrane integrity is crucial for proper renal function and for avoiding leakage of intracellular components. Some phytochemicals present in the plant might be able to stabilize biological membranes and decrease their liability to toxic damage.

Another crucial part of nephroprotection is the prevention of the programmed cell death or apoptosis, which is often seen in kidney diseases. High rates of apoptosis can lead to a loss of functioning renal cells and impaired renal function. Anti-apoptotic properties of *B. variegata* antioxidants are believed to be mediated by their ability to decrease the oxidative stress and preserve mitochondrial function. These effects may play a role in the better survival of renal cells in pathological conditions.

The renoprotective effect of *Bauhinia variegata* may also be mediated through its antidiabetic effects. Diabetes mellitus is one of the major causes of chronic kidney disease in the world. The continuous exposure of kidneys to hyperglycemia results in the onset of microvascular damage, which is the origin of diabetic nephropathy. The chronically high blood glucose causes oxidative stress, inflammation and



microvascular damage in kidneys, ultimately leading to diabetic nephropathy. The benefits of *Bauhinia variegata* could be due to its ability to modulate the glucose metabolism and to minimise oxidative stress, preventing renal complications associated with diabetes. In research work carried out with medicinal plants, it has been shown that agents which protect the kidneys in the plant are also effective in reducing the degrees of tubular degeneration, glomerular damage, inflammatory infiltration and interstitial edema. Based on their pharmacological profile, the same protective effects can be expected after the use of extracts from *Bauhinia variegata*. The changes in renal histopathology, along with biochemical markers, would be good indications of nephroprotective activity.

Although promising results have been reported, the majority of the current studies on the nephroprotective property of *Bauhinia variegata* is from the preclinical level. Further research is needed to determine the effect of these phytoconstituents in protecting the kidneys and for their long-term safety, as well as for specific dosage regimens. In addition, clinical trials are required to verify the efficacy in humans and to enable its inclusion in evidence-based treatment. From the available scientific literature, it can be concluded that *Bauhinia variegata* has a high nephroprotective potential in general. It is rich in antioxidant and anti-inflammatory phytochemical content, has traditional medicinal applications and a variety of pharmacological activities, all of which make it a potential natural renal protection agent. On-going studies could potentially help lead to new plant-based treatments for kidney diseases, prevention, and the alleviation of kidney toxicity.

7. Renal Protection by *Bauhinia variegata* - Mechanisms.

The mechanism of renal protection is an important factor that needs to be understood for the evaluation of the therapeutic potential of medicinal plants. Kidney injury is a multistage process that includes oxidative stress, inflammation, apoptosis, mitochondrial dysfunction and cellular damage. The nephroprotective activities of *Bauhinia variegata* may be attributed to its multi-target action against renal injury pathways due to the combination of its various

phytochemicals. While additional mechanistic investigations are needed, there is some science that helps to explain how the plant might work to protect.

7.1 Antioxidant Mechanism

Oxidative stress is one of the main causes for the onset and development of kidney damage. A high amount of production of reactive oxygen species (ROS) may overwhelm the antioxidant defense mechanisms of the body and cause lipid peroxidation, protein oxidation, DNA damage and cell dysfunction. Gentamicin is a nephrotoxic drug that is known to cause oxidative stress that leads to structural and functional damage in kidney tissue. The *Bauhinia variegata* is rich in flavonoids, phenolics, tannins etc. which have the ability to neutralise free radicals and diminish oxidative load. These phytochemicals neutralize the ROS to prevent them from damaging the cellular components. They can also boost the activity of antioxidant enzymes like superoxide dismutase, catalase and glutathione peroxidase, which help to boost the body's own antioxidant capacity. By these actions, the plant helps to maintain the redox balance and to prevent oxidative damage in the renal cells.

7.2 Anti-inflammatory Mechanism

Kidney disease has a lot to do with inflammation. Nephrotoxic agents can often induce release of inflammatory mediators that recruit immune cells to damaged tissues. This inflammatory reaction can lead to further deterioration of the tissues and reduced kidney function.

In different experimental tests bioactive compounds present in *Bauhinia variegata* have shown anti-inflammatory properties. These compounds could inhibit production of pro-inflammatory cytokines and decrease infiltration of inflammatory cells into renal tissues. The plant may help to reduce inflammation, which can help prevent any further damage to the tissue and aid in healing of the damaged kidneys.

7.3 Inhibition of Lipid Peroxidation

Lipid peroxidation is defined as the oxidation of membrane lipids, and is believed to be one of the more important effects of oxidative stress. The process will



disrupt the integrity of the membrane, change the permeability of the cell, and finally cause the cell death. Renal tubular cells are especially vulnerable to lipid peroxidation due to their high metabolic activity and consumption of oxygen. The antioxidant properties of *Bauhinia variegata* may help to suppress the lipid peroxidation process through free radical scavenging and blocking the chain reaction in lipid peroxidation. Maintaining the integrity of the membrane is important for normal cell function and safeguarding kidney tissues against toxic insults.

7.4 Cytoprotective and Membrane-Stabilizing Effects

The normal kidney function depends on the integrity of the cell membrane. The damage can lead to leakage of intracellular contents, disturbances of intracellular ion balance and loss of cell viability. The membrane stabilizing effect of *Bauhinia variegata* may be due to the protective effect of phytochemicals on the phospholipid structure from oxidative damage. This protective effect will increase the resistance of the cells to toxic agents and help to maintain the renal architecture. In specific contexts, such as chronic oxidative or inflammatory insult to the renal cell, it is crucial that cytoprotection takes place.

7.5 Anti-apoptotic Activity

One of the important mechanisms involved in nephrotoxicity and progression of kidney disease is Apoptosis (programmed cell death). A high rate of apoptosis will decrease the number of working kidney cells and affect kidney function. Apoptosis is mediated by several nephrotoxic agents, which cause oxidative stress, mitochondrial dysfunction and activation of the intracellular signaling pathways.

Oxidative stress and mitochondrial stability could help to mediate the antioxidant activity of *Bauhinia variegata* to inhibit the apoptosis. The inhibition of apoptosis cell death protects the integrity of renal tissue and helps achieve normal kidney function.

7.6 The Cellular Signalling Pathways are controlled by modulation.

A variety of pathological mechanisms related to kidney injury are controlled by intricate cellular

signaling pathways. Signaling cascades as a result of oxidative stress and inflammation have effects on gene expression, cytokine production, and cell survival. Medicinal plants contain phytochemicals that can affect these pathways and their activity. While further studies are still underway, the presence of bioactive substances in *Bauhinia variegata* leaves could potentially affect pathways related to oxidative stress and inflammatory responses. This modulation might help to minimize tissue damage and increase cell adaptability to pathological situations.

7.7 Drug Induced Nephrotoxicity Prevention

Nephrotoxicity is an important concern in the clinic regarding drug-induced nephrotoxicity. Significant renal damage has been induced by medications like aminoglycoside antibiotics, chemotherapeutic agents and nonsteroidal anti-inflammatory drugs in an oxidative and inflammatory manner.

The pharmacological activity of *Bauhinia variegata* indicates that it could potentially be used to prevent kidney damage caused by drugs through the mechanisms of free radical scavenging, anti-inflammatory, membrane stabilizing and renal protecting effects. These protective effects may be useful to reduce the structural damage and recovery after nephrotoxic substance exposure.

8. Experimental evidence of the Nephroprotective Activity of *Bauhinia variegata*

Traditional validation of medicinal plants through scientific studies can facilitate the determination of their therapeutic relevance and possible clinical use. Recently, there has been a growing focus on assessing plant-derived ingredients as protective agents against renal injury. The studies focused on the nephroprotective effects of *Bauhinia variegata* are still very limited compared to those of other medicinal plants, but the available experimental data and its well established pharmacological effects suggest that it could be a promising plant for renal protection.

Preliminary experimental studies on different extracts of *Bauhinia variegata* have shown both antioxidant and anti-inflammatory activity, which are basic mechanisms underlying nephroprotective properties. Treatment with plant extracts demonstrated the ability



to decrease oxidative damage, increase antioxidant enzyme activity and maintain tissue architecture in experimental models of organ toxicity. The result indicated that *Bauhinia variegata* might be beneficial in safeguarding the renal tissues against toxic agents and pathological conditions. Animal models continue to be the most commonly used method to test nephroprotective agents. In such studies, nephrotoxicity is usually induced by chemicals or drugs that are known to be toxic to the kidneys such as gentamicin, cisplatin, paracetamol and heavy metals. Plant extracts are then evaluated for their preventive/reversive potential against biochemical, hematological and histopathological changes in kidney damage.

A key parameter of nephroprotective activity is the normalisation of renal function parameters. High blood urea nitrogen and blood creatinine are hallmarks of poor renal function. The antioxidant activity of the extract of these medicinal plants has been consistently shown in experimental studies to significantly decrease these markers and hence improve glomerular filtration and renal function. *Bauhinia variegata* has a similar phytochemical makeup and has significant potential to impart similar benefits. Oxidative stress markers are also often assessed in nephrotoxicity studies as well as biochemical parameters. Renal injury is characterized by elevated lipid peroxidation and the fall in endogenous antioxidants. Extracts of *Bauhinia variegata* have been shown to exert antioxidant properties and to limit the oxidative damage in several experimental systems. The results obtained support indirectly its nephroprotective efficacy and indicate the central role of antioxidant mechanisms as a mechanism of renal protection.

Another important aspect of nephroprotective studies is histopathological examination. Histological examination of the kidney can be done on a microscopic level to determine structural changes including tubular degeneration, glomerular damage, inflammatory infiltration, necrosis and interstitial edema. This pathological changes shall be reduced by protective agents and normal renal architecture shall be preserved. The antioxidant and anti-inflammatory properties of *Bauhinia variegata* suggest its ability to reduce tissue damage and facilitate tissue healing after

toxic exposure. In several studies conducted on medicinal plants that are rich in flavonoids and phenolics, the authors observed marked improvement in renal histopathology, which included restoration of tubular integrity and reduction in inflammatory lesions. Due to the presence of comparable phytochemicals, it would be reasonable to suggest that *Bauhinia variegata* could have similar protective effects. Nevertheless, additional studies need to be conducted with specific nephrotoxicity models to confirm its effectiveness in those models.

The nephroprotective activity of *Bauhinia variegata* is also reinforced with the reported antidiabetic and hepatoprotective activities. Diabetes is often accompanied by elevated oxidative stress and systemic inflammation, which are known factors of renal injury, and liver dysfunction is also linked with these two factors. Renal dysfunction is often associated with elevated systemic inflammation and increased oxidative stress, both of which are associated with diabetes, and liver dysfunction is also associated with these two factors. The plant's ability to balance metabolism and lower oxidative stress can indirectly promote kidney health and ward off kidney-related problems.

In recent years, the importance of phytopharmacology in identifying the individual bioactive compounds with therapeutic effect has been highlighted. Biological activities of some constituents which have been isolated from *Bauhinia variegata* have been shown to be relevant to nephroprotection. Further research on these compounds could aid in the development of standardisation of formulations, as well as increase knowledge of how these compounds act.

While there are some positive results, there are certain drawbacks. There is limited evidence available from preclinical studies and extensive clinical work is lacking. The outcome of studies can also vary depending on how the plants are extracted, dosages, experimental design, etc. Thus, further studies are needed to provide standardized procedures and to assess the safety and efficacy of *Bauhinia variegata* in humans.



9. The Need for New Research and the Safety and Toxicological Considerations of Existing Studies

Interest in medicinal plants as alternative therapeutic agents has grown, and there is a need to comprehensively evaluate their safety and toxicological profile. *Bauhinia variegata* is a century-old traditional medicine plant and is safe for general use, but scientific evaluation of its toxicity is still necessary in order to achieve its widespread therapeutic use. Because of the chronicity of some health issues like kidney diseases, it is very important to assess the safety of herbals to be used for extended periods. The potential for safety of *Bauhinia variegata* has been suggested as relatively favorable based on traditional use in various traditional and modern systems of medicine. Leaves, bark, flowers and flower buds of the plant have been used medicinally or as food, and there has been no report of any serious adverse effects. The flower buds are generally accepted and appear to be non-toxic in moderate quantities and are used as a vegetable in several parts of India.

In experimental toxicological studies, plant extracts of *Bauhinia variegata* have been shown to be generally safe at therapeutic doses. Laboratory animal tests of acute toxicity have indicated low death rates and little behavioral abnormalities after intake of plant extracts. The results indicate a fairly high margin of safety but toxicity may depend on the method of extraction, plant part employed, dose, period of exposure, and mode of administration. Although these products seem safe, the risks of consuming too much of a herbal product could still exist. In some cases, high doses of phytochemicals may have unwanted side effects, especially if taken long-term. Some bioactive compounds can interact with metabolic enzymes, affect drug absorption, or affect physiological functions. Thus, caution should be exercised when standardizing doses so that they are effective and safe.

Herb-drug interactions are another important factor to take into consideration. Patients suffering from kidney disease are prescribed many drugs such as anti-hypertensive, anti-diabetic, anti-bacterial and anti-diuretic drugs. Theoretically, phytochemicals found in *Bauhinia variegata* can affect the pharmacokinetics or pharmacological actions of these

drugs. As of yet, documented interactions are sparse but healthcare professionals need to be cautious of concurrent use of herbal with conventional therapies.

Another significant hurdle in herbal medicine research is its quality control. The phytochemical content of medicinal plants can be influenced by geographical variation, climatic factors, period of harvesting, soil texture, and extraction methods. Therefore, therapeutic activity and safety of the various preparations of *Bauhinia variegata* can vary. Standardisation of plant material and extraction procedures is thus critical to ensure reproducibility of results. One major drawback of existing studies is that only a few studies focused on the nephroprotective effects of *Bauhinia variegata*. Although many studies have shown its antioxidant, anti-inflammatory, antidiabetic, and hepatoprotective effects, there is only limited data existing on renal protection. The information available is mostly from the indirect observations and based on the biological activities of the phytoconstituents found in it, which are derived from preclinical investigations.

In addition, many experimental studies have small sample sizes and brief treatment periods, and may have limited applicability. Comparing studies may be difficult to do because of inter-species differences, different nephrotoxicity models, different extraction methods, and different dosage regimens. The methodological variations emphasize the need for standardized experimental procedures. Another important research gap is the lack of detailed mechanistic studies. Antioxidant and anti-inflammatory mechanisms are thought to be major mechanisms involved in nephroprotection, but the exact molecular pathways are not fully understood. The elucidation of the exact mechanism of action of the identified active compounds in the plant and the understanding of its therapeutic potential would be greatly improved.

Perhaps most important, there are no well-designed clinical trials. The medicinal effects of *Bauhinia variegata* have been mostly studied in the laboratory and on animals. There are few human trials to assess the effectiveness, safety, pharmacokinetics and long-term effects of its use. Clinical investigations are



crucial to bring the promising preclinical results into the clinical realm and into therapeutic applications.

Chronic toxicity, reproductive toxicity, genotoxicity and detailed pharmacokinetic studies should be conducted in future toxicological assessment. These would be helpful for safety information on long-term administration of these as well as evidence based dosage recommendations. Based on current evidence, the safety of *Bauhinia variegata* can be considered relatively good and there is little evidence of adverse effects. However, additional toxicological research and clinical trials are needed to establish its long-term safety and efficacy. Research gaps related to the development of *Bauhinia variegata* as a scientifically proven nephroprotective agent will be essential.

10. CONCLUSION

Bauhinia variegata is a medicinal plant that has been used for hundreds of years in traditional medicine. The plant contains various bioactive phytochemicals, such as flavonoids, phenolic compounds, tannin, saponin, glycosides, and terpenoids, supporting its ability to have many pharmacological activities. The scientific studies have shown that *Bauhinia variegata* possesses considerable antioxidant, anti-inflammatory, antidiabetic, antimicrobial, hepatoprotective, wound healing, and nephroprotective activity. Many of its traditional therapeutic uses are scientifically supported by these biological activities, and suggest its potential as a source of natural medicine agents.

From the available evidence, the nephroprotective properties of *B. variegata* seem to be related to its anti-oxidative, anti-inflammatory, anti-cytotoxic and maintaining renal structure and function properties. The results are encouraging, but the majority of research carried out to date consisted of preclinical studies, and additional research is needed to isolate bioactive ingredients, uncover molecular mechanisms, standardise herbal preparations, and confirm the clinical effectiveness. The lack of adequate experimental and clinical studies in the future will be crucial for assessing its therapeutic potential and for the safe, effective and evidence-based development of nephroprotective formulation. In general, *B. variegata* is a significant natural resource for protecting the kidneys and seems to be

useful in the prevention and management of kidney diseases in the future.

REFERENCES

1. Gudavalli, D., Pandey, K., Ede, V. G., Sable, D., Ghagare, A. S., & Kate, A. S. (2024). Phytochemistry and pharmacological activities of five species of *Bauhinia* genus: A review. *Fitoterapia*, 174, 105830.
2. Sharma, A., Chandel, V., & Roy, S. (2025). Exploring the bioactive properties and diverse uses of *Bauhinia variegata*: A comprehensive review. *Discover Plants*, 2, 86.
3. Sahu, G., & Gupta, P. K. (2012). A review on *Bauhinia variegata* Linn. *International Research Journal of Pharmacy*, 3(1), 61–64.
4. Negi, A., & Sharma, N. (2012). Spectrum of pharmacological activities from *Bauhinia variegata*: A review.
5. Kamal, Y., Khan, T., Haq, I., Zahra, S. S., Asim, M. H., Shahzadi, I., Mannan, A., & Fatima, N. (2022). Phytochemical and biological attributes of *Bauhinia variegata* L. *Brazilian Journal of Biology*, 82, e257990.
6. Deswal, G., & Arora, K. (2015). Ethnobotany and phytopharmacology of *Bauhinia variegata*. *International Journal of Pharmaceutics and Drug Analysis*, 3(9), 261–263.
7. Prabhu, S., Vijayakumar, S., Ramasubbu, R., Praseetha, P. K., Karthikeyan, K., Thiyagarajan, G., & Sureshkumar, J. (2021). Traditional uses, phytochemistry and pharmacology of *Bauhinia racemosa* Lam.: A comprehensive review. *Future Journal of Pharmaceutical Sciences*, 7, 101.
8. Kirtikar, K. R., & Basu, B. D. (2006). Indian medicinal plants (Vol. 2). International Book Distributors.
9. Chopra, R. N., Nayar, S. L., & Chopra, I. C. (2006). Glossary of Indian medicinal plants. National Institute of Science Communication.
10. Nadkarni, K. M. (2009). Indian materia medica (Vol. 1). Popular Prakashan.
11. Harborne, J. B. (1998). *Phytochemical methods: A guide to modern techniques of plant analysis* (3rd ed.). Chapman & Hall.
12. Trease, G. E., & Evans, W. C. (2009). *Pharmacognosy* (16th ed.). Saunders Elsevier.



13. Kokate, C. K., Purohit, A. P., & Gokhale, S. B. (2010). *Pharmacognosy*. Nirali Prakashan.
14. Evans, W. C. (2009). *Trease and Evans pharmacognosy* (16th ed.). Saunders.
15. Sofowora, A. (2008). *Medicinal plants and traditional medicine in Africa* (3rd ed.). Spectrum Books.
16. Parekh, J., & Chanda, S. (2007). In vitro antimicrobial activity and phytochemical analysis of some Indian medicinal plants. *Turkish Journal of Biology*, 31(1), 53–58.
17. Halliwell, B., & Gutteridge, J. M. C. (2015). *Free radicals in biology and medicine* (5th ed.). Oxford University Press.
18. Rang, H. P., Dale, M. M., Ritter, J. M., & Flower, R. J. (2016). *Rang and Dale's pharmacology* (8th ed.). Elsevier.
19. Brunton, L. L., Hilal-Dandan, R., & Knollmann, B. C. (2018). *Goodman & Gilman's the pharmacological basis of therapeutics* (13th ed.). McGraw-Hill.
20. Katzung, B. G. (2021). *Basic and clinical pharmacology* (15th ed.). McGraw-Hill.
21. Mali, R. G., Mahajan, S. G., & Mehta, A. A. (2007). Rakta Kanchan (*Bauhinia variegata*): Chemistry, traditional and medicinal uses—A review. *Pharmacognosy Reviews*, 1(2), 314–319.
22. Deswal, G., & Arora, K. (2015). Ethnobotany and phytopharmacology of *Bauhinia variegata*. *International Journal of Pharmaceutics and Drug Analysis*, 3(9), 261–263.
23. Negi, A., & Sharma, N. (2012). Spectrum of pharmacological activities from *Bauhinia variegata*: A review. *Journal of Pharmacy Research*, 5(2), 1000–1005.
24. Al-Snafi, A. E. (2014). The pharmacological importance of *Bauhinia variegata*: A review. *International Journal of Pharma Sciences and Research*, 5(12), 160–164.
25. Evans, W. C. (2009). *Trease and Evans pharmacognosy* (16th ed.). Saunders Elsevier.
26. Kokate, C. K., Purohit, A. P., & Gokhale, S. B. (2010). *Pharmacognosy*. Nirali Prakashan.
27. Harborne, J. B. (1998). *Phytochemical methods: A guide to modern techniques of plant analysis* (3rd ed.). Chapman & Hall.
28. Sofowora, A. (2008). *Medicinal plants and traditional medicine in Africa* (3rd ed.). Spectrum Books.
29. Khandelwal, K. R. (2008). *Practical pharmacognosy* (19th ed.). Nirali Prakashan.
30. Wallis, T. E. (2005). *Textbook of pharmacognosy* (5th ed.). CBS Publishers.
31. Farnsworth, N. R. (1994). *Ethnopharmacology and drug development*. Ciba Foundation Symposium, 185, 42–59.
32. Fabricant, D. S., & Farnsworth, N. R. (2001). The value of plants used in traditional medicine for drug discovery. *Environmental Health Perspectives*, 109(Suppl. 1), 69–75.
33. Cowan, M. M. (1999). Plant products as antimicrobial agents. *Clinical Microbiology Reviews*, 12(4), 564–582.
34. Halliwell, B. (2007). Oxidative stress and cancer: Have we moved forward? *Biochemical Journal*, 401(1), 1–11.
35. Halliwell, B., & Gutteridge, J. M. C. (2015). *Free radicals in biology and medicine* (5th ed.). Oxford University Press.
36. Valko, M., Leibfritz, D., Moncol, J., Cronin, M. T. D., Mazur, M., & Telser, J. (2007). Free radicals and antioxidants in normal physiological functions and human disease. *International Journal of Biochemistry & Cell Biology*, 39(1), 44–84.
37. Sies, H. (2015). Oxidative stress: A concept in redox biology and medicine. *Redox Biology*, 4, 180–183.
38. Forman, H. J., Zhang, H., & Rinna, A. (2009). Glutathione: Overview of its protective roles. *Molecular Aspects of Medicine*, 30(1–2), 1–12.
39. Li, S., Tan, H. Y., Wang, N., Zhang, Z. J., Lao, L., Wong, C. W., & Feng, Y. (2015). The role of oxidative stress and antioxidants in liver diseases. *International Journal of Molecular Sciences*, 16(11), 26087–26124.
40. Kumar, V., Abbas, A. K., & Aster, J. C. (2020). *Robbins and Cotran pathologic basis of disease* (10th ed.). Elsevier.
41. Brunton, L. L., Hilal-Dandan, R., & Knollmann, B. C. (2018). *Goodman & Gilman's the pharmacological basis of therapeutics* (13th ed.). McGraw-Hill.
42. Katzung, B. G. (2021). *Basic and clinical pharmacology* (15th ed.). McGraw-Hill.
43. Rang, H. P., Ritter, J. M., Flower, R. J., & Henderson, G. (2019). *Rang and Dale's pharmacology* (9th ed.). Elsevier.



44. Dipiro, J. T., Talbert, R. L., Yee, G. C., Matzke, G. R., Wells, B. G., & Posey, L. M. (2020). *Pharmacotherapy: A pathophysiologic approach* (11th ed.). McGraw-Hill.
45. Tietz, N. W. (2018). *Fundamentals of clinical chemistry and molecular diagnostics* (8th ed.). Elsevier.
46. Gilman, A. G. (2018). *Goodman & Gilman's manual of pharmacology and therapeutics* (2nd ed.). McGraw-Hill.
47. Brenner, B. M., & Rector, F. C. (2016). *The kidney* (10th ed.). Elsevier.
48. Johnson, R. J., Feehally, J., & Floege, J. (2019). *Comprehensive clinical nephrology* (6th ed.). Elsevier.
49. Chatterjee, A., & Pakrashi, S. C. (2003). *The treatise on Indian medicinal plants* (Vol. 5). National Institute of Science Communication.
50. Gupta, A. K. (2010). *Quality standards of Indian medicinal plants* (Vol. 8). Indian Council of Medical Research.
51. Wagner, H., & Bladt, S. (2009). *Plant drug analysis: A thin layer chromatography atlas* (2nd ed.). Springer.
52. Heinrich, M., Barnes, J., Gibbons, S., & Williamson, E. M. (2012). *Fundamentals of pharmacognosy and phytotherapy* (2nd ed.). Elsevier.
53. Bruneton, J. (1999). *Pharmacognosy, phytochemistry, medicinal plants* (2nd ed.). Lavoisier Publishing.
54. Dewick, P. M. (2009). *Medicinal natural products: A biosynthetic approach* (3rd ed.). Wiley.
55. Evans, W. C. (2002). *Pharmacognosy and phytochemistry*. Saunders.
56. Duke, J. A. (2002). *Handbook of medicinal herbs* (2nd ed.). CRC Press.
57. Ekor, M. (2014). The growing use of herbal medicines: Issues relating to adverse reactions and challenges in monitoring safety. *Frontiers in Pharmacology*, 4, 177.
58. Rates, S. M. K. (2001). Plants as source of drugs. *Toxicon*, 39(5), 603–613.
59. Newman, D. J., & Cragg, G. M. (2020). Natural products as sources of new drugs over nearly four decades. *Journal of Natural Products*, 83(3), 770–803.
60. Cragg, G. M., & Newman, D. J. (2013). Natural products: A continuing source of novel drug leads. *Biochimica et Biophysica Acta*, 1830(6), 3670–3695.

HOW TO CITE: Sarika More, Ansari MD Huzaifa Siraj Ahmad, Taarique Deshmukh, Rashid Akhtar Nehal Ahmad, Shripad Bairagi, Nephroprotective Potential of *Bauhinia variegata*: Phytochemistry, Pharmacological Activities, and Therapeutic Perspectives, *J. Pharm. Sci.*, 2026, 2 (6), 415-429. <https://doi.org/10.5281/zenodo.20766980>

