

The Rise of *Centella Asiatica*: A Botanical Powerhouse Reshaping the Cosmetic and Wellness Industries

Author: Grishma Patil

Affiliation: Independent Researcher

Contact: grishmapatil14@gmail.com

ORCID: 0009-0000-3885-2290

Date: February 01, 2026

Abstract

Centella asiatica (CA), commonly known as Gotu Kola or Tiger Grass, has emerged as a key botanical ingredient in modern cosmetic and wellness formulations, supported by both traditional medicinal use and a growing body of scientific evidence. Extensive pharmacological research has identified CA's primary triterpenoid constituents, asiaticoside, madecassoside, asiatic acid, and madecassic acid, as central to its biological activity (Brinkhaus et al., 2000; Gohil et al., 2010). These compounds have been shown to influence wound healing, collagen synthesis, inflammatory modulation, and antioxidant defense, making CA particularly relevant for skin-repair, barrier-support, and soothing applications (Shukla et al., 1999; Somboonwong et al., 2012).

Beyond dermatological use, emerging studies suggest that CA may exert neuroprotective, anxiolytic, and cognitive-supportive effects, expanding its potential utility within the wellness and nutraceutical space (Orhan, 2012; Hashim et al., 2011; Puttarak et al., 2017). Mechanistic insights into fibroblast activation, extracellular matrix remodeling, and oxidative stress regulation further support its inclusion in both cosmetic and drug-adjacent formulations.

As consumer demand increasingly favors plant-derived ingredients with demonstrated efficacy and mechanistic plausibility, *Centella asiatica* has become a versatile active across serums, creams, wound-care products, and functional wellness formulations. This review synthesizes current evidence on CA's bioactive profile and therapeutic relevance, highlighting its growing role at the intersection of cosmetic science and wellness innovation.

1. Introduction

Centella asiatica (L.) Urban is a perennial creeping herb belonging to the Apiaceae family and is widely distributed across tropical and subtropical regions of Asia and Africa (Brinkhaus et al., 2000; James & Dubery, 2009). The plant has been used for centuries in multiple traditional medical systems, including Ayurveda, Traditional Chinese Medicine, and various African ethnomedicinal practices, where it has been valued for its effects on wound healing, inflammation, circulation, and cognitive support (Brinkhaus et al., 2000; Gohil et al., 2010). These long-standing applications reflect the plant's broad therapeutic relevance well before its integration into modern cosmetic and pharmaceutical research.

Renewed scientific interest in *Centella asiatica* is largely driven by its rich content of biologically active triterpenoid compounds, including asiaticoside, madecassoside, asiatic acid, and madecassic acid, which are responsible for many of its documented pharmacological effects (Gohil et al., 2010; Hashim et al., 2011). Extensive experimental studies have demonstrated anti-inflammatory, antioxidant, wound-healing, angiogenic, and neuroprotective properties associated with these constituents, supporting both topical and systemic applications (Shukla et al., 1999; Somboonwong et al., 2012; Bylka et al., 2014).

As understanding of these mechanisms has expanded, *Centella asiatica* has become increasingly incorporated into dermatological, cosmetic, and wellness formulations. Its demonstrated effects on fibroblast activity, collagen synthesis, extracellular matrix remodeling, and oxidative stress regulation have positioned it as a valuable ingredient for skin-repair, barrier-support, soothing, and anti-aging applications (Shukla et al., 1999; Brinkhaus et al., 2000). In parallel, growing consumer demand for plant-derived ingredients with scientific substantiation has accelerated its adoption across cosmeceutical, wound-care, and nutraceutical products. As a result, *Centella asiatica* now occupies a unique position at the intersection of traditional medicine, modern cosmetic science, and wellness innovation, with continued relevance in both research and commercial formulation development.

2. Phytochemical Composition and Bioactive Mechanisms

Centella asiatica is a chemically complex botanical rich in triterpenoid saponins, phenolic compounds, flavonoids, tannins, and volatile constituents that collectively contribute to its pharmacological profile (James & Dubery, 2009; Bylka et al., 2014). Among these, pentacyclic triterpenoids represent the primary bioactive class responsible for the plant's therapeutic relevance in dermatological, cosmetic, and wellness applications. The relative abundance of these compounds varies with plant maturity, geographic origin, and extraction methodology, underscoring the importance of standardized processing for reproducible cosmetic and pharmaceutical use (Bylka et al., 2014; Hashim et al., 2011).

2.1 Major Triterpenoids and Their Biological Actions

The principal pharmacologically active constituents of *Centella asiatica* include asiaticoside, madecassoside, asiatic acid, and madecassic acid. These compounds have been extensively investigated for their wound-healing, anti-inflammatory, antioxidant, and collagen-modulating properties (Brinkhaus et al., 2000; Gohil et al., 2010).

Asiaticoside has been shown to stimulate fibroblast proliferation, enhance collagen type I synthesis, and accelerate re-epithelialization, making it one of the most studied actives in burn, incision, and chronic wound models (Shukla et al., 1999; Somboonwong et al., 2012). Madecassoside exhibits complementary anti-inflammatory activity, contributing to reduced inflammatory mediator expression and improved tissue remodeling outcomes (Gohil et al., 2010; Bylka et al., 2014).

The aglycone forms, asiatic acid and madecassic acid, contribute significantly to antioxidant defense and cellular protection. These molecules are associated with modulation of oxidative stress pathways and protection against reactive oxygen species, supporting tissue resilience and barrier recovery in stressed or inflamed skin (Hashim et al., 2011).

Together, these triterpenoids support a coordinated wound-repair process involving fibroblast activation, extracellular matrix remodeling, and angiogenic support, forming the mechanistic foundation for CA's widespread dermatological use (Brinkhaus et al., 2000; Somboonwong et al., 2012).

2.2 Expanded Phytochemical Profile

Beyond triterpenoids, *Centella asiatica* contains flavonoids such as quercetin and kaempferol, which contribute additional antioxidant and anti-inflammatory activity through radical scavenging and metal-chelation mechanisms (James & Dubery, 2009; Bylka et al., 2014). Phytosterols and other lipid-soluble constituents further support barrier integrity and lipid regulation, reinforcing CA's role in skin-repair and moisturizing formulations (Bylka et al., 2014).

Minor constituents, including polyacetylenes, amino acids, and volatile compounds, have been associated with antimicrobial, soothing, and hydration-supportive effects, enhancing the plant's multifunctional therapeutic profile (James & Dubery, 2009). The synergistic interaction among these phytochemicals broadens CA's biological relevance across dermatological, neurological, and vascular applications (Gohil et al., 2010).

2.3 Mechanistic Insights Into Skin Repair and Anti-Inflammatory Activity

The wound-healing and anti-aging effects of *Centella asiatica* are mediated through multiple overlapping biological pathways:

2.3.1 Collagen Synthesis and Extracellular Matrix Remodeling

CA triterpenoids have been shown to enhance fibroblast activity and promote collagen synthesis, contributing to improved dermal structure, tensile strength, and scar quality (Shukla et al., 1999; Brinkhaus et al., 2000). Regulation of extracellular matrix turnover supports both wound resolution and anti-aging effects in cosmetic applications (Gohil et al., 2010).

2.3.2 Angiogenesis and Microcirculation Support

Experimental models demonstrate that CA extracts enhance granulation tissue formation and vascularization, improving oxygen and nutrient delivery to healing tissue (Somboonwong et al., 2012). These effects are particularly relevant for compromised or chronic wounds and contribute to improved tissue regeneration outcomes.

2.3.3 Anti-Inflammatory and Barrier-Recovery Pathways

CA reduces inflammatory signaling and oxidative damage, promoting faster re-epithelialization and barrier restoration. These properties support its widespread use in soothing, barrier-repair, and post-procedure skincare formulations (Brinkhaus et al., 2000; Bylka et al., 2014).

2.3.4 Antioxidant Defense and Oxidative Stress Regulation

CA enhances endogenous antioxidant defenses and mitigates oxidative stress, contributing to protection against photoaging and inflammation-associated tissue damage (Hashim et al., 2011).

2.4 Neuroprotective and Cognitive Mechanisms

In addition to its dermatological relevance, *Centella asiatica* has demonstrated neuroprotective activity in preclinical models. Triterpenoids and flavonoids have been

associated with improved mitochondrial function, synaptic support, and cognitive performance, aligning with the plant's traditional use as a cognitive tonic and its emerging role in wellness and nootropic research (Orhan, 2012; Puttarak et al., 2017).

2.5 Vascular and Anti-Cellulite Applications

CA has been shown to support microcirculation and vascular integrity by improving venous tone and capillary resistance. Clinical studies have demonstrated benefits in conditions related to venous insufficiency, supporting its use in anti-cellulite formulations and oral venotonic supplements (Cesarone et al., 2001; De Sanctis et al., 2001).

2.6 Extraction Technologies and Standardization

The biological activity of CA extracts is highly dependent on extraction method. Ethanolic and hydroalcoholic extractions effectively preserve triterpenoid saponins, while advanced techniques such as ultrasonic-assisted extraction improve yield through enhanced cell-wall disruption (Hashim et al., 2011; Khursheed et al., 2024). Standardized extracts, including titrated extract of *Centella asiatica* (TECA), ensure consistent ratios of key triterpenoids and are widely used in clinical, cosmetic, and pharmaceutical formulations (Brinkhaus et al., 2000).

3. Dermatological and Cosmetic Applications

Centella asiatica has become an important ingredient in dermatological and cosmetic formulations due to its ability to address inflammation, compromised barrier function, and visible signs of skin aging, three central concerns in modern skincare (Bylka et al., 2014). The plant's triterpenoid constituents support epithelial repair, stimulate collagen synthesis, and promote dermal remodeling, making CA valuable for both clinical dermatology and consumer cosmetic products (Shukla et al., 1999; Brinkhaus et al., 2000).

Clinical and experimental findings support the role of *Centella asiatica* in improving skin firmness, elasticity, hydration, and overall texture. Human clinical evidence indicates that CA based skincare formulations support barrier function and reduce irritation in sensitive skin populations (Su et al., 2025), while pilot clinical studies have demonstrated improvements in hydration, elasticity, and skin texture following application of CA extracellular vesicle based formulations (Chang et al., 2025). In addition, key triterpenoid constituents such as asiaticoside and madecassoside exhibit anti-inflammatory activity that contributes to reduced redness, irritation, and inflammatory signaling, supporting CA's suitability for sensitive, acne prone, and barrier impaired skin types (Somboonwong et al., 2012; Bylka et al., 2014).

CA's calming and reparative properties have also led to its widespread use in post-procedure skincare, including formulations intended for use after laser treatments, chemical exfoliation, and other dermatological interventions. Its ability to accelerate barrier recovery and reduce visible irritation has made it a preferred botanical in barrier-repair creams, soothing serums, and recovery balms used in both professional and at-home settings (Brinkhaus et al., 2000; Bylka et al., 2014).

Commercial adoption further underscores CA's dermatological relevance. The ingredient is widely marketed under the term "CICA" and appears across multiple global skincare product categories, including creams, serums, toners, masks, and intensive repair treatments. This trend reflects sustained consumer demand for botanicals that combine soothing effects with demonstrated biological activity and good tolerability.

Within the United States, the growing use of *Centella asiatica* also aligns with evolving regulatory expectations under the Modernization of Cosmetics Regulation Act of 2022 (MoCRA), which expanded FDA authority over cosmetic product safety, substantiation, and post-market oversight. MoCRA emphasizes improved ingredient safety documentation, adverse event monitoring, and the development of formulations that minimize irritation and support skin barrier integrity, particularly for products intended for sensitive or compromised skin. Botanical ingredients such as *Centella asiatica*, which demonstrate anti-inflammatory activity, barrier-supportive effects, and favorable tolerability profiles, are increasingly relevant within this regulatory framework, as they enable manufacturers to meet heightened safety and quality expectations while maintaining formulation efficacy (U.S. Food and Drug Administration, 2022).

4. Internal Benefits and Systemic Applications

While *Centella asiatica* is widely recognized for its topical dermatological benefits, its systemic therapeutic potential has also been extensively explored. In traditional Ayurvedic and Chinese medicine, CA has long been used to support cognitive function, emotional balance, digestive comfort, and vascular health, reflecting its broad physiological relevance (Brinkhaus et al., 2000; Gohil et al., 2010). In modern contexts, CA is commonly incorporated into nutraceutical formulations such as capsules, teas, and standardized extracts aimed at stress modulation, cognitive support, and circulatory health.

Experimental and translational research suggests that CA exhibits adaptogenic and neuroprotective properties, in part through modulation of oxidative stress pathways and neurotransmitter systems associated with cognition and emotional regulation (Orhan, 2012). Preclinical studies indicate that CA extracts may influence cholinergic signaling and mitochondrial function while reducing stress related oxidative damage in neuronal tissues, mechanisms that support its traditional use as a cognitive tonic (Puttarak et al., 2017). Although large scale human trials remain limited, these findings provide a mechanistic basis for its growing inclusion in wellness and nootropic formulations.

The systemic vascular benefits of *Centella asiatica* are among its most clinically substantiated applications. Standardized extracts rich in triterpenoids have been shown to improve venous tone, enhance microcirculation, reduce capillary permeability, and strengthen connective tissue integrity (Cesarone et al., 2001). These effects have positioned CA as a well-recognized phytotherapeutic option for chronic venous insufficiency, varicose veins, and edema, particularly within European clinical practice.

In addition, CA has demonstrated relevance in systemic wound healing contexts. Oral administration of CA triterpenoids has been associated with reduced oxidative stress and

improved tissue repair capacity in experimental models, supporting its adjunctive role in conditions where healing is impaired (Shukla et al., 1999). The combined topical and internal applicability of CA distinguishes it as a versatile botanical with cross category relevance spanning dermatology, vascular medicine, and wellness science.

4.1 Clinical Evidence Supporting Systemic and Dermatological Use

Centella asiatica has been evaluated in a range of clinical and translational studies examining its effects on skin structure, vascular function, and overall skin tolerance. In dermatological research, human studies have demonstrated that topical formulations containing *Centella asiatica* contribute to improvements in skin barrier function, hydration, elasticity, and overall skin comfort. A prospective clinical study published in the *Journal of Cosmetic Dermatology* reported that a skincare product formulated with *Centella asiatica* leaf extract, ceramide NP, and panthenol significantly improved skin tolerance, reduced irritation, and supported barrier recovery in subjects with sensitive skin following regular use (Su et al., 2025).

In addition, a recent pilot clinical study evaluating a *Centella asiatica* extracellular vesicle based formulation demonstrated measurable improvements in facial skin hydration, elasticity, and texture after 28 days of application, further supporting the ingredient's relevance in cosmetic formulations targeting skin quality and visible aging parameters (Chang et al., 2025).

Clinical evidence supporting CA's vascular applications is particularly robust. Placebo controlled human trials conducted in Italy have evaluated standardized *Centella asiatica* extracts, including titrated extract of *Centella asiatica* (TECA), in individuals with venous hypertension and microcirculatory disorders. These studies demonstrated significant improvements in venous tone, capillary filtration, edema reduction, and microcirculatory function following supplementation, with good tolerability and no significant adverse effects reported (Cesarone et al., 2001; De Sanctis et al., 2001). Such findings underpin CA's established role in the management of chronic venous insufficiency.

With respect to cognitive and neurological outcomes, human clinical data remain limited; however, reviews integrating preclinical and early translational evidence suggest that CA extracts may support cognitive performance, stress resilience, and neuroprotection through antioxidant and mitochondrial-supportive mechanisms (Orhan, 2012; Puttarak et al., 2017). While further clinical studies are needed to define dosage and efficacy in human populations, the existing evidence aligns with CA's traditional use and supports its continued investigation within wellness and nutraceutical research.

Collectively, these clinical and translational findings validate *Centella asiatica* as a multi system botanical with demonstrated relevance to dermatological care, vascular health, and wellness focused applications. The convergence of topical and systemic evidence supports its ongoing use in both cosmetic and therapeutic-adjacent formulations.

5. Formulation Challenges and Ingredient Compatibility

Despite its well documented therapeutic potential, incorporating *Centella asiatica* into stable and effective cosmetic and pharmaceutical formulations presents several technical challenges. The primary bioactive constituents, asiaticoside, madecassoside, asiatic acid, and madecassic acid are chemically sensitive to heat, oxidative stress, and extreme pH conditions. Improper processing or incompatible formulation environments can lead to degradation of these triterpenoids, compromising product stability, shelf life, and biological efficacy (Hashim et al., 2011; James & Dubery, 2009).

To mitigate these limitations, modern formulation strategies increasingly rely on protective delivery systems. Liposomal encapsulation, nanoemulsions, and polymer-based carriers are widely used strategies to improve stability and delivery of sensitive actives, enhance skin penetration, and support controlled release within the epidermis and dermis (Bylka et al., 2014; Khursheed et al., 2024). Such systems are particularly valuable for preserving bioactivity during manufacturing and storage while improving bioavailability at target sites associated with collagen synthesis and wound repair.

pH control represents another critical formulation parameter. The triterpenoid components of *Centella asiatica* demonstrate optimal stability within mildly acidic environments, generally aligning with the physiological skin pH range of approximately 4.5–5.5 (Hashim et al., 2011). Formulations outside this range, particularly those containing strong alkalis or high concentrations of acidic exfoliants may accelerate degradation or reduce efficacy. Consequently, careful buffering and excipient selection are essential when CA is incorporated into multifunctional skincare systems.

Ingredient compatibility also plays a significant role in formulation success. *Centella asiatica* is frequently combined with soothing and barrier supporting ingredients such as niacinamide, panthenol, hyaluronic acid, and ceramides, which complement its anti-inflammatory and reparative effects (Bylka et al., 2014). These combinations are especially well suited for products targeting sensitive, post-procedure, or compromised skin. In contrast, caution is warranted when pairing CA with highly acidic exfoliating agents or aggressive retinoids, as such combinations may increase irritation potential or negatively impact triterpenoid stability.

Beyond topical applications, formulation research is expanding into nutricosmetic and ingestible formats. Emerging studies are exploring the synergistic use of *Centella asiatica* with other adaptogenic botanicals, such as *Withania somnifera* (ashwagandha) and *Panax ginseng*, to support systemic stress modulation, oxidative balance, and skin health from within (Orhan, 2012). As CA continues to be integrated into increasingly complex multi-ingredient systems, advanced formulation science remains essential to ensure stability, compatibility, and consistent therapeutic performance across cosmetic, pharmaceutical, and wellness applications.

6. Displacement and Complementation of Conventional Ingredients

The growing adoption of *Centella asiatica* in modern skincare and wellness formulations reflects a broader industry shift toward multifunctional, well tolerated botanicals. Rather than fully replacing conventional ingredients, CA increasingly serves as an alternative or

complementary active in formulations designed for sensitive, compromised, or post-procedure skin.

Historically, botanicals such as *Aloe vera* and *Calendula officinalis* have been widely used for their soothing and anti-inflammatory properties. While these ingredients remain valuable, CA offers a broader functional profile. In addition to calming inflammation, CA triterpenoids stimulate fibroblast activity, enhance collagen synthesis, and support extracellular matrix remodeling, mechanisms not strongly associated with aloe or calendula (Shukla et al., 1999; Bylka et al., 2014). As a result, CA is often selected in formulations intended to support wound healing, scar modulation, and barrier recovery alongside soothing benefits.

Niacinamide (vitamin B3) continues to be a cornerstone ingredient for barrier strengthening, tone improvement, and inflammation control. However, in clinical and cosmetic contexts involving ultra-sensitive or post-procedure skin, CA is frequently favored or used in combination due to its lower irritation potential and reparative activity (Brinkhaus et al., 2000; Bylka et al., 2014). In such cases, CA complements niacinamide by supporting skin recovery without provoking transient flushing or tingling that some users experience with higher niacinamide concentrations.

Hyaluronic acid remains unmatched as a primary humectant for hydration and skin plumping. However, CA contributes additional barrier-supportive and anti-inflammatory effects that extend beyond hydration alone. Its ability to reduce oxidative stress, promote dermal repair, and support structural integrity positions CA as a multifunctional adjunct in formulations where barrier repair and inflammation control are prioritized over moisture replenishment alone (Hashim et al., 2011; Shukla et al., 1999).

The versatility of *Centella asiatica* across concerns such as acne, aging, sensitivity, and post-inflammatory recovery has encouraged formulators to favor it when designing streamlined, efficacy driven skincare systems. This trend reflects a shift toward fewer, more biologically relevant actives rather than ingredient redundancy.

6.1 Comparative Perspective with Other Botanicals

Compared with other commonly used botanicals, *Centella asiatica* demonstrates a uniquely broad therapeutic spectrum. *Aloe vera*, while effective for hydration and surface level soothing, lacks the documented collagen stimulating and angiogenic properties associated with CA triterpenoids (Shukla et al., 1999). Similarly, chamomile and calendula are valued for their calming effects but do not exhibit the same influence on fibroblast proliferation, extracellular matrix remodeling, or microcirculatory support (Bylka et al., 2014).

Licorice root (*Glycyrrhiza glabra*) is widely used for brightening and inflammation reduction via tyrosinase inhibition; however, its primary activity is pigment modulation rather than tissue repair. In contrast, CA provides gentler, reparative support suitable for long-term use across a wider range of skin types, including sensitive and compromised skin (Brinkhaus et al., 2000).

Among adaptogenic botanicals used in wellness formulations, such as *Withania somnifera* (ashwagandha) and *Panax ginseng*, CA is distinctive in its compatibility with both topical and oral applications. While these adaptogens are primarily used systemically, CA's demonstrated efficacy in topical dermatology, vascular support, and cognitive research positions it as a rare cross-category botanical (Orhan, 2012).

As a result, CA is increasingly selected by formulators as a central active, either replacing single function botanicals or enhancing existing formulations, to meet growing demand for gentle, multifunctional, and scientifically supported ingredients in cosmetic and wellness products.

7. Global Market Trends and Economic Significance

The growing prominence of *Centella asiatica* reflects broader global trends favoring plant-derived, multifunctional ingredients across cosmetic, pharmaceutical, and wellness industries. Demand for CA has increased steadily over the past decade, driven by consumer preference for ingredients that combine efficacy, tolerability, and alignment with clean label and science backed formulation principles.

The global rise of Korean skincare ("K-beauty") played a central role in accelerating awareness of CA, particularly under the widely adopted "CICA" designation. Korean brands were among the first to position CA as a core ingredient for barrier repair, post-inflammatory care, and sensitive skin support. As consumer awareness expanded beyond Asia, multinational beauty and personal care companies incorporated CA into dermatologist tested, hypoallergenic, and recovery focused product lines, reinforcing its credibility and mainstream acceptance.

Beyond cosmetics, CA has gained visibility in pharmaceutical adjacent and nutraceutical applications, particularly in formulations targeting wound healing, vascular health, and cognitive wellness. Its multifunctional biological profile has enabled adoption across multiple product categories, reducing reliance on single function synthetic actives and supporting the development of streamlined, efficacy driven formulations.

From a supply chain perspective, *Centella asiatica* is primarily cultivated in regions including India, Sri Lanka, Madagascar, and parts of Southeast Asia. Increased global demand has encouraged more structured cultivation practices, including controlled harvesting, improved post harvest processing, and the adoption of organic and sustainability certifications. These developments have economic significance for producing regions, where CA cultivation contributes to agricultural diversification and income generation for smallholder farming communities.

The visibility of CA as a highlighted ingredient on product packaging and marketing materials underscores its dual scientific and commercial success. Its trajectory illustrates how a traditionally used botanical can transition into a globally recognized functional ingredient when supported by pharmacological research, formulation science, and evolving consumer expectations. As demand for evidence based natural ingredients

continues to rise, *Centella asiatica* serves as a representative case study of successful botanical integration into modern consumer health markets.

8. Sustainability and Ethical Sourcing

As global demand for *Centella asiatica* continues to rise, ensuring sustainable and ethical sourcing has become an increasingly important consideration. Historically, CA was often collected through wild harvesting; however, growing commercial demand has raised concerns regarding environmental degradation, biodiversity loss, and long term supply stability. In certain regions of South Asia and Africa, unmanaged harvesting has contributed to declining wild populations, underscoring the need for regulated and responsible cultivation practices.

In response, structured agricultural initiatives have emerged to support sustainable CA production. In India, cultivation programs led by cooperatives and non governmental organizations have promoted controlled farming and agroecological methods, enabling small scale farmers to transition from wild collection to cultivated supply. Similar efforts have been observed in Madagascar and Sri Lanka, where brands increasingly engage in direct partnerships with local growers to support fair trade practices, improve crop management, and invest in long-term soil and ecosystem health.

Certification frameworks such as ECOCERT, COSMOS, and USDA Organic are now commonly pursued by manufacturers seeking to demonstrate compliance with environmental and ethical standards. In parallel, traceability has become a growing priority within the cosmetic and wellness industries. Some supply chains now incorporate digital traceability tools, including batch level documentation and origin tracking, to provide transparency regarding sourcing practices and ingredient provenance.

Sustainable and ethical sourcing of *Centella asiatica* not only contributes to environmental conservation but also supports economic resilience within producing communities. As botanical ingredients become increasingly central to modern formulation science, the integration of sustainability principles alongside efficacy and safety considerations is essential. The case of CA highlights how botanical innovation can align scientific advancement with ecological responsibility and social accountability.

9. Conclusion

Centella asiatica's evolution from a traditional medicinal herb to a globally recognized cosmetic and wellness ingredient reflects the successful integration of ethnobotanical knowledge with modern scientific research. Its well characterized bioactive compounds, broad pharmacological activity, and growing body of clinical and translational evidence have established CA as a multifunctional botanical with relevance across dermatological, pharmaceutical adjacent, and wellness applications.

The widespread incorporation of CA into formulations addressing acne, skin aging, barrier dysfunction, post-inflammatory recovery, vascular health, and cognitive wellness underscores its adaptability within an evolving consumer and regulatory landscape. As demand continues to shift toward evidence based, well tolerated, and ethically sourced

ingredients, CA exemplifies how traditional botanicals can meet modern expectations for efficacy, safety, and sustainability.

Ongoing research is expected to further refine delivery systems, optimize formulation strategies, and explore additional clinical indications for *Centella asiatica*. These efforts will likely expand its therapeutic scope while reinforcing its role as a model ingredient for the future of functional botanicals, one that bridges heritage, science, and global health innovation.

10. Future Directions and Innovations

The future development of *Centella asiatica* lies at the convergence of traditional botanical knowledge and emerging scientific technologies. Advances in molecular biology, formulation science, and delivery systems are expected to further expand the clinical and cosmetic potential of CA. Novel topical delivery platforms, including nanoencapsulated gels, lipid based carriers, and dissolvable microneedle systems, are under investigation for their ability to enhance penetration, improve bioavailability, and enable more targeted delivery of triterpenoid saponins within the skin (Bylka et al., 2014; Khursheed et al., 2024).

Growing interest in psychodermatology, the study of interactions between psychological stress and skin health, represents another promising area for CA research. Given its documented neuroprotective and anxiolytic activity, CA is being explored as a supportive ingredient in formulations addressing stress associated skin conditions such as eczema, psoriasis, and stress exacerbated inflammatory disorders. Early experimental findings suggest that CA may influence cutaneous neuroendocrine signaling pathways involved in stress induced inflammation, although further mechanistic and clinical studies are required to substantiate these effects (Orhan, 2012; Puttarak et al., 2017).

Microbiome focused skincare represents an additional frontier for CA innovation. Research is evaluating how CA-derived triterpenoids interact with skin associated microbes and downstream inflammatory pathways. For example, madecassoside (from *Centella asiatica*) showed Propionibacterium (Cutibacterium) acnes related anti-inflammatory activity and supported skin hydration/barrier associated markers in vitro, suggesting potential relevance for acne-prone and sensitive skin contexts (Shen et al., 2019).

Renewed interest is also emerging in wound management applications, including post surgical recovery and chronic wounds such as diabetic ulcers. CA's angiogenic, collagen stimulating, and anti scarring properties position it as a valuable component of next generation wound-care strategies (Shukla et al., 1999; Somboonwong et al., 2012). Current exploratory work includes polyherbal and multi active formulations combining CA with complementary bioactives such as panthenol, peptides, and other plant-derived compounds to enhance reparative outcomes.

Finally, sustainability driven innovation is shaping the future of CA production. Advances in green chemistry and extraction technologies are enabling more

environmentally responsible methods that reduce solvent use, improve yield efficiency, and preserve bioactive integrity. These developments support a holistic approach to CA utilization, one that integrates efficacy, safety, and environmental stewardship, aligning botanical innovation with the principles of future-ready cosmeceutical and wellness design.

References

- Brinkhaus, B., Lindner, M., Schuppan, D., & Hahn, E. G. (2000). Chemical, pharmacological and clinical profile of the East Asian medical plant *Centella asiatica*. *Phytomedicine*, 7(5), 427–448.
[https://doi.org/10.1016/S0944-7113\(00\)80065-3](https://doi.org/10.1016/S0944-7113(00)80065-3)
- Bylka, W., Znajdek-Awizen, P., Studzińska-Sroka, E., Dańczak-Pazdrowska, A., & Brzezińska, M. (2014). *Centella asiatica* in dermatology: An overview. *Phytotherapy Research*, 28(8), 1117–1124.
<https://doi.org/10.1002/ptr.5110>
- Cesarone, M. R., Belcaro, G., De Sanctis, M. T., Incandela, L., Bavera, P., Ippolito, E., Bucci, M., & Griffin, M. (2001). Effects of the total triterpenic fraction of *Centella asiatica* in venous hypertensive microangiopathy: A prospective, placebo-controlled, randomized trial. *Angiology*, 52(Suppl 2), S15–S18.
PMID: 11666117
- De Sanctis, M. T., Belcaro, G., Incandela, L., & Cesarone, M. R. (2001). Treatment of edema and increased capillary filtration in venous hypertension with total triterpenic fraction of *Centella asiatica*. *Angiology*, 52(Suppl 2), S55–S59.
PMID: 11666125
- Khursheed, T., Khalil, A. A., Akhtar, M. N., Alsulami, T., Mugabi, R. M., & Nayik, G. A. (2024). Ultrasound-assisted solvent extraction of phenolics, flavonoids, and major triterpenoids from *Centella asiatica* leaves: A comparative study. *Ultrasonics Sonochemistry*, 111, 107091.
<https://doi.org/10.1016/j.ultsonch.2024.107091>
- Gohil, K. J., Patel, J. A., & Gajjar, A. K. (2010). Pharmacological review on *Centella asiatica*: A potential herbal cure-all. *Indian Journal of Pharmaceutical Sciences*, 72(5), 546–556.
PMCID: PMC3210009; PMID: 21188013
- Puttarak, P., Dilokthornsakul, P., Saokaew, S., Dhippayom, T., Kongkaew, C., Sruamsiri, R., & Chuthaputti, A. (2017). Effects of *Centella asiatica* on cognitive function and mood: A systematic review and meta-analysis. *Scientific Reports*, 7, 10646.

<https://doi.org/10.1038/s41598-017-09823-9>

Hashim, P., Sidek, H., Helan, M. H. M., Sabery, A., Palanisamy, U. D., & Ilham, M. (2011).

Triterpene composition and bioactivities of *Centella asiatica*. *Molecules*, 16(2), 1310–1322.

<https://doi.org/10.3390/molecules16021310>

James, J. T., & Dubery, I. A. (2009).

Pentacyclic triterpenoids from the medicinal herb *Centella asiatica* (L.) Urban. *South African Journal of Botany*, 75(3), 417–426.

<https://doi.org/10.1016/j.sajb.2009.02.002>

Orhan, I. E. (2012).

Centella asiatica (L.) Urban: From traditional medicine to modern medicine with neuroprotective potential. *Evidence-Based Complementary and Alternative Medicine*, 2012, Article ID 946259.

<https://doi.org/10.1155/2012/946259>

Su, Z., Zheng, Y., Yi, J., Lai, W., & Ye, C. (2025).

The effectiveness and safety of a skin care product with *Centella asiatica* leaf extract, ceramide NP, and panthenol in subjects with sensitive skin: A prospective, observational study. *Journal of Cosmetic Dermatology*, 24(7), e70324.

<https://doi.org/10.1111/jocd.70324>

Chang, T.-M., Wu, C.-C., Huang, H.-C., Lu, J.-Y., Chuang, C.-H., Kao, P.-L., Tang, W.-H., Liu, L.-T.-C., Qiu, W.-Y., Percec, I., Chen, C., & Kuo, T.-Y. (2025).

A 28-day pilot study of the effects on facial skin hydration, elasticity, and texture of a *Centella asiatica* extracellular vesicle-based skin care formulation. *Cosmetics*, 12(5), 186.

<https://doi.org/10.3390/cosmetics12050186>

Shukla, A., Rasik, A. M., & Dhawan, B. N. (1999).

Asiaticoside-induced elevation of antioxidant levels in healing wounds. *Phytotherapy Research*, 13(1), 50–54.

Somboonwong, J., Kankaisre, M., Tantisira, B., & Tantisira, M. H. (2012).

Wound healing activities of different extracts of *Centella asiatica* in incision and burn wound models: An experimental animal study. *BMC Complementary and Alternative Medicine*, 12, Article 103.

<https://doi.org/10.1186/1472-6882-12-103>

U.S. Food and Drug Administration (FDA). (2022).

Modernization of Cosmetics Regulation Act of 2022 (MoCRA). U.S. Department of Health and Human Services. <https://www.fda.gov/cosmetics/cosmetics-laws->

[regulations/modernization-cosmetics-regulation-act-2022](#)

Shen, X., Guo, M., Yu, H., Liu, D., Lu, Z., & Lu, Y. (2019). *Propionibacterium acnes* related anti-inflammation and skin hydration activities of madecassoside, a pentacyclic triterpene saponin from *Centella asiatica*. **Bioscience, Biotechnology, and Biochemistry**, 83(3), 561–568. <https://doi.org/10.1080/09168451.2018.1547627>