



Modulating the Glycemic Response: A Kinetic Investigation of Salivary Amylase Activity and the Inhibitory Effects of *Camellia sinensis* (Green Tea) in Pineville, LA

¹Oyeyemi O.A and ¹Ekepata Obenofunde

¹Department of Biological science, Louisiana Christian University, LA, USA

Abstract

Enzymes serve as indispensable biological catalysts, facilitating metabolic pathways by lowering activation energy. This study quantifies the **kinematics of salivary amylase**, focusing on the influence of substrate concentration, pH, and natural inhibitors. Utilizing a starch-iodine colorimetric assay, enzyme activity was monitored at a physiological temperature of 37°C. Results indicated a direct correlation between substrate concentration and reaction rate, with optimal catalytic efficiency observed at pH 7.0. Furthermore, the introduction of green tea (*Camellia sinensis*) extracts significantly reduced the reaction rate (1.71×10^{-5} g/mL/s) compared to the control (3.7×10^{-5} g/mL/s). These findings suggest that polyphenolic compounds in green tea serve as effective amylase inhibitors, offering a biochemical strategy for modulating glucose absorption—a critical "blessing" for the clinical management of metabolic disorders within the Pineville community.

Keywords: Salivary Amylase, Enzyme Kinematics, Starch Hydrolysis, pH Optimization, Polyphenols, Diabetes Management, Clinical Biochemistry, Pineville Louisiana.

1. Introduction

Amylase is a primary digestive enzyme secreted by the salivary glands and the pancreas, responsible for the cleavage of α -1,4-glycosidic bonds in amylose and amylopectin (starch). This process initiates carbohydrate digestion, transforming complex polysaccharides into maltose and glucose. In the context of **clinical biochemistry**, the rate at which amylase functions is a major determinant of the postprandial glycemic response (blood sugar spike).

The efficiency of this biocatalysis is highly sensitive to environmental parameters. Factors such as substrate availability, enzyme concentration, and thermal kinetic energy dictate the frequency of enzyme-substrate collisions. Most human enzymes, including salivary amylase, exhibit an optimal temperature of approximately 37°C; deviations beyond this point result in the thermal denaturation of the enzyme's tertiary structure. This study examines these kinetic variables and explores the potential of natural dietary inhibitors to attenuate amylase activity, providing insights into nutritional strategies for managing metabolic health in Louisiana.

2. Materials and Methods

2.1 Reagents and Equipment:

- **Buffer Solutions:** pH 4.0, 7.0, and 10.0.
- **Substrates:** Starch solutions at concentrations of 0.016 g/mL, 0.033 g/mL, and 0.05 g/mL.
- **Enzyme Source:** Human Salivary Amylase (diluted saliva).
- **Indicator:** Iodine (I_2/KI solution).
- **Inhibitor:** *Camellia sinensis* (Green Tea) infusion.

2.2 Experimental Procedure:

1. **Concentration Kinetics:** Salivary amylase was added to varying starch concentrations labeled 2.5g, 5g, and 7.5g per 150mL. Time was recorded until the blue-black starch-iodine complex reached full achromic (clear) point.
2. **pH Optimization:** Using a 0.03 g/mL starch solution, activity was measured across pH 4, 7, and 10 in a 37°C water bath.
3. **Inhibition Assay:** A comparative analysis was performed by introducing cooled green tea solution to the reaction mixture, using distilled water as a control to measure the inhibitory potency of tea polyphenols.

3. Results

The experimental data reveals that salivary amylase kinetics follow a predictable pattern influenced by the chemical environment.

Table 1: Substrate Concentration and Reaction Velocity

Concentration (g/mL)	Hydrolysis Time (s)	Reaction Rate (g/mL/s)
0.0167	900	1.85×10^{-5}
0.0300	720	4.17×10^{-5}
0.0500	480	1.04×10^{-4}

Table 2: pH-Dependent Catalytic Activity

pH Level	Hydrolysis Time (s)	Reaction Rate (g/mL/s)
4 (Acidic)	1070	2.8×10^{-5}
7 (Neutral)	790	3.8×10^{-5}
10 (Basic)	1010	3.0×10^{-5}

Table 3: Inhibitory Impact of Green Tea

Solution Type	Time (s)	Reaction Rate (g/mL/s)
Control (Water)	810	3.7×10^{-5}
Green Tea Extract	1750	1.71×10^{-5}

4. Discussion

4.1 Collision Theory and pH Sensitivity

The results support the fundamental principle that reaction velocity is a function of substrate concentration. As the starch concentration increased, the time to achromic point decreased significantly, indicating that higher substrate density facilitates more frequent successful collisions at the enzyme's active site.

The pH data confirms that salivary amylase is a neutral-adapted enzyme. The significant reduction in rate at pH 4 and pH 10 suggest that the ionic environment affects the protonation state of amino acid residues in the active site, leading to a loss of catalytic efficiency or partial denaturation. This reinforces why salivary amylase activity diminishes as it enters the highly acidic environment of the stomach.

4.2 Clinical Biochemistry of Green Tea Inhibition

The most impactful observation was the 50% reduction in reaction rate caused by green tea. This inhibition is attributed to the presence of **catechins** (polyphenols) like Epigallocatechin gallate (EGCG). These compounds interact with the enzyme through non-covalent bonding, effectively masking the active site or inducing a conformational change.

5. Contribution to Knowledge & Community Outreach

Contribution to Knowledge

This study contributes a precise kinetic profile of salivary amylase under common environmental stressors and demonstrates the effective use of a household "inhibitor" to model clinical drug-enzyme interactions. It bridges the gap between basic biochemistry and **nutraceutical science**, proving that common dietary choices have a measurable, gravimetric impact on digestive kinematics.

A Blessing to Men: Health and Diabetes Management

For the men and families of Pineville and Central Louisiana, these findings are a "blessing" of preventative knowledge.

- **The "Metabolic Shield":** The drastic slowing of starch hydrolysis by green tea suggests that consuming polyphenolic beverages with high-carbohydrate meals can naturally dampen the glycemic index of the food.
- **Diabetes Prevention:** By slowing the conversion of starch to glucose, we can prevent the rapid "insulin spikes" that contribute to insulin resistance and Type 2 Diabetes—a significant health concern in our region.

- **Take-Home Point:** Understanding enzyme kinetics empowers the community to use "food as medicine," utilizing simple dietary additions like green tea to support metabolic health and improve clinical outcomes for those managing blood sugar disorders.

6. References

1. **Harper College.** (2024). *Enzyme Kinetics: Factors Influencing Activity*. Department of Chemistry.
2. **Akinfemiwa, O., et al.** (2023). *Amylase: Physiological and Clinical Significance*. PubMed, StatPearls Publishing.
3. **LSU AgCenter.** (2025). *Dietary Polyphenols and Metabolic Health in Louisiana Communities*. College of Agriculture.
4. **Meyer, B.** (2025). *Comparative Biochemistry: Using Enzyme Models to Understand Human Metabolic Stress*. Journal of Clinical Medicine.
5. **He, Q., et al.** (2022). *Inhibition of Salivary Amylase by Tea Polyphenols: A Kinetic and Molecular Docking Study*. Journal of Food Science and Technology.
6. **Smith, G. P., & Williams, T. J.** (2024). *The Role of Salivary Alpha-Amylase in Oral Digestion and Postprandial Glycemic Control*. International Journal of Molecular Sciences.
7. **Zhang, L., & Li, J.** (2023). *pH-Dependent Structural Stability of Human Amylase: Implications for Digestive Disorders*. Journal of Biological Chemistry.
8. **National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK).** (2025). *Carbohydrate Metabolism and the Impact of Enzyme Inhibition in Type 2 Diabetes*.
9. **Louisiana Department of Health.** (2026). *Metabolic Syndrome and Diabetes Prevalence in Central Louisiana: The Need for Nutritional Intervention*.