Discover Agriculture

RESEARCH Open Access

Mineral composition of neglected and underutilized food crops and their role in nutrition security in Sierra Leone

Sahr Foday[1*], Richard Wadsworth[2], Tamba S. Sonda[3] and Samuel Maxwell Tom Williams[2,4]

*Correspondence:
Sahr Foday
sfoday@njala.edu.sl
[1]Njala University Hospital, Njala University, Moyamba District, Bo, Sierra Leone
[2]Department of Biological Sciences, Njala University, Moyamba District, Bo, Sierra Leone
[3]Institute of Food Technology, Nutrition & Consumer StudiesMoyamba District, Bo, Sierra Leone
[4]Laboratory of Public Health, Graduate School of Infectious Disease, Hokkaido University, Hokkaido, Japan

Abstract

Background Sierra Leone's food system faces persistent nutritional challenges, largely due to limited crop diversity and reliance on imported staples. Neglected and Underutilized Food Crops (NUFCs), though culturally embedded and ecologically resilient, remain underexplored in formal nutrition and agricultural strategies. This study evaluates the mineral composition of selected NUFCs cultivated in Bo and Kenema districts regions known for their agricultural activity and vulnerability to micronutrient deficiencies.

Methodology A mixed-methods approach was employed, combining ethnobotanical surveys with laboratory-based nutritional analysis. Plant samples were collected through guided field walks and community engagement, then identified and authenticated at the National Herbarium, Njala University. Mineral profiling was conducted using standardized procedures: Kjeldahl digestion for nitrogen (converted to protein), Bray I extraction for phosphorus, and Atomic Absorption Spectrophotometry (AAS) and flame photometry for trace elements including iron (Fe), zinc (Zn), copper (Cu), calcium (Ca), magnesium (Mg), potassium (K), sodium (Na), and phosphorus (P). All concentrations were standardized to mg/100 g for comparability.

Result The results revealed that *Piper umbellatum* exhibited the most diverse mineral profile, with high concentrations of zinc (12.49 mg/100 g), phosphorus (103.1 mg/100 g), magnesium (8.94 mg/100 g), calcium (11.62 mg/100 g), and sodium (1.93 mg/100 g). *Celosia argentea* and *Luffa aegyptiaca* were iron-rich, while *Ipomoea aquatica* showed the highest protein content (7.00%) and notable potassium levels. These findings affirm the nutritional relevance of NUFCs and their potential to address micronutrient deficiencies, particularly in rural communities with limited access to nutrient rich foods.

Conclusion This study confirms the nutritional value of selected indigenous NUFCs that are widely cultivated, long-naturalized, and traditionally consumed in Sierra Leone. Species such as *Piper umbellatum*, *Celosia argentea*, and *Ipomoea aquatica* demonstrate strong potential to enhance dietary diversity, public health, and food system resilience. The findings lay a foundation for future research, policy action, and community-level nutrition initiatives.

© The Author(s) 2026. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/.

Keywords Neglected plant species, Leafy vegetables, Micronutrient composition, Nutrition secutrity, Sierra Leone

1 Introduction

Sierra Leone's agricultural landscape is rich in biodiversity, yet its food system remains heavily reliant on a narrow range of staple crops primarily rice, cassava, and maize that offer limited nutritional diversity and are increasingly vulnerable to climate shocks. Despite having over 5.36 million hectares of cultivable land [12, 41], the country continues to import approximately 80% of its food [38], underscoring a systemic gap between agricultural potential and nutritional outcomes. This reliance on imported staples and low-diversity cropping systems has contributed to widespread micronutrient deficiencies, particularly among women and children, with iron-deficiency anemia, zinc deficiency, and protein-energy malnutrition remaining prevalent [34]. According to the World Food Programme's 2024 Food Security Monitoring Report, 82% of the population in Sierra Leone is food insecure, of which 18% are classified as severely food insecurity [38].

Neglected and Underutilized Food Crops (NUFCs) represent a promising yet overlooked solution to these challenges. These crops, often indigenous, climate-resilient, and culturally embedded have historically contributed to local diets and traditional medicine but remain marginalized in formal agricultural and nutrition programs [16, 28]. In Sierra Leone, species such as *Piper umbellatum L.*, *Celosia argentea (L.) T. Anderson*, *Ipomoea aquatica* Forssk and *Vernonia amygdalina Delile* are widely known and consumed in rural communities, yet their nutritional profiles are poorly documented and underappreciated in policy and research circles. The lack of empirical data on the mineral composition of NUFCs has hindered their integration into national food security strategies. While anecdotal and ethnobotanical evidence supports their nutritional value, systematic laboratory-based assessments are scarce. Comparative studies with global benchmarks reveal inconsistencies in nutrient values due to environmental, genetic, and methodological factors [3, 29]. Without standardized data and validated nutritional profiles, NUFCs remain excluded from agricultural extension services, dietary guidelines, and public health interventions.

This study addresses that gap by evaluating the mineral composition of selected NUFCs cultivated in Bo and Kenema districts regions known for their agricultural engagement and vulnerability to nutritional deficiencies. By quantifying key minerals such as iron (Fe), zinc (Zn), copper (Cu), phosphorus (P), potassium (K), calcium (Ca), magnesium (Mg), and sodium (Na), the research aims to scientifically validate the nutritional potential of these crops.

The study aligns with national development priorities, including the Feed Salone initiative, which seeks to reduce dependency on imported staples and promote indigenous crops for food sovereignty. It also contributes to global efforts to mainstream underutilized crops in sustainable food systems [22, 37]. By bridging scientific analysis with traditional knowledge, this research positions NUFCs as vital assets in the fight against hidden hunger and agricultural vulnerability in Sierra Leone.

We aim to evaluate the mineral composition of selected NUFCs cultivated in Bo and Kenema districts to assess their potential contribution to food and nutrition security

in Sierra Leone. In this study, we hypothesize that these crops possess mineral profiles capable of addressing micronutrient deficiencies prevalent in the region. In this study, we hypothesize that these crops possess mineral profiles capable of addressing micronutrient deficiencies prevalent in the region. Because these crops were collected from different localities, their nutritional contribution is influenced not only by inherent mineral composition but also by factors such as bioavailability, antinutrients, dietary practices, cultural patterns of consumption, and environmental or varietal differences; these aspects were beyond the scope of the present study.

2 Methodology

2.1 Study area

The study was conducted in Bo and Kenema districts of Sierra Leone, two agriculturally active regions located in the Southern and Eastern Provinces, respectively. Bo District spans approximately 5473 km^2 and is characterized by a tropical climate with fertile soils suitable for rice, cassava, maize, and leafy vegetables [26]. Kenema District, covering about 6053 km^2, shares similar agro-ecological features but is also known for its inland valley swamp (IVS) systems and cash crops like cocoa and coffee [27]. Both districts experience a monsoon-type humid tropical climate, with annual rainfall ranging from 2,000 mm in the north to 4000 mm in the south [27]. These conditions support diverse cropping systems and make the regions ideal for studying NUFCs, which are often cultivated in marginal or mixed farming environments (Fig. 1).

2.2 Study design

This research employed a mixed-methods design, integrating both quantitative laboratory analysis and qualitative ethnobotanical inquiry. The study focused on assessing the mineral composition of selected NUFCs to determine their nutritional potential. A purposive sampling strategy was used to select plant species based on local consumption patterns, traditional knowledge, and availability in the study area. The design included structured questionnaires with an in-depth interview, guided field walks, and laboratory-based chemical assays. This triangulated approach ensured a comprehensive understanding of both the biochemical properties and socio-cultural relevance of NUFCs.

2.3 Sample size and species selection

The study collected data on 45 neglected and underutilized food crops (NUFCs) comprising 23 species. Species were categorized into three groups based on binomial statistical analysis: crops with increasing utilization, crops with decreasing utilization, and crops with no significant change.

For the present study on proximal mineral composition, eight NUFC species were purposely selected from the latter two categories, those with declining usage or no significant change ($p > 0.05$). Selection was guided not only by statistical classification but also by practical considerations including local availability, adaptability to extreme environmental conditions, cultural values, and potential nutritional relevance. The final sample included species such as *Luffa aegyptic Mill*, *Justicia insularis T. Anderson*, *Triumfetta cordifolia A. Rich*, *Piper umbellatum L.*, *Asystasia gangetica (L.)*, *Celosia argentinae (L.) T. Anderson*, *Vernoniamy amygdalina Delile*, and *Ipomoea aquatica* Forssk. These crops were identified through community consultations with experienced local farmers,




Fig. 1 Satellite map of Sierra Leone showing the study districts of Bo and Kenema, highlighted in yellow. Major cities and district boundaries are labeled for geographic reference

women, hunters, and farm traders, and represent a subset of NUFCs at risk of underutilization despite their nutritional promise.

2.4 Data collection

2.4.1 Fieldwork and ethnobotanical survey

Data collection began with the training of research assistants in digital data-capture tools such as Kobo Toolbox, followed by a pilot survey to refine the questionnaire and ensure clarity. As part of a broader research project, an ethnobotanical survey was conducted to document indigenous knowledge, cultivation practices, and community perceptions of NUFCs. Although the detailed ethnobotanical findings have been published separately [13, 14], this component informed species selection and contextual understanding for the present nutritional analysis.

The ethnobotanical survey employed semi-structured interviews, focus group discussions (FGDs), and guided field walks. To ensure broad representation of knowledge holders, the sample universe was expanded using a snowball sampling approach. Initial key informants identified through local agricultural extension officers recommended additional participants with expertise in NUFC cultivation and use. This iterative process enabled the inclusion of experienced farmers, market vendors, traditional healers,

and community elders across multiple villages and markets in Bo and Kenema districts. Snowball sampling is widely used in ethnobotanical research to capture community-embedded and often dispersed traditional knowledge systems.

The design and implementation of the ethnobotanical survey followed established methodological guidelines, including those outlined by Martin [19], Alexiades and Sheldon [4], and Cotton [11]. These references informed the structure of interview guides, participant selection procedures, and field documentation techniques. Guided field walks allowed for direct observation, identification, and photographic documentation of species in both cultivated and wild settings. In the context of this manuscript, these procedures ensured accurate species identification and relevance to local food systems, while the detailed cultural and medicinal insights are reported in the previously published ethnobotanical study.

2.4.2 Plant specimen collection and identification

Plant samples were collected from home gardens, farms, and forest margins. Priority was given to fertile specimens bearing flowers or fruits to ensure accurate identification. Young leaves were harvested using knives or cutlasses and stored in sealed transparent plastic bags to preserve freshness. With guidance from herbarium experts, samples were treated to prevent early dehydration and transported to the Quality Control Laboratory at Njala University for analysis.

Voucher specimens were prepared and authenticated by experts at the National Herbarium, Njala University, using standard botanical literature and morphological keys. Photographs and local names were recorded for each species, and samples were preserved using alcohol and pressing techniques for reference and laboratory analysis. The plant parts analyzed (proximal mineral composition) in this study were the young leaves sourced from the following locations and institutions:

1. *Piper umbellatum L.*: Collected from Kasewe Forest Reserve (coordinates: 8.336248 -12.170405), under guidance of District Forest Officer, Moyamba District.
2. *Triumfetta cordifolia A.Rich*: Collected from Kasewe Forest Reserve (coordinates: 8.342758 -12.180665), under guidance of District Forest Officer, Moyamba District.
3. *Vernoniamy amygdalina Delile*: Obtained from Small Bo chiefdom, Kenema District under the guidance of the town chiefs (coordinates: 7.8502519 -11.2486597).
4. *Justicia insularis (L.) T. Anderson*: Obtained from Tikonko chiefdom, Bo District under the guidance of the town chief (coordinates: 7.8877352 -11.7858397).
5. *Asystasia gangetica*: Collected from Gbo chiefdom, Bo District under the guidance of the town chief (coordinates: 8.070604210636244, -11.826134556167704).
6. *Luffa aegyptica Mill*: Collected from Gbo chiefdom, Bo District under the guidance of the town chief (coordinates: 8.070604210636244, -11.826134556167704).
7. *Celosia argentea L.*: Collected from Kandu Leppiam chiefdom, Kenema District under the guidance of the town chief (coordinates: 8.070604210636244, -11.826134556167704).
8. *Ipomoea aquatica*: Obtained from Small Bo chiefdom, Kenema District under the guidance of the town chief (coordinates: 7.8502519 -11.2486597).

2.4.3 Laboratory analysis

Proximal mineral analysis was conducted on eight selected neglected and underutilized food crops (NUFCs) identified through prior statistical and ethnobotanical screening. Fresh leaf samples of each species were collected from Bo and Kenema districts, air-dried at ambient temperature, and subsequently oven-dried at 60 °C to constant weight. The dried samples were ground into fine powder using a stainless-steel mill and stored in airtight containers prior to analysis. The chemical analysis was conducted at the Environmental Chemistry Laboratory, Njala University. The Kjeldahl method was used to estimate nitrogen content, which was converted to protein using the [6] standard factor of 6.25.

2.5 Sample preparation and digestion

Following field collection and processing, each plant species was represented by one composite sample, prepared from the edible portion of a single representative plant. To ensure analytical reliability, each composite sample was analyzed in duplicate, resulting in two independent laboratory repetitions per species.

All samples were oven-dried, finely ground, and sieved prior to mineral analysis. Each powdered sample was subjected to acid digestion using a mixture of nitric acid (HNO_3) and perchloric acid ($HClO_4$) in a 2:1 ratio. Approximately 0.5 g of sample was digested in a fume hood until a clear solution was obtained. The digested samples were then filtered and diluted to a final volume of 50 mL with deionized water. The resulting solutions were used for quantification of mineral elements.

Mineral elements were quantified using:

• Nitrogen content was determined using the Kjeldahl method, which involves digestion with concentrated sulfuric acid (H_2SO_4) in the presence of a catalyst, followed by distillation and titration. Crude protein was calculated by multiplying the total nitrogen content by a conversion factor of 6.25, based on the assumption that protein contains approximately 16% nitrogen [6].

• Iron (Fe), cooper (Cu), and Zinc (Zn) were determined using Atomic Absorption Spectrophotometry (AAS), following the Ammonium Bicarbonate DTPA method [32].

• Phosphorus (P) was quantified using the Bray I extraction method, followed by colorimetric determination with a spectrophotometer [9].

• Flame photometry for potassium (K), sodium (Na), calcium (Ca), and magnesium (Mg).

All concentrations were initially recorded in mg/kg and converted to mg/100 g using the Nutraceuticals Group's online conversion scale [23] to ensure consistency with global nutritional reporting standards.

2.6 Statistical analysis

Quantitative data from laboratory assays were analyzed using Microsoft Excel, SPSS v26, and R. Descriptive statistics were used to summarize mineral concentrations across NUFC species. Comparative analysis was performed against global literature to assess alignment and variability in nutrient profiles.

Qualitative data from interviews and FGDs were thematically analyzed to contextualize the nutritional findings within traditional knowledge systems and community

practices. This dual analysis provided a holistic understanding of the nutritional value and cultural relevance of NUFCs in Sierra Leone's food system.

2.7 Ethical considerations

Although the study was primarily based in Bo and Kenema districts, two plant species were also collected from the Kasewe Forest Reserve in Moyamba District. Prior to data collection, participants were informed about the purpose and nature of the research. Ethical approval was obtained from the Directorate of Research and Development, Njala University, Sierra Leone, and the study adhered to the ethical principles of the Declaration of Helsinki. Informed consent was obtained from all participants, including consent for the publication of anonymized data. The study does not involve a clinical trial; hence, a clinical trial number is not applicable.

In addition to human subject protocols, ethical clearance also covered the collection of plant specimens. All plant materials were collected in accordance with national guidelines and institutional protocols. No endangered or protected species were involved. Permissions were obtained from the Sierra Leone Ministry of Agriculture and Forestry and local authorities, including district forest officers and town chiefs. While no formal reference number was issued, written consent and permission letters were provided by Njala University to the Ministry of Agriculture and local authorities prior to specimen collection. Documentation of these permissions is available upon request.

3 Results

3.1 Proximal mineral composition of selected NUFCs

The mineral analysis revealed considerable variation in nutrient profiles across the eight selected neglected and underutilized food crops (NUFCs). Table 1 summarizes the concentrations of key minerals calcium (Ca), magnesium (Mg), potassium (K), sodium (Na), phosphorus (P), iron (Fe), zinc (Zn) as well as crude protein content, expressed as a percentage of dry weight.

3.1.1 Iron (Fe)

The crops with the highest iron content are *Celosia argentea* (15.226 mg/100 g) and *L.aegyptica* (12.962 mg/100 g) making them an excellent candidate for addressing iron deficiency anemia, a significant public health issue in Sierra Leone. Iron is crucial for oxygen transport and energy production, and its deficiency can lead to fatigue, weakened immunity, and developmental delays in children (Obeagu, 2025). *C. argentea* and *L. aegyptica* rich iron content highlights its potential to improve dietary iron intake, particularly in populations with limited access to fortified or animal-based iron sources.

Table 1 N/Protein and mineral composition of the selected NUFC (mg/100 g)

Species	Fe	Cu	Zn	*N*	*P*	Na	K	Mg	Ca
Luffa aegyptic	129.62	2.59	9.59	0.66	105.40	3.47	12.46	14.51	20.88
Justicia insularis	99.82	2.00	13.09	0.28	62.80	8.93	3.93	16.07	53.61
Triumfetta cordifolia	15.21	0.30	12.30	0.10	70.90	13.20	3.20	6.68	25.20
Piper umbellatum	24.74	0.49	12.49	0.31	103.10	1.930	24.93	8.94	11.62
Asystasia gangetica	77.18	1.54	5.54	0.24	105.40	5.63	28.67	12.56	34.00
Celosia argentinae	152.26	3.05	15.05	0.38	111.50	3.41	86.41	8.40	20.80
Vernoniamy cordiafolia	7.65	221.30	17.55	2.84	11.80	7.65	221.30	55.41	76.89
Ipomoea aquatica	0.14	179.90	6.55	1.12	23.52	0.14	179.90	33.20	45.59

This crop could be integrated into local diets to enhance iron availability and overall nutritional status. It is also important to note that the nutritional impact of iron depends on factors influencing its absorption such as bioavailability and the presence of dietary inhibitors or enhancers which were not assessed in this study.

3.1.2 Copper (Cu)

Vernonia amygdalina stands out with the highest copper content (22.13 mg/100 g), followed by *Ipomoea aquatica* (17.99 mg/100 g), far exceeding other crops in the study. Copper is essential for red blood cell formation, immune system function, and the maintenance of healthy nerves and bones. Its high concentration in *Vernonia amygdalina* and *Ipomoea acquatica* underscores the crop's potential role in preventing copper deficiency, which can cause anemia and cardiovascular issues. This exceptional copper content makes it an ideal addition to diets in areas where trace mineral deficiencies are prevalent.

3.1.3 Zinc (Zn)

Piper umbellatum leads in zinc content with 12.49 mg/100 g, making it a valuable source for supporting immune health, wound healing, and cellular metabolism. Zinc deficiencies are associated with stunted growth, increased susceptibility to infections, and delayed wound healing. By incorporating *Piper umbellatum* into local diets, particularly in regions with zinc-deficient soils, communities can address this critical nutrient gap and improve overall health outcomes, especially in children and pregnant women.

3.1.4 Protein

The crop with the highest protein content is *Ipomoea aquatica* (7.00%), positioning it as a critical resource for addressing protein-energy malnutrition. Protein is vital for growth, tissue repair, and enzymatic functions, making it a cornerstone of human nutrition. *Ipomoea aquatica* has relatively high protein levels offers an affordable plant-based option for supplementing protein intake, particularly in low-resource settings where animal protein sources are scarce. This crop's role in improving dietary diversity and nutrition is significant, especially for vulnerable populations.

3.1.5 Phosphorus (P)

Piper umbellatum exhibits the highest phosphorus content (103.1 mg/100 g), making it an essential crop for supporting bone health, energy production, and cellular repair. Phosphorus is a key component of DNA, RNA, and ATP, underscoring its importance in both structural and metabolic functions. By integrating *Piper umbellatum* into diets, communities can enhance their phosphorus intake and support overall growth and development, particularly in children and adolescents.

3.1.6 Sodium (Na)

Among the crops analyzed, *Piper umbellatum* also leads in sodium content (1.93 mg/100 g), followed by *Triumfetta cordiafolia* (1.32 mg/100 g). While sodium is commonly present in diets, its moderate levels in this crop suggest it can contribute to maintaining electrolyte balance without posing risks of excessive sodium intake. This makes *Piper umbellatum* and *Triumfetta cordiafolia* suitable for complementing

low-sodium diets while still supporting essential functions such as nerve transmission and muscle contraction.

3.1.7 Potassium (K)

The highest potassium levels are found in *Vernonia amygdalina* (22.13 mg/100 g) followed by *Ipomoea acquatica* (17.99 mg/100 g), highlighting its role in supporting cardiovascular health and maintaining fluid balance. Potassium helps regulate blood pressure, making it a key nutrient for preventing hypertension and reducing the risk of heart disease. By incorporating *Vernonia amygdalina* and *Ipomoea acquatica* into daily diets, individuals can improve their potassium intake and promote overall cardiovascular health, especially in regions with a high prevalence of hypertension.

3.1.8 Magnesium (Mg)

With a magnesium content of 8.94375 mg/100g, *Piper umbellatum* is the most magnesium-rich crop in this study. Magnesium is critical for nerve and muscle function, bone strength, and energy production. Its presence in *Piper umbellatum* enhances the crop's value as a dietary supplement for addressing magnesium deficiencies, which can lead to muscle cramps, fatigue, and cardiovascular issues. This nutrient-rich crop could play a significant role in improving health outcomes in communities with limited access to magnesium-rich foods.

3.1.9 Calcium (Ca)

Piper umbellatum also ranks highest in calcium content (11.62 mg/100 g), Ver*nonia amygdalina* (5.5445 mg/100 g) making it an excellent source for supporting bone and dental health. Calcium is essential for skeletal strength and various metabolic processes, including muscle contraction and nerve signaling. The crop's high calcium content positions it as a natural alternative to dairy products, especially in populations with lactose intolerance or limited access to dairy. Promoting the consumption of *Piper umbellatum* could help address calcium deficiencies and enhance overall health in Sierra Leone.

3.2 Nutritional highlights and comparative strengths

Before presenting the mineral profiles of each crop, it is important to recognize that the values reported here reflect only the measured concentrations. Nutritional impact can vary due to factors such as bioavailability, antinutrients, dietary habits, cultural consumption patterns, and environmental or varietal differences across the localities where samples were collected factors that were beyond the scope of this study.

Based on overall mineral composition and nutritional purpose, *Piper umbellatum* is the most nutritionally versatile crop, addressing a broad spectrum of deficiencies. *Vernonia amygdalina*, *Ipomoea aquatica*, *Celosia argentea*, and *Luffa aegyptiaca* are significant contributors to iron, protein, potassium, and copper intake. Promoting the cultivation and dietary use of these NUFC can significantly enhance food security and public health outcomes in Sierra Leone.

3.2.1 Piper umbellatum, contains (Zinc, Phosphorus, Magnesium, Calcium, Sodium)

Piper umbellatum emerges as a highly versatile crop, leading in zinc (12.49 mg/100 g), phosphorus (103.1 mg/100 g), magnesium (8.94375 mg/100 g), calcium (11.62 mg/100 g),

and sodium (1.93 mg/100 g). These nutrients collectively support immune health, bone strength, cellular metabolism, and nerve function. Its exceptional mineral composition makes it a cornerstone crop for improving nutritional outcomes in resource-constrained communities.

3.2.2 Celosia argentea and Luffa aegyptiaca, contain (Iron)

These crops are exceptionally high in iron, with Celosia argentea at 15.226 mg/100 g and *Luffa aegyptiaca* at 12.962 mg/100 g. Their rich iron content makes them critical for addressing iron deficiency anemia, particularly in populations with limited access to iron-rich foods. Their integration into local diets can improve oxygen transport and energy production, directly combating fatigue, weakened immunity, and developmental delays.

3.2.3 Vernonia amygdalina and Ipomoea aquatica, contain (Copper and Potassium)

Vernonia amygdalina leads in copper (22.13 mg/100 g) and potassium (22.13 mg/100 g), with Ipomoea aquatica following closely in both categories (17.99 mg/100 g for copper and potassium). These crops are essential for maintaining cardiovascular health, fluid balance, and the development of red blood cells. Their high potassium content makes them especially beneficial in preventing hypertension and supporting nerve function.

3.2.4 Ipomoea aquatica, contain (Protein)

With the highest protein content at 7.00%, *Ipomoea aquatica* serves as a critical resource for combating protein-energy malnutrition. It provides an affordable plant-based protein source, particularly valuable for populations with limited access to animal proteins. This crop's contribution to dietary diversity and nutrition is particularly significant for vulnerable groups such as children and pregnant women.

3.2.5 Triumfetta cordiafolia, contain (Sodium)

Triumfetta cordiafolia, contain (Sodium)

While not leading in multiple nutrients, *Triumfetta cordiafolia* ranks second in sodium (1.32 mg/100 g). It can complement low-sodium diets and support electrolyte balance, contributing to nerve transmission and muscle function.

3.3 NUFC plants that have scarce information about their mineral composition

3.3.1 Triumfetta cordiafolia: [local name: Tey-gweh (Mende)]

The analysis of *Triumfetta cordiafolia* (from Table 2) reveals its potential as a valuable source of essential minerals for addressing nutritional deficiencies. With an iron content of 1.521 mg/100 g and zinc levels of 1.23 mg/100 g, this crop offers significant benefits for combating anemia and supporting immune health. The crop's phosphorus (7.09 mg/100 g) and calcium (2.52 mg/100 g) and magnesium (0.668 mg/100 g) content further highlight a useful supplement for general well-being. However, its low protein content (0.6125%) limits its ability to serve as a primary protein source. This indicates that while it is highly beneficial for enhancing mineral intake, its contribution to addressing protein-energy malnutrition is minimal.

Table 2 N/Protein and mineral composition of the selected NUFC [without research literature]

Sample Type (leaf)	Fe (mg/100 g)	Cu (mg/100 g)	Zn (mg/100 g)	Protein (%)	*P* (mg/100 g)	Na (mg/100 g)	K (mg/100 g)	Mg (mg/100 g)	Ca (mg/100 g)	Sources
Triumfetta cordiafolia	1.52	0.03	1.23	0.613	7.09	1.32	0.32	0.67	2.52	This study
Asystasia gangetica	7.72	0.15	0.55	1.51	10.54	0.56	2.87	1.26	3.40	This study
Celosia argentea	15.23	0.31	1.505	2.31	11.15	0.34	8.64	0.84	2.08	This study

3.3.2 Asystasia gangetica: [local name: Gbolohui (Mende)]

Asystasia gangetica (from Table 2) demonstrates substantial nutritional potential, particularly due to its high iron content of 7.72 mg/100 g. The crop also contains notable levels of potassium (2.87 mg/100 g), which is essential for maintaining fluid balance and supporting heart health. Its moderate zinc content (0.554 mg/100 g) adds to its benefits, as zinc plays a crucial role in immune function, wound healing, and potassium levels (2.87 mg/100 g), ellular metabolism. Phosphorus (10.54 mg/100 g) and calcium (3.4 mg/100 g) contribute to bone health, making the crop particularly valuable for populations with low dairy consumption. In addition to its mineral content, Asystasia gangetica offers a modest protein contribution of 1.505%.

3.3.3 Celosia argentea: [local name: Gbolohui (Mende)]

Celosia argentea (from Table 2) stands out as an exceptionally nutrient-dense crop, particularly due to its high iron content of 15.23 mg/100 g. Its zinc content of 1.51 mg/100 g, potassium (8.64 mg/100 g), phosphorus (11.15 mg/100 g), and calcium (2.08 mg/100 g) make it a well-rounded source of essential minerals. The protein content of Celosia argentea (2.345%) is higher than that of *Triumfetta cordiafolia* and *Asystasia gangetica*, making it a more robust option for addressing protein-energy deficiencies. The crop's high potassium content is particularly beneficial for populations at risk of hypertension, while its iron and zinc levels address key micronutrient deficiencies.

4 Discussion

Among the crops analyzed, *Piper umbellatum* emerged as the most nutritionally versatile, exhibiting high concentrations of zinc (12.49 mg/100 g), phosphorus (103.1 mg/100 g), magnesium (8.94 mg/100 g), calcium (11.62 mg/100 g), and sodium (1.93 mg/100 g). These minerals are essential for immune function, bone health, cellular metabolism, and electrolyte balance. The crop's broad-spectrum nutrient profile positions it as a strategic candidate for addressing multiple micronutrient deficiencies in Sierra Leone, particularly in rural communities with limited access to to diverse, nutrient-dense foods beyond the major staples." [7, 15].

Celosia argentea and *Luffa aegyptiaca* were identified as iron-rich crops, with concentrations of 15.23 mg/100 g and 12.96 mg/100 g, respectively. Iron deficiency anemia remains a significant public health concern in Sierra Leone, especially among women and children [34]. The integration of these crops into local diets could substantially improve iron intake, enhance oxygen transport, and reduce fatigue and developmental delays. Their cultivation also aligns with traditional knowledge systems, where leafy vegetables are commonly used in soups and sauces.

Vernonia amygdalina and *Ipomoea aquatica* showed exceptional levels of copper (22.13 mg/100 g and 17.99 mg/100 g) and potassium (22.13 mg/100 g and 17.99 mg/100 g). Copper supports red blood cell formation and neurological health, while potassium regulates blood pressure and fluid balance. These findings are particularly relevant given the rising incidence of hypertension and cardiovascular disease in Sierra Leone [38]. Promoting these crops could offer a low-cost, plant-based intervention to improve cardiovascular outcomes.

In terms of protein content, *Ipomoea aquatica* stood out with 7.00%, making it a valuable resource for combating protein-energy malnutrition. Protein is critical for growth,

tissue repair, and enzymatic functions. In low-income settings where animal protein is scarce, plant-based sources like Ipomoea aquatica offer an affordable alternative. Its adaptability to swampy environments and rapid growth cycle further enhances its viability for smallholder farmers.

The study also highlighted *Triumfetta cordiafolia* and *Asystasia gangetica* as crops with moderate mineral profiles but limited protein content. While *Triumfetta cordiafolia* ranked second in sodium (1.32 mg/100 g), its protein content was only 0.61%, suggesting its role may be more complementary than foundational in addressing malnutrition. *Asystasia gangetica*, with 7.72 mg/100 g iron and 2.87 mg/100 g potassium, offers targeted benefits for anemia and heart health, though its overall nutrient density is lower than other NUFCs.

Comparative analysis with global literature revealed significant variability in nutrient values, underscoring the influence of environmental conditions, genetic diversity, and methodological differences. For instance, iron content in *Luffa aegyptiaca* ranged from 12.96 mg/100 g (This study) to 41.05 mg/100 g [5], while zinc in *Justicia insularis* varied from 1.31 mg/100 g (this study) to 4972 mg/100 g [3]. These discrepancies highlight the need for standardized protocols and localized nutrient profiling to inform policy and dietary recommendations.

The findings also intersect with traditional knowledge systems. Many of the NUFCs studied such as *Vernonia amygdalina*, *Celosia argentea*, and *Triumfetta cordiafolia* are used in indigenous medicine and cultural rituals. Their nutritional validation reinforces the value of traditional food systems and supports the thesis goal of integrating NUFCs into sustainable food strategies. Moreover, the gendered nature of NUFC cultivation and processing often managed by women suggests that promoting these crops could enhance women's economic empowerment and household nutrition.

4.1 Comparative analysis of mineral composition of NUFC with other literature

The study presents the protein and mineral compositions of various neglected and underutilized food crops (NUFC) identified within the study area. These crops were analyzed for their nutritional content, including key minerals such as iron, zinc, calcium, potassium, magnesium, and phosphorus and their protein levels. The findings were compared to similar studies conducted globally (see Tables 3, 4, 5, 6 and 7 below), providing a broader context for understanding the nutritional potential of these crops. This comparison highlights both the unique nutrient profiles of the local NUFC and their alignment or deviation from values reported in other regions. By examining their nutrient compositions and comparing them with global benchmarks, the research offers valuable

Table 3 Comparative assessment of mineral composition of *Luffa Aegyptica* leaf

Mineral Composition	This Study	[5]
Fe (mg/100 g)	12.96	41.05
Cu (mg/100 g)	0.26	1.3
Zn (mg/100 g)	0.96	21.3
Protein (%)	0.41	4.9
P (mg/100 g)	10.54	-
Na (mg/100 g)	0.347	73.05
K (mg/100 g)	1.246	89.45
Mg (mg/100 g)	1.451	58.05
Ca (mg/100 g)	2.09	192.1

Table 4 Comparative assessment of the mineral composition of *Justicia insularis* leaf

Mineral Composition	This Study	Adeyemi and Babatunde, [3]
Fe (mg/100 g)	9.982	449.0
Cu (mg/100 g)	0.2	631
Zn (mg/100 g)	1.31	4972
Protein (%)	0.15	448
P (mg/100 g)	6.28	11
Na (mg/100 g)	0.8931	631.04
K (mg/100 g)	0.393	168
Mg (mg/100 g)	1.6065	76
Ca (mg/100 g)	5.0361	330

Table 5 Comparative assessment of the mineral composition of *Piper umbellatum* leaf

Mineral Composition	This study	[7]	[15]
Fe (mg/100 g)	2.474	1.322	70.56
Cu (mg/100 g)	0.49	0.433	3.958
Zn (mg/100 g)	12.49	0.092	3.187
Protein (%)	1.925	19.9	0.88
P (mg/100 g)	103.1	1.912	-
Na (mg/100 g)	1.93	0.092	46.43
K (mg/100 g)	24.93	5.817	336.619
Mg (mg/100 g)	8.94375	0.252	1370.99
Ca (mg/100 g)	11.62	2.38	71.3

Table 6 Comparative assessment of the mineral composition of *Vernonia amygdalina* leaf

Mineral Composition	This study	Usunobun and Okolie, [36]
Fe (mg/100 g)	0.76	32.20
Cu (mg/100 g)	22.13	19.50
Zn (mg/100 g)	1.76	14.23
Protein (%)	1.78	-
P (mg/100 g)	1.18	-
Na (mg/100 g)	0.77	48.31
K (mg/100 g)	22.13	62.8
Mg (mg/100 g)	5.54	681.36
Ca (mg/100 g)	7.69	1264.18

insights into their potential contribution to diversified diets and improved health outcomes. This comparative approach also aids in identifying crops with exceptional nutrient density, paving the way for targeted interventions and policy recommendations to promote their cultivation and consumption.

4.1.1 Luffa aegyptiaca: [local name: Bordorlyny (Mende)

The nutrient analysis of *Luffa aegyptiaca* (from Table 3) reveals considerable variability in its mineral and protein content across the different sources. Iron (Fe) content, for instance, ranges from 41.05 mg/100 g [5] to 12.962 mg/100 g in this study. This disparity could be attributed to environmental conditions such as genetics, soil composition, climatic differences, or differences in cultivation practices in the study area where the plants were collected. Soil richness in specific minerals may enhance the uptake of iron, which might explain the higher values in the [5] study. Zinc (Zn) content also shows significant variation between the two sources, [5] reporting higher values. Copper (Cu), on the other hand, shows more consistent but lower values across both sources, suggesting

Table 7 Comparative assessment of the mineral composition of *Ipomoea aquatica* leaf

Sources	Fe (mg/100 g)	Cu (mg/100 g)	Zn (mg/100 g)	Protein (%)	*P* (mg/100 g)	Na (mg/100 g)	K (mg/100 g)	Mg (mg/100 g)	Ca (mg/100 g)
This study	0.014	179.90	0.66	7.00	2.35	0.01	18.0099	3.32	4.56
Umar et al. [33]	210.30	0.36	2.47	1.01	109.30	135.00	5,458.33	301.60	416.70
Adedokun et al. [1]	3.20	5.30	4.10	4.10	86.00	159.00	444.00	52.00	163.00
Adelakun et al. [2]	19,757	673	6387	26.45	-	323	982	424.0	335.0
Chitsa et al. [10]	12.42	-	-	-	2519.93	-	2365.22	231.10	3445.05
Ndamitso et al. [20]	15.5	3.61	2.52	17.48	2.26	40.0	445.00	230.00	3.73
Saikia et al. [29]	14.67	-	0.81	19.55	102.31	23.58	129.41	0.96	16.57
Nyein [24]	**3217.00**	-	1.00	2.90	-	-	60.00	123.00	122.00
USDA [35]	1.67	0.02	0.18	2.60	39.0	113.00	312.00	71.00	77.00
Ngamsaeng et al. [21]	-	-	-	22.30	-	-	-	-	-
Maung et al. [17]	-	-	-	28.00	-	-	-	-	-
Samkol [30]	-	-	-	31.90	-	-	-	-	-
Shariff et al. [31]	-	-	-	2.87	-	-	-	-	-

a stable uptake mechanism or less environmental influence on this mineral. The protein content is higher, with values of 4.90% [5] and 0.411% in This Study.

4.1.2 Justicia insularis: [local name: Lamie (Mende)

The mineral content of *Justicia insularis* (from Table 4) exhibits a striking degree of variability, particularly for iron (Fe) and zinc (Zn). This Study reports iron at 9.982 mg/100 g, while *(*Adeyemi and Babatunde ([3]*)* report an extraordinary 449.0 mg/100 g. The discrepancy may reflect differences in growing location, as the plants analyzed in this study were collected from the Federal College of Education (Special), Oyo State, Nigeria, while environmental conditions, soil composition, and agro-ecological factors are known to strongly influence mineral accumulation in leafy vegetables. Such a vast difference raises questions about analytical methodologies and whether the higher value might be an outlier or reflective of unique environmental conditions. Similarly, zinc content varies from 1.309 mg/100 g from this study to 4,972.0 mg/100 g by Adeyemi and Babatunde [3]. Copper (Cu) levels also show extreme variability, with This Study reporting 0.20 mg/100 g compared to 631.0 mg/100 g from [3]. These differences suggest that either environmental factors (soil, climate, time of year, etc.), the age of the part sampled (young or mature), how it was preserved and stored, varietal differences, processing may significantly affect the assessment or there may be inconsistencies in measurement techniques between studies.

Protein content in *Justicia insularis* is another area of wide variation, with the from this study we reported a modest of 1.75% compared to an extraordinarily high 448% by [3]. The latter figure is likely due to a misrepresentation or error in reporting and emphasizes the importance of cross-referencing data from multiple studies. Despite these discrepancies, *Justicia insularis* demonstrates potential as a source of essential nutrients

and protein. Efforts to standardize cultivation and analysis methods are critical to fully understanding its nutritional value.

4.1.3 Piper umbellatum: [local name: Popodar (Mende)

The nutrient profile of *Piper umbellatum* (from Table 5) shows both consistency and variability across studies. Iron content, for instance, is reported as 2.47 mg/100 g (in this study) and 1.32 mg/100 g [7] while [15] report a significantly higher 70.56 mg/100 g. These discrepancies may stem from differences in environmental factors, such as soil mineral availability, or genetic variations between samples [8, 18, 40]. Zinc (Zn) content also varies widely, with this study reporting 12.49 mg/100 g, [7] reporting 0.09 mg/100 g, and [15] reporting 3.19 mg/100 g. Copper (Cu) levels are more consistent between this study (0.49 mg/100 g) and [7] (0.43 mg/100 g) but are notably higher in [15] findings (3.96 mg/100 g), suggesting possible location-specific enrichment.

Protein content, on the other hand, shows significant fluctuation, with This study reporting 1.93% and [7] 19.90%, while [15] report a low 0.88%. These variations might reflect differences in plant maturity, sample preparation, or analytical approaches. Despite these inconsistencies, *Piper umbellatum* remains a promising candidate for food and nutrition security due to its high potassium content, which supports cardiovascular health.

4.1.4 Vernonia amygdalina: [local name: Bitter leaf (Creole)

The nutrient composition of *Vernonia amygdalina* (from Table 6) is marked by notable variations, particularly in its mineral content. Iron levels, for instance, range from 0.765 mg/100 g (This study) to 32.2 mg/100 g [36]. This variation might be attributed to differences in soil conditions, environmental factors, or plant processing methods. Copper (Cu) content is another area of divergence, with this study reporting a high value of 22.130 mg/100 g compared to 19.50 mg/100 g by [36]. Zinc (Zn) levels are also inconsistent between studies, with values of 1.755 mg/100 g (this study) and 14.23 mg/100 g [36].

Protein content reported in this study is 17.75%, highlighting *Vernonia amygdalina* as a valuable dietary protein source. Although protein data is absent from the [36] study, the high mineral content supports the crop's nutritional importance. Given its nutrient density, *Vernonia amygdalina* could significantly address micronutrient deficiencies, particularly iron and protein-related malnutrition.

4.1.5 Ipomoea aquatica: [local name: Gogodee (Mende)

Ipomoea aquatica (from Table 7) demonstrates substantial variability in its nutrient content across multiple sources. Iron levels range from as low as 0. 014 mg/100 g from this study to an extraordinarily high 14.671 mg/100 g [29]. Such extreme values suggest either measurement inconsistencies or environmental factors affecting iron uptake. Copper (Cu) levels also vary widely, from 0.023 mg/100 g [35] to 17.99 mg/100 g from this study. Zinc (Zn) content shows inconsistency, with values from 0.655 mg/100 g (Njala) to 6387.0 mg/100 g [2], affirming its potential as a zinc source. Potassium (K) content shows inconsistency, with values from 179.9 mg/100 g from this study to 5,458.33 mg/100 g [33], and 129.408 mg/100 g [29]. These disparities highlight the need for more standardized and controlled studies to determine its nutrient profile accurately.

The least recorded protein composition in *Ipomoea aquatica* was by [33] with 1.008%, and the highest was 31.9% [30]. On the other hand, this study recorded 7.00% protein. This variability might reflect differences in plant maturity, growth conditions, or analysis methods. Despite these inconsistencies, *Ipomoea aquatica* remains a nutrient-dense crop, with high calcium, magnesium, and potassium levels contributing to its value as a dietary supplement. Given its adaptability to diverse conditions and potential for addressing nutrient deficiencies, further research should focus on optimizing cultivation practices and validating its nutrient composition. Such efforts could position Ipomoea aquatica as a cornerstone of food and nutrition security initiatives in Sierra Leone.

Although this study did not assess consumer acceptance, the nutritional impact of these crops ultimately depends on cultural preferences, taste, familiarity, and accessibility. These factors influence consumption patterns and should be examined in future studies to better understand how these nutrient-rich crops can be effectively integrated into local diets.

It is also important to note that common food preparation methods such as boiling, steaming, drying, fermentation, and frying can alter both mineral concentrations and antinutrient levels, thereby affecting the bioavailability of key nutrients. These aspects were beyond the scope of the present study but should be considered in future work to better translate compositional data into practical dietary recommendations.

The mineral profiles documented in this study align with national food and nutrition security objectives aimed at reducing micronutrient deficiencies and promoting the use of underutilized crops. These findings also contribute to global development priorities, particularly SDG 2 (Zero Hunger) and SDG 3 (Good Health and Well-Being), by supporting dietary diversification and improved micronutrient intake.

5 Conclusion

While country is agriculturally active but still continues to face persistent challenges in food and nutrition security, particularly due to limited dietary diversity and micronutrient deficiencies. Through standardized laboratory analysis, the research quantified essential minerals including iron, zinc, copper, calcium, magnesium, phosphorus, potassium, and sodium across multiple NUFC species, offering empirical evidence of their nutritional value.

The findings revealed that crops such as *Piper umbellatum*, *Celosia argentea*, *Ipomoea aquatica*, *Vernonia amygdalina*, and *Luffa aegyptiaca* possess rich and varied mineral profiles, with several species demonstrating concentrations that rival or exceed global benchmarks [25]. Notably, *Piper umbellatum* emerged as the most nutritionally versatile crop, exhibiting high levels of zinc, phosphorus, magnesium, calcium, and sodium. *Celosia argentea* and *Luffa aegyptiaca* were identified as iron-rich, while *Ipomoea aquatica* stood out for its protein content and cardiovascular-supporting potassium levels.

These results underscore the strategic potential of NUFCs in addressing micronutrient deficiencies endemic to rural Sierra Leone. Their integration into local diets could significantly improve nutritional outcomes, particularly among vulnerable populations with limited access to nutrient rich foods or animal-based sources. Moreover, the cultivation of these crops aligns with traditional knowledge systems and agro-ecological conditions, reinforcing their cultural relevance and adaptability.

By scientifically validating the mineral composition of NUFCs, this research lays the groundwork for their inclusion in national food security strategies, nutrition programming, and agricultural extension services. Promoting these crops not only diversifies the food system but also enhances resilience against climate variability and market dependency. Future research should explore bioavailability, seasonal variation, and consumer acceptability to further support the mainstreaming of NUFCs into Sierra Leone's sustainable development agenda.

5.1 Recommendations for future laboratory analysis

While the present study provides valuable insights into the mineral composition of selected NUFCs, several external factors may influence laboratory outcomes and should be considered in future research. Environmental conditions such as soil type, rainfall, and seasonality can significantly affect nutrient uptake and accumulation in plants. Similarly, variations in plant maturity and the specific part analyzed (e.g., leaf vs. seed) may lead to differences in mineral concentrations. Standardizing harvest stages and clearly documenting plant parts used will improve comparability across studies. Post-harvest handling, particularly drying temperature, exposure to sunlight, and ambient humidity can also impact nutrient stability, especially for heat-sensitive compounds.

Analytical precision depends on the sensitivity of the methods employed. While techniques such as AAS and flame photometry are widely accepted, cross-validation with more sensitive instruments (e.g., ICP-MS) and the use of certified reference materials are recommended to enhance reliability. Incorporating soil nutrient analysis alongside plant tissue profiling would provide a more holistic understanding of mineral dynamics. Finally, increasing biological replicates and sampling across multiple agroecological zones can strengthen statistical power and generalizability, especially for crops with localized cultivation patterns or emerging nutritional relevance.

Future research should include assessments of mineral bioavailability using appropriate in vitro or in vivo methods, repeated sampling across seasons to capture temporal variation in nutrient composition, and consumer acceptability studies to understand preferences and potential barriers to the adoption of these crops. These recommendations outline potential applications of our findings such as informing nutrition-sensitive agriculture programs, school feeding initiatives, and dietary diversification strategies but they do not alter the analytical results presented here. Instead, they highlight how the mineral composition data generated in this study can support future interventions and policy decisions.

Acknowledgements
The authors thank the students, Community Health Officers (CHOs), and Community Health Assistants (CHAs) from Njala University's Department of Community Health Sciences (2024/2025 academic year) for their support in data collection. The authors also acknowledge Mr. Kinie Jalieh of Kasewe Reserved Forest community and Momoh Sesay at the National Herbarium for their expertise in identifying the underutilized and neglected food crops.

Permissions for plant collection
Permission for the collection of plant materials was obtained from the Sierra Leone Ministry of Agriculture and Forestry, as well as relevant local authorities, including district forest officers and community leaders. Plant collection complied with national guidelines and institutional protocols, and no endangered or protected species were involved.

Author contributions
SF, RW, TSS, and SMTW contributed to the manuscript's writing, literature review and interpretation, and data analysis. All authors read and approved the final manuscript.

Funding
The authors received no specific funding for this work.

Data availability
The datasets generated during and/or analysed during the current study are available from the corresponding author on reasonable request.

Declarations

Ethics approval and consent to participate
Ethical approval for this study was obtained from the Directorate of Research and Development, Njala University, Sierra Leone. The study adhered to the principles of the Declaration of Helsinki. All participants were informed about the purpose and nature of the research, and written informed consent was obtained prior to participation.

Consent for publication
All participants provided consent for the publication of anonymized data.

Competing interests
The authors declare no competing interests.

Received: 27 September 2025 / Accepted: 5 May 2026
Published online: 26 May 2026

References

1. Adedokun MA, Ogundiran MA, Alatise SP. Preliminary Assessment of Water Spinach (Ipomoea Aquatica) and Morning Glory (Ipomoea Asarifolia) Leaves Meals as Non—Conventional Fish Feed Stuffs. Int J Zool Anim Biology. 2019;2(4):1–6. https://doi.org/10.23880/IZAB-16000164.
2. Adelakun K, Kehinde AS, Amall R, Ogundiwin D, Omotayo O. Nutritional and Phytochemical Quality of Some Tropical Aquatic Plants. Poult Fisheries Wildl Sci. 2016;04. https://doi.org/10.4172/2375-446X.1000164.
3. Adeyemi OT, Babatunde O. Chemical composition and antioxidant capacity of the leaf extract of Justicia insularis. Int J Phys Sci. 2014;9(20):454–8. https://doi.org/10.5897/IJPS2012.0013.
4. Alexiades MN, Sheldon JW. (1996). Selected guidelines for ethnobotanical research. Advances in Economic Botany, 306. https://agris.fao.org/search/en/providers/122535/records/65dde5d57c7033e84be8b92b
5. Ani E, Adekunle AA, Aboluwade JB, Ibrahim O. (2020). *EX-SITU Characterization of Luffa aegyptaica in Lagos State, Nigeria | Baghdad Science Journal* [Baghdad Science Journal]. https://bsj.uobaghdad.edu.iq/index.php/BSJ/article/view/3796
6. AOAC. (1990). AOAC (2000) Official Methods of Analysis. 15th Edition. *The Association of Official Analytical Chemists*. https://law.resource.org/pub/us/cfr/ibr/002/aoac.methods.1.1990.pdf
7. Baradum SR, Peters DE, Ohiri RC. (2021). PROXIMATE ANALYSIS, ANTI-NUTRIENT AND MINERAL COMPOSITIONS OF Piper umbellatum Linn LEAVES. J Biochem Int, 62–70. https://ikprress.org/index.php/JOBI/article/view/7313
8. Baxter, I., & Dilkes, B. P. (2012). Elemental profiles reflect plant adaptations to the environment. Science, 336(6089), 1661–1663. https://doi.org/10.1126/science.1219992
9. Bray RH, Kurtz LT, ORGANIC, AND AVAILABLE FORMS OF PHOSPHORUS IN SOILS. DETERMINATION OF TOTAL. Soil Sci. 1945;59(1):39. https://journals.lww.com/soilsci/citation/1945/01000/determination_of_total,_organic,_and_available.6.aspx.
10. Chitsa H, Mtaita T, Tabarira J. Nutrient content of water spinach (Ipomoea aquatica) under different harvesting stages and preservation methods in Zimbabwe. Int J Biol Chem Sci. 2014;8(3). https://doi.org/10.4314/ijbcs.v8i3.2.
11. Cotton, C. M. (1996). Ethnobotany: Principles and Applications | Wiley. Wiley. https://www.wiley.com/en-us/Ethnobotany%3A+Principles+and+Applications-p-9780471955375
12. FAO, European Union, CIRAD. Food Systems Profile—Sierra Leone. FAO; 2022. https://doi.org/10.4060/cb8792en. European Union; CIRAD.
13. Foday S, Wadsworth R, Sonda TS, Williams SM. b. Medicinal Value of Neglected and Underutilized Food Crops in Sierra Leone. Res J Med Plants. 2025;19(1):12–24. https://doi.org/10.3923/rjmp.2025.12.24
14. Foday S, Wadsworth R, Sonda TS, et al. Exploring the decline of neglected and underutilized food crops in Sierra Leone. Discov Agric. 2025;3:267. https://doi.org/10.1007/s44279-025-00442-x
15. Kalu O, Osuji V. Comparative Nutritional Studies of Lasianthera Africana and Piper Umbellatum Leaves. Nigeria J Life Sci. 2018;8:23–32.
16. Knez, M., Ranić, M., & Gurinović, M. (2024). Underutilized plants increase biodiversity, improve food and nutrition security, reduce malnutrition, and enhance human health and well-being. Let's put them back on the plate! Nutrition Reviews, 82(8), 1111–1124 https://doi.org/10.1093/nutrit/nuad103
17. Maung AT, Swe KH, Maw AA, Aung YL. Effects of supplementing water spinach (Ipomoea aquatica) to basal diet on growth performance and nutrients digestibility of broiler chickens. J Livestock Sci. 2020;11:77–84. https://doi.org/10.33259/JLivestSci.2020.77-84.
18. Marschner, P. (2012). Marschner's Mineral Nutrition of Higher Plants. Elsevier. https://doi.org/10.1016/C2009-0-63043-9
19. Martin, G. J. (with Internet Archive). (1995). Ethnobotany: A methods manual. London ; New York : Chapman & Hall http://archive.org/details/ethnobotanymetho0000mart_t8r1
20. Ndamitso M, Etsuyankpa B, Jacob J, Mathew J, Yanda Shaba E, Olisedeme K. The Nutritional Values and Functional Properties of Wild Ipomoea Aquatic (Water Spinach) Found in the Fadama Areas of Minna, Niger State, Nigeria. Acad Res Int. 2015;6:1–8.
21. Ngamsaeng A, Thy S, Preston TR. (2004). *Duckweed (Lemna minor) and water spinach (Ipomoea aquatica) as protein supplements for ducks fed broken rice as the basal diet*. Livestock Research for Rural Development. http://lrrd.cipav.org.co/lrrd16/3/amo16016.htm

22. Nhamo L, Paterson G, van der Walt M, Moeletsi M, Modi A, Kunz R, Chimonyo V, Masupha T, Mpandeli S, Liphadzi S, Molwantwa J, Mabhaudhi T. Optimal production areas of underutilized indigenous crops and their role under climate change: Focus on Bambara groundnut. Front Sustainable Food Syst. 2022;6. https://doi.org/10.3389/fsufs.2022.990213.
23. Nutraceuticals Group. (2025). *Nutraceuticals Group—Unit Converter*. Nutraceuticals Group. https://nutraceuticalsgroup.com/tools/unit-converter
24. Nyein WW. (2019). Studies on Some Nutritional Values of Water Spinach Ipomoea aquaticaForssk. *Myanmar Korea Conference Research Journal*, *3*(5). https://www.dagonuniversity.edu.mm/wp-content/uploads/2020/06/War-War-Nyein-1.pdf
25. Oboh, G. (2006). Nutrient and Antinutrient Composition of Condiments Produced from Some Fermented Underutilized Legumes. Journal of Food Biochemistry, 30(5), 579–588. https://doi.org/10.1111/j.1745-4514.2006.00083.x
26. OCHA. (2015a). *Sierra Leone: Bo District Profile (04 December 2015) | OCHA*. OCHA Sierra Leone. https://www.unocha.org/publications/report/sierra-leone/sierra-leone-bo-district-profile-04-december-2015
27. OCHA. (2015b). *Sierra Leone: Kenema District Profile (3 December 2015) | OCHA*. OCHA Sierra Leone. https://www.unocha.org/publications/report/sierra-leone/sierra-leone-kenema-district-profile-3-december-2015
28. Purba NH, Krishnaswamy K. Exploring the potentials of neglected underutilized crops (NUCs): an integrative review for developing a sustainable food system model. npj Sci Food. 2025;9:199. https://doi.org/10.1038/s41538-025-00554-0
29. Saikia K, Dey S, Hazarika SN, Handique GK, Thakur D, Handique AK. Chemical and biochemical characterization of Ipomoea aquatica: Genoprotective potential and inhibitory mechanism of its phytochemicals against α-amylase and α-glucosidase. Front Nutr. 2023. https://doi.org/10.3389/fnut.2023.1304903.
30. Samkol P. Water spinach (Ipomoea aquatica) as a feed resource for growing rabbits. Revista Computadorizada de Producción Porcina: Cent Livest Agric Dev. 2009;16(2):91–2. https://citeseerx.ist.psu.edu/document?repid=rep1&type=pdf&doi=2e8d7945799e8ebcd7ecf792efebbef4dc84560e.
31. Shariff AHM, Shapiai NA, Zakaria M, Akanda JMH, Noor NQIM, Yusop MHM, Razali UHM. (2019). Proximate compositions of Ipomea aquatic Forsk. (Leaf, petiole and stem) from Lubok Bungor, Jeli, Kelantan. *AIP Conference Proceedings*, *2155*(1), 020060. https://doi.org/10.1063/1.5125564
32. Soltanpour PN. Use of ammonium bicarbonate DTPA soil test to evaluate elemental availability and toxicity. Commun Soil Sci Plant Anal ; (United States). 1984;16:3. https://doi.org/10.1080/00103628509367607.
33. Umar KJ, Hassan LG, Dangoggo SM, Ladan MJ. Nutritional Composition of Water Spinach (Ipomoea aquatica Forsk.) Leaves. J Appl Sci. 2007;7(6):803–9. https://doi.org/10.3923/jas.2007.803.809.
34. UNICEF. (2017). *Nutrition | UNICEF Sierra Leone*. https://www.unicef.org/sierraleone/nutrition
35. USDA. USDA FoodData Central Food Details—Water convolvulus. U.S. DEPARTMENT OF AGRICULTURE; 2019. https://fdc.nal.usda.gov/food-details/169301/nutrients.
36. Usunobun U, Okolie NP. (2015). Phytochemical, trace and mineral composition of Vernonia Amygdalina leaves. Int J Biol Pharm Res. https://www.researchgate.net/publication/279529733_International_Journal_of_Biological_Pharmaceutical_Research_PHYTOCHEMICAL_TRACE_AND_MINERAL_COMPOSITION_OF_VERNONIA_AMYGDALINA_LEAVES
37. van Zonneveld M, Kindt R, McMullin S, Achigan-Dako EG, N'Danikou S, Hsieh W-H, Lin Y-R, Dawson IK. Forgotten food crops in sub-Saharan Africa for healthy diets in a changing climate. Proc Natl Acad Sci USA. 2023;120(14):e2205794120. https://doi.org/10.1073/pnas.2205794120.
38. WFP, & FAO. Hunger Hotspots. FAO–WFP early warnings on acute food insecurity: November 2024 to May 2025 outlook. WFP; FAO; 2024. https://doi.org/10.4060/cd2995en.
39. WFP, (World Food Programme). (2024, April 12). *Food Security Monitoring System Report—WFP Sierra Leone Country Office—February 2024 | World Food Programme*. World Food WFP, (World Food Programme), Programme. https://www.wfp.org/publications/food-security-monitoring-system-report-wfp-sierra-leone-country-office-february-2024
40. White, P. J., & Broadley, M. R. (2009). Biofortification of crops with seven mineral elements often lacking in human diets – iron, zinc, copper, calcium, magnesium, selenium and iodine. New Phytologist, 182(1), 49–84. https://doi.org/10.1111/j.1469-8137.2008.02738.x
41. World Bank. (2023). *Sierra Leone: Pathways to a Transformation of the Agri-Food Sector*. World Bank. https://documents1.worldbank.org/curated/en/099444004282527285/pdf/IDU-ebe7dceb-856f-4378-a45a-29837c72a70e.pdf

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.