



Back to the origins: biopesticides as promising alternatives to conventional agrochemicals

Ahmed A. A. Aioub · Soumya Ghosh · Ammar AL-Farga ·
Allah Nawaz Khan · Rabia Bibi · Abdullah M. Elwakeel · Asma Nawaz ·
Naglaa T. Sherif · Sohaila A. Elmasry · Esraa E. Ammar

Accepted: 15 April 2024 / Published online: 15 May 2024
© The Author(s) 2024

Abstract Presently, the world is using eco-friendly products to limit pollution in soil, air, water, and marine environments and to mitigate rapid climate change according to the sustainable development goals of the United Nations Development Programme. As a result, most countries attempt to produce environmentally friendly herbicides, fertilizers, and pesticides from plants, algae (e.g., *Cladophora glomerata*, *Laurencia pinnata*, *Plocamium cartilagineum*,

Polcamium spp.) or animal manure. Plants, such as *Anethum sowa*, *Thymus vulgaris*, *Foeniculum vulgare*, *Syzygium aromaticum*, *Pinus sylvestris*, *Citrus* spp., *Piper* spp. and *Mentha spicata*, are ecofriendly sources of essential oils, containing safe components, which can resist harmful pests. This review evaluates the common plants and algae used for extracting biopesticides, geographical distribution, target pests, mode of action, and commercial viability.

A. A. A. Aioub (✉)
Plant Protection Department, Faculty of Agriculture,
Zagazig University, Zagazig 44511, Egypt
e-mail: a.aioub@zu.edu.eg; ahmedaioub1991@gmail.com

S. Ghosh
Natural and Medical Sciences Research Center, University
of Nizwa, Nizwa 616, Oman

A. AL-Farga
Department of Biochemistry, Faculty of Science
University of Jeddah, Jeddah, Saudi Arabia

A. N. Khan
State Key Laboratory of Vegetation and Environmental
Change, Institute of Botany, Chinese Academy
of Sciences, University of Chinese Academy of Sciences,
Nanxincun, Xiangshan, Beijing, China

A. N. Khan
Department of Botany, University of Agriculture,
Faisalabad, Pakistan

R. Bibi
Centre of Excellence in Marine Biology, University
of Karachi, Karachi 75270, Pakistan

A. M. Elwakeel
Chemistry Department, Faculty of Science, Tanta
University, Tanta 31527, Egypt

A. Nawaz
Government College University, Faisalabad, Pakistan

N. T. Sherif
Faculty of Science, Tanta University, Tanta 31527, Egypt

S. A. Elmasry
Faculty of Science, Damanhur University,
Damanhur 5845171, Egypt

E. E. Ammar
School of Food and Biological Engineering, Jiangsu
University, Zhenjiang 212013, China

E. E. Ammar
Plant Ecology Sector, Botany Department, Faculty
of Science, Tanta University, Tanta 31527, Egypt

Keywords Biopesticides · Plants · Algae · Nano pesticides · Sustainable development

Introduction

The world's population is increasing tremendously and is expected to reach 8.5 billion by 2030. Conversely, food will decrease due to various reasons including harmful pests (Desa, 2011) and land for agriculture becoming less (Jhala et al., 2020). Annually, many crops are spoiled by an estimated 67,000 pests like fungi, weeds, bacteria, and insects (Agrios, 2005; Meena & Mishra, 2020). Using plants and algae for controlling pests is not new and is a traditional method (Choi et al., 2004). Biopesticides, originating from natural products (plant, algae, fungi) are eco-friendly, less toxic, and have less resistance build-up among insects and pathogens (Regnault-Roger et al., 2012). Biopesticides are effective because they have specific targets, and inhibit or kill harmful pests without negative effects on humans and beneficial organisms. Pests are a major factor in the declining yield of crops, which influence economies and feed populations (Sinha & Biswas, 2011).

Biopesticides are biodegradable and promote eco-friendly environments (Mazid et al., 2011). Safe plant and algae biopesticides are commercially available (Alam et al., 2002). In the United States of America, almost 200 biopesticides are used against harmful pests (Kumar & Singh, 2015), while in India 2.89% of the total market are biopesticides, which are increasingly used (Thakore, 2006). Biopesticides mainly derived from plants and algae have gained attention and publicity due to potentially healthier benefits in agriculture and for human health (Gupta & Dikshit, 2010). It possesses antifungal, antibacterial, antiviral, and insecticidal biopesticides (Akinbode & Ikotun, 2008; Hao et al., 2021; Harlapur et al., 2010; Yasmin et al., 2008).

Plant-based pesticides are commonly derived from herbs and plant-based pesticides are known as herbal pesticides (Pal et al., 2013). Almost 6000 plant species are used as insecticides (Dimetry et al., 1993). Different forms of algae are potential biopesticides. Microalgae are a microscopic group of algae that have bioactive compounds, which are effective against harmful organisms (Costa et al., 2019). Interest and awareness are increasing to use medicinal plants and

algae for controlling and inhibiting plant diseases caused by pests. For this purpose, almost 2400 plant species have been screened for potential phytoactivity (Khafagi & Dewedar, 2000).


In this review we will discuss both biopesticides active against plant pathogens and biopesticides active against insect pests.

Assessment of plants as biopesticide

Plant derived pesticides are obtained from leaves, roots, barks, seeds, fruits and bulbs. The plant parts are dried and bioactive compounds extracted (Chougule & Andoji, 2016). Different parts of plants are used for making biopesticides and extracts taken for assessing potential activity. (Ahmad et al., 2009). For example, cinnamon essential oil extract from cinnamon bark showed antifungal activity against *Phytophthora colocasiae* which causes acute decline, and up to 100% crop loss. The main component of the oil is cinnamaldehyde. At 0.625 mg/mL, the maximum inhibition of mycelial growth, zoospore germination, and sporulation of the fungus were observed, whereas leaf necrosis due to phytotoxicity was 100% at 1.25 mg/mL (Hong et al., 2021).

Essential oils can be used as biopesticide alternatives to monocrotophos. The eugenol derivative, 4-allyl-2-methoxy-1-(4-trifluoromethyl-benzyloxy)-benzene with a feeding deterrent index (FDI) of 54.14% and the thymol derivative, 2-isopropyl-4-methyl-2-(4-nitrobenzyl) oxy) benzene with an FDI of 53.88% are extracted from clove and thyme essential oils respectively. Both showed as antifeedant agents against the fourth instar larvae of *Spodoptera littoralis* through oral administration for 14 days. This study revealed that the ether compound was more effective than the ester compound regardless of the essential oil (Yan et al., 2021). Commercially available herbaceous plant *Nepeta cataria* extracts are effective repellents against mites, mosquitoes and cockroaches. The essential oil and hydrosol from *Cuminum cyminum* (cumin) seeds, characterized by γ -terpinene-7-al (34.95%), cumin aldehydes (26.48%), and α -terpinene-7-al (12.77%), were evaluated against second-stage juvenile (J2s) root-knot nematodes *Meloidogyne incognita* and *M. javanica* in the soil. The motility, hatching, and survival of J2s was affected after immersion in the essential oil at

Fig. 1 Bioactive chemical and compounds found in plants that has potential as bio-pesticide



1000 and 2000 $\mu\text{L/L}$. The lowest numbers of hatched J2s were recorded. More than 70% of paralyzed J2s were recorded after immersion for 48 h, which increased to more than 90% after 96 h (Pardavella et al., 2021).

Azadirachta indica L., *Thuja orientalis* L., *Nicotiana tabacum* L., *Nicotiana rustica* L., and *Melia azedarach* L. aqueous and powder extracts are used as alternatives to synthetic insecticides for the control of *C. maculatus* in stored pulses. Their toxicity was evaluated under specific conditions at concentrations of 0.50%, 1%, 1.5%, 2%, 2.5%, and 3%, and the following was the most effective biopesticide, *N. tabacum*, *A. indica*, and *M. azedarach*. The mortality rate of *C. maculatus* was significantly higher with all aqueous phytochemicals and all *N. tabacum* concentrations as well as powdered *A. indica*. *T. orientalis* recorded lower mortalities with the least number of phytochemicals (Akbar & Khan, 2021). Azadirachtin, a component of NeemAzal, is a plant-based insecticide, which is bio-efficient and biodegradable. Azadirachtin suppresses sugar beet moths (*Scrobipalpa ocellatella*) by reducing feeding, delays in larval and nymphal development, incomplete ecdysis, sterile eggs, deformed pupae and adults, and reduced fecundity. The optimal period to spray azadirachtin was 5–6 days after the moth deposited eggs, and concentrations of 2 and 2.5-ml L^{-1} had a substantial impact on *S. ocellatella* mortality. Thus, defined concentrations of a sugar beet moth insecticide applied at

the end of the growing season significantly reduced crop loss (Allahvaisi et al., 2021). The Asteraceae family of plants are used as botanical pesticides as they have effective chemicals for killing and controlling harmful fungi. Bioactive compounds include alkaloids, terpenoids and flavonoids (Ferreira et al., 2013). *Tarhonianthus camphorats* is capable of killing and inhibiting many bacteria (Sunder et al., 2011). Current trends are towards the extraction and use of plant materials for pest control (Villaverde et al., 2016) (Fig. 1).

Assessment of algae as eco-friendly biopesticides

Marine macro-algae (seaweed), are present on the seafloor attached to rocks or boulders, floating over the seawater, or sometimes drifting to shore. Seaweeds are found in the inter-tidal zones of tropical to temperate to cold waters (Coppejans et al., 2009). More than 10,000 species of macro-algae are known. Seaweeds are categorized into three main groups i.e., red (Rhodophyta), brown (Ochrophyta), and green (Chlorophyta), based on pigmentation, morphological, and anatomical features (Guiry & Guiry, 2017). Seaweeds have been used as food, forage, fertilizer, medicine, and other therapeutic applications benefiting human health (Blunden, 1993; Smit, 2004). Marine algae synthesize and emit bioactive compounds as a defence mechanism, which have been

identified and isolated (Perez et al., 2016). More than 2700 natural products have been reported. The secondary metabolites belong to terpenoids, alkaloids, phenols, esters, triglycerides, hydrocarbons, isoprenoids, alcohols, fatty alcohols, steroids, and waxy constituents. The largest variety of natural products has been reported from the red algae (Rhodophyta) mainly from genus *Laurencia* (Ioannou & Roussis, 2009). *Laurencia* is a complex genus comprising 383 species, of which 209 are taxonomically accepted (Cassano et al., 2019). The newly discovered *Laurencia karachiana* from Pakistan also possess insecticidal potential (Bibi et al., 2021) enriched with alcohols, fatty acids, sesquiterpenes, diterpenes, and steroids (Bibi et al., 2020).

The insecticidal potential of seaweeds is described (Table 1). Chemical composition analysis of seaweed extracts revealed that halogenated compounds such as triterpenoids and lipophilic compounds such as fatty acids are involved in insecticidal activity (Gressler et al., 2009). Oil-based compounds penetrate insects' bodies through cuticles and respiratory tracts and cause cellular and haemolymph acidosis, which leads to death (Sugiura et al., 2008). *Fucus spiralis* brown algae essential oil has an insecticidal effect against the Mediterranean fruit fly, *Ceratitis capitata* (Diptera: Tephritidae) (Gressler et al., 2009). Insecticidal efficiency was tested on the pupae and adults using different concentrations of *F. spiralis* essential oils. The

activity of the essential oils against adults was higher, with an LD₅₀ and an LD₉₀ of 0.239 and 2.467 ppm, respectively (Boutjagualt et al., 2022). Gas chromatography–mass spectrometry revealed the presence of borneol (13.13%), hentriacontane (6.13%), bornyl salicylate (5.48%), docosane (6.93%), octacosane (5.62%), heptacosane (7.22%), and limonene (4.9%), which made up the essential oil (Boutjagualt et al., 2022).

Widespread distribution of seaweeds can be utilized as an alternative approach to develop eco-friendly pesticides and repellents against mosquito species (Bibi et al., 2020). Further volatile compounds that are synthesized by seaweeds for their defence against herbivores and predators induce feeding deterrents and toxic effects against agricultural pests (Roviroso et al., 2013). These antifeedants can be isolated and applied against harmful sap sucking pests and stored grain commodities. Thus, huge biomass of seaweeds, enriched with potential bioactive compounds, is wasted every year along the shore and is the most alternative and safe source of biopesticides that can replace harmful synthetic pesticides.

Mechanism of action of biopesticides

Despite the variety of phytochemical pesticides, the mode of action of all compounds is unknown (Singh, 2017). Plants' bioactive compounds have different

Table 1 Efficacy of plants and algae as a biopesticide against pests

Plants and Algae source	Target pest	Reference
<i>Azadirachta indica</i>	<i>Aphis craccivora</i> , <i>Amrasca devastans</i> , <i>Sitobion avenae</i> <i>Bemisia tabaci</i> , and <i>Sciothrips cardamom</i> (Insects)	(Baidoo et al., 2012; Stanley et al., 2014)
<i>Allium sativum</i>	<i>Rhizoctonia solani</i> , <i>Colletotrichum</i> sp., <i>Staphylococcus aureus</i> , <i>Staphylococcus epidermidis</i> , (Fungi) and <i>Spodoptera littorals</i> (Insects)	(Baidoo & Mochiah, 2016; Suleiman & Abdallah, 2014)
<i>Euphorbia</i> spp.	<i>Escherichia coli</i> , <i>Salmonella typhi</i> (Bacteria), <i>Tribolium castaneum</i> (Insect) and <i>Aspergillus flavus</i> (Fungi)	(Abubakar, 2009; Gayathri & Ramesh, 2013; Mohammadi et al., 2016; Voukeng et al., 2017)
<i>Curcuma longa</i>	<i>Streptococcus pyogenes</i> , <i>Ralstonia solanacearum</i> , <i>Escherichia coli</i> , <i>Listeria monocytogenes</i> , and <i>Bacillus subtilis</i> (Bacteria)	(Abubakar, 2009; Gayathri & Ramesh, 2013; Mohammadi et al., 2016; Voukeng et al., 2017)
<i>Cinnamomum zeylanicum</i>	<i>Escherichia coli</i> (Bacteria), <i>Bursaphelenchus xylophilus</i> , <i>Meloidogyne</i> sp., (Nematode) and <i>Penicillium</i> spp. (Fungi)	(Nikkhah et al., 2017; Shirurkar & Wahegaonkar, 2012; Zhang et al., 2016)
<i>Zingiber officinale</i>	<i>Drosicha mangiferae</i> , <i>Trichoplusia binotalis</i> , and <i>Necrobial rufipes</i> (Insect)	(Al-Rahmah et al., 2013)

actions that may be physical or biological towards plant pathogens, insects, nematodes, and interactions depending on the type of compound and the pathogen (Grdiša and Kristina 2013; Aioub et al., 2022).

Mode of action of biopesticides

Plant extracts are lethal or toxic and affect the physiological activities of insect pests like preventing feeding and decreasing oviposition (Chengala & Singh, 2017). Antifeedant plant extracts prevent feeding by making the meal taste bad and the larvae die of starvation. Examples include *M. alternifolia* oil on *Helicoverpa armigera* (Liao et al., 2017); *Gliricidium sepium* methanol extract against coffee Mealybugs (*Planococcus citri* Risso) (Nukmal et al., 2017); and *Polyalthia longifolia* (methanol extract), *Polyalthia longifolia* (petroleum ether), *Limonia acidissima* (methanol) against *Spodoptera litura* (Arora, 2017). The eco-friendly essential oil of *Ageratina adenophora* has an antifeedant activity of 87.46% against *Plutella xylostella* (L.) (Mayanglambam et al., 2022).

Botanical pesticides have a toxic effect on insects infesting stored products by poisoning the mitochondrion, resulting in electron transport chain disruption and energy production inhibition (Stevenson et al., 2017). The crude phenolic compounds (methanolic extract) of *Acorus calamus* poisons the tobacco cutworm (*Spodoptera litura*) larvae and affect the hormonal balance by decreasing carboxylesterase, which affects the neurotransmitter acetylcholine (Wiwattanawanichakun et al., 2022). Essential oils from *Mentha arvensis* are highly toxic and penetrate the thick, hard exoskeleton, leading to disorders in the biological systems of *Sitophilus granaries* (Renoz et al., 2022). *Thaumastocoris peregrinus* is controlled by toxic essential oil from *Eugenia uniflora* (L.), which affects the hatching and increases the mortality of adults and nymphs (Stenger et al., 2021).

Botanical insecticides harm the pathogen by decreasing or inhibiting its stages of development (Isman, 2017). Essential oils of *Teucrium quadrifarium* and *Boenninghausenia albiflora* have growth inhibitor properties against *Spilarctia obliqua*, causing morphological and physiological effects, leading to a decrease in larval and pupal weight and increased mortality (Tandon & Mittal, 2018). Plant extracts from *Lantana camara*, *Catharanthus roseus*, *Coleus amboinicus*, *Curcuma longa*, and *Alpinia pyramidata* provide

toxicity, growth regulator properties, and antifeedant activity towards *Plutella xylostella* Linn diamondback moth (2nd larval instar) (Javier et al., 2019).

Extracted plant metabolites like alcohols, terpenes, alkaloids, phenols and tannins inhibit fungal growth by inhibiting spore germination and development of mycelia, or by degenerating hypha and mycelia (Yoon et al., 2013). The pathogenic fungus *Valsa mali* was affected by wood tar (EC₅₀ ranged from 69.54 to 92.81 µg/mL), leading to hyphal breaks, increased cell membrane permeability, and unbalanced enzyme production related to the decrease in pectinase and related genes. (Chen et al., 2022). A series of Michael-type amino derivatives were prepared from xanthatin, a secondary metabolite extract from *Xanthium strumarium*, with antifungal activity and inhibition of mycelium growth on *C. mandshurica* and spore germination against *F. solani* (Zhi et al., 2022). *Uvaria grandiflora* extract (zeylenone) inhibited growth of cucumber powdery mildew, and sporangium formation of *Phytophthora capsica* by destroying the cell wall structure and energy metabolism. Also, zeylenone was toxic to normal cell lines (He et al., 2021).

Similarly, botanical pesticide exposure affects bacterial growth especially Gram-negative organisms through its effect on cellular metabolism like protein synthesis (Djeussi et al., 2013). Pesticide exposure causes microbial death by increasing plasma membrane permeability (Khan et al., 2009). For example, hydroalcoholic extract from *Larrea tridentata* provides bactericidal activity to multidrug-resistant bacteria like *Pseudomonas syringae*, *Michiganensis*, *Clavibacter michiganensis* sbsp., especially *Xanthomonas campestris* (Morales-ubaldo et al., 2021). Parthenolide from *Magnolia grandiflora* Linn causes cell death in *Xanthomonas oryzae* pv. *oryzae* (Xoo) by accumulation of reactive oxygen species, decreased glutathione reductase and increased glutathione peroxidase activity (Xu et al., 2018).

Plant extracts with nematocidal activity inhibits egg hatching and reduces nematode populations (Abd El-Aal et al., 2021; Eldeeb et al., 2022). For example, methanolic extracts of *Anaphalis virgata* and *Anaphalis nepalensis* were tested against root-knot nematodes (Ismail et al., 2021). Extract of *W. indica* roots provide high nematocidal activity towards J2 of *M. incognita* by three isolated compounds i.e., waltherione C, 5'-methoxywaltherione A and waltherione A (Jang et al., 2019).

Some phytochemicals inhibit viral infection by inducing host resistance (Montanha & Moellerke, 2004), inhibiting virus penetration and virus replication (Rajasekaran et al., 2013). Benzosuberene compound extracted from *Cedrus deodara* oils inhibit self-assembly of tobacco mosaic virus (Sharma et al., 2021). The liberation of nucleic acids of watermelon mosaic virus is blocked by *Thuja orientalis* extract, leading to inhibition of multiplication (Elbeshehy et al., 2015).

Nano-biopesticides

Nanopesticides are nanostructures with two to three dimensions between 1 to 200 nm, used to carry agro-chemical ingredients. Nano-biopesticides are appealing because of their small size, high surface-area-to-volume ratio, stability, increased effectiveness, improved solubility, mobility, and importantly low toxicity (Fig. 2) (Lade et al., 2019). Toxins released into the food chain by chemical pesticides sprayed directly on plants can cause environmental problems.

Pesticides that contain nanoparticle formulations, such as micelles and nano-composites, lower the risk of environmental and health problems (Aioub et al., 2022). Clay-based nanotubes similarly transmit chemical insecticides (Sasson et al., 2007). Nano-biopesticides, like nano-fertilizers, are packaged in carriers that allow for the controlled release of active ingredients to achieve the intended effects in each environment.

The addition of nano-biopesticides to biopolymers improves stiffness, penetrability, crystallinity, thermal stability, solubility, and biodegradability (Manjunatha et al., 2016).

Nano-biopesticides comprising nano-materials that are applied to soils, results in the establishment of mutualistic bacteria that boost plant growth (Lu et al., 2010). Toxicity can be caused by silver-based nanoparticle coatings, which can be reversed by biocompatible coatings, improving the likelihood of seed germination in plants. Nano-emulsions, nano-encapsulates, nano-containers, and nano-cages have recently been described as nano-pesticide delivery strategies for plant protection with




Fig. 2 The importance of nanotechnology for the formulation of nano-based bio-pesticides. This figure is reproduced from Lade et al. (2019)

various capabilities (Bouwmeester et al., 2009; Hazafa et al., 2022). Cationic polymers may adhere to polyanionic surfaces of bacteria, rupture cell membranes, and kill pests. Biopesticides, such as nano-biopesticides, can be used on plants to reduce microbial resistance, but chemicals applied directly to plants are unable to suppress a wide spectrum of bacterial growth (Li et al., 2019). Long amino acid chains including tertiary ammonium groups can be found in nanoparticles. Depending on their structure, these groups may combat a variety of pests and ailments including bacteria (Hazafa et al., 2022; Ng et al., 2014), because of their strong activity in a range of environmental and chemical conditions. Many polymers with this property have been discovered and studied. For example, amphiphilic copolymers, functionalized cationic polycarbonates, poly(amidoamines), polyethylenimine, poly (methyl methacrylates), amino celluloses and chlorinated cellulose acetates are available commercially (Ng et al., 2014).

Plants containing nano-emulsions of essential oils can be employed to reduce insect larval infections. Nano-emulsions from *Tagetes minuta*, *Ageratum conyzoides*, and *Achillea fragrantissima* are used to inhibit the growth of *Callosobruchus maculatus*. The nano-emulsions can be used in fumigation to kill or restrict the formation of eggs and larvae, with treatment ranging from 16.1–40.5 $\mu\text{L/L}$ and 4.5–243 $\mu\text{L/L}$, respectively (Nenaah et al., 2015). Encapsulation of essential oils can be accomplished with nanoparticles made of liposomes and solid lipids. Inverse gelation and oil emulsion can be used to encapsulate essential oils (Lopez et al., 2012). Encapsulation in liposomes is beneficial against bacteria because the cell membrane is protected from the impact of essential oils (Sherry et al., 2013). However, many forms of nanoparticles are employed for pest control. Microparticles are also employed to keep the effects of essential oils stable. Encapsulation of carvacrol with the cell wall of *Saccharomyces cerevisiae* was conducted with a diameter of 0.5 μm to control the larvae of *Rhipicephalus microplus* with LC_{50} of 0.71 mg/mL. *Saccharomyces cerevisiae* cell wall appears to be beneficial in preserving the low volatility of encapsulated carvacrol, which retains acaricidal action for up to 60 h (Da Silva et al., 2017).

Commercial applications of biopesticides derived from plants and algae

Commercially made pesticides derived from higher plants and algae are biodegradable and eco-friendly. Compounds extracted from biological origins contain chemicals like terpenes, alkaloids, quinines, aldehydes, amino acids, saccharides, and flavonoids. These compounds are ecologically important as they are used commercially for antifeedant, nematocide, insecticide and fungicide products (Souto et al., 2021). From 1939 to 1962, the world focused on synthetic pesticides like 1,1,1-trichloro-2-propanol and 2-bis(4-chlorophenyl)-ethane (Jarman & Ballschmiter, 2012). As these chemicals are harmful and remain in contaminated soil, water and air for long periods and have adverse environmental effects (Multigner et al., 2010).

Formation of commercial biopesticides with algae is simple and low cost, as different varieties of microalgae can be obtained through wastewater (Ranglová et al., 2021). Duckweed (*Lemna minor*), water hyacinth (*Eichhornia crassipes*) and filamentous algae (*Lyngbya wollei*) produce allelopathic compounds that have potential commercial chemical application to inhibit the growth of pests (Fu et al., 2020). Biopesticides are evaluated and checked with the same regulations and rules as synthetic pesticides. Products from plants and algae are required to be prepared in safe environments according to strict guidelines before registration (Wang et al., 2000). It is believed that the commercial market of biopesticides will completely replace the market of synthetic pesticides by 2050 (Damalas, 2018) (Table 2).

Conclusion and future prospective

Synthetic pesticides' quick action makes them the first line of treatment, but their large-scale use has resulted in environmental hazards and the development of resistance. Therefore, environmentally safe plant products with fungicidal and insecticidal properties are a potential alternative. They can be as effective as conventional pesticides when applied as part of an integrated pest management strategy, especially for crops such as fruits, vegetables, nuts, and flowers. Application of chemical pesticides leads to the destruction of beneficial microorganisms and residues may accumulate. Bioactive compounds present in plants

Table 2 Insecticidal activities of marine seaweeds

Marine seaweeds	Extracts Compounds	Pests	Results	Ref
<i>Lyngbya wollei</i>	-	<i>Escherichia coli</i> , <i>Bacillus subtilis</i> , <i>Salmonella typhi</i>	Inhibition of growth rate	(Fu et al., 2020; Kumar & Singh, 2015)
Micro and macro algae spp.	-	<i>Stiobion avenae</i> , <i>Staphylococcus epidermidis</i>	Inhibition of growth rate	(Asimakis et al., 2022)
<i>Cladophora glomerata</i>	Saturated/unsaturated fatty acids	Mosquito, <i>Aedes triseriatus</i>	Larvicidal (LC ₅₀ 3 – 14 ppm)	(LaLonde et al., 1979)
<i>Laurencia pinnata</i>	Isolaurepinnacin	Azuki bean beetle, <i>Callosobruchus chinensis</i>	97% mortality after 24 h	(Fukuzawa & Masamune, 1981)
<i>L. pinnata</i>	Laurepinnacin	Cabbage army worm, <i>Manestra brassicae</i>	Larvicidal	(Fukuzawa & Masamune, 1981)
<i>Plocamium cartilagineum</i>	Monoterpene, Violacene	Tobacco hornworm, <i>Manduca sexta</i>	Larvicidal	(Crews et al., 1984)
<i>Chondria armata</i>	Domoic acid	American cockroach, <i>Periplaneta americana</i> ,	100% mortality at 1 µg/g,	(Maeda et al., 1984)
		House fly, <i>Musca domestica</i>	40% adult mortality at 0.1 µg / insect	
		German cockroach, <i>Blattella germanica</i>	60% adult mortality at 0.6 µg / insect	
<i>Digenea simplex</i>	Kainic acid	American cockroach, <i>P. americana</i>	100% adult mortality at 100 µg/ g	(Maeda et al., 1984)
<i>C. armata</i>	Isodomic acid-B	American cockroach, <i>P. americana</i>	Strong insecticidal activity	(Maeda et al., 1984)
<i>Polcamium</i> sp.	γ- BHC	Mosquito, <i>Anopheles gambiae</i>	0.0096 µg/mL	(Watanabe et al., 1990)
<i>Polcamium</i> sp.	γ- BHC and telfairine	German cockroach, <i>B. germanica</i>	Excitatory effect on central nervous system	(Watanabe et al., 1990)
<i>Caulerpa scalpelliformis</i>	2:8 Methanol acetone	Mosquito, <i>Ae. aegypti</i>	LC ₅₀ 15.85 mg/L	(Thangam & Kathiresan, 1991)
<i>Plocamium telairiae</i>	Aplysia terpenoid A and telfairine	Mosquito, <i>An. Gambiae</i>	Neurotoxic effect, convulsion, death	(Bloomquist, 1993)
<i>Caulerpa racemose</i>	Petroleum ether-acetone	Mosquito, <i>Ae. aegypti</i> , <i>Cx. quinquefasciatus</i>	LC ₅₀ < 100 mg/L	(Thangam & Kathiresan, 1996)
<i>C. peltata</i>	13-Epilaurencienyne	Ant	Knockdown effect	(Iliopoulou et al., 2002)
<i>Laurencia obtusa</i>	Methanol	Red flour beetle, <i>T. castaneum</i>	40% mortality after 24 h	(Rizvi & Shameel, 2006)
<i>Jolya laminarioides</i>	Monoterpene	Cereal aphid, <i>Schizaphis graminum</i>	92% adult mortality at 100 ppm after 48 h	(Argandoña et al., 2000)
<i>P. cartilagineum</i>	Violacene	Aphids <i>Myzus persicae</i> and <i>Ropalosiphum padi</i>	Deterrent	(Argandoña et al., 2002)
<i>P. cartilagineum</i>	Polyhalogenated monoterpenes	Mosquito, <i>Culex</i> sp	60.2% mortality at 6 mg/mL	(Selvin & Lipton, 2004)
<i>Ulva fasciata</i>	1:1 Dichloromethane- methanol	Mosquito, <i>Cx. quinquefasciatus</i>	62.2% mortality at 100 µg/mL	(Manilal et al., 2009)
<i>Acrosiphonia orientalis</i>	1:1 Dichloromethane-methanol	Mosquito, <i>Cx. pipiens</i>	Second instar (LC ₅₀ 1.42 ppm)	(Alarif et al., 2010)
<i>C. racemose</i>	Alkaloid caulerpin	Mosquito, <i>Cx. pipiens</i>		

Table 2 (continued)

Marine seaweeds	Extracts Compounds	Pests	Results	Ref
<i>C. scalpelliformis</i>	Chloroform	Cotton bug, <i>Dysdercus cingulatus</i>	(LC ₅₀ 256 µg/L)	(Rajesh et al., 2011)
<i>Laurencia papillosa</i>	Acetogenin	Red flour beetle, <i>Tribolium confusum</i>	LD ₅₀ 0.21% after 6 h	(Abou-elnaga et al., 2011)
<i>C. prolifera</i> , <i>C. serrulata</i> <i>Padina pavonica</i> , <i>Cystoseira myrica</i> , <i>Nitophyllum punctatum</i>	Dried ground sample	Mosquito, <i>Culex pipiens</i>	Prolonged larval duration	(Elbanna & Hegazi, 2011)
<i>Padina pavonica</i>	Chloroform, benzene	Red cotton bug, <i>D. cingulatus</i>	Prolonged nymphal duration	(Sahayaraj & Kalidas, 2011)
<i>Lobophora variegata</i>		Mosquito sp., <i>Ae. Aegypti</i> , <i>Cx. quinquefasciatus</i>	LC ₅₀ 70.3 µg/ml, 79.4 µg/ml	(Kumar & Sujith, 2011)
<i>Sargassum wightii</i>	Methanol	Mosquito, <i>An. sudaicus</i>	Prolonged developmental stages, reduced adult emergence	(Kumar et al., 2012)
<i>Ulva fasciata</i> , <i>U. lactuca</i>	Methanol, chloroform	Red cotton bug <i>D. cingulatus</i>	Reduced longevity, fecundity, eggs hatchability, whole body weight	(Asha et al., 2012)
<i>C. scalpelliformis</i>	Chloroform	<i>Spodoptera litura</i> Red cotton bug <i>D. cingulatus</i>	Repellent effects with black colouration in insects body	(Kombiah & Sahayaraj, 2012)
<i>Sargassum tenerrium</i>	Benzene	Cotton Pest, <i>D. cingulatus</i>	Anti-insect activity (LC ₅₀ 0.009%)	(Sahayaraj & Jeeva, 2012)
<i>Laurencia dendroidea</i>	Sesquiterpene, Elatol	Mosquito, <i>Ae. aegypti</i>	LC ₅₀ 10.7 ppm	(Bianco et al., 2013)
<i>C. racemose</i>	Ethanol	Mosquito, <i>Ae. Aegypti</i> , Mosquito, <i>Cx. quinquefasciatus</i>	(LC ₅₀ 0.05 µg/mL), (LC ₅₀ 0.05 µg/mL), (LC ₅₀ 0.06 µg/mL)	(Ali et al., 2011)
<i>P. cartilagineum</i>	Polyhalogenated monoterpenes	Mosquito, <i>An. stephensi</i> Tobacco budworm, <i>Heliothis virescens</i> ,	Anti-insect activity	(Rovirosa et al., 2013)
<i>Bryopsis pennata</i>	Chloroform	Mosquito, <i>Ae. aegypti</i>	Larvicidal (LC ₅₀ 82.5 µg/mL) with morphological abnormalities	(Yu et al., 2015)
<i>Jania rubens</i>	Hexane	Mosquito, <i>Ae. aegypti</i>	Larvicidal (60% mortality after 24 h at 40 mg/ml)	(Jagadeesan et al., 2015)
<i>L. obtusa</i>	halogenated sesquiterpene (-)-elatol and (+)-obtusol	Mosquito, <i>Ae. aegypti</i>	Larvicidal with reported destruction in the intestinal epithelium	(Salvador-Neto et al., 2016)
<i>B. pennata</i>	Methanol	Mosquito, <i>Ae. aegypti</i> (L) <i>Ae. albopictus</i>	Larvicidal LC ₅₀ 156 µg/ml LC ₅₀ 177 µg/ml,	(Ahmad et al., 2016)
		Mosquito, <i>Ae. aegypti</i> (L) <i>Ae. albopictus</i>	Adulticidal LC ₅₀ 17.5 µg/mL LC ₅₀ 35.4 µg/mL	
<i>P. cartilagineum</i>	Mortensene, Dibromomertensene, dihydromertensene	Tomato moth, <i>Tuta absoluta</i> , Cereal aphid, <i>Schizaphis graminum</i>	Larvicidal, adulticidal	(Argandoña et al., 2000)

Table 2 (continued)

Marine seaweeds	Extracts Compounds	Pests	Results	Ref
<i>L. brandenii</i>	Fraction	Rice weevil, <i>Sitophilus, Oryzae</i>	LD ₅₀ 3.7 mg/cm ²	(Manilal et al., 2011)
<i>Dictyota dichotoma</i>	3:1 Ethanol and water	<i>Ae. aegypti</i>	LC ₅₀ 0.068 mg/mL	(Beula et al., 2011)
<i>L. papillosa</i>	Acetogenin, (12E)-cis maneo-nene-E	Mosquito, <i>Culex pipiens pallens</i>	Second instar LC ₅₀ 30.7 µg/mL	(Abou-elnaga et al., 2011)
<i>Chondria dasyphylla</i>	Ethyl acetate fraction	Mosquito, <i>An. stephensi</i>	LC ₅₀ 10.6 µg/mL	(Khanavi et al., 2011)
<i>Padina minor</i>	95% Ethanol	Mosquito, <i>Ae. aegypti</i>	LC ₅₀ 50.8 mg/mL	(Bantoto & Dy, 2013)
<i>Gracilaria coricata</i>	Ethanol	Mosquito, <i>Ae. Aegypti</i>	LC ₅₀ 0.08 µg/mL	(Ali et al., 2013)
		Mosquito, <i>Cx. quinquefasciatus</i>	LC ₅₀ 0.09 µg /mL	
		<i>An. stephensi</i>	LC ₅₀ 0.09 µg/mL	
Seaweeds	Extracts	Mosquito, <i>Aedes aegypti</i> (L)	Larvicidal	(Hira et al., 2017)
<i>Turbinaria ornate</i>	Hexane	Mosquito, <i>Aedes aegypti</i> (L)	Larvicidal LC ₅₀ 1.38 µg/mL	(Sowmiya et al., 2017)
<i>L. nidifica</i>	Laurinterol	Maize weevil, <i>Sitophilus zeamais</i> , Termites	Repellent	(Ishii et al., 2017)
<i>S. cristaeifolium, C. Agardh</i>	Methanol	Tobacco caterpillar, <i>Spodoptera litura</i> Fab	Antifeedant, larvicidal, growth inhibitor	(Gowthish et al., 2018)
<i>L. intricata</i>	(+)-Cyclocolorenone	Maize weevil, <i>S. zeamais</i> , Termites	Repellent	(Ishii et al., 2019)
<i>S. polycystum</i>	Silver nanoparticles	Mosquito, <i>Aedes aegypti</i>	LC50 0.03 µg/mL after 72 h	(Vinoth et al., 2019)
		<i>An. stephensi</i>	LC50 1.15 µg/mL	
		<i>Cx. quinquefasciatus</i>	(0.23 µg/mL)	
<i>C. veravalensis</i>	Tetradecanoic acid, 10, 13-dimethyl-, methyl ester	Cotton pest, <i>D. cingulatus</i>	Growth, development and oviposition	(Sahayaraj et al., 2019)
<i>Gracilaria filiformis</i>	Fraction polysaccharide	Mosquito, <i>An. stephensi</i>	LC ₅₀ 0.23 µg/mL	(Venkatesan et al., 2019)
<i>Turbinaria conoides</i>	Fraction polysaccharide	Mosquito, <i>Ae. Aegypti</i>	LC ₅₀ 0.17 µg/mL	(Venkatesan et al., 2019)
<i>Enteromorpha compressa</i>	Fraction polysaccharide	Mosquito, <i>Cx. quinquefasciatus</i>	LC ₅₀ 0.1117 µg/mL	(Venkatesan et al., 2019)
<i>Gracilaria edulis</i>	Methanol	Tobacco cutworm, <i>Spodoptera litura</i>	53% larval mortality after 72 h with growth inhibiting effects	(Gowthish & Kannan, 2019)
<i>Jania rubens</i>	Hexane	Green peach aphid, <i>Myzus persicae</i>	Antifeedant	(Bibi et al., 2020)
		Mosquito, <i>Ae. aegypti</i> L	Neurotoxic (EC ₅₀ 37.4 µg/mL), Larvicidal (LC ₅₀ 32 µg/mL), 24 h	(Bibi et al., 2020)
<i>Laurenciai karachiana</i>	Hexane	Mosquito, <i>Ae. aegypti</i> L	Larvicidal LC ₅₀ 89 µg/mL, 24 h	(Bibi et al., 2020)
<i>Asparagopsis taxiformis</i>	Dichloromethane	Cowpea weevil, <i>Callosobruchus maculatus</i>	LC ₅₀ 1.14 after 24 h, reduced oviposition	(Bibi et al., 2021)
<i>Gracilaria foliifera</i>	Hexane	Mosquito, <i>Ae. aegypti</i> L	Larvicidal LC ₅₀ 76.8 µg/mL, 24 h	(Bibi et al., 2021)

Table 2 (continued)

Marine seaweeds	Extracts Compounds	Pests	Results	Ref
<i>Chaetomorpha antennina</i>	Isolated fraction 5 enriched with methyl ester of chloroacetic acid	Tobacco cutworm, <i>S. litura</i>	LC ₅₀ 38.73 ppm, reduce haemocyte count (69.8%) and some major defence enzyme	(Chanthini et al., 2021)
<i>Lobophora variegata</i>	Silver nanoparticle	Tobacco cutworm, <i>S. litura</i>	Affect development, moulting, morphology	(Kitherian et al., 2021)

and algae have pesticidal effects. Biopesticides derived from them are eco-friendly, safe and beneficial for the environment. Therefore, development of eco-friendly plant and algal-based pesticides is needed for pest management with minimal environmental impact. Biopesticides along with fewer application constraints, enhanced resistance management potential, and environmental safety make them more efficient. This supports the United Nations Development Program's 2030–2050 objectives, with the preservation of the global climate, which protects life on land and in water. Thus, a clean environment is secured where with adequate food for every human being, a reduction in poverty, and enhanced scientific and cultural development.

Acknowledgements The authors are thankful to Miss/ Yathreb M. Hashim for her preliminary literature searches about the mode of action of biopesticides, and Miss/ Nouran A. El-shershaby for paraphrasing some parts of the preliminary manuscript.

Authors Contributions Ahmed A.A. Aioub: Collected and summarized the results and revise the final manuscript.

Soumya Ghosh: participated in writing the manuscript. The authors read and approved the final manuscript.

Ammar AL-Farga: Collected and summarized the results and revise the final manuscript.

Abdullah M. Elwakeele; conducted a systematic literature review and wrote the primary manuscript.

Allah Nawaz Khan: conducted a systematic literature review and wrote the primary manuscript.

Rabia Bibi: conducted a systematic literature review.

Soumya Ghosh: participated in writing the manuscript. The authors read and approved the final manuscript.

Asma Nawaz: participated in writing the manuscript. The authors read and approved the final manuscript.

Naglaa T. Sherif: participated in writing the manuscript. The authors read and approved the final manuscript.

Sohaila A. Elmasry: participated in writing the manuscript and References.

Esraa E. Ammar: Collected and summarized the results, design graphics, and revise the final manuscript.

Funding Open access funding provided by The Science, Technology & Innovation Funding Authority (STDF) in cooperation with The Egyptian Knowledge Bank (EKB).

Data availability Data available on request.

Declarations

Ethical approval Not applicable.

Consent to participate All authors are consent to participate.

Consent to publish All authors are consent to publish.

Competing interests There are no conflict of interest.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Abd El-Aal, E. M., Shahan, M., Sayed, S., et al. (2021). In vivo and in vitro management of *Meloidogyne incognita* (Tylenchida: Heteroderidae) using rhizosphere bacteria, *Pseudomonas* spp. and *Serratia* spp. compared with oxamyl. *Saudi Journal of Biological Sciences*, 28, 4876–4883.
- Abou-elnaga, Z. S., Alarif, W. M., Al-lihaibi, S. S., & Arabia, S. (2011). New Larvicidal Acetogenin from the Red Alga *Laurencia papillosa*. *CLEAN–Soil, Air, Water*, 39, 787–794. <https://doi.org/10.1002/clen.201000597>
- Abubakar, E. M. (2009). Antibacterial activity of crude extracts of *Euphorbia hirta* against some bacteria associated with enteric infections. *Journal of Medicinal Plants Research*, 3, 498–505.
- Agrios, G. N. (2005). *Plant Pathology* (5th ed.). Amsterdam: Elsevier Academic Press.
- Ahmad, A., Alkarkhi, A. F., Hena, S., & Khim, L. H. (2009). Extraction, separation and identification of chemical ingredients of *Elephantopus scaber* L. using factorial design of experiment. *International Journal of Chemistry*, 1, 36.
- Ahmad, R., Yu, K. X., Wong, C. L., & Jantan, I. (2016). Larvicidal and adulticidal activities of Malaysian seaweeds against *Aedes aegypti* (L.) and *Aedes albopictus* Skuse (Diptera: Culicidae). *The Southeast Asian Journal of Tropical Medicine and Public Health*, 47, 719–730.
- Aioub, A. A. A., Elesawy, A. E., & Ammar, E. E. (2022). Plant growth promoting rhizobacteria (PGPR) and their role in plant-parasitic nematodes control: A fresh look at an old issue. *Abstract Journal of Plant Diseases and Protection*, 129(6), 1305–1321. <https://doi.org/10.1007/s41348-022-00642-3>
- Akbar, R., & Khan, I. A. (2021). *Callosobruchus Maculatus* Fab. (Coleoptera Bruchidae) a major insect pest of stored pulses. *Fresenius Environmental Bulletin*, 30, 5098–5107.
- Akinbode, O. A., & Ikotun, T. (2008). Evaluation of some bioagents and botanicals in in vitro control of *Colletotrichum destructivum*. *African Journal of Biotechnology*, 7, 868–872.
- Alam, S., Akhter, M., Begum, F., et al. (2002). Antifungal activities (in vitro) of some plant extracts and smoke on four fungal pathogens of different hosts. *Pakistan Journal of Biological Sciences*, 5, 307–309.
- Alarif, W. M., Ayyad, S.-E.N., & Al-lihaibi, S. S. (2010). Acyclic Diterpenoid from the Red Alga *Gracilaria foliifera*. *Revista Latinoamericana de Química*, 38, 52–57.
- Ali, M. Y. S., Ravikumar, S., & Beulaohanson, M. (2011). Mosquito larvicidal activity of seaweeds extracts against *Anopheles stephensi*, *Aedes aegypti* and *Culex quinquefasciatus*. *Asian Pacific Journal of Tropical Medicine*, 3, 196–201. [https://doi.org/10.1016/S2222-1808\(13\)60040-7](https://doi.org/10.1016/S2222-1808(13)60040-7)
- Ali, A., Murphy, C. C., Demirci, B., et al. (2013). Insecticidal and biting deterrent activity of rose-scented geranium (*Pelargonium* spp.) essential oils and individual compounds against *Stephanitis pyrioides* and *Aedes aegypti*. *Pest Management Science*, 69, 1385–1392.
- Allahvaisi, S., Hassani, M., & Heidari, B. (2021). Bioactivity of azadirachtin against *Scrobipalpa ocellatella* Boyd. (Lepidoptera: Gelechiidae) on sugar beet. *Journal of Plant Protection Research*, 61(3), 280–289. <https://doi.org/10.24425/jppr.2021.137954>
- Al-Rahmah, A. N., Mostafa, A. A., Abdel-Megeed, A., et al. (2013). Fungicidal activities of certain methanolic plant extracts against tomato phytopathogenic fungi. *African Journal of Microbiology Research*, 7, 517–524.
- Argandoña, V., Del Pozo, T., San-Martín, A., & Roviroso, J. (2000). Insecticidal activity of *Plocamium cartilagineum* monoterpenes. *Boletín La Soc Chil Química*, 45, 371–376.
- Argandoña, V. H., Roviroso, J., San-Martín, A., et al. (2002). Antifeedant effects of marine halogenated monoterpenes. *Journal of Agriculture and Food Chemistry*, 50, 7029–7033.
- Arora, S. (2017). Antifeedant and Insecticidal Activity of Plant Extracts Against *Spodoptera litura* (Fab.) and *Lipaphis erysimi*. *Proceedings of the National Academy of Sciences, India Section B: Biological Sciences*, 87, 1229–1236. <https://doi.org/10.1007/s40011-015-0697-4>
- Asha, A., Rath, J. M., Raja, D. P., & Sahayaraj, K. (2012). Biocidal activity of two marine green algal extracts against third instar nymph of *Dysdercus cingulatus* (Fab.) (Hemiptera: Pyrrhocoridae). *Journal of Biopesticides*, 5, 129.
- Asimakis, E., Shehata, A. A., Eisenreich, W., et al. (2022). Algae and Their Metabolites as Potential Bio-Pesticides. *Microorganisms*, 10, 307.
- Baidoo, P. K., & Mochiah, M. B. (2016). Comparing the Effectiveness of Garlic (*Allium sativum* L.) and Hot Pepper (*Capsicum frutescens* L.) in the Management of the Major Pests of Cabbage Brassica oleracea (L.). *Sustainable Agriculture Research*, 5, 83–91. <https://doi.org/10.5539/sar.v5n2p83>
- Baidoo, P. K., Baidoe-Ansah, D., & Agbonu, I. (2012). Effects of Neem (*Azadirachta indica* A. Juss) Products on *Aphis craccivora* and its Predator *Harmonia axyridis* on Cowpea. *American Journal of Experimental Agriculture*, 2, 198–206.
- Bantoto, V., & Dy, D. (2013). The larvicidal activity of brown algae *Padina minor* (Yamada 1925) and *Dicyota linearis*

- (Greville 1830) against the dengue vector, *Aedes aegypti* (Linn 1762)(Diptera: Culicidae). *Journal of Vector Borne Diseases*, 50, 68.
- Beula, J. M., Ravikumar, S., & Ali, M. S. (2011). Mosquito larvicidal efficacy of seaweed extracts against dengue vector of *Aedes aegypti*. *Asian Pacific Journal of Tropical Biomedicine*, 1, S143–S146.
- Bianco, E. M., Pires, L., Santos, G. K. N., et al. (2013). Larvicidal activity of seaweeds from northeastern Brazil and of a halogenated sesquiterpene against the dengue mosquito (*Aedes aegypti*). *Industrial Crops and Products*, 43, 270–275. <https://doi.org/10.1016/j.indcrop.2012.07.032>
- Bibi, R., Muhammad, R., & Rasheed, M. (2020). Toxic assessment, growth disrupting and neurotoxic effects of red seaweeds' botanicals against the dengue vector mosquito *Aedes aegypti* L. *Ecotoxicology and Environmental Safety*, 195, 110451. <https://doi.org/10.1016/j.ecoenv.2020.110451>
- Bibi, R., Tariq, R. M., Abdelgaleil, S. A., & Rasheed, M. (2021). Insecticidal potential of botanicals from red seaweeds against stored grain pests, rice weevil (*Sitophilus oryzae* L.) and cowpea weevil (*Callosobruchus maculatus* Fab.). *Pakistan Journal of Zoology*, 54, 1657–1664. <https://doi.org/10.17582/journal.pjz/20190819070846>
- Bloomquist, J. R. (1993). Toxicology, mode of action and target site-mediated resistance to insecticides acting on chloride channels. *Comparative Biochemistry and Physiology Part C: Pharmacology, Toxicology and Endocrinology*, 106, 301–314.
- Blunden, G. (1993). Marine Algae as Sources of Biologically Active Compounds. *Interdisciplinary Science Reviews*, 18, 73–80.
- Boutjagualt, I., Hmimid, F., Errami, A., & Bouharroud, R. (2022). Biocatalysis and Agricultural Biotechnology Chemical composition and insecticidal effects of brown algae (*Fucus spiralis*) essential oil against *Ceratitis capitata* Wiedemann (Diptera : Tephritidae) pupae and adults. *Biocatalysis and Agricultural Biotechnology*, 40, 102308. <https://doi.org/10.1016/j.cbab.2022.102308>
- Bouwmeester, H., Dekkers, S., Noordam, M. Y., et al. (2009). Review of health safety aspects of nanotechnologies in food production. *Regulatory Toxicology and Pharmacology*, 53, 52–62. <https://doi.org/10.1016/j.yrtph.2008.10.008>
- Cassano, V., Santos, N., Maria, E., et al. (2019). *Laurencia longiramea* sp. nov. for Brazil and an emendation of the generic delineation of *Corynecladia* (Ceramiales, Rhodophyta). *Phycologia*, 58, 115–127. <https://doi.org/10.1080/00318884.2018.1523519>
- Chanthini, K. M. P., Senthil-Nathan, S., Stanley-Raja, V., et al. (2021). Biologically active toxin from macroalgae *Chaetomorpha antennina* Bory, against the lepidopteran *Spodoptera litura* Fab. and evaluation of toxicity to earthworm, *Eudrilus eugeniae* Kinb. *Chemical and Biological Technologies in Agriculture*, 8, 1–15.
- Chen, Y., Lv, M., Zhou, J., et al. (2022). Potential Value of Wood Tar as a Natural Fungicide against *Valsa mali*. *Molecules*, 27, 1531.
- Chengala, L., & Singh, N. (2017). Botanical pesticides – a major alternative to chemical pesticides : A review. *International Journal of Life Sciences*, 5, 722–729.
- Choi, G. J., Jang, K. S., Kim, J. S., et al. (2004). In Vivo Anti-fungal Activities of 57 Plant Extracts Against Six Plant Pathogenic Fungi. *Plant Pathology Journal*, 20, 184–191. <https://doi.org/10.5423/PPJ.2004.20.3.184>
- Chougule, P. M., & Andoji, Y. S. (2016). Antifungal activity of some common medicinal plant extracts against soil borne phytopathogenic fungi *Fusarium oxysporum* causing wilt of tomato. *International Journal of Development Research*, 3, 7030–7033.
- Coppejans, E., Leiaert, F., Dargent, O., Gunasekara, E., Clerk, O. (2009). Sri Lankan seaweeds: Methodologies and Field guide to the dominant species (Vol. 6, pp. 265). Belgian Development Cooperation. [https://doi.org/10.1002/1526-4998\(200008\)56:8<651:AID-PS201>3.0.CO;2-U](https://doi.org/10.1002/1526-4998(200008)56:8<651:AID-PS201>3.0.CO;2-U)
- Costa, J. A. V., Freitas, B. C. B., Cruz, C. G., et al. (2019). Potential of microalgae as biopesticides to contribute to sustainable agriculture and environmental development. *Journal of Environmental Science and Health, Part B, Pesticides, Food Contaminants, and Agricultural Wastes*, 54, 366–375. <https://doi.org/10.1080/03601234.2019.1571366>
- Crews, P., Myers, B. L., Naylor, S., et al. (1984). Bio-active monoterpenes from red seaweeds. *Phytochemistry*, 23, 1449–1451.
- Da Silva, L., Macie, A., Mendonça, C., & Unior, L. (2017). Use of encapsulated carvacrol with yeast cell walls to control resistant strains of *Rhipicephalus microplus* (Acari : Ixodidae). *Industrial Crops and Products*, 108, 190–194. <https://doi.org/10.1016/j.indcrop.2017.06.037>
- Damalas, C. A. (2018). Current status and recent developments in biopesticide use. *Agriculture*, 8, 13. <https://doi.org/10.3390/agriculture8010013>
- Desa UN. (2011). World population prospects: The 2010 revision, highlights and advance tables. United Nations.
- Dimetry, N. Z., Amer, S. A. A., & Reda, A. S. (1993). Biological activity of two neem seed kernel extracts against the two-spotted spider mite *Tetranychus urticae* Koch. *Journal of Applied Entomology*, 116, 308–312. <https://doi.org/10.1111/j.1439-0418.1993.tb01201.x>
- Djeussi, D. E., Noumedem, J. A. K., Seukep, J. A., et al. (2013). Antibacterial activities of selected edible plants extracts against multidrug-resistant Gram-negative bacteria. *BMC Complementary and Alternative Medicine*, 13, 1–8.
- Elbanna, S. M., & Hegazi, M. M. (2011). Screening of some seaweeds species from South Sinai, Red Sea as potential bioinsecticide against mosquito larvae, *Culex pipens*. *Egyptian Academic Journal of Biological Sciences. A, Entomology*, 4, 21–30.
- Elbeshehy, E. K. F., Metwali, E. M. R., & Almaghrabi, O. A. (2015). Antiviral activity of *Thuja orientalis* extracts against watermelon mosaic virus (WMV) on *Citrus lanatus*. *Saudi Journal of Biological Sciences*, 22, 211–219. <https://doi.org/10.1016/j.sjbs.2014.09.012>
- Eldeeb, A. M., Farag, A. A. G., Al-Harbi, M. S., et al. (2022). Controlling of *Meloidogyne incognita* (Tylenchida: Heteroderidae) using nematicides, *Linum usitatissimum* extract and certain organic acids on four peppers cultivars under greenhouse conditions. *Saudi Journal of Biological Sciences*, 29, 3107–3113.

- Ferreira, I., De Oliveira, R. G., Soares, I. M., et al. (2013). Evaluation of acute toxicity , antibacterial activity , and mode of action of the hydroethanolic extract of *Piper umbellatum* L. *Journal of Ethnopharmacology* 1–7. <https://doi.org/10.1016/j.jep.2013.10.011>
- Fu, Y., Bhadha, J. H., Rott, P., et al. (2020). Investigating the use of aquatic weeds as biopesticides towards promoting sustainable agriculture. *PLoS ONE*, 15, 1–23. <https://doi.org/10.1371/journal.pone.0237258>
- Fukuzawa, A., & Masamune, T. (1981). Laurepinnacin and isolaurepinnacin, new acetylenic cyclic ethers from the marine red alga *Laurencia pinnata* Yamada. *Tetrahedron Letters*, 22, 4081–4084.
- Gayathri, A., & Ramesh, K. V. (2013). Antifungal activity of *Euphorbia hirta* L. inflorescence extract against *Aspergillus flavus* A mode of action study. *International Journal of Current Microbiology and Applied Sciences*, 2, 31–37.
- Gowthish, K., & Kannan, R. (2019). Pesticidal potentials of some red algal seaweeds from Tuticorin coast against the tobacco cutworm *Spodoptera litura* Fab. (Lepidoptera: Noctuidae). *International Journal of Scientific & Technology Research*, 8, 502–506.
- Gowthish, K., Muthukumaran, N., & Selvanarayanan. (2018). Antixenosis resistance in black gram against leaf caterpillar *Spodoptera litura* fab. *Multilogic in Science*, 8, 62–65.
- Grdisa, M., & Kristina, G. (2013). Botanical Insecticides in Plant Protection. *Agriculturae Conspectus Scientificus*, 78, 85–93.
- Gressler, V., Coplepicolo, P., & Pinto, E. (2009). Useful strategies for algal volatile analysis. *Current Analytical Chemistry*, 5, 271–292. <https://doi.org/10.2174/157341109788680255>
- Guiry, M. D., & Guiry, G. M. (2017). *AlgaeBase*. World-wide electronic publication, National University of Ireland, Galway. <https://www.algaebase.org>. Accessed 20 Oct 2021
- Gupta, S., & Dikshit, A. K. (2010). Biopesticides: An ecofriendly approach for pest control. *Journal of Biopesticides*, 3, 186–188.
- Hao, H., Zuo, Y., Fang, J., et al. (2021). Transcriptome Analysis of *Aedes albopictus* (Diptera: Culicidae) Larvae Exposed With a Sublethal Dose of Haedoxan A. *Journal of Medical Entomology*, 58, 2284–2291.
- Harlapur, S. I., Kulkarni, M. S., Wali, M. C., Kulkarni, S. (2010). Evaluation of plant extracts, bio-agents and fungicides against *Exserohilum turcicum* (Pass.) Leonard and Suggs. causing *Turcicum* leaf blight of maize. *Karnataka Journal of Agricultural Sciences* 20.
- Hazafa, A., Murad, M., Umer Masood, M., Bilal, S., Nasir Khan, M., Farooq, Q., ... Naeem, M. (2022). Nano-biopesticides as an emerging technology for pest management. *IntechOpen*. <https://doi.org/10.5772/intechopen.101285>
- He, J., Dou, M., Xie, J., et al. (2021). Discovery of zeylenone from *Uvaria grandiflora* as a potential botanical fungicide. *Pest Management Science*, 5407–5417. <https://doi.org/10.1002/ps.6580>
- Hira, K., Tariq, R. M., Sultana, V., et al. (2017). Effect of seaweeds occurring at Karachi coast on mosquito larvae and liver function in rats. *Pharmaceutical Sciences*, 30, 387–391.
- Hong, Z., Talib, K. M., Mujtaba, K. G., et al. (2021). Antifungal potential of cinnamon essential oils against *Phytophthora colocasiae* causing taro leaf blight. *Chemical and Biological Technologies in Agriculture*, 8, 1–9. <https://doi.org/10.1186/s40538-021-00238-3>
- Iliopoulou, D., Vagias, C., Harvala, C., & Roussis, V. (2002). C 15 Acetogenins from the red alga *Laurencia obtusa*. *Phytochemistry*, 59, 111–116.
- Ioannou, E., & Roussis, V. (2009). Natural products from seaweeds. In A. E. Osbourn & V. Lanzotti (Eds.), *Plant-derived natural products* (pp. 51–81). Springer-Verlag. https://doi.org/10.1007/978-0-387-85498-4_2
- Ishii, T., Nagamine, T., Cao, B., et al. (2017). Insecticidal and Repellent Activities of Laurinterol from the Okinawan Red Alga *Laurencia nidifica*. *Records of Natural Products*, 1, 63–68.
- Ishii, T., Abe, T., & Kikuchi, N. (2019). Investigation of Insect Repellent Activity of Cyclocolorenone Obtained from the Red Alga *Laurencia intricata* and Minoru Suzuki. *Records of Natural Products*, 1, 81–84.
- Ismail, M., Javed, S., Batool, Z., et al. (2021). Nematocidal, Insecticidal, Anti-Inflammatory And Cytotoxic Activities Of Selected Medicinal Plants. *Analele Univ Din Oradea, Fasc Biol*, 28, 1–13.
- Isman, M. B. (2017). Industrial Crops & Products Bridging the gap : Moving botanical insecticides from the laboratory to the farm. *Industrial Crops and Products*, 110, 10–14. <https://doi.org/10.1016/j.indcrop.2017.07.012>
- Jagadeesan, Y., Singh, V. P., & Balaiah, A. (2015). In-Vitro assessment of larvicidal activity of marine macro algae collected from South Indian coastal area. *Asian Journal of Microbiology, Biotechnology & Environmental Sciences*, 17, 171–176.
- Jang, J. Y., Dang, L., Choi, J., et al. (2019). Control of root-knot nematodes using *Waltheria indica* producing 4-quinolone alkaloids. *Pest Management Science*, 75, 2264–2270. <https://doi.org/10.1002/ps.5363>
- Jarman, W. M., & Ballschmiter, K. (2012). From coal to DDT : The history of the development of the pesticide DDT from synthetic dyes till Silent Spring. *Endeavour*, 36, 131–142. <https://doi.org/10.1016/j.endeavour.2012.10.003>
- Javier, A. M. V., Ocampo, V. R., Ceballo, F. A., et al. (2019). Insecticidal Activity of Crude Ethanolic Extracts of Selected Philippine Plants against Diamondback Moth, *Plutella xylostella* Linnaeus. *The Philippine Journal of Science*, 148, 33–43.
- Jhala, J., Baloda, A. S., & Rajput, V. S. (2020). Role of bio-pesticides in recent trends of insect pest management: A review. *Journal of Pharmacognosy and Phytochemistry*, 9, 2237–2240.
- Khafagi, I. K., & Dewedar, A. (2000). The efficiency of random versus ethno-directed research in the evaluation of Sinai medicinal plants for bioactive compounds. *Journal of Ethnopharmacology*, 71, 365–376. [https://doi.org/10.1016/S0378-8741\(00\)00164-1](https://doi.org/10.1016/S0378-8741(00)00164-1)
- Khan, R., Islam, B., Akram, M., et al. (2009). Antimicrobial Activity of Five Herbal Extracts Against Multi Drug Resistant (MDR) Strains of Bacteria and Fungus of

- Clinical Origin. *Molecules*, 14, 586–597. <https://doi.org/10.3390/molecules14020586>
- Khanavi, M., Toulabi, P. B., Abai, M. R., et al. (2011). Larvicidal activity of marine algae, *Sargassum swartzii* and *Chondria dasyphylla*, against malaria vector *Anopheles stephensi*. *Journal of Vector Borne Diseases*, 48, 241.
- Kitherian, S., Thangapandi, V., Rath, M., & Antony, J. (2021). Environmental Nanotechnology, Monitoring & Management Seaweed *Lobophora variegata* -based Silver Nanopesticide for environmental friendly management of economically important pest, *Spodoptera litura*. *Environmental Nanotechnology, Monitoring & Management*, 16, 100531. <https://doi.org/10.1016/j.enmm.2021.100531>
- Kombiah, P., & Sahayaraj, K. (2012). Repellent activity of *Caulerpa scalpelliformis* extracts and its formulations against *Spodoptera litura* and *Dysdercus cingulatus* (Fab.). *Journal of Biopesticides*, 5, 145.
- Kumar, R. S., & Sujith, S. (2011). In vitro mosquito larvicidal activity of marine algae against the human vectors, *Culex quinquefasciatus* (Say) and *Aedes aegypti* (Linnaeus) (Diptera: Culicidae). *International Journal of Zoological Research*, 7, 272–278.
- Kumar, K. P., Murugan, K., Kovendan, K., et al. (2012). Combined effect of seaweed (*Sargassum wightii*) and *Bacillus thuringiensis* var. *israelensis* on the coastal mosquito, *Anopheles sundanicus*, Tamil Nadu, India. *Science Asia*, 38, 41–46.
- Kumar, S., Singh, A. (2015). Biopesticides: Present Status and the Future Prospects. *Journal of Biofertilizers & Biopesticides*, 06. <https://doi.org/10.4172/jbfbp.1000e129>
- Lade, B., Gogle, D. P., Lade, D. B., Moon, G. M., Nandeshwar, S. B., & Kumbhare, S. D. (2019). Nanobiopesticide formulations: Application strategies today and future perspectives. *Nano-Biopesticides Today and Future Perspectives*.
- LaLonde, R. T., Morris, C. D., Wong, C. F., et al. (1979). Response of *Aedes triseriatus* larvae to fatty acids of *Cladophora*. *Journal of Chemical Ecology*, 5, 371–381.
- Li, Y., Aioub, A. A., Lv, B., et al. (2019). Antifungal activity of pregnane glycosides isolated from *Periplocia sepium* root barks against various phytopathogenic fungi. *Industrial Crops and Products*, 132, 150–155.
- Liao, M., Xiao, J. J., Zhou, L. J., et al. (2017). Chemical composition, insecticidal and biochemical effects of *Melaleuca alternifolia* essential oil on the *Helicoverpa armigera*. *Journal of Applied Entomology*, 141, 721–728.
- Lopez, M. D., Maudhuit, A., & Poncelet, D. (2012). Development of Formulations to Improve the Controlled-Release of Linalool to Be Applied As an Insecticide. *Journal of Agriculture and Food Chemistry*, 60, 1187–1192.
- Lu, W., Senapati, D., Wang, S., et al. (2010). Effect of surface coating on the toxicity of silver nanomaterials on human skin keratinocytes. *Chemical Physics Letters*, 487, 92–96. <https://doi.org/10.1016/j.cplett.2010.01.027>
- Maeda, M., Kodamat, T., Tanaka, T., et al. (1984). Insecticidal and neuromuscular activities of domoic acid and its related compounds. *Journal of Pesticide Science*, 9, 27–32.
- Manilal, A., Sujith, S., Kiran, G. S., et al. (2009). Antimicrobial potential and seasonality of red algae collected from the southwest coast of India tested against shrimp, human and phytopathogens. *Annales De Microbiologie*, 59, 207–219.
- Manilal, A., Sujith, S., Sabarathnam, B., et al. (2011). Biological activity of red alga *Laurencia brandenii*. *Acta Botanica Croatica*, 70, 81–90.
- Manjunatha, S., Biradar, D., & Aladakatti, Y. (2016). Nanotechnology and its applications in agriculture : A review. *Journal of Farm Sciences*, 29, 1–13.
- Mayanglambam, S., Raghavendra, A., & Rajashekar, Y. (2022). Use of *Ageratina adenophora* (Spreng.) essential oil as insecticidal and antifeedant agents against diamondback moth, *Plutella xylostella* (L.). *Journal of Plant Diseases and Protection*, 129, 439–448. <https://doi.org/10.1007/s41348-022-00573-z>
- Mazid, S., Kalita, J. C., & Rajkhowa, R. C. (2011). A review on the use of biopesticides in insect pest management. *International Journal of Advanced Science and Technology*, 7, 169–178.
- Meena, R. K., Mishra, P. (2020). Bio-pesticides for Agriculture and Environment Sustainability. *Resource Use Efficiency in Agriculture*, 85–107. https://doi.org/10.1007/978-981-15-6953-1_3
- Mohammadi, S., Ataollah, S., Shandiz, S., et al. (2016). Evaluation of Anti-Bacterial Properties of *Euphorbia Condylolcarpa* Methanol Extract. *Archives of Microbiology & Immunology*, 1, 12–20.
- Montanha, J. A., & Moellerke, P. (2004). Antiviral Activity of Brazilian Plant Extracts. *Acta Farmaceutica Bonaerense*, 23, 183–186.
- Morales-ubaldo, A. L., Rivero-perez, N., Avila-ramos, F., et al. (2021). Bactericidal Activity of *Larrea tridentata* Hydroalcoholic Extract against Phytopathogenic Bacteria. *Agronomy*, 11, 1–11.
- Multigner, L., Ndong, J. R., Giusti, A., et al. (2010). Chlordecone Exposure and Risk of Prostate Cancer. *Journal of Clinical Oncology*, 28, 3457–3462. <https://doi.org/10.1200/JCO.2009.27.2153>
- Nenaah, G. E., Ibrahim, S. I. A., & Al-assiuty, B. A. (2015). Chemical composition, insecticidal activity and persistence of three Asteraceae essential oils and their nanoemulsions against *Callosobruchus maculatus* (F.). *Journal of Stored Products Research*, 61, 9–16. <https://doi.org/10.1016/j.jspr.2014.12.007>
- Ng, V. W. L., Tan, J. P. K., Leong, J., et al. (2014). Antimicrobial polycarbonates: Investigating the impact of nitrogen-containing heterocycles as quaternizing agents. *Macromolecules*, 47, 1285–1291. <https://doi.org/10.1021/ma402641p>
- Nikkhah, M., Hashemi, M., Habibi, M. B., & Farhoosh, R. (2017). Synergistic effects of some essential oils against fungal spoilage on pear fruit. *International Journal of Food Microbiology*, 257, 285–294. <https://doi.org/10.1016/j.ijfoodmicro.2017.06.021>
- Nukmal, N., Rosa, E., & Apriliyani, K. M. (2017). Insecticidal effects of the flavonoid-rich fraction of leaves extract of gamal (*Gliricidia sepium*) on the coffee mealybugs (*Planococcus citri* risso.). *Annual Research & Review in Biology*, 16, 1–9. <https://doi.org/10.9734/ARRB/2017/36209>
- Pal, G. K., Kumar, B., & Shahi, S. K. (2013). Antifungal activity of some common weed extracts against

- phytopathogenic fungi *Alternaria* spp. *International Journal of Universal Pharmacy and Life Sciences*, 3, 6–14.
- Pardavella, I., Daferera, D., Tselios, T., Skiada, P., & Gianakou, I. (2021). The use of essential oil and hydrosol extracted from cuminum cyminum seeds for the control of meloidogyne incognita and meloidogyne javanica. *Plants*, 10(1), 46. <https://doi.org/10.3390/plants10010046>
- Perez, M., Falque, E., & Dominguez, H. (2016). Antimicrobial action of compounds from marine seaweed. *Marine Drugs*, 14(3), 52. <https://doi.org/10.3390/md14030052>
- Rajasekaran, D., Palombo, E. A., Yeo, T. C., et al. (2013). Identification of Traditional Medicinal Plant Extracts with Novel Anti-Influenza Activity. *PLoS ONE*, 8, 1–15. <https://doi.org/10.1371/journal.pone.0079293>
- Rajesh, S., Asha, A., Kombiah, P., Sahayaraj, K. (2011). Biocidal activity of algal seaweed on insect pest and fungal plant pathogen. Proc Natl Semin Harmful/Beneficial Insects Agric Importance held 17–18 Febr 2011 St Joseph's Coll Calicut, Kerala 86–91.
- Ranglová, K., Lakatos, G. E., Manoel, J. A. C., et al. (2021). Growth, biostimulant and biopesticide activity of the MACC-1 *Chlorella* strain cultivated outdoors in inorganic medium and wastewater. *Algal Research*, 53, 102136. <https://doi.org/10.1016/j.algal.2020.102136>
- Regnault-Roger, C., Vincent, C., & Arnason, J. T. (2012). Essential oils in insect control: low-risk products in a high-stakes world. *Annu Rev Entomol*, 57, 405–424.
- Reno, F., Demeter, S., Degand, H., et al. (2022). The modes of action of *Mentha arvensis* essential oil on the granary weevil *Sitophilus granarius* revealed by a label-free quantitative proteomic analysis. *J Pest Sci* (2004), 95, 381–395. <https://doi.org/10.1007/s10340-021-01381-4>
- Rizvi, M. A., & Shameel, M. (2006). In vitro nematocidal activities of seaweed extracts from Karachi coast. *Pakistan Journal of Botany*, 38, 1245–1248.
- Roviro, J., Soler, A., Blanc, V., et al. (2013). Bioactive monoterpenes from antarctic plocamium cartilagineum. *Journal of the Chilean Chemical Society*, 58, 2025–2026. <https://doi.org/10.4067/S0717-97072013000400026>
- Sahayaraj, K., & Jeeva, Y. M. (2012). Nymphicidal and Ovipositional Efficacy of Seaweed *Sargassum tenerrimum* (J. Agardh) against *Dysdercus cingulatus* (Fab.) (Pyrrhocoridae). *Chilean Journal of Agricultural Research*, 72, 152–156. <https://doi.org/10.4067/s0718-58392012000100024>
- Sahayaraj, K., & Kalidas, S. (2011). Evaluation of nymphicidal and ovicidal effect of a seaweed, *Padina pavonica* (Linn.) (Phaeophyceae) on cotton pest, *Dysdercus cingulatus* (Fab.). *Indian Journal of Marine Sciences*, 40, 125–129.
- Sahayaraj, K., Asharaja, A., Ponsankar, A., et al. (2019). Behavioral response and relative toxicity for the active compounds of *Caulerpa veravalensis* (Thivy and Chauhan) against nymph of *Dysdercus cingulatus* (Fab.) (Hemiptera : Pyrrhocoridae). *Journal of Asia-Pacific Entomology*, 22, 417–426. <https://doi.org/10.1016/j.aspen.2019.02.009>
- Salvador-Neto, O., Gomes, S. A., Soares, A. R., et al. (2016). Larvicidal potential of the halogenated sesquiterpene (+)-obtusol, isolated from the alga *Laurencia dendroidea* J. Agardh (Ceramiales: Rhodomelaceae), against the Dengue Vector Mosquito *Aedes aegypti* (Linnaeus) (Diptera: Culicidae). *Marine Drugs*, 14, 20. <https://doi.org/10.3390/md14020020>
- Sasson, Y., Levy-Ruso, G., Toledano, O., & Ishaaya, I. (2007). Nanosuspensions: Emerging novel agrochemical formulations. In I. Ishaaya, A. R. Horowitz, & R. Nauen (Eds.), *Insecticides design using advanced technologies*. Springer. https://doi.org/10.1007/978-3-540-46907-0_1
- Selvin, J., & Lipton, A. P. (2004). Biopotentials of ulva fasciata and hypnea musciformis collected from the peninsular coast of India. *Journal of Marine Science and Technology*, 12, 1–6. <https://doi.org/10.51400/2709-6998.2214>
- Sharma, J., Bhardwaj, V. K., Das, P., & Purohit, R. (2021). Plant-based analogues identified as potential inhibitor against tobacco mosaic virus: A biosimulation approach. *Pesticide Biochemistry and Physiology*, 175, 104858. <https://doi.org/10.1016/j.pestbp.2021.104858>
- Sherry, M., Charcosset, C., Fessi, H., & Greige-Gerges, H. (2013). Essential oils encapsulated in liposomes: A review. *Journal of Liposome Research*, 23, 268–275.
- Shirurkar, D. D., & Wahegaonkar, N. K. (2012). Antifungal activity of selected plant derived oils and some fungicides against seed borne fungi of maize. *European Journal of Experimental Biology*, 2, 1693–1696.
- Singh, I. (2017). Antimicrobials in higher plants: Classification, mode of action and bioactivities. *Chemical Biology Letters*, 4, 48–62.
- Sinha, B., & Biswas, I. (2011). Potential of Bio-Pesticides in Indian Agriculture Vis-a-Vis Rural Development. *SSRN Electronic Journal*. <https://doi.org/10.2139/ssrn.1472371>
- Smit, A. J. (2004). Medicinal and pharmaceutical uses of seaweed natural products: A review. *Journal of Applied Phycology*, 16, 245–262.
- Souto, A. L., Sylvestre, M., Tölke, E. D., et al. (2021). Plant-derived pesticides as an alternative to pest management and sustainable agricultural production: Prospects, applications and challenges. *Molecules*, 26, 4835.
- Sowmiya, R., Balasubramani, G., Deepak, P., et al. (2017). Characterization and screening of in vitro antimalarial and larvicidal activities of selected seaweeds from south-east coast of India against *Plasmodium falciparum* and *Anopheles stephensi*. *Journal of Coastal Life Medicine*, 5, 242–248.
- Stanley, J., Preetha, G., Chandrasekaran, S., et al. (2014). Efficacy of Neem oil on cardamom thrips, sciothrips cardamomi ramk., and organoleptic studies. *Psyche* (London) 2014. <https://doi.org/10.1155/2014/930584>
- Stenger, L. D., Abati, R., Pawlak, I. G., et al. (2021). Toxicity of essential oil of *Eugenia uniflora* (L.) to *Thaumastocoris peregrinus* (Hemiptera: Thaumastocoridae) and selectivity to the parasitoid *Cleruchoides noackae* (Lin & Hubert) (Hymenoptera: Mymaridae). *Crop Protection*, 147, 105693. <https://doi.org/10.1016/j.cropro.2021.105693>
- Stevenson, P. C., Isman, M. B., & Belmain, S. R. (2017). Industrial Crops & Products Pesticidal plants in Africa : A global vision of new biological control products from local uses. *Industrial Crops and Products*, 110, 2–9. <https://doi.org/10.1016/j.indcrop.2017.08.034>

- Sugiura, M., Horibe, Y., Kawada, H., & Takagi, M. (2008). Insect spiracle as the main penetration route of pyrethroids. *Pesticide Biochemistry and Physiology*, 91, 135–140. <https://doi.org/10.1016/j.pestbp.2008.03.001>
- Suleiman, E., & Abdallah, W. (2014). In vitro Activity of Garlic (*Allium sativum*) on Some Pathogenic Fungi. *European J Med Plants*, 4, 1240–1250. <https://doi.org/10.9734/ejmp/2014/10132>
- Sunder, J., Singh, D. R., Jeyakumar, S., et al. (2011). Antibacterial activity in solvent extract of different parts of *Morinda citrifolia* plant. *Journal of Pharmaceutical Sciences and Research*, 3, 1404–1407.
- Tandon, S., & Mittal, A. K. (2018). Insecticidal and growth inhibitory activity of essential oils of *Boenninghausenia albi fl ora* and *Teucrium quadrifarium* against *Spilarcia obliqua*. *Biochemical Systematics and Ecology*, 81, 70–73. <https://doi.org/10.1016/j.bse.2018.09.010>
- Thakore, Y. (2006). The biopesticide market for global agricultural use. *Industrial Biotechnology*, 3, 194–208.
- Thangam, T. S., & Kathiresan, K. (1991). Mosquito Larvicidal Effect of Seaweed Extracts. *Botanica Marina*, 34, 433–436. <https://doi.org/10.1515/botm.1991.34.5.433>
- Thangam, T. S., & Kathiresan, K. (1996). The prey consumption and prey preference of the larvae of the mosquito *Culex* (Lutzia) raptor on the larvae of *Culex quinquefasciatus*. *Experientia*, 52, 380–382. <https://doi.org/10.1007/BF01919544>
- Venkatesan, M., Arumugam, V., Pugalendi, R., et al. (2019). Antioxidant, anticoagulant and mosquitocidal properties of water soluble polysaccharides (WSPs) from Indian seaweeds. *Process Biochemistry*, 84, 196–204. <https://doi.org/10.1016/j.procbio.2019.05.029>
- Villaverde, J. J., Sandín-españa, P., Sevilla-morán, B., et al. (2016). Biopesticides from Natural Products: Current Development, Legislative Framework, and Future Trends. *BioResources*, 11, 5618–5640.
- Vinoth, S., Shankar, S. G., Gurusaravanan, P., et al. (2019). Anti-larvicidal Activity of Silver Nanoparticles Synthesized from *Sargassum polycystum* Against Mosquito Vectors. *Journal of Cluster Science*, 30, 171–180. <https://doi.org/10.1007/s10876-018-1473-4>
- Voukeng, I. K., Veronique, P., Kuete, V. (2017). Multidrug resistant bacteria are sensitive to *Euphorbia prostrata* and six others Cameroonian medicinal plants extracts. *BMC Research Notes* 1–8. <https://doi.org/10.1186/s13104-017-2665-y>
- Wang, M. B., Abbott, D. C., & Waterhouse, P. M. (2000). A single copy of a virus-derived transgene encoding hairpin RNA gives immunity to barley yellow dwarf virus. *Molecular Plant Pathology*, 1, 347–356. <https://doi.org/10.1046/j.1364-3703.2000.00038.x>
- Watanabe, K., Umeda, K., Kurita, Y., et al. (1990). Two insecticidal monoterpenes, telfairine and aplysiaterpenoid A, from the red alga *Plocamium telfairiae*: Structure elucidation, biological activity, and molecular topographical consideration by a semiempirical molecular orbital study. *Pesticide Biochemistry and Physiology*, 37, 275–286. [https://doi.org/10.1016/0048-3575\(90\)90134-N](https://doi.org/10.1016/0048-3575(90)90134-N)
- Wiwattanawanichakun P, Saehlee S, Yooboon T, Kumrunsee N (2022) Toxicity of isolated phenolic compounds from *Acorus calamus* L. to control *Spodoptera litura* (Lepidoptera : Noctuidae) under laboratory conditions. *Chemical and Biological Technologies in Agriculture*, 1–9. <https://doi.org/10.1186/s40538-021-00274-z>
- Xu, S., Zhao, X., Liu, F., et al. (2018). Crucial role of oxidative stress in bactericidal effect of parthenolide against *Xanthomonas oryzae* pv. *oryzae*. *Pest Management Science*, 74, 2716–2723. <https://doi.org/10.1002/ps.5091>
- Yan, T. K., Asari, A., Salleh, S. A., & Azmi, W. A. (2021). Eugenol and thymol derivatives as antifeedant agents against red palm weevil, *Rhynchophorus ferrugineus* (Coleoptera: Dryophthoridae) larvae. *Insects*, 12, 551. <https://doi.org/10.3390/insects12060551>
- Yasmin, M., Hossain, K. M., & Bashar, M. A. (2008). Effects of some angiospermic plant extracts on in vitro vegetative growth of *Fusarium moniliforme*. *Bangladesh Journal of Botany*, 37, 85–88.
- Yoon, M., Cha, B., & Kim, J. (2013). Recent Trends in Studies on Botanical Fungicides in Agriculture. *The Plant Pathology Journal*, 29, 1–9.
- Yu, K., Wong, C., Ahmad, R., & Jantan, I. (2015). Mosquitocidal and Oviposition Repellent Activities of the Extracts of Seaweed *Bryopsis pennata* on *Aedes aegypti* and *Aedes albopictus*. *Molecules*, 20, 14082–14102. <https://doi.org/10.3390/molecules200814082>
- Zhang, Y., Liu, X., Wang, Y., et al. (2016). Antibacterial activity and mechanism of cinnamon essential oil against *Escherichia coli* and *Staphylococcus aureus*. *Food Control*, 59, 282–289.
- Zhi, X., Song, L., Liang, J., et al. (2022). Bioorganic & Medicinal Chemistry Letters Synthesis and in vitro antifungal activity of new Michael-type amino derivatives of xanthatin, a natural sesquiterpene lactone from *Xanthium*. *Bioorganic & Medicinal Chemistry Letters*, 55, 128481. <https://doi.org/10.1016/j.bmcl.2021.128481>