

***Lindernia dubia* (L.) PENNELL: A NEW INVASIVE IN THE ROMANIAN BANAT**

Alina NEACȘU, Ilinca IMBREA, Alina LAȚO, Gicu-Gabriel ARSENE

Banat's University of Agricultural Sciences and Veterinary Medicine "King Michael I of Romania"
from Timisoara, Faculty of Agriculture, 119 Aradului Street,
Timisoara, 300645, Romania

Corresponding author email: botanix.timagro@gmail.com

Abstract

In this paper we report the presence of the species Lindernia dubia (L.) Pennell in the flora of Banat (SW Romania) and at the same time we present a new place for this species in Romania. We found the species on the lake Surduc area. This is a 362 ha artificial lake, a protected area (nature reserve of national interest). We found Lindernia dubia here for the first time in 2005, in flooded areas periodically (on clay soils), on the shore of the lake as well as in the nearby arable fields. We suppose he arrived here with the fish he brought to populate his lake or by seeds brought through the birds. The size of the population depends on the fluctuation of water level. Being an invasive species, we believe it is necessary to monitor it; Sirbu & Oprea (2011) say it extends into characteristic habitats while Doroftei & Anastasiu (2014) consider it one of the most abundant invasive species from the Danube Delta.

Key words: *Lindernia dubia* (L.) Pennell, Romanian Banat, invasive.

INTRODUCTION

The genus *Lindernia* Allioni encompasses about 70 species distributed throughout warm tropical and warm-temperate regions of New and Old World (Flora of China, 1998); recently, based on cpDNA sequences, many *Lindernia* species have been transferred to different genera within *Linderniaceae* (Les, 2017). Formerly included in *Scrophulariaceae*, this genus belongs to *Linderniaceae* Borsch et al., K. Müller, & Eb. Fischer (Stevens, 2017). In Europe, the genus *Lindernia* is represented mainly by two species: *Lindernia procumbens* (Krock.) Philcox, native and *Lindernia dubia* (L.) Pennell, from North America, naturalized (Tutin et al., 1972). Marhold (2017, The Euro+Med PlantBase¹) mentions also *Lindernia anagallidea* (Michx.) Pennell (*Lindernia dubia* var. *anagallidea* (Michx.) Cooperr.) from Italy, a species native from North America (Cook, 1985). The most significant morphological differences between *Lindernia procumbens* (Krock.) Philcox and *Lindernia dubia* (L.) Pennell are documented by Molnár et al. (2000): length of leaves, length of pedicels, ratio of pedicels/leaf length. These

two species may occur simultaneously in a habitat and may interbreed (Yoshino et al., 2006, in Schmotzer, 2015). *Lindernia dubia* (L.) Pennell is characterized by: serrated leaves, floral pedicels shorter than the bracts, corolla longer than the calyx, two fertile stamens (Figure 1). *Lindernia procumbens* (Krock.) Philcox (Natura 2000 species code: 1725) is included in the Annex IV of the Council Directive 92/43/EEC among the species of community interest in need of strict protection. *Lindernia dubia* (L.) Pennell is an invasive, accidentally introduced in France in 1850, by ships on the Loire river banks (Fournier, 1940²; Simons & Jansen, 2018). According to Marhold (2017), the species is to be found in: Bulgaria, France, Germany, Switzerland, Italy, Portugal, Slovenia, while DAISIE Database³ mentions *Lindernia dubia* (L.) Pennell present in: Belgium, Bulgaria, Czech Republic, France, Germany, Greece, Hungary, Italy, Macedonia, Portugal, Romania, Spain. The Euro+Med PlantBase presents the following countries as a distribution for genus *Lindernia*: Armenia, Austria, Bulgaria, Czech Republic, Croatia, Egypt, France, Germany,

¹ <https://ww2.bgbm.org/EuroPlusMed/PTaxonDetail.asp?NameCache=Lindernia&PTRefFk=7200000>

² http://www.fcbrn.fr/sites/fcbrn.fr/files/ressource_telechargeable/fiche_lindernia_dubia_v2.pdf


³ <http://www.europe-aliens.org/speciesFactsheet.do?speciesId=19525#>

Greece, Switzerland, Spain, Hungary, Italy, Portugal, Poland, Central European Russia, South European Russia, Romania, Slovakia, Slovenia, Serbia (including Kosovo and Vojvodina), Asiatic Turkey, Ukraine and for


the species *Lindernia dubia* (L.) Pennell, the following: Bulgaria, France, Germany, Spain, Italy, Portugal, Slovenia. The European area of the species included also Poland (first mention: 2003, Drobnik & Buchalik, 2004).



a) Lake shore with *Lindernia dubia*



b) Corolla longer than the calyx



c) Pedicel shorter than leaf length



d) The two fertile stamens

Figure 1. *Lindernia dubia* (L.) Pennell - overview of the habitat (a) and morphological aspects (b, c, d)

In Romania, *Lindernia dubia* (L.) Pennell have been identified for the first time in the Danube Delta by Ciocârlan & Costea, 1994, and is now one of the most abundant invasive species in the Danube Delta Nature Reserve (Doroftei & Anastasiu, 2014; Sîrbu et al., 2011; Anastasiu et al., 2014). *Lindernia* species are among the few annual species whose germination doesn't diminish in floodable conditions; *Lindernia dubia* (L.) Pennell germinates very well in the riparian areas of the rivers Loire and Allier

(France), but if the water layer persists the plants do not grow (Abernethy & Willby, 1999). *Lindernia dubia*'s capacity to germinate both in floodable conditions and in non-flooded soil is also highlighted by Neff et al. (2009), Šumberová et al. (2012) and http://beta.floranorthamerica.org/Lindernia_dubia⁴. In seasonally flooded habitats the ability to preempt the growth of flooding-sensitive annuals by germinating before water levels

⁴http://beta.floranorthamerica.org/Lindernia_dubia

have fully receded will bestow a considerable competitive advantage and may account for the present abundance of *Lindernia dubia* (L.) Pennell in the riparian zone of sections of the Loire and Allier (Abernethy & Willby, 1999). In the Czech Republic, Šumberová et al. (2005) discussing the two species, they appear in the marginal vegetation; *Lindernia dubia* (L.) Pennell regenerates earlier from the stock of seeds, while *Lindernia procumbens* (Krock.) Philcox appears at the end of June when there are high temperatures necessary for germination.

In this country, the species is also study by Horáková et al. (2005), Kaplan et al. (2016). In 1995, Seliškar et al. revised the *Lindernia* genus and first mentioned the *Lindernia dubia* (L.) Pennell species in the flora of Slovenia. In Slovakia, Hrivnák et al. (2016) discussing *Lindernia dubia* (L.) Pennell in its Central European context. In Bulgaria, both species grow in humid sandy places, in ponds and rice paddies. *Lindernia procumbens* (Krock.) Philcox is cited in 1889, *Lindernia dubia* (L.) Pennell appearing much later in 1984. Stojchev & Cheshmedziev (2005) investigate the anatomy of the two species and complete their chorology in Bulgaria with new data. The

anatomical characteristics remain in close parameters, the observed differences refer only to a greater density of the epidermal cells and the stomata from the *Lindernia procumbens* (Krock.) Philcox leaf and the higher number of cavities and medullary rays from the *Lindernia dubia* (L.) Pennell stem.

Pignatti (2005) considers as main ecological differences between the two species of *Lindernia* the fact that *Lindernia procumbens* (Krock.) Philcox has slightly higher requirements for light, slightly lower for temperature, and slightly lower for humidity, compared to *Lindernia dubia* (L.) Pennell. Ellenberg et al. (1992) present the two species as having the same light requirements (high) and *Lindernia procumbens* (Krock.) Philcox slightly more thermophilic.

Julve (2020a, 2020b) presents *Lindernia dubia* (L.) Pennell as a more thermophilic and more trophic soils. Both species are intolerant to salinity. Sanda et al. (2003), consider that there are no ecological differences between the populations of the two species in Romania, while Sârbu et al. (2013) say that *Lindernia dubia* (L.) Pennell needs less light and slightly higher temperature.




Figure 2. Geographical position of Surduc Lake

During some researches on the flora and vegetation around Surduc Lake (Timiș County,

W Romania, Figure 2), during 2005-2019 we followed the presence of invasive species

populations, given that this area is subject to a chaotic tourist development (Neacșu et al., 2017). *Lindernia dubia* (L.) Pennell is among the species we found here.

MATERIALS AND METHODS

We conducted field research in July-September, during 2005-2019. The identification of the species was made according to the current determinants in Romania (Ciocârlan, 2009, Sârbu et al., 2013), and according to Molnar et al. (2000) and Flora of China, 1998 (online edition). In the areas with populations of *Lindernia dubia* (L.) Pennell we also noticed the presence of other species of cormophytes. We also made observations on anthropogenic pressures. Nomenclature is according to Euro+Med PlantBase. Voucher specimens are stored in the Herbarium of Banat's University of

Agricultural Sciences and Veterinary Medicine "King Michael I of Romania" from Timisoara.

RESULTS AND DISCUSSIONS

We found populations of *Lindernia dubia* (L.) Pennell in six areas of the shores of Surduc Lake (Figure 3). Given the characteristics of the habitat (accumulation lake on the edge of a small river, with a narrow valley), two ways in which *Lindernia dubia* (L.) Pennell got here seem more likely: together with the fish brought here for breeding or by seeds in the mud encrusted in birds' feathers and legs (natural zoochory).

Both anthropic and natural *Lindernia dubia* (L.) Pennell dispersion is considered by various authors (e.g. Thiébault, 2007; Schmotzer, 2015). Simons & Jansen (2018) present as a way of introduction the zoochory.



Figure 3. Zones with *Lindernia dubia* (L.) Pennell populations on the Lake Surduc shore

We noticed that the morphology of *Lindernia dubia* (L.) Pennell specimens depends on the characteristics of the microhabitat:

a) Along the shore with clay soil we met abundant populations, with vigorous specimens, ample, ascending or slightly procumbent stems, which take root at the

nodes. When the soil is dry, it blooms. In autumn, the stems turn reddish. In these microhabitats are also found: *Eleocharis palustris* (L.) R. Br., *Eleocharis acicularis* (L.) Roemer et Schultes, *Lythrum portula* (L.) D. A. Webb, *Echinochloa crus-gali* (L.) Beauv., *Gypsophilla muralis* L., *Gnaphalium*

uliginosum L., *Polygonum aviculare* L., *Polygonum lapathifolium* L., *Xanthium orientale* subsp. *italicum* (Moretti) Greuter;

b) On the waterfront, in the sandy areas, *Lindernia dubia* (L.) Pennell grows together with: *Polygonum aviculare* L., *Trifolium repens* L., *Potentilla reptans* L., *Plantago media* L., *Bidens tripartita* L., *Alisma plantago-aquatica* L., *Echinochloa crus-galli* (L.) Beauv. These forms are vigorous, with ascending stems.

c) In the agricultural crops near the shore we found isolated specimens, small in size, with weak rooting.

In the whole studied area, the shores of the lake are occupied by populations of the following species: *Alisma plantago-aquatica* L., *Bidens tripartita* L., *Calystegia sepium* (L.) R.Br., *Carex riparia* Curtis, *Cyperus fuscus* L., *Echinochloa crus-galli* (L.) Beauv., *Eleocharis acicularis* (L.) Roemer et Schultes, *Eleocharis palustris* (L.) R. Br., *Gnaphalium uliginosum* L., *Gypsophila muralis* L., *Impatiens noli-tangere* L., *Juncus bufonius* L., *Juncus effusus* L., *Juncus tenuis* Willd., *Leersia oryzoides* (L.) Sw., *Leontodon autumnalis* L., *Lotus corniculatus* L., *Lysimachia vulgaris* L., *Lythrum hyssopifolia* L., *Lythrum salicaria* L., *Mentha pulegium* L., *Oenanthe aquatica* (L.) Poir., *Plantago media* L., *Polygonum aviculare* L., *Polygonum hydropiper* L., *Polygonum lapathifolium* L., *Polygonum persicaria* L., *Pulicaria vulgaris* Gaertn., *Ranunculus repens* L., *Rorippa amphibia* L. (Besser), *Rorippa sylvestris* L. (Besser), *Salix alba* L., *Salix cinerea* L., *Stachys palustris* L., *Trifolium pratense* L., *Trifolium repens* L., *Xanthium orientale* subsp. *italicum* (Moretti) Greuter.

In our research (Neacșu, 2008), we encountered on the shores of Lake Surduc and *Lindernia procumbens* (Krock.) Philcox and we included these phytocenoses in the vegetal association *Eleocharidetum acicularis* W. Koch 1926 em. Oberd. 1957, which we did not find later. Near the lake, *Lindernia procumbens* (Krock.) Philcox is also reported by Karácsonyi & Negrean (2012). From a phytosociological point of view, *Lindernia procumbens* (Krock.) Philcox is presented by Sârbu et al. (2013) as a characteristic species for *Isoëto-Nanojuncetea*, while *Lindernia dubia* (L.) Pennell is considered characteristic for *Nanocyperion*.

Mucina et al. (2016) presents the unit *Cypero-Lindernion dubiae* Müller-Stoll et Pietsch in T. Müller 1963 as synonym for *Eleocharition soloniensis* Philippi 1968, described as pioneer ephemeral rush-vegetation in temporarily flooded mesotrophic habitats of Central and Western Europe. *Lindernia dubia* (L.) Pennell is considered in Mucina et al. (2016) classification as a diagnostic species for the class *Oryzetea sativae* Miyawaki 1960 (www.synbioosys.alterra.nl/evc/).

CONCLUSIONS

In the paper, we signal a new place in Romania for *Lindernia dubia* (L.) Pennell: the shores of Surduc Lake. This habitat is represented by depression areas, flooded in spring, whose existence is conditioned by the water levels. When these areas are covered in water the species takes refuge on the nearby arable lands. In the habitat near Surduc Lake, *Lindernia dubia* (L.) Pennell grows alongside other cormophyte species, most frequently being accompanied by: *Echinochloa crus-galli* (L.) Beauv., *Eleocharis acicularis* (L.) Roemer et Schultes, *Gypsophila muralis* L., *Polygonum lapathifolium* L., *Alisma plantago-aquatica* L., *Bidens tripartita* L.

Due to the area occupied by the species on the lake shores and large population fluctuations from one year to another, as well as the status of the lake (protected and tourist area), we do not recommend drastic measures to manage the species, especially since there is a probability of reappears *Lindernia procumbens* (Krock.) Philcox. However, it is necessary to monitor the area and the lakes and ponds in the area, where *Lindernia dubia* (L.) Pennell could be spread by birds.

REFERENCES

- Abernethy, V.J., Willby, N.J. (1999). Changes along a disturbance gradient in the density and composition of propagule banks in floodplain aquatic habitats. *Plant Ecology*, 140, 177–190.
- Anastasiu, P., Negrean, G., Smarandache, D., Lițescu, S., Basnou, C. (2014). Neophytes in protected areas. Case study: The Danube Delta Biosphere Reserve. *Acta Horti Bot. Bucurest.*, 41, 41-68, at: <https://ahbb.unibuc.ro/wp-content/uploads/2019/04/Ahbb-2014-41-41-68.pdf>.

- Burkart, M. (2001). River corridor plants (Stromtalpflanzen) in Central European lowland: a review of a poorly understood plant distribution pattern. *Global Ecology & Biogeography*, 10, 449–468.
- Ciocârlan, V. (2009). Flora ilustrată a României: *Pteridophyta* et *Spermatophyta*. Bucharest, RO: Ceres Publishing House, 689.
- Ciocârlan, V., Costea, M. (1994). *Lindernia dubia* (L.) Pennell, specie nouă în flora României. *Analele științifice ale Institutului de Cercetare și Proiectare Delta Dunării*, 7-8.
- Cook, C.D.K. (1985). Range Extensions of Aquatic Vascular Plant Species. *Journal of Aquatic Plant Management*, 1-6.
- Csiky, J., Purger, D. (2008). Monitoring of plant species along the Drava river and in Baranja (Croatia). In: Purger, J. J. (ed.), *Biodiversity studies along the Drava river*, University of Pécs, Hungary, 13-56.
- Doroftei, M., Anastasiu, P. (2014). Potential Impacts of Climate Change on Habitats and Their Effects on Invasive Plant Species in Danube Delta Biosphere Reserve, Romania. In: Rannow, S., Neubert, M. (eds.), *Managing Protected Areas in Central and Eastern Europe Under Climate Change*, Springer, Dordrecht - Heidelberg - New York - London, 267-279.
- Drobnik J., Buchalik M. (2004). *Lindernia dubia* (Scrophulariaceae) – nowy gatunek we florze Polski. *Fragmenta Floristica et Geobotanica Polonica*, 11(1), 5-14.
- Ellenberg, H., Weber, H.E., Düll, W., Werner, W., Paulißen, D. (1992). Indicator values of plants in Central Europe. *Scripta Geobotanica*, 18, 119.
- Horáková, V., Šumberová, K., Lososová, Z., Fabšicová, M. (2005). *Lindernia dubia* in the Czech Republic: Distribution and Ecology. *8th International Conference on the Ecology and Management of Alien Plant Invasions*, “Alien Plant Species Biology And Distribution”, Katowice, Poland, at: http://www.sci.muni.cz/botany/vz/pdf/Horakova_Lindernia%20dubia%20in%20the%20Czech%20Republic_distribution%20and%20ecology.pdf.
- Hrivnák, R., Kochjarová, J., Šumberová, K., Schmotzer, A. (2016). Alien wetland annual *Lindernia dubia* (Scrophulariaceae): the first recently mentioned localities in Slovakia and their central European context. *Biologia*, Section Botany, 71/3, 281-286.
- Julve, Ph. (2020). *Lindernia dubia* (L.) Pennell. *Baseflor. Index botanique, écologique et chorologique de la flore de France*. Version 27 avril 2020. <https://telabotanica.org/projets/psytosociologie>.
- Julve, Ph. (2020). *Lindernia palustris* Hartmann. *Baseflor. Index botanique, écologique et chorologique de la flore de France*. Version 27 avril 2020. <https://telabotanica.org/projets/psytosociologie>.
- Kaplan, Z., Danihelka, J., Štěpánková, J., Ekrt, L., Chrtěk Jr., J., Zázvorka, J., Grulich, V., Řepka, R., Pranč, J., Ducháček, M., Kúr, P., Šumberová, K., Brůna, J. (2016). Distributions of vascular plants in the Czech Republic. Part 2, *Preslia*, 88, 229–322.
- Karácsonyi, K., Negrean, G. (2012). O pădure de stejar pedunculat cu *Molinia caerulea* din Câmpia de Vest a României. *Studii și Comunicări*, seria Științele Naturii, Satu-Mare, XIII, 7-14.
- Les, D.H. (2017). Aquatic Dicotyledons of North America: Ecology, Life History, and Systematics. *CRC Press Taylor & Francis Group*, Boca Raton – London – New York, 704-707.
- Marhold, K. (2017). *Linderniaceae*. In: *Euro+Med Plantbase-the information resource for Euro-Mediterranean plant diversity. (Lindernia procumbens)* <http://ww2.bgbm.org/EuroPlusMed/PTaxonDetail.asp?NameCache=Lindernia%20procumbens&PTRefFk=7200000>; *Lindernia dubia* - <http://ww2.bgbm.org/EuroPlusMed/PTaxonDetail.asp?NameCache=Lindernia%20dubia&PTRefFk=7200000>.
- Molnár, A., Pfeiffer, N., Ristow, M. (2000). Adatok hazai *Nanocyperon*-fajok ismeretéhez IV. A *Lindernia dubia* (L.) Pennel [Scrophulariaceae] Magyarországon / Data to the knowledge of *Nanocyperion*-species in Hungary IV. *Lindernia dubia* (L.) Pennel in Hungary, *Kitaibelia*, V(2), 279-287.
- Mucina, L., Bültmann, H., Dierssen, K., Theurillat, J.-P., Raus, T., Čarni, A., Šumberová, K., Willner, W., Dengler, J., Gavilán García, R., Chytrý, M., Hájek, M., Di Pietro, R., Iakushenko, D., Pallas, J., Daniëls, F.J.A., Bergmeier, E., Santos Guerra, A., Ermakov, N., Valachovič, M., Schaminée, J.H.J., Lysenko, T., Didukh, Y.P., Pignatti, S., Rodwell, J. S., Capelo, J., Weber, H. E., Solomeshch, A., Dimopoulos, P., Aguiar, C., Hennekens, S. M., Tichý, L. (2016). Vegetation of Europe: hierarchical floristic classification system of vascular plant, bryophyte, lichen, and algal communities. *Applied Vegetation Science*. Wiley, vol. 19, supp.1, 3-264.
- Neacșu, A., Arsene, G.-G., Arsene, A. (2017). Notes on the vascular flora of the Lake Surduc area. *Research Journal of Agricultural Science*, 49(3), 145-154.
- Neacșu, A.-G. (2008). Cercetări asupra florei și vegetației unor lacuri de acumulare din județul Timiș. (Research on the biodiversity of flora and vegetation in some accumulation lakes in Timis county). Banat University of Agricultural Sciences and Veterinary Medicine from Timisoara, Romania, Faculty of Agriculture. *Doctoral thesis*.
- Neff, K.P., Rusello, K., Baldwin, A.H. (2009). Rapid Seed Bank Development in Restored Tidal Freshwater Wetlands. *Restoration Ecology* 7(4), 539–548.
- Pignatti, S. (2005). Valori di bioindicazione delle piante vascolari della flora d'Italia. *Braun-Blanquetia*, 39, 58.
- Sanda, V., Biță-Nicolae, C.D., Barabaș, N. (2003). *Flora cormofitelor spontane și cultivate din România*. Bacău, RO: Ion Borcea Publishing House, 169.
- Sârbu, I., Ștefan, N., Oprea, A. (2013). *Plante vasculare din România. Determinator ilustrat de teren*. Bucharest, RO: Victor B Victor Publishing House, 700.

- Schmotzer, A. (2015). Occurrence of *Lindernia dubia* in the Ipoly Valley (Hungary and Slovakia). *Studia Botanica Hungarica*, 46(1), 77-89.
- Seliškar, A., Trpin, D., Vreš, B. (1995). Flora in vegetacija vlažnih rastišč Slovenije - I. Rod *Lindernia* All. (Flora and vegetation on wet habitats in Slovenia - I. Genus *Lindernia* All.), *Biološki vestnik*, 40(3-4), 45-58.
- Simons & Jansen (2018). Ecology of naturalized invasive species *Lindernia dubia* (L.). Pennell in the Netherlands, *Gorteria - Dutch Botanical Archives* 40. 001-010
- Sîrbu, C. (coord.), Oprea, A. (2011). *Plantele adventive în flora României*. Iași, RO: Ion Ionescu de la Brad Publishing House, 368-369.
- Sîrbu, C., Oprea, A., Eliáš, P., Jr. Ferus, P. (2011). New contribution to the study of alien Flora in Romania. *Journal of Plant Development*, 18, 121-134, at: <https://plant-journal.uaic.ro/docs/2011/17.pdf>
- Stevens, P.F. (2017). *Angiosperm Phylogeny. Website*, Version 14, July 2017, at: <http://www.mobot.org/MOBOT/research/APweb>, (2001 onwards).
- Stojchev, G., Cheshmedziev, I. (2005). Horological and anatomical investigation on *Lindernia procumbens* and *Lindernia dubia* (*Scrophulariaceae*). In: Gruev, B., Nikolova, M., Donev, A. (eds.), *Proceedings of the Balkan Scientific Conference of Biology*, Plovdiv (Bulgaria), 248-256.
- Šumberová, K., Horáková, V., Lososová, Z. (2005). Vegetation dynamics on exposed pond bottoms in the Českbudějovická basin (Czech Republic). *Phytocoenologia*, 35(2-3), 421-448.
- Šumberová, K., Lososová, Z., Ducháček, M., Horáková, V., Fabšičová, M. (2012). Distribution, habitat ecology, soil seed bank and seed dispersal of threatened *Lindernia procumbens* and alien *Lindernia dubia* (*Antirrhinaceae*) in the Czech Republic. *Phyton (Horn)*, 52(1), 39-72.
- Thiébaud, G. (2007). Non-indigenous aquatic and semi-aquatic plant species in France. In: Gherardi, F. (ed.), *Biological invaders in inland waters: Profiles, distribution, and threats*, Springer, Dordrecht, 209-229.
- Tomović, G., Vukojičić, S., Niketić, M., Lakušić, D. (2007). New chorological data on some threatened and rare plants in Serbia, *Archives of Biological Sciences*, Belgrade, 59(1), 3-73.
- Tutin, T.G., Heywood, V.H., Burges, N.A., Moore, D.M., Valentine, D.H., Walters, S.M., Webb, D.A. (eds.) (1972). *Flora Europaea*, 3, Cambridge University Press, 203-204.
- ****Lindernia dubia* (L.) Pennell - Lindernie fausse-gratiolle, Lindernie douteuse. Fiche réalisée par la Fédération des Conservatoires botaniques nationaux, Version 2, 2012, at: http://www.fcbn.fr/sites/fcbn.fr/files/ressource_telechargeable/fiche_-_lindernia_dubia_v2.pdf
- ****Lindernia dubia*. Species Factsheet, Delivering Alien Invasive Species Inventory in Europe (DAISIE), at: <http://www.europe-alien.org/speciesFactsheet.do?speciesId=19525#>
- ****Lindernia* Allioni, Flora of China, 18, 30-37, at: <http://flora.huh.harvard.edu/china/PDF/PDF18/lindernia.pdf>.
- ***Council Directive 92/43/EEC of 21 May 1992 on the conservation of natural habitats and of wild fauna and flora, at: <http://eur-lex.europa.eu/legal-content/EN/TXT/?uri=CELEX:01992L0043-20130701>.
- ***European Vegetation Classification at: www.synbiosys.alterra.nl/evc/
- ***Flora of North America, at: http://beta.floranorthamerica.org/Lindernia_dubia.
- ***The Euro+Med PlantBase, at: <https://ww2.bgbm.org/EuroPlusMed/PTaxonDetail.asp?NameCache=Lindernia&PTRefFk=7200000>.