



Allometric relations between DBH and sapwood area for predicting stand transpiration: lessons learned from the *Quercus* genus

R. Schoppach^{1,2} · K. P. Chun³ · J. Klaus⁴

Received: 15 June 2022 / Revised: 6 March 2023 / Accepted: 15 March 2023 / Published online: 29 March 2023
© The Author(s) 2023

Abstract

Catchment-scale transpiration is commonly determined by the use of sap-flow sensors, and its quantification, which is critical for water and forest management, relies crucially on the total catchment's sapwood area (A_s). Species-specific allometric relationships between the trees A_s and the trees diameter at breast height (DBH) are widely used for determining stand or catchment A_s . However, substantial differences between studies challenge the robustness of these relationships between sites displaying various topographical and environmental characteristics. Our objectives for this study are to compare the parameters of these relationships between species of the *Quercus* genus from different sites across the globe and to test the role of topographical factors on the A_s -DBH relationship in *Quercus petraea*. Using 145 trees sampled within a 0.455 km² catchment, we found that topography (slope, flow accumulation, aspect, curvature, and topographic wetness index) does not modulate the A_s -DBH relationship in *Q. petraea*, within our catchment. We compared our curve parameters with those from 16 studies on oak trees and found that the A_s -DBH relationship is not only species-specific, but depends on the site's conditions. The use of species-specific parameters from other sites may lead to more than 100% difference in the calculation of A_s , and therefore in forest transpiration. In the light of these results, we recommend building site- and species-specific A_s -DBH relationships for determining stand or catchment transpiration, using a minimum of nine, randomly sampled trees, and different methods and azimuthal directions for determining sapwood depth.

Keywords Sapwood area · Allometry · DBH · Topography · *Quercus* · Transpiration

Introduction

The estimation of catchment-scale transpiration remains critical for forest and water management. To date, a widely applied method is based on the use of sap-flow sensors

(Hassler et al. 2018; Tsuruta et al. 2019; Mitra et al. 2020). Different types of sensors exist, but most rely on the measurement of sap-flux density in a subset of trees that is most commonly multiplied by the sapwood area of the trees to obtain catchment-scale transpiration (Granier et al. 1996; Flo et al. 2019). Sapwood area therefore plays a crucial role in estimating tree and catchment transpiration (Cermak and Nadezhdina 1998; Hölscher et al. 2005; Meinzer et al. 2005), which motivated numerous studies on sapwood allometric relations (e.g. Wang et al. 2010; Quiñonez-Piñón and Valeo 2017; Lubczynski et al. 2017; Mitra et al. 2020). Commonly, stand sapwood area is derived from empirical relationships with the DBH of the stand's trees, because sampling all the trees in a catchment/stand is too resource intense (e.g. Hassler et al. 2018; Tsuruta et al. 2019). Given the importance of these empirical relationships for determining forest water use, they have been studied in a wide range of forested ecosystems and on various tree species like temperate broad-leaved species (Gebauer et al. 2008), boreal tree species (Quiñonez-Piñón and Valeo 2017),

Communicated by Ruediger Grote.

✉ R. Schoppach
remy.schoppach@gmail.com

- ¹ Catchment and Eco-Hydrology Research Group, Environmental Research and Innovation Department, Luxembourg Institute of Science and Technology, Belvaux, Luxembourg
- ² Earth and Life Institute, UCLouvain, Louvain-la-Neuve, Belgium
- ³ Department of Geography and Environmental Management, University of the West of England, Bristol, UK
- ⁴ Department of Geography, University of Bonn, Bonn, Germany

snow-dependent tree species (Mitra et al. 2020), tropical rain forest species (Granier et al. 1996; Motzer et al. 2005; Horna et al. 2011; Aparecido et al. 2016; Kunert et al. 2015; Moore et al. 2018), Amazonian tree species (Parolin et al. 2008; Aparecido et al. 2019), desert tree species (Lubczynski et al. 2017), mountainous tree species (Vertessy et al. 1995; Tsuruta et al. 2019) or Australian tree species (Wang et al. 2016) among others. The relationships can take the form of different mathematical equations (e.g. linear, exponential, and hyperbolic). For most species from the Fagaceae family, including the *Quercus* genus, the relation commonly takes the form of a power-law function (Eq. 1),

$$A_s = B_0 * DBH^{B_1} \quad (1)$$

where A_s is the sapwood area of the tree (cm²), DBH is the diameter at breast height (cm), and B_0 and B_1 are curve parameters.

Methods for determining these empirical relationships can be destructive, like the use of a wood disc taken at breast height (Miranda et al. 2009), or invasive, such as core analysis (Gebauer et al. 2008; Horna et al. 2011), dye-immersed (Parolin et al. 2008; Aparecido et al. 2019) or thermo-imaged fallen trees (Granier et al. 1994). Non-destructive methods also exist and are commonly based on electromagnetic imaging techniques (Čermák et al. 2004; Bieker and Rust 2010a, b; Wang et al. 2016; Benson et al. 2019; Salomón et al. 2020). The allometric relations of trees from the *Quercus* genus have been extensively studied (Table 1), due to its high natural abundance (Rüther and Walentowski 2008; Petritan et al. 2012) and a broad range of suitable growing conditions (Mette et al. 2013; Pretzsch et al. 2013). Additionally, the sapwood-heartwood limit from most *Quercus* trees can be distinguished straightforwardly by the darker colour of the heartwood (Mosedale et al. 1996; Taylor et al. 2002), which facilitates the quantification of A_s within this genus by direct inspection after coring.

Many eco-hydrological studies aiming at quantifying tree or stand water uses rely on so-called species-specific relationships (e.g. Chiu et al. 2016; Hassler et al. 2018; Schoppach et al. 2021). However, different studies on the same species have shown contrasting values for the B_0 and B_1 parameters (e.g. Aranda et al. 2005; Schmidt 2007; Grossiord et al. 2014), questioning the robustness of these relationships between sites (Horna et al. 2011) and raising concern on the level of difference in sapwood area, when measurements are up-scaled to the stand or catchment level using these parameter values. In the past decades, studies showed that the A_s -DBH allometric relationships of Japanese cedar and Japanese cypress may be affected by topographical factors as they vary with the slope position of the trees (Kumagai et al. 2007; Kume et al. 2016). These topographical effects may explain the inconsistencies of B_0 and

B_1 between studies. However, topographical effects on A_s -DBH relationships have never been explored in the *Quercus* genus, despite the wide use of this relationship for quantifying stand-scale transpiration, especially across Europe. To date, most forest transpiration studies at the catchment scale rely on B_0 and B_1 parameters from the literature. There is a need for testing the robustness of these parameters between sites, specific species of a given genus, and topographical factors, to provide guidance for researchers and practitioners to build strong A_s -DBH allometric relationships.

In this study, we aim to fill these gaps by (i) comparing A_s -DBH allometric relationships between species of the *Quercus* genus growing in diverse site conditions from 17 field studies, (ii) investigating the role of topographical factors (i.e. slope, flow accumulation, aspect, curvature, and topographic wetness index) on A_s -DBH allometric relationships of *Q. petraea*, and (iii) determining the minimum number of samples required for a robust quantification of the A_s -DBH power-law function for *Q. petraea* in a temperate European forest.

Material and methods

Study site

The Weierbach is a forested catchment of 0.455 km² located in north-western Luxembourg (lat 49.827, long 5.795). The forest is a mix of European beech (*Fagus sylvatica* L.) and pedunculate (*Q. robur* L.) and sessile oak (*Q. petraea* (Matt.) Liebl.) (Fabiani et al. 2021; Schoppach et al. 2021). Catchment's elevation ranges between 458 and 514 m asl, and the geology is composed of Devonian slate and phyllites covered by 60 cm of silt and rock fragments (Juilleret et al. 2011). Distributions of the catchment topographical characteristics can be found in Schoppach et al. (2021).

Measurement of sapwood depth in oak trees and landscape characteristics analysis

During the 2019 and 2020 growing seasons, we measured sapwood depth of oak trees within 145 randomly selected trees in the catchment (Fig. 1b). Sapwood depth was determined by drilling one single core per tree directly coring horizontally, in a randomized azimuthal direction, at breast height. All sampled oak trees are mature trees in the co-dominant strata. Sapwood and heartwood were discriminated based on colour variation (Mosedale et al. 1996; Quiñonez-Piñón and Valeo 2018). Bark was removed from the cores before the sapwood depth was measured. Here, we assumed that the sapwood depth was constant on all azimuthal directions, and the tree trunks were considered as perfect cylinder for calculating A_s (Gebauer et al. 2008; Hassler et al. 2018).

Table 1 Equation 1 parameters measured on oak trees (*Quercus* genus).

Study	Binomial name	B ₀	B ₁	R ²	RMSE	Covered area (m ²)	Trees	Data
Our study	<i>Quercus petraea</i>	0.3226	1.872	0.65	149	Within a 4.55 10 ⁶ m ² catchment	145	Raw
	<i>Quercus prinus</i> L.	0.375	1.751	0.93	–	Within a 1.92 10 ⁴ m ² plot	46	Eq
	<i>Quercus alba</i> L.	0.979	1.514	0.81	–	Within a 1.92 10 ⁴ m ² plot	45	Eq
	<i>Quercus rubra</i> L.	1.122	1.440	0.75	–	Within a 1.92 10 ⁴ m ² plot	60	Eq
Wullschlegel and Hanson (2006)*	<i>Quercus alba</i> L.	0.1021	1.5483	0.89	–	Within a 1.92 10 ⁴ m ² plot	24	Eq
	<i>Quercus prinus</i> L.	0.1146	1.7154	0.92	–	Within a 1.92 10 ⁴ m ² plot	24	Eq
	<i>Quercus rubra</i> L.	0.0875	1.5075	0.77	–	Within a 1.92 10 ⁴ m ² plot	24	Eq
	<i>Quercus garryana</i>	0.37	1.42	–	–	–	6	Eq
Poyatos et al. (2007)**	<i>Quercus pubescens</i>	4.16	0.66	–	–	78.5 m ² (Within a 10 m circle)	10	Eq
Limousin et al. (2009)	<i>Quercus ilex</i>	0.363	2.149	0.97	–	4.2 10 ² m ² (Taken outside of the three 140 m ² plots)	18	Eq
Grossiord et al. (2014)*	<i>Quercus cerris</i>	1.843	1.6351	0.90	–	2.7 10 ³ m ² (Selected right outside the three 30 * 30 m study plots)	21	Eq
Jung et al. (2011)	<i>Quercus petraea</i>	0.3653	1.8483	0.94	–	2.7 10 ³ m ² (idem)	48	Eq
	<i>Quercus mongolica</i>	0.1846	2.1	0.94	–	2.0 10 ³ m ² (Taken in the 40 m * 50 m plot)	9	Eq
Aranda et al. (2005)	<i>Quercus petraea</i>	0.90	1.7	0.92	–	(In a plot near the experimental site)	–	Eq
Salomón et al. (2020)	<i>Quercus ilex</i>	0.682	1.911	–	–	3.0 10 ² m ² (Within three plots of 10 × 10 m)	20	Eq
Benson et al. (2019)***	<i>Quercus virginia</i> Mill	0.1808	0.4788	0.97	5.99	–	19	Raw
	<i>Quercus nigra</i> L.	2.320	1.995	0.80	20.59	–	7	Raw
Martin et al. (1998)	<i>Quercus. coccinea</i> Muenchh	0.2544	1.7	0.80	21.66	(Within the Coweeta Basin)	5	Raw
Schmidt (2007)	<i>Quercus alba</i> L.	0.1148	1.973	0.88	39.03	(Idem)	10	Raw
	<i>Quercus prinus</i> L.	1.496	1.372	0.95	25.08	(Idem)	10	Raw
	<i>Quercus rubra</i> L.	0.1876	1.801	0.79	41.63	(Idem)	9	Raw
	<i>Quercus petraea</i>	0.0627	2.25	0.97	12.78	1.58 10 ⁴ m ² (Within a 1.29 and a 0.23 ha plot)	20	Raw
Study	Binomial name	Equation	R ²	Covered area (m ²)	Trees	Data		
Other relationship conformation than power-law function								
Bovard et al. (2005)	<i>Quercus rubra</i> L.	3.24 * DBH—10.24	0.90	2.3 10 ³ m ² (Within a 0.23 ha plot)	11	Eq		
Jonard et al. (2011)	<i>Quercus petraea</i>	9.442 * DBH –129.11	0.73	2.5 10 ³ m ² (Within a 0.23 ha plot)	16	Eq		

Table 1 (continued)

Study	Binomial name	Equation	R ²	Covered area (m ²)	Trees	Data
Wang et al. (2010)	<i>Quercus mongolica</i>	$\log_{10} A_s = -1.05(\log_{10} \text{DBH}) + 2.1$	0.95	–	8	Eq
Zhu et al. (2017)	<i>Quercus variabilis</i> <i>Blume</i>	$0.0083 * e^{0.0495 \text{DBH}}$	0.95	(Outside the experimental plots)	10	Eq

*These studies used Eq. 1 with A_s in m² and DBH in m

**This study used basal area instead of DBH in Eq. 1

***Not measured at breast height but at various heights in each tree (2.2 different heights per tree, on average)

“ B_0 ” and “ B_1 ” parameters correspond to Eq. 1 parameters. “RMSE” = Root Mean Square Error. “Covered area (m²)” indicates the area in which trees have been sampled, using the original papers formulation for clarifying when it is ambiguously or vaguely described. “Trees” indicates the number of sampled trees within each study. “Data” indicates if the equations parameters are provided in the study (Eq.) or if only raw data are provided (Raw), so we calculated B_0 and B_1 parameters

We used a high-resolution (1 m) digital elevation model (Luxembourgish air navigation administration 2017) and the Spatial Analysis Toolbox of ArcGIS Desktop 10.5 to determine the topographical characteristics at each sampling location.

We smoothed our raster maps by averaging topographical values of each one-square-metre grid plot with the values of the plots encompassed within a 10 m circle around it, to avoid the influence of small differences from the tree actual position to the analysis. The topographical characteristics within those 10 m circles were extracted for each sampled tree. Two relationships between sapwood depth and DBH were established based on a value criterion for each topographical characteristic. We selected these criteria to obtain roughly balanced (Table 2) and normally distributed groups in terms of DBH (Fig. 2), without significant differences in the means and variances (Table 2). Criteria, number of trees composing each group, and mean $\pm$ SE are displayed in Table 2. The slope values represent the steepness of the land surface. The flow accumulation represents the number of m² of converging area flowing into each downslope area. The aspect is the orientation of the down-hill slope. The curvature indicates if the surface is upwardly convex or concave. The topographic wetness index (TWI) was calculated as follows (Eq. 2):

$$\text{TWI} = \ln \left(\frac{(\text{flow ac.} + 1) * a}{\tan \left(\text{slope} * \frac{\pi}{180} \right) + 0.001} \right) \quad (2)$$

with *flow ac.* being the flow accumulation of a m², *a* being the size (1 m²), and slope being the *slope* value in degrees.

We tested the statistical differences between the A_s -DBH relationships derived from all pairs of datasets (two groups per topographical characteristic) using extra-sum-of-squares *F* tests and Akaike’s information criterions. For both approaches, we tested the probability that one model with identical B_0 and B_1 parameters fits the data as well as two models with different B_0 and B_1 values.

Comparison between A_s -DBH allometric relationships from the literature

For determining the significance of the difference in A_s produced by the application of the various B_1 and B_0 parameters from the literature, we applied Eq. 1 to our dataset of 145 DBH and compared the resulting A_s . We used one-way analysis of variance (ANOVA) to determine the significance ($p < 0.05$) of the species and site effect on the produced A_s . Given the obvious differences between relationships of different species, we focused our analysis on *Q. Petraea* datasets, which present




Fig. 1 Location of the study site within Luxembourg (a), location of the wood core samplings within the catchment (b), and 5 m contour lines (c). Panel B: Red circles represent the location of the trees sam-

pled in 2019, and blue circles represent the location of the trees sampled in 2020. Light green line delineates the catchment boundaries. (Color figure online)

Table 2 Groups characteristics for topographical investigation.

Criteria	Sample size	Mean DBH ($\pm$ SE)	p value t test ^a	p value F test ^b
Slope $< 5^\circ$	66	52.90 $\pm$ 1.34	0.83	0.36
Slope $> 5^\circ$	79	52.54 $\pm$ 1.10		
Aspect $< 160^\circ$	72	52.43 $\pm$ 1.29	0.75	0.22
Aspect $> 160^\circ$	73	52.97 $\pm$ 1.11		
Curvature $< 0^\circ$	60	53.77 $\pm$ 1.49	0.29	0.05
Curvature $> 0^\circ$	85	51.95 $\pm$ 1.00		
Flow accumulation $< 100 \text{ m}^b$	70	53.29 $\pm$ 1.31	0.51	0.25
Flow accumulation $> 100 \text{ m}^2$	75	52.16 $\pm$ 1.10		
TWI ^c	68	52.57 $\pm$ 1.22	0.88	0.77
TWI	77	52.82 $\pm$ 1.19		

^a p value of the unpaired t test between both groups

^b p value of the F test for comparing variances between both groups

^cTWI topographic wetness index

relatively similar curves, and ran a Tukey's multiple comparison test in order to test the significance of the site effect alone.

Power analysis to determine the minimum number of samples required

We used a power analysis to estimate the minimum sample size required to reasonably detect an effect of a given size (Maxwell 2000). The effect size is a number measuring the strength of the relationship between two variables in a sampling panel. In our case, the relationship links A_s to DBH. Our effect size estimation is based on Lehr's rule of thumb which states that the sample size n for a two-sided two-sample t test with power 80% ($\beta=0.2$) and significance level ($\alpha=0.05$) should be (Lehr 1992; Van Belle 2011):

$$n = \frac{16}{\Delta^2} \quad (3)$$

where Δ is the treatment difference to be detected in units of the standard deviation—the standardized difference. We used “*pwr*” R package (Champely et al. 2018) to calculate the number of samples needed to get at least the same level of coefficient of determination that we obtained with our 145 samples. As Eq. 1 is not linear, *pwr* was applied on the relation between A_s and pA_s (the predicted A_s based on Eq. 1). The coefficient of determination is based on a linear assumption which is not technically appropriate for our nonlinear study, but we use them for a rough performance reference.




Fig. 2 Topographical characteristics associated with each group of trees. Left panels display boxplots of the topographical characteristics with horizontal lines representing the minimum, mean and maximum values of DBH. Right panels display the DBH distribution within each group of trees. Solid line filled with red and dashed line not filled delineate the distribution of the different groups. Threshold values defining each group are displayed on each pane. (Color figure online)

Results

Effect of topographical characteristics on A_s -DBH relationships

According to the extra-sum-of-squares F test, among all tested topographical characteristics (i.e. slope, flow

accumulation, aspect, curvature, and TWI), none produced a significant effect on the A_s -DBH relationship (Fig. 3). The strongest effect of topographical factor in our dataset was the aspect, but with only a p value of 0.1 between the two models. Results are similar using the Akaike's information criterion. Except for the aspect, none of the topographical characteristics produced a likely effect.

Variations of the B_1 and B_0 coefficients between species of the *Quercus* genus and sites

We identified 16 studies in the literature in which the allometric A_s -DBH relationship was determined for oak trees (Table 1). Within these studies and our additional study, 14 relationships have the shape of a power-law function (Fig. 4), while two others were linear and one exponential (Table 1). The A_s -DBH relationship was established for 13 different species from the *Quercus* genus (Fig. 4a, Table 1). We found a highly significant (p value < 0.0001) effect of the studies, which includes species and sites effects, on the different A_s datasets calculated using B_1 and B_0 coefficients from the literature and our DBH dataset. The mean A_s calculated varies by two orders of magnitude (between 56.7 and 6592 cm²) using the most extreme study-specific parameters. Beside our study, four other studies determined the A_s -DBH relationship on *Q. petraea* (Aranda et al. 2005; Schmidt 2007; Jonard et al. 2011; Grossiord et al. 2014). Our study has, on average, seven times more samples than the previous studies (145 vs. an average of 20). Our A_s -DBH relationship presents a R^2 36% lower (0.65 vs an average of 0.88). The area covered by the sampling is almost a thousand time larger (904) than the mean area of all the other studies (Table 1). We found a highly significant site effect (p value < 0.0001—Fig. 5) on the calculated A_s of *Q. Petraea*. The mean A_s ranged from 368.5 ± 8.03 cm² when applying Jonard's B_0 and B_1 parameters to 1151 ± 35.0 cm² when applying Grossiord's parameters (Table 3).

Number of samples required for a reliable quantification of A_s -DBH relationship

The power tests indicated that at least nine samples are needed to reach the same level of correlation, or even better, than what we obtained with our 145 samples ($n = 8.7$, $r = 0.81$).

Fig. 3 Effect of the topographical characteristics on the A_s -DBH allometric relationship. Each circle represents one tree and the colours denote their group within each topographical characteristic. Black and red curves represent the A_s -DBH allometric relationship (Eq. 1) fitted to each group of data. (Color figure online)





Fig. 4 A_s -DBH allometric curves determined on *Quercus* trees. Panel **a** displays the A_s -DBH relationships of all *Quercus* species from literature, while panel **b** displays the A_s -DBH relationship on *Quercus petraea* only. Different colours represent different studies and species (see legend). In panel **b**, each grey circle represents one of our core measurements. (Color figure online)



Discussion

Topography does not modulate the allometric relation between DBH and sapwood area in *Q. petraea* at our study site

Topography does not influence the A_s -DBH allometric relationship in *Q. petraea* at the Weierbach study site. This contrasts with the results of Kumagai et al. (2007) and Kume et al. (2016) in Japanese cedar and Japanese cypress who found different relationships at different slope positions. Based on our analysis, we can argue that A_s -DBH




Fig. 5 Sapwood areas (A_s) calculated via Eq. 1 using different B_0 and B_1 parameters from *Q. petraea* and our 145 DBH dataset. The box extends from the 25th to 75th percentiles. The line in the middle of the box is plotted at the median. Whiskers represent 5th and 95th percentiles. Dots represent data outside of the 5–95 percentiles. Letters denote significant differences between groups

relationship in *Q. petraea* is robust, independent of the topographical position of the sampled trees at the study site. This is a major finding for the robustness of future upscaling analysis relying on this allometric relation. Further research on different species and in different topographical settings is undoubtedly needed as a way forward.

Allometric curves vary between species of the *Quercus* genus

The allometric A_s -DBH relationships are commonly described as “species-specific”. In our study, we show that

determining A_s using different *Quercus*’s B_0 and B_1 parameters from the literature produce highly significant differences. This site and species specificity of parameters led to a difference of two orders of magnitude in the calculated mean A_s . Tropical forests typically display a wide variety of species which commonly imposes the use of a single pair of parameters for determining A_s of a stand (e.g. Moore et al. 2018) with Eq. 1. For the species from the *Quercus* genus in temperate forests, we demonstrated that this method leads to under- or over-estimate A_s by up to hundred times. This massive difference confirms and reinforces the requirement of species-specific relationships when upscaling sap-flux density data from a subset of trees to stand or catchment scale.

Quercus petraea’s B_0 and B_1 parameters vary between sites

Our analysis showed that applying B_0 and B_1 parameters determined in different studies and at different sites on pedunculate oaks resulted in significantly different values of A_s (Table 3, Fig. 5). Using the parameters from other studies would induce a difference ranging from an overestimation of 107% (Grossiord et al. 2014) to an underestimation of 44% (Jonard et al. 2011) compared to the A_s determined with our own B_0 and B_1 . These major differences highlight the need for, not only species-specific, but also site-specific allometric relations when estimating sap-flow at stand or catchment level. Similar investigations on larger catchments would help evaluating how far B_0 and B_1 remain valid.

Interestingly, the use of B_0 and B_1 parameters from our study and Schmidt (2007) did not result in significant differences in A_s (Fig. 5). In Jonard et al. (2011), authors found a linear relationship, probably because of the too low DBH range of the sample trees (29–48 cm). Schmidt’s study was carried out in northern Bavaria, less than 300 km from Weierbach. Jonard’s study was carried out in the Belgian Ardennes, the northern site among the five, located within less than 100 km from Weierbach. Aranda’s study was carried out in Spain, and Grossiord’s study took place in Tuscany, both being much further south and in a clearly

Table 3 Meteorological and stand characteristics of the studies on *Quercus petraea*

Study	Mean annual temperature (°C)	Total annual precipitation (mm)	Tree density (Tree/ha)	Mean A_s ($\pm$ SE)**
Our study	8.9	815	301	555.9 $\pm$ 17.1
Aranda et al. (2005)	8.1	1105	867	777.8 $\pm$ 21.7
Schmidt (2007)	7.7	800	350 and 636*	494.0 $\pm$ 18.6
Jonard et al. (2011)	9.8	1044	468	368.5 $\pm$ 8.03
Grossiord et al. (2014)	13.5	950	867	1151 $\pm$ 35.0

*Two plots with two densities mentioned

**Mean A_s determined by applying B_0 and B_1 parameters from each study to our database of 145 DBH values

different climate than the two first cited. One explanation for the significantly different curve parameters within the same species could therefore be the growing conditions effect on the trees allometry. Indeed, higher water demand and lower water availability have been pointed out as a driver of defoliation, which subsequently affects the proportion of sapwood area (e.g. Bréda et al. 2006; Limousin et al. 2012). In Scots pine, summer vapour pressure deficit and maximum temperature have been shown to be negatively correlated with the leaf area/sapwood area ratio across Europe (Mencuccini and Bonosi 2001). Sapwood proportion is suspected to depend on the availability of light (Sellin 1994; Thurner et al. 2019) and the water demand (Gebauer et al. 2008; Horna et al. 2011). Trees exposed to higher level of light and potential evapotranspiration require to maximize sapwood area for sustaining the higher evaporative demand and thus preventing leaf overheating and dehydration (Aparecio et al. 2019). Importantly, the hydraulic conductivity of sapwood is also influenced by growth rate which depends on resources availability and demand (Medhurst and Beadle 2002). The total annual precipitation amount, the mean annual temperature, and the tree densities varies between the five above-mentioned studies on *Q. petraea* (Table 2). Total radiation received by each site and the associated potential evapotranspiration are not available in the original papers. However, Spanish and Tuscan sites are likely to receive higher level of annual radiation and are exposed to higher evaporative demands due to their southern locations compared with Luxembourgish, Belgian, and German sites (Cornes et al. 2018). Sites with lower tree densities present lower sapwood area for a given DBH, which contrasts with the findings of Benyon et al. (2015) in eucalypt forests, showing a decrease in sapwood thickness with an increase in forest density. Higher forest density is supposed to reduce the exposition to high radiation and VPD conditions, which influences the development of sapwood. Here, we cannot decipher the various effects of the growing conditions on the A_s -DBH relationship based on these annual data. Further research is certainly needed to unravel the role of temperature, radiation, water demand, water availability, and forest density on the A_s -DBH allometric relationship on *Q. petraea*.

Number of samples required for a reliable quantification of A_s -DBH relationship

For upscaling purposes, a reliable quantification of the A_s -DBH allometric relationship remains crucial as it appeared that Eq. 1 parameters are constrained not only by species, but also by sites/stand characteristics. We thus recommend to experimentally determine Eq. 1 parameters for each site rather than using values from literature. According to our power test, at least nine samples are needed for building a robust A_s -DBH relationship. Importantly, the sampled trees

should cover a range of DBH that is as wide as possible to avoid misinterpretation of the curve shape (e.g. Jonard et al. 2011). This number of samples is close to or lower than the number of samples most studies on *Quercus* genus used (Table 1). Therefore, Eq. 1 parameters extracted from these studies can be considered as robust for the species and sites concerned. However, the A_s -DBH allometric relation calculated in this study displays a substantially lower R^2 compared to previous studies, which potentially pools down the required number of samples.

Confounding effects and methodological caveats

In this study, we were not able to highlight significant topographical effects on the A_s -DBH relationship of *Q. petraea*. Also, we determined an A_s -DBH relationship displaying a substantially lower R^2 compared to literature. Confounding effects may be responsible for the lack of significance of the topographical effects and the lower R^2 . Confounding effects may include a much larger sampling area than other studies with higher unexplained variances and random errors, a higher number of sampled trees, a wider range of tree's DBH, or methodological limitations.

A methodological limitation that may introduce some deviation of the measurement from the A_s -DBH curve is that A_s of each tree was inferred from a single core sampled at a random azimuth. We assumed the sapwood depth to be constant on all azimuths (Čermák et al. 2004; Tsuruta et al. 2010; Benson et al. 2019). Sapwood area was indeed inferred from a single core taken at random azimuth because we had no authorization from the forest owner to core each tree more than once during the experiment. However, the radial profile of sapwood thickness is suspected to be variable (e.g. Looker et al. 2016), especially in regard to slope (Pilate et al. 2004). Indeed, trees growing on a slope will produce tension wood to counteract the forces of gravity (Scurfield 1973; Pilate et al. 2004). This alters the production of wood, and presumably the sapwood thickness around the circumference of the tree, which could be a substantial confounding effect. The variability in radial sap-flux densities was found to be considerable in temperate broad-leaved species (Gebauer et al. 2008). For boreal trees, radial profile of sapwood thickness was shown to be dependent on the species; some species displaying constant sapwood depth, while others are growing thicker on the North-East side (Quiñonez-Piñón and Valeo 2017). For trembling aspen and white spruce, the influence of azimuthal direction on sapwood depth had a low impact (Merlin et al. 2020). In oak trees, the xylem vessel size was found to be larger on the northern side of *Quercus suber*, compared with the southern side (Barij et al. 2011), but, to our knowledge, the influence of azimuthal direction on sapwood depth itself remains unexplored in *Q. petraea*.

Topographical effect may also be confounded by the threshold values selected for comparing groups, which was imposed by catchment characteristics and done in such a way to force similar sample sizes, but was not necessarily tied to meaningful physical attributes. Further research on catchments displaying larger ranges of topographical characteristics could help untangling these effects.

Within studies on *Q. petraea* displaying power-law functions, Aranda et al. (2005), Schmidt (2007), and Grossiord et al. (2014) had a difference between their smaller and larger DBH of 14, 20, and 36 cm, respectively. Our dataset displayed a difference of 61 cm and our number of sampled trees was substantially higher. Also, our panel encompassed many trees with large DBH, while both Schmidt (2007) and Grossiord et al. (2014) had no trees larger than 50 cm. This value falls at 19 cm for Aranda et al. (2005). However, excluding the 85 trees larger than 50 cm in DBH from our dataset led to a substantially lower R^2 (0.28) of the A_s -DBH relationship. Larger range of DBH within the sampled trees therefore did not explain the low R^2 value displayed by our relationship.

Our dataset presents sapwood depth values varying between 10 and 86 mm with a mean of 42 mm. If we consider a deviation of 1 mm due to a recurrent measurement error, this one mm deviation would result in a deviation of 2.17% (14 cm²) of the A_s of a tree with average DBH (53 cm). For our smallest tree (DBH = 29.3 cm), the deviation increases to 9% (8.5 cm²). However, the potential effect of a systematic measurement mistake remains far too weak for explaining the observed spread of the points around the allometric curve (Fig. 3b). Dyeing experiments suggest that A_s is likely susceptible to be overestimated by the determination solely on heartwood coloration (Aparecido et al. 2019), despite other studies contradicting this suggestion (e.g. Githiomi and Dougal 2012). Experiments using dye are able to differentiate between observable sapwood and the functional conducting sapwood, the latter being typically lower. For this reason, we recommend the use of variable methods to validate the delimitation of sapwood.

The influence of the sampling area cannot be ignored. Most studies presented in Table 1 aimed at quantifying stand transpiration, and their main objectives was not to determine the A_s -DBH allometric relationship, which is commonly done only on few trees outside the field setup. In our experiment, we sampled within a catchment of 0.455 km², which is by far wider than all previous studies on oak trees. Therefore, our A_s -DBH curve is likely to be influenced by non-accounted micro-climatic conditions, influencing the resources availability (i.e. water and radiation) and the evaporative demand.

Conclusion

Sapwood area plays a key role for determining sap-flow of a tree and upscaling sap-flow measurements from tree to stand or catchment level. A reliable quantification of sapwood area is therefore crucial for water and forest management. In this study, we confirmed that the A_s -DBH allometric relationship of the *Quercus* genus is not only species-specific, but depends on the site conditions. We showed that topography does not modulate the A_s -DBH relationship in *Q. petraea*, within our catchment. In the light of these results, we emphasize that each upscaling study should rely on its own site- and species-specific relationship and recommend a minimum of nine, randomly sampled trees, in order to build a robust A_s -DBH relationship. Due to the potentially high variability in radial profile of sapwood thickness and the caveats related to colour distinction, we recommend coring at different azimuthal locations in the trees and validating the delimitation of sapwood via different methods. Further research is certainly needed for evaluation on how far B_0 and B_1 parameters remain valid.

Acknowledgements This work was supported by the Luxembourg National Research Fund (FNR/CORE/C17/SR/11702136/EFFECT). KP Chun is supported by the Accelerator Programme (AP) 2022-24 and the Starter Scheme by the University of West of England, Bristol. We thank the members of the Catchment and Eco-hydrology group at LIST, and especially Jean-François Iffly and Jérôme Juilleret for their support in field work. We thank the Luxembourgish air navigation administration for providing the Digital Elevation Model.

Author contributions RS contributed to conceptualization, investigation, data analysis, data visualization, and writing original draft and editing. KPC performed data analysis and reviewing and editing. JK contributed to funding acquisition, supervision, reviewing and editing, and project administration.

Funding This work was supported by the Luxembourg National Research Fund (FNR/CORE/C17/SR/11702136/EFFECT).

Data availability The data used in this study are the property of the Luxembourg Institute of Science and Technology (LIST) and can be obtained upon request to the corresponding author, after approval by LIST.

Code availability Not applicable

Declarations

Competing interests The authors declare no competing interests.

Conflict of interest The authors declare that they have no known competing financial interests or personal relationships that could appear to influence the work reported in this paper.

Consent for publication Not applicable

Ethical approval Not applicable

Consent to participate Not applicable

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Aparecido LMT, Miller GR, Cahill AT, Moore GW (2016) Comparison of tree transpiration under wet and dry canopy conditions in a Costa Rican premontane tropical forest. *Hydrol Process* 30(26):5000–5011. <https://doi.org/10.1002/hyp.10960>
- Aparecido LMT, Santos JD, Higuchi N, Kunert N (2019) Relevance of wood anatomy and size of Amazonian trees in the determination and allometry of sapwood area. *Acta Amazon* 49:1–10. <https://doi.org/10.1590/1809-4392201800961>
- Aranda I, Gil L, Pardos JA (2005) Seasonal changes in apparent hydraulic conductance and their implications for water use of European beech (*Fagus sylvatica* L.) and sessile oak [*Quercus petraea* (Matt.) Liebl] in South Europe. *Plant Ecol* 179(2):155–167. <https://doi.org/10.1007/s11258-004-7007-1>
- Barij N, Čermák J, Stokes A (2011) Azimuthal variations in xylem structure and water relations in cork oak (*Quercus suber*). *IAWA J* 32(1):25–40. <https://doi.org/10.1163/22941932-90000040>
- Benson AR, Koeser AK, Morgenroth J (2019) Estimating conductive sapwood area in diffuse and ring porous trees with electronic resistance tomography. *Tree Physiol* 39(3):484–494. <https://doi.org/10.1093/treephys/tpy092>
- Benyon RG, Lane PNJ, Jaskierniak D, Kuczera G, Haydon SR (2015) Use of a forest sapwood area index to explain long-term variability in mean annual evapotranspiration and streamflow in moist eucalypt forests. *Water Resour Res* 51(7):5318–5331. <https://doi.org/10.1002/2015WR017321>
- Bieker D, Rust S (2010a) Non-destructive estimation of sapwood and heartwood width in Scots pine (*Pinus sylvestris* L.). *Silva Fenn* 44(2):267–273. <https://doi.org/10.14214/sf.153>
- Bieker D, Rust S (2010b) Electric resistivity tomography shows radial variation of electrolytes in *Quercus robur*. *Can J for Res* 40(6):1189–1193. <https://doi.org/10.1139/x10-076>
- Bovard BD, Curtis PS, Vogel CS, Su HB, Schmid HP (2005) Environmental controls on sap flow in a northern hardwood forest. *Tree Physiol* 25(1):31–38. <https://doi.org/10.1093/treephys/25.1.31>
- Bréda N, Huc R, Granier A, Dreyer E (2006) Temperate forest trees and stands under severe drought: a review of ecophysiological responses, adaptation processes and long-term consequences. *Ann For Sci* 63(6):625–644. <https://doi.org/10.1051/forest:2006042>
- Čermák J, Kučera J, Nadezhdina N (2004) Sap flow measurements with some thermodynamic methods, flow integration within trees and scaling up from sample trees to entire forest stands. *Trees* 18(5):529–546. <https://doi.org/10.1007/s00468-004-0339-6>
- Cermak J, Nadezhdina N (1998) Sapwood as the scaling parameter-defining according to xylem water content or radial pattern of sap flow? In: *Annales des sciences forestieres*, vol 55, No. 5. EDP Sciences, pp 509–552. <https://doi.org/10.1051/forest:19980501>
- Champely S, Ekstrom C, Dalgaard P, Gill J, Weibelzahl S, Anandkumar A, De Rosario MH (2018) Package ‘pwr’. R package version. 1(2)
- Chiu CW, Komatsu H, Katayama A, Otsuki K (2016) Scaling-up from tree to stand transpiration for a warm-temperate multi-specific broadleaved forest with a wide variation in stem diameter. *J For Res* 21(4):161–169. <https://doi.org/10.1007/s10310-016-0532-7>
- Cornes RC, van der Schrier G, van den Besselaar EJ, Jones PD (2018) An ensemble version of the E-OBS temperature and precipitation data sets. *J Geophys Res Atmos* 123(17):9391–9409. <https://doi.org/10.1029/2017JD028200>
- Fabiani G, Schoppach R, Penna D, Klaus J (2021) Transpiration patterns and water use strategies of beech and oak trees along a hillslope. *Ecohydrology* 15(2):e2382. <https://doi.org/10.1002/eco.2382>
- Flo V, Martinez-Vilalta J, Steppe K, Schuldt B, Poyatos R (2019) A synthesis of bias and uncertainty in sap flow methods. *Agric For Meteorol* 271:362–374. <https://doi.org/10.1016/j.agrformet.2019.03.012>
- Gebauer T, Horna V, Leuschner C (2008) Variability in radial sap flux density patterns and sapwood area among seven co-occurring temperate broad-leaved tree species. *Tree Physiol* 28(12):1821–1830. <https://doi.org/10.1093/treephys/28.12.1821>
- Githiomi JK, Dougal E (2012) Analysis of heartwood–sapwood demarcation methods and variation of sapwood and heartwood within and between 15 year old plantation grown *Eucalyptus regnans*. *Int J Appl Sci Technol* 2(8)
- Granier A, Anfodillo T, Sabatti M, Cochard H, Dreyer E, Tomasi M, Bréda N (1994) Axial and radial water flow in the trunks of oak trees: a quantitative and qualitative analysis. *Tree Physiol* 14(12):1383–1396. <https://doi.org/10.1093/treephys/14.12.1383>
- Granier A, Biron P, Bréda N, Pontallier JY, Saugier B (1996) Transpiration of trees and forest stands: short and long-term monitoring using sapflow methods. *Glob Change Biol* 2(3):265–274. <https://doi.org/10.1111/j.1365-2486.1996.tb00078.x>
- Grossiord C, Gessler A, Granier A, Pollastrini M, Bussotti F, Bonal D (2014) Interspecific competition influences the response of oak transpiration to increasing drought stress in a mixed Mediterranean forest. *For Ecol Manage* 318:54–61. <https://doi.org/10.1016/j.foreco.2014.01.004>
- Hassler SK, Weiler M, Blume T (2018) Tree-, stand- and site-specific controls on landscape-scale patterns of transpiration. *Hydrol Earth Syst Sci* 22(1):13–30. <https://doi.org/10.5194/hess-22-13-2018>
- Hölscher D, Koch O, Korn S, Leuschner C (2005) Sap flux of five co-occurring tree species in a temperate broad-leaved forest during seasonal soil drought. *Trees* 19(6):628–637. <https://doi.org/10.1007/s00468-005-0426-3>
- Horna V, Schuldt B, Brix S, Leuschner C (2011) Environment and tree size controlling stem sap flux in a perhumid tropical forest of Central Sulawesi Indonesia. *Ann For Sci* 68(5):1027–1038. <https://doi.org/10.1007/s13595-011-0110-2>
- Jonard F, André F, Ponette Q, Vincke C, Jonard M (2011) Sap flux density and stomatal conductance of European beech and common oak trees in pure and mixed stands during the summer drought of 2003. *J Hydrol* 409(1–2):371–381. <https://doi.org/10.1016/j.jhydrol.2011.08.032>
- Juilleret J, Ifly JF, Pfister L, Hissler C (2011) Remarkable Pleistocene periglacial slope deposits in Luxembourg (Oesling): pedological implication and geosite potential. *Bull Soc Nat Luxemb* 112:125–130
- Jung EY, Otieno D, Lee B, Lim JH, Kang SK, Schmidt MWT, Tenhunen J (2011) Up-scaling to stand transpiration of an Asian temperate mixed-deciduous forest from single tree sapflow measurements. *Plant Ecol* 212(3):383–395. <https://doi.org/10.1007/s11258-010-9829-3>

- Kumagai TO, Aoki S, Shimizu T, Otsuki K (2007) Sap flow estimates of stand transpiration at two slope positions in a Japanese cedar forest watershed. *Tree Physiol* 27(2):161–168. <https://doi.org/10.1093/treephys/27.2.161>
- Kume T, Tsuruta K, Komatsu H, Shinohara Y, Katayama A, Ide JI, Otsuki K (2016) Differences in sap flux-based stand transpiration between upper and lower slope positions in a Japanese cypress plantation watershed. *Ecohydrology* 9(6):1105–1116. <https://doi.org/10.1002/eco.1709>
- Kunert N, Aparecido LMT, Higuchi N, dos Santos J, Trumbore S (2015) Higher tree transpiration due to road-associated edge effects in a tropical moist lowland forest. *Agric For Meteorol* 213:183–192. <https://doi.org/10.1016/j.agrformet.2015.06.009>
- Lehr R (1992) Sixteen S-squared over D-squared: a relation for crude sample size estimates. *Stat Med* 11(8):1099–1102. <https://doi.org/10.1002/sim.4780110811>
- Limousin JM, Rambal S, Ourcival JM, Rocheteau A, Joffre R, Rodriguez-Cortina R (2009) Long-term transpiration change with rainfall decline in a Mediterranean *Quercus ilex* forest. *Glob Change Biol* 15(9):2163–2175. <https://doi.org/10.1111/j.1365-2486.2009.01852.x>
- Limousin JM, Rambal S, Ourcival JM, Rodriguez-Calcerrada J, Perez-Ramos I, Rodriguez-Cortina R et al (2012) Morphological and phenological shoot plasticity in a Mediterranean evergreen oak facing long-term increased drought. *Oecologia* 169(2):565–577. <https://doi.org/10.1007/s00442-011-2221-8>
- Looker N, Martin J, Jencso K, Hu J (2016) Contribution of sapwood traits to uncertainty in conifer sap flow as estimated with the heat-ratio method. *Agric For Meteorol* 223:60–71. <https://doi.org/10.1016/j.agrformet.2016.03.014>
- Lubczynski MW, Chavarro-Rincon DC, Rossiter DG (2017) Conductive sapwood area prediction from stem and canopy areas—allometric equations of Kalahari trees Botswana. *Ecohydrology* 10(6):e1856. <https://doi.org/10.1002/eco.1856>
- Luxembourgish air navigation administration (2017) <https://data.public.lu/fr/datasets/digital-surface-model-high-dem-resolution/> Downloaded. Accessed 15 May 2020
- Martin JG, Kloeppel BD, Schaefer TL, Kimbler DL, McNulty SG (1998) Aboveground biomass and nitrogen allocation of ten deciduous southern Appalachian tree species. *Can J For Res* 28(11):1648–1659. <https://doi.org/10.1139/x98-146>
- Maxwell SE (2000) Sample size and multiple regression analysis. *Psychol Methods* 5(4):434. <https://doi.org/10.1037/1082-989X.5.4.434>
- Medhurst JL, Beadle CL (2002) Sapwood hydraulic conductivity and leaf area—sapwood area relationships following thinning of a *Eucalyptus nitens* plantation. *Plant Cell Environ* 25:1011–1019. <https://doi.org/10.1046/j.1365-3040.2002.00880.x>
- Meinzer FC, Bond BJ, Warren JM, Woodruff DR (2005) Does water transport scale universally with tree size? *Funct Ecol* 19(4):558–565. <https://doi.org/10.1111/j.1365-2435.2005.01017.x>
- Mencuccini M, Bonosi L (2001) Leaf/sapwood area ratios in Scots pine show acclimation across Europe. *Can J For Res* 31:442–456. <https://doi.org/10.1139/x00-173>
- Merlin M, Solarik KA, Landhäusser SM (2020) Quantification of uncertainties introduced by data-processing procedures of sap flow measurements using the cut-tree method on a large mature tree. *Agric For Meteorol* 287:107926. <https://doi.org/10.1016/j.agrformet.2020.107926>
- Mette T, Dolos K, Meinardus C, Bräuning A, Reineking B, Blaschke M, Wellstein C (2013) Climatic turning point for beech and oak under climate change in Central Europe. *Ecosphere* 4(12):1–19. <https://doi.org/10.1890/ES13-00115.1>
- Miranda I, Gominho J, Pereira H (2009) Variation of heartwood and sapwood in 18-year-old *Eucalyptus globulus* trees grown with different spacings. *Trees* 23(2):367–372. <https://doi.org/10.1007/s00468-008-0285-9>
- Mitra B, Papuga SA, Alexander MR, Swetnam TL, Abramson N (2020) Allometric relationships between primary size measures and sapwood area for six common tree species in snow-dependent ecosystems in the Southwest United States. *J Res* 31(6):2171–2180. <https://doi.org/10.1007/s11676-019-01048-y>
- Moore GW, Orozco G, Aparecido LM, Miller GR (2018) Upscaling transpiration in diverse forests: insights from a tropical premontane site. *Ecohydrology* 11(3):e1920. <https://doi.org/10.1002/eco.1920>
- Mosedale JR, Charrier B, Janin G (1996) Genetic control of wood colour, density and heartwood ellagitannin concentration in European oak (*Quercus petraea* and *Q. robur*). *For an Int J for Res* 69(2):111–124. <https://doi.org/10.1093/forestry/69.2.111>
- Motzer T, Munz N, Küppers M, Schmitt D, Anhof D (2005) Stomatal conductance, transpiration and sap flow of tropical montane rain forest trees in the southern Ecuadorian Andes. *Tree Physiol* 25(10):1283–1293. <https://doi.org/10.1093/treephys/25.10.1283>
- Parolin P, Müller E, Junk WJ (2008) Sapwood area in seven common tree species of Central Amazon floodplains. *Area Pesqui Bot* 59:277–286
- Petrutan AM, Biris IA, Merce O, Turcu DO, Petrutan IC (2012) Structure and diversity of a natural temperate sessile oak (*Quercus petraea* L.)—European Beech (*Fagus sylvatica* L.) forest. *For Ecol Manage* 280:140–149. <https://doi.org/10.1016/j.foreco.2012.06.007>
- Pilate G, Chabbert B, Cathala B, Yoshinaga A, Leplé JC, Laurans F, Ruel K (2004) Lignification and tension wood. *CR Biol* 327(9–10):889–901. <https://doi.org/10.1016/j.crv.2004.07.006>
- Poyatos R, Čermák J, Llorens P (2007) Variation in the radial patterns of sap flux density in pubescent oak (*Quercus pubescens*) and its implications for tree and stand transpiration measurements. *Tree Physiol* 27(4):537–548. <https://doi.org/10.1093/treephys/27.4.537>
- Pretzsch H, Schütze G, Uhl E (2013) Resistance of European tree species to drought stress in mixed versus pure forests: evidence of stress release by inter-specific facilitation. *Plant Biol* 15(3):483–495. <https://doi.org/10.1111/j.1438-8677.2012.00670.x>
- Quiñonez-Piñón MR, Valeo C (2017) Allometry of sapwood depth in five boreal trees. *Forests* 8(11):457. <https://doi.org/10.3390/f8110457>
- Quiñonez-Piñón MR, Valeo C (2018) Assessing the translucence and color-change methods for estimating sapwood depth in three boreal species. *Forests* 9(11):686. <https://doi.org/10.3390/f9110686>
- Rüther C, Walentowski H (2008) Tree species composition and historic changes of the Central European oak/beech region. In: *Canopy arthropod research in Europe*, pp 61–88
- Salomón RL, Steppe K, Ourcival JM, Villers S, Rodríguez-Calcerrada J, Schapman R, Limousin JM (2020) Hydraulic acclimation in a Mediterranean oak subjected to permanent throughfall exclusion results in increased stem hydraulic capacitance. *Plant Cell Environ* 43(6):1528–1544. <https://doi.org/10.1111/pce.13751>
- Schmidt M (2007) Canopy transpiration of beech forests in Northern Bavaria—Structure and function in pure and mixed stands with oak at colline and montane sites. Doctoral dissertation
- Schoppach R, Chun KP, He Q, Fabiani G, Klaus J (2021) Species-specific control of DBH and landscape characteristics on tree-to-tree variability of sap velocity. *Agric For Meteorol* 307:108533. <https://doi.org/10.1016/j.agrformet.2021.108533>
- Scurfield G (1973) Reaction Wood: its Structure and Function: lignification may generate the force active in restoring the trunks of leaning trees to the vertical. *Science* 179(4074):647–655. <https://doi.org/10.1126/science.179.4074.647>

- Sellin A (1994) Sapwood–heartwood proportion related to tree diameter, age, and growth rate in *Picea abies*. *Can J For Res* 24(5):1022–1028. <https://doi.org/10.1139/x94-133>
- Taylor AM, Gartner BL, Morrell JJ (2002) Heartwood formation and natural durability—a review. *Wood Fiber Sci* 34:587–611
- Thurner M, Beer C, Crowther T, Falster D, Manzoni S, Prokushkin A, Schulze ED (2019) Sapwood biomass carbon in northern boreal and temperate forests. *Glob Ecol Biogeogr* 28(5):640–660. <https://doi.org/10.1111/geb.12883>
- Tsuruta K, Kume T, Komatsu H, Higashi N, Umebayashi T, Kumagai TO, Otsuki K (2010) Azimuthal variations of sap flux density within Japanese cypress xylem trunks and their effects on tree transpiration estimates. *J For Res* 15(6):398–403. <https://doi.org/10.1007/s10310-010-0202-0>
- Tsuruta K, Komatsu H, Kume T, Otsuki K, Kosugi Y, Kosugi KI (2019) Relationship between stem diameter and transpiration for Japanese cypress trees: implications for estimating canopy transpiration. *Ecohydrology* 12(5):e2097. <https://doi.org/10.1002/eco.2097>
- Van Belle G (2011) Statistical rules of thumb, vol 699. Wiley
- Vertessy RA, Benyon RG, O'sullivan SK, Gribben PR (1995) Relationships between stem diameter, sapwood area, leaf area and transpiration in a young mountain ash forest. *Tree Physiol* 15(9):559–567. <https://doi.org/10.1093/treephys/15.9.559>
- Wang X, Wang C, Zhang Q, Quan X (2010) Heartwood and sapwood allometry of seven Chinese temperate tree species. *Ann For Sci* 67(4):410. <https://doi.org/10.1051/forest/2009131>
- Wang H, Guan H, Guyot A, Simmons CT, Lockington DA (2016) Quantifying sapwood width for three Australian native species using electrical resistivity tomography. *Ecohydrology* 9(1):83–92. <https://doi.org/10.1002/eco.1612>
- Wullschlegel SD, Hanson PJ (2006) Sensitivity of canopy transpiration to altered precipitation in an upland oak forest: evidence from a long-term field manipulation study. *Glob Change Biol* 12(1):97–109. <https://doi.org/10.1111/j.1365-2486.2005.001082.x>
- Wullschlegel SD, Hanson PJ, Todd DE (2001) Transpiration from a multi-species deciduous forest as estimated by xylem sap flow techniques. *For Ecol Manag* 143(1–3):205–213. [https://doi.org/10.1016/S0378-1127\(00\)00518-1](https://doi.org/10.1016/S0378-1127(00)00518-1)
- Zhu L, Hu Y, Zhao X, Zeng X, Zhao P, Zhang Z, Ju Y (2017) The impact of drought on sap flow of cooccurring Liquidambar formosana Hance and *Quercus variabilis* Blume in a temperate forest. *Cent China Ecohydrol* 10(4):e1828. <https://doi.org/10.1002/eco.1828>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.