



Sorting *F. graminearum* core effector candidates shows multiple fungal proteins that target the wheat cell nucleus during Fusarium Head Blight

Shimlal Ayilalath, Lilian Faurie, Emmanuel Vanrobays, Florian Rocher, Loriane Loizeau, Géraldine Philippe, Marie Javelle, Mickaël Bosio, Christophe Sallaud, Christophe Tatout, et al.

► To cite this version:

Shimlal Ayilalath, Lilian Faurie, Emmanuel Vanrobays, Florian Rocher, Loriane Loizeau, et al.. Sorting *F. graminearum* core effector candidates shows multiple fungal proteins that target the wheat cell nucleus during Fusarium Head Blight. *Journal of Experimental Botany*, 2026, <10.1093/jxb/erag067>. <hal-05508565>

HAL Id: hal-05508565

<https://hal.inrae.fr/hal-05508565v1>

Submitted on 13 Feb 2026

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons CC BY 4.0 - Attribution - International License

Title

Sorting *F. graminearum* core effector candidates shows multiple fungal proteins that target the wheat cell nucleus during Fusarium Head Blight

Authors

Shimlal Ayilalath¹, Lilian Faurie^{1,2}, Emmanuel Vanrobays², Florian Rocher¹, Loriane Loizeau¹,
Géraldine Philippe¹, Marie Javelle³, Mickaël Bosio³, Christophe Sallaud³, Christophe Tatout² and
Ludovic Bonhomme¹

Affiliations

¹ Université Clermont Auvergne, INRAE, GDEC, F-63000 Clermont-Ferrand, France

² iGReD, CNRS, Inserm, Université Clermont Auvergne, 63000 Clermont-Ferrand, France

³ Limagrain, Chappes Research Center, Route d'Ennezat, 63720 Chappes, France.

Corresponding authors: christophe.tatout@uca.fr and ludovic.bonhomme@inrae.fr

Highlight

This study proposes a method based on *in silico* and *in vivo* screens to identify interacting pairs between core pathogenicity effectors localized in the nucleus and susceptibility factors in plants.

Abstract

Effectors are small molecules secreted by microbial pathogens that disrupt host basal functioning and responses during infection by targeting various plant susceptibility factors. This study reports a candidate selection approach for identifying novel, potential plant nuclear localized effectors from *Fusarium graminearum* secretory proteins. From a dataset of core secretory proteins conserved across several *Fusarium* strains, candidates were selected based on predicted nuclear localization, structural characteristics, and expression profiles during infection. Transient expression in *Nicotiana benthamiana* confirmed accumulation in the plant nucleus, that were further confirmed in wheat protoplasts. One of these proteins was selected for yeast two-hybrid (Y2H) screening to identify wheat protein targets, using a *Fusarium*-infected wheat spike cDNA library specifically generated for this study. The screening identified a high confident interaction with a nuclear-localized wheat beta-amylase 2. The structural modeling of the protein complex between beta-amylase 2 and the putative effector was used to predict interacting amino acid

© The Author(s) 2026. Published by Oxford University Press on behalf of Society for Experimental Biology. All rights reserved. For commercial re-use, please contact reprints@oup.com for reprints and translation rights for reprints. All other permissions can be obtained through our RightsLink service via the Permissions link on the article page on our site—for further information please contact journals.permissions@oup.com.

residues and informed a deletion analysis to disrupt the interaction. This research identifies a *Fusarium graminearum* secretory core protein that interacts with a conserved wheat beta-amylase 2, showcasing a method to select pathogenicity factors conserved across multiple pathogens and host plants, with implications for developing broad-spectrum resistance strategies.

Keywords

Fusarium graminearum, secretory protein, protein structure, plant nucleus.

Abbreviations

Alphafold-Multimer Local Interaction Score (AFM-LIS)

AGAMOUS-LIKE 62 (AGL62)

AGPase (ADP-glucose pyrophosphorylase)

Conserved Domain Database (CDD)

Fusarium graminearum (*F. graminearum*)

Fusarium head blight (FHB)

Green Fluorescent Protein (GFP)

hours post inoculation (hpi)

Interface Predicted Template Modeling (ipTM)

Local Interaction Area (LIA)

Local Interaction Score (LIS)

Monomeric red fluorescent protein (mRFP)

MUTS HOMOLOG 1 (MSH1)

Nicotiana benthamiana (*N. benthamiana*)

Nuclear Localization Signals (NLS)

Possible Biological Score (PBS)

Predicted Aligned Error (PAE)

Predicted Template Modeling (pTM)

Replication factor A protein 1 (RPA)

- 1 Position-Specific Iterated BLAST (RPS-BLAST)
- 2 Sucrose non-fermenting-1-related protein Kinase 1 α (TaSnRK1 α)
- 3 Synthetic defined (SD)
- 4 Yeast two-hybrid (Y2H)
- 5

6 **Introduction**

7 Effectors are small molecules produced by microbial pathogens that interfere with a wide array of
 8 host cellular responses to promote conducive conditions for pathogen development and enhance
 9 infection processes. Secreted and then released into the host tissues, the effectors reach precise
 10 subcellular targets to achieve their roles through direct or indirect interactions with molecular
 11 functions and/or cellular components, the so-called plant susceptibility factors (Zhang *et al.*, 2022).
 12 In plant-pathogen interactions, these targets are remarkably diverse, depicting the need of the
 13 pathogen to control multiple specialized host functions at specific stages of the infection process
 14 (Deslandes and Rivas, 2011). The nucleus is a prime target for many effectors due to its central
 15 role in regulating plant cell functions and responses. Accordingly, a wide range of plant pathogens,
 16 including bacteria, viruses, nematodes, fungi, and oomycetes, produce effectors that specifically
 17 localize to the nucleus (Harris *et al.*, 2023). Many studies have demonstrated their remarkable
 18 actions in manipulating plant cell responses, especially through direct or indirect effects on the
 19 expression of a diverse set of host genes to manipulate plant immunity, to suppress the activation
 20 of genes responsible for plant defense, and even to interfere with generic and specific pathogen
 21 recognition signals (Jiang *et al.*, 2007; Zhang *et al.*, 2022; Cai *et al.*, 2023). For instance, in
 22 *Nicotiana benthamiana* (*N. benthamiana*), the bacterial effector Pi23226 from *Phytophthora*
 23 *infestans* has been proven to accumulate in the nucleus and to interfere with the 25S rRNA
 24 biogenesis. This ultimately results in the disruption of protein translation and promotes cell death,
 25 thereby enabling the pathogen to circumvent the plant's defensive mechanisms and facilitate its
 26 proliferation within the plant (Lee *et al.*, 2023). Similarly, during the rice blast disease, the fungal
 27 effector AvrBs3, secreted by *Magnaporthe oryzae*, has been shown to inhibit the action of specific
 28 transcription factors by directly binding to the host DNA, which results in the suppression of the
 29 defense response (Zhang *et al.*, 2022). As such, increasing our knowledge on nuclear-targeted
 30 microbial effectors will lead to a better understanding of plant pathogen interactions and help to

1 further discover the range of susceptibility factors that pathogens are able to manipulate (Todd *et*
2 *al.*, 2022). By screening their precise actions, this can leverage new avenues to unveil the host
3 vulnerabilities that pathogens exploit during infection and to identify new targeted strategies for
4 crop protection.

5 *Fusarium* head blight (FHB) is one of the most devastating plant diseases in small grains such as
6 wheat and barley (Dean *et al.*, 2012). Primarily caused by the fungus *Fusarium graminearum* (*F.*
7 *graminearum*), FHB not only reduces grain yield and quality but also contaminates grains with
8 harmful mycotoxins leading to serious health risk to humans and animals (Ekwomadu *et al.*, 2021;
9 Xu *et al.*, 2023). Over the past 30 years, genetic approaches have yielded more than 650 QTLs for
10 FHB resistance but providing overall small improvements in the fields with high dependency on
11 environmental conditions (Steiner *et al.*, 2017). Searching for fungal effectors is opening up new
12 opportunities to identify novel key determinants of plant disease, further exploring related wheat
13 susceptibility factors to identify new sources of resistance (Fabre *et al.*, 2020). To date, about a
14 dozen *F. graminearum* effectors have been identified with diverse functions and plant subcellular
15 targets (Voigt *et al.*, 2005; Jiang *et al.*, 2020; Hao *et al.*, 2023; Jin *et al.*, 2024). For instance, OSP24
16 interacts with the *Triticum aestivum* Sucrose non-fermenting-1-related protein Kinase 1 α
17 (TaSnRK1 α) to promote the degradation of the kinase by the proteasome, resulting in a decreased
18 FHB resistance in wheat (Jiang *et al.*, 2020). TaSnRK1 α modulates the activity of the enzyme
19 AGPase (ADP-glucose pyrophosphorylase), which is a key enzyme in starch biosynthesis linking
20 essential metabolic functions such as sugar metabolism and resistance to fungal pathogens.
21 Another effector, called FgNls1, contains several Nuclear Localization Signals (NLS) and its
22 transient expression in *N. benthamiana* as a FgNls1-GFP fusion protein shows its accumulation in
23 the nuclei of plant cells (Hao *et al.*, 2023). Consistent with its nuclear localization, affinity
24 chromatography experiments identified a wheat histone H2B as a main target protein that interacts
25 with FgNls1. FgNls1 expression is significantly upregulated during wheat spike infection while
26 FgNLS1 silencing reduced FHB severity, clearly linking a nuclear-targeted effector to the FHB
27 disease process (Hao *et al.*, 2023). These few examples demonstrate the unequivocal role of *F.*
28 *graminearum* effectors in wheat susceptibility to FHB and suggest that the nucleus can be a
29 primary target to control an intricate set of plant responses. As such, the identification of interacting
30 pairs between effectors and susceptibility factors is a promising strategy to weaken the FHB
31 disease and to improve wheat tolerance.

In our previous works, we identified a core secretome of 357 proteins found to be expressed by different *F. graminearum* strains of varying aggressiveness in different wheat cultivars of contrasting susceptibility (Rocher *et al.*, 2022). Displaying close expression patterns that depicted two early waves of secretory protein delivery over a 96-hour period, these data showed that the genetic program developed by *F. graminearum* during FHB is well conserved across different strains and independent from the host's genetic background. One set of secretory proteins, called early secretory proteins, is typically expressed during the very early stages of infection (i.e. until 48 hours after fungal inoculation) while late secretory proteins are expressed at later stages of infection (i.e. starting from 72 hours after inoculation) (Rocher *et al.*, 2022). The conservation of these expression dynamics in different *F. graminearum* strains suggests that these secretory proteins play an active role in promoting infection by targeting essential host functions at specific times of the infection. Interestingly, among these core secretory proteins, 46 harbored plant NLS and 41 were found to be co-regulated with regulatory hubs of the complex gene regulation network of FHB responses in wheat (Rocher *et al.*, 2024), suggesting their potential role in interfering with massive gene expression regulation. Based on this dataset, we aim to demonstrate that multiple *F. graminearum* secretory proteins effectively accumulate in the plant nucleus to fulfill their function. First, starting from the 357 core secretory proteins identified in Rocher *et al.* (2022), we selected a set of candidates based on their structural features predicted by *in silico* tools and their early and late expression profiles during infection. Nuclear accumulation was assayed using transient expression in *N. benthamiana* and subsequently confirmed in wheat protoplasts for a subset of them. Bringing all this information together allowed the selection of one specific secreted protein hereafter called Effector18 (EFF18) to determine whether it could physically interact with wheat proteins, using the yeast two-hybrid (Y2H) system. For that we designed a specific wheat spike cDNA library and identified four targets with high confidence scores. While the identified interacting host proteins were further validated by yeast two hybrid interaction assay, we focused our attention to EFF18 and LOC123105923, a beta-amylase 2 protein, because their interaction was observed not only in yeast but also predicted by the *in-silico* prediction tool ColabFold_Multimer. Finally, we took advantage of the predicted interacting complex to define the putative interacting residues and used these predictions to design a deletion analysis to confirm the predicted interaction domain. This study identifies a core secretory protein from *F. graminearum* that interacts with a well-conserved beta-amylase 2 and illustrates a strategy to select

pathogenicity factors conserved across multiple pathogens and their host plants to develop promising generalized broad-spectrum resistance.

Materials and Methods

Microbial strains, plant materials, growth conditions and plant assays

The *F. graminearum* strain MDC_Fg1 considered aggressive on different European cultivars (Alouane *et al.*, 2018; Fabre *et al.*, 2021; Rocher *et al.*, 2022) was used in this study. Spores were obtained as previously detailed (Rocher *et al.*, 2022), kept frozen until use and adjusted to a concentration of 1×10^5 spores/ml in distilled water before inoculation. The 'Recital' wheat cultivar, an FHB-susceptible winter wheat (Rocher *et al.*, 2022, 2024), was used for transient expression assay and to prepare a specific cDNA library. For wheat protoplast preparation, surface sterilized seeds of the 'Recital' wheat cultivar were germinated on a filter paper in petri dish in a germination chamber at 25°C with a 16:8 light cycle for 3-5 days. Seedlings were then transferred in pots containing sterilized soil in a growth chamber with same growth conditions as above, for 7 days until plants reached the two-leaf stage. *Agrobacterium tumefaciens* strain GV3101 infiltration assay was performed using *N. benthamiana* (lab line) grown on sterilized soil and incubated in a germination chamber maintained at 25°C with a 16:8 light cycle. Seedlings were transferred to a growth chamber with the same conditions for 3-4 weeks until plants reached the 5-7 leaves stage appropriate for infiltration.

Transient expression of *F. graminearum* secretory proteins in plant cells

Genomic DNA sequences from *F. graminearum* MDC_Fg1 encoding the 14 secretory proteins excluding their signal peptide sequences were used for *de novo* gene synthesis (Genscript USA Inc). The synthesized sequences were cloned into the pGWB605 destination vector (Nakagawa *et al.*, 2007) to express the proteins as C-terminal GFP fusions under the control of the p35S promoter. A second pGWB605 destination vector was designed to express a NLS fused to the mRFP under the control of the pDon promoter. In addition, a third vector expressing the P19 suppressor of RNA silencing was used in all transient expression assay to increase the expression level of co-expressed proteins (Petersen *et al.*, 2019). Transient expression assays were performed by *A. tumefaciens* strain GV3101 infiltration of *N. benthamiana* leaves and observed after 72 to

96 hours after infiltration as described in Sparkes *et al.* (2006). Localization of proteins was analyzed using a Zeiss Cell Observer Spinning Disk microscope under 40X objective. A second transient expression assay was performed in wheat protoplast using the Recital wheat cultivar and the polyethylene glycol-mediated method as previously described (Luo *et al.*, 2022). Protoplast viability was assessed with fluorescent visibility stain (EVANS Blue) and observed under a wide field microscope under 40X objective, and 50,000 protoplasts were mixed with the secretory protein-GFP fusion plasmid and NLS-mRFP. The transfected protoplasts were incubated at 25°C for 2 days and observed using a Zeiss LSM 800 confocal microscope under a 63X oil immersion objective. The NLS-mRFP nuclear marker was used to determine the nuclear localization of the secretory proteins.

Bimolecular Fluorescence Complementation (BiFC) Assay

To validate the interaction between EFF18 and LOC123105923 in planta, bimolecular fluorescence complementation (BiFC) assays were performed. The full-length coding sequence of EFF18 (excluding the signal peptide) was cloned into the pB4cYGW BiFC destination vector to generate a fusion with the C-terminal half of the yellow fluorescent protein (YFP), resulting in the construct EFF18_pCT366_pB4cYGW. Similarly, LOC123105923 was cloned into the pB4nYGW BiFC vector to fuse with the N-terminal half of YFP, generating LOC123105923_pCT367_pB4nYGW.

Constructs were introduced into *A. tumefaciens* strain GV3101, and equal volumes of bacterial suspensions carrying each BiFC construct ($OD_{600} = 0.8$) were co-infiltrated into *N. benthamiana* leaves as described by Sparkes *et al.* (2006). A silencing suppressor (P19) was co-infiltrated in all experiments to enhance protein expression. Leaves were incubated under controlled conditions (25°C, 16:8 light: dark) for 72 hours before imaging.

YFP fluorescence was detected using a Zeiss LSM 800 confocal laser scanning microscope with a 63X oil immersion objective. Excitation was performed at 514 nm, and emission was collected between 525–550 nm. The nuclear localization of the BiFC signal was confirmed by co-expression with the NLS-mRFP marker. Negative controls included co-expression of each BiFC construct with an unrelated protein partner or empty vector (Mermet *et al.*, 2023). No YFP signal was observed in these combinations, confirming the specificity of the EFF18–LOC123105923 interaction. KAKU4-KAKU4 interaction was used as a positive control (Mermet *et al.*, 2023).

Yeast 2-Hybrid assays

Protein-protein interactions were performed by Hybrigenics services (Evry, France) using the ULTIMate Yeast 2-hybrid (Y2H) system and a dedicated cDNA library obtained from Recital wheat cultivar inoculated with the *F. graminearum* strain MDC_Fg1. To prepare the cDNA library, FHB-infected and mock spikes at different stages of the infection dynamics were produced as previously detailed (Rocher *et al.*, 2022). Briefly, plants were sown in buckets, vernalized during six weeks and transferred in a growth chamber with controlled conditions (16:8 light cycle with 21°C/17°C (day/night) temperatures and relative humidity of 80%). Infection was performed at mid-anthesis by inoculating the floral cavity of three synchronous flowering spikes. Spikelets inoculated with water served as controls. For each individual, the inoculated spikelets were collected and immediately frozen in liquid nitrogen. The cDNA library was constructed using inoculated spikelets collected from four plants at 48 and 72hpi as in Sasi *et al.* (Sasi *et al.*, 2023). The resulting library was cloned into the Y2H vector as a prey construct. Coding sequencing of the secretory proteins without the signal peptide were cloned into Y2H vectors as bait constructs and expressed as a fusion protein with the LexA and/ or Gal4 DNA-binding domain. The bait constructs were transformed into yeast strain AH109 and the Y2H screening was performed by mating with the yeast strain Y187 containing the prey, which corresponds to the wheat cDNA fused to the DNA activation domain. Interaction-dependent activation of the reporter genes allowed for the identification of positive interactions. Identified binding partners in Y2H screens were classified in 4 grades (A, B, C and D) using the Hybrigenics Possible Biological Score (PBS) scoring system reflecting the reliability of the interactions where grade A represents the strongest interactions that exhibit high specificity, and which have been observed in independent experiments (Velásquez-Zapata *et al.*, 2021). Selected interactions of grade A and B were subsequently validated using the Yeast 2-hybrid from Clontech, MATCHMAKER GAL4 Two-Hybrid System. Entry vectors containing bait and prey were synthesized by Genscript USA Inc and subsequently subcloned into the bait vector pDEST-GBKT7 or the prey vector pDEST-GADT7 (Rossignol *et al.*, 2007). Yeast cultures were grown at 30 °C on Yeast Extract Peptone Dextrose (YPD) or on selective Synthetic Defined (SD) media. *Saccharomyces cerevisiae* haploid strains AH109 Gold and Y187 were transformed according to the classical protocol (Gietz and Woods, 2002), respectively with the secretory

proteins cloned into the bait vector pDEST-GBKT7 or with the effectors cloned into the prey vector pDEST-GADT7 and grown on appropriate medium (SD-Trp or SD-Leu plates). After mating on YPD and selection of diploids on SD-Leu-Trp medium, interactions were tested on stringent medium SD-Leu-Trp-His-Ade. Empty pDEST-GBKT7 or pDEST-GADT7 vectors were used as negative controls. Two deletion derivatives were designed for EFF18, a C-terminal deletion and deletion of 45 amino acids corresponding to interacting residues of EFF18 with LOC123105923. C-terminal deletion called EFF18_DC was carried out with Phusion PCR DNA Polymerase (Thermo Fischer Scientific) on pUC57_EFF18 with primers C-term_EFF18_Forward (5'-GGGGACAAGTTTGTACAAAAAAGCAGGCTTCATGGTGCCCAAGGCTGACA-3') and C-term-EFF18_Reverse (5'-CGGGGACCACTTTGTACAAGAAAGCTGGGTGTTAGTAGCCGAAAGGCGAAAGG-3'). Deletion of interacting residues (EFF18_DM) was carried out on pUC57_EFF18 by using QuikChange II Site-Directed Mutagenesis Kit (Agilent) with primers Interaction_EFF18_Forward (5'-AATACATCCTTTCGCCTTTCGGCTACACCGGCGCGGCGCCTGACCA-3') and Interaction_EFF18_Reverse (5'-TGGTCAGGCGCCGCGCCGGTGTAGCCGAAAGGCGAAAGGATGTATT-3') designed by overlapping the deletion area (underlined). PCR products were subsequently introduced in pGADT7 and used as bait for Y2H assay.

***In silico* analyses**

Subcellular localization prediction of secreted proteins was carried out using the web tool LOCALIZER1.0.4 (Sperschneider *et al.*, 2017) from mature protein sequences. SignalP5.0 was used to define the signal peptide in the secreted proteins (Nielsen *et al.*, 2019). Protein features of sequences without the signal peptide were evaluated through the EffectorP-fungi 3.0 web tool (Sperschneider and Dodds, 2022). Depicter2 server was used to predict Intrinsically Disordered Region (IDR) (Basu *et al.*, 2023). Monomeric and multimeric structural predictions of candidate proteins were computed with ColabFold Batch (v.1.5.5) (Mirdita *et al.*, 2022). Configuration of ColabFold settings were maintained as default, with 5 predicted structural models, each comprising 3 cycles/reiterations per model. ColabFold scores calculated from monomeric or

multimeric protein structures called pTM (Predicted Template Modeling) and ipTM (Interface Predicted Template Modeling) respectively, were used as global indicators of the prediction confidence. AlphaFold-Multimer Local Interaction Score (AFM-LIS) (Kim *et al.*, 2024) was used to predict protein interactions at the residual scale using the Local Interaction Score (LIS) and the Local Interaction Area (LIA). Probability of interaction of individual amino acids was based on the Predicted Aligned Error (PAE) scores calculated from the PAE matrix with a threshold of 12 Ångström (distance between amino acids). To predict positive protein protein interaction, the LIS and LIA were evaluated according to the following thresholds: a best LIS ≥ 0.203 and a best LIA ≥ 3432 (Kim *et al.*, 2024). Visualization of 3D structures and identification of interacting residues at less than 5 Ångström of distance were performed using ChimeraX (v.1.7.1) (Goddard *et al.*, 2018). Clustering analyses were performed with FoldSeek (Van Kempen *et al.*, 2024) using predicted protein structures as .pdb files available at RCSB Protein Data Bank (RCSB PDB) or predicted by ColabFold. FoldSeek was used with default parameters except for coverage set at 0.8, Predicted Local Distance Difference Test (pLDDT) threshold set at 0.7 and TM score threshold that quantifies the similarity between two protein structures set at 0.7. Clustering was performed either on the 351 core secreted proteins (pdf files were available for 351/357 accessions) alone or in combination with 3,991 additional putative effectors collected from the literature (Derbyshire and Raffaele, 2023; Seong and Krasileva, 2023; Rocafort *et al.*, 2022). Genetic diversity was assessed using a set of 22 fungi species including basidiomycetes and ascomycetes and 22 plant species including unicellular algae, bryophytes and angiosperms. For each of the plant and fungi species, the protein database was retrieved from the NCBI/Taxonomy browser (**Supplementary Tables 1 and 2**). BLASTp search of the PHI-Base database was performed using the Uniprot IDs of secreted proteins predicted to be localized in the nucleus and included into a cluster of ≥ 2 members. Using the same accessions, 3D structure annotation analyses was performed using FoldSeek and a local version of the full AlphaFold database. Hits were selected for LDDT ≥ 0.5 and with a coverage of $\geq 70\%$ to avoid selection based on small fragment similarities. The corresponding Uniprot IDs were then used to search PHI-Base database. The search for orthologous sequences was carried out against downloaded databases of representative species of fungi (for secretory proteins) or plants (for wheat targets). For *F. graminearum* proteins, a tyrosine kinase protein (kinase, PF07714) (Alouane *et al.*, 2021) and the Replication factor A protein 1 (RPA, YAR007C) (Cappela-Gutierrez *et al.*, 2014) were selected as controls for their high and low

percentage of identity, respectively, while AGAMOUS-LIKE 62 (AGL62, Q9FKK2) (Nam et al., 2004) and MUTS HOMOLOG 1 (MSH1, Q84LK0) (Bai and Guo, 2023) were chosen for similar reasons in wheat. The MMseqs2 tool (v.15.6f452) (Steinegger and Söding, 2017) was used with the reciprocal best hits (easy-rbh) option to select the best potential ortholog. Protein sequences were aligned using the mafft tool (v.7.525) (Katoh and Standley, 2013) and the output files were used as input for IQ-TREE (v.2.3.3) by generating 1000 bootstraps, with the ModelFinder tool to identify the best-fit model. The treefile format was imported and modified using the iTOL web service (v.6) (Letunic and Bork, 2024). The Conserved Domain Database (CDD) web server (Lu et al., 2020) was used to identify conserved protein domains in protein sequences via Position-Specific Iterated BLAST (RPS-BLAST). The conserved domains present in the secreted proteins and their targets were identified using the CD-search batch option restricted to protein domains annotated in the pfam database (Mistry et al., 2021). Positions of conserved domains were used to draw the protein structures of secretory proteins and their putative targets identified by Y2H using the drawProtein R package. Bioinformatic scripts are available at https://gitlab.com/lilian_fau/fhb_secure/-/wikis/home.

RESULTS

In silico* characterization of the core secretome of 357 proteins from *Fusarium graminearum

As a starting point for this study, the 357 putative core secretory proteins previously identified by Rocher et al. (2022) (**Supplementary Table 3**) were studied further to refine their characteristics and analyze their similarities both to one another and to those described in available published studies (Rocafort et al., 2022; Derbyshire and Raffaele, 2023; Seong and Krasileva, 2023). The 357 *F. graminearum* genes were systematically expressed and regulated during FHB infection in three strains of contrasting aggressiveness or when facing five different wheat host cultivars of contrasting susceptibility (Rocher et al., 2022). Overall, this gene set encodes small proteins, with an average size of 384 amino acids (AA) (**Figure 1A**), and containing an N-terminal signal peptide for 293/357 (80%) of them, as predicted by SignalP5.0 (**Supplementary Table 4**). Overall, 212 (~60%) did not contain the known specific effector features predicted by EFFECTORP-fungi 3.0 (**Figure 1B, Supplementary Table 5**), suggesting that they may represent novel structural features not yet implemented in the EFFECTORP-fungi 3.0 model. At the primary sequence level, 187 were unique, while 170 shared similarities with another member of the 357

core secretory proteins (**Figure 1C, Supplementary Table 6**). We also observed that while 113 did not yield any functional annotation, 244 (68%) harboured conserved domains referenced in the pfam database (**Supplementary Table 6 and 7**).

We then used FoldSeek (van Kempen *et al.*, 2024) to cluster the 357 proteins according to their protein structure similarities. First, predicted structures were successfully collected for 351/357 and looking at their pLDDT (Predicted Local Distance Difference Test) used as a metric to assess the reliability of predictions (**Supplementary Table 8**), 92/351 have pLDDT below 0.7 and 26/351 below 0.5. While low values may impair the clustering efficiency, 47 clusters and 245 singletons (~70%) were detected, depicting substantial heterogeneity in their structural folds. Among evidenced identified clusters, 39, 5, 2 and 1 gathered 2, 3, 4 or 5 secretory proteins, respectively (**Supplementary Table 8**). In a second strategy, clustering was performed between the 351 core secretory proteins supplemented with 3,991 known secreted proteins identified in three recent studies from different pathogens (Rocafort *et al.*, 2022; Derbyshire and Raffaele, 2023; Seong and Krasileva 2023). This analysis narrowed down to 81 singletons (22%), while the remaining proteins were grouped into clusters of varying sizes, including a large cluster of 180 members and 52 others belonging to clusters containing 2 to 87 members (**Supplementary Table 9**). Comparing the different approaches of domain analysis, similarity searches between the protein sequences of the 357 proteins and their clustering with other published fungi secretomes, 43 proteins (12% of our 351 core secretome) remained unannotated, highlighting the distance still to be covered to fully characterize the *F. graminearum* secretome, while 113 (32%) contained protein domains shared between different members of the secretome and were included in FoldSeek clusters already described (**Figure 1D, Supplementary Table 8 and 9**). To show that our secreted proteins share significant characteristics with secreted proteins identified in previous studies, we then conducted a new analysis to annotate the 56 secretory proteins predicted by LOCALIZER to be localized in the nucleus (see **Figure 1B and Supplementary Table 5**) by searching the PHI-Base database for sequence or structural similarities. Of these, 51/56 proteins were assigned to one of our FoldSeek clusters (≥ 2 members), indicating that our predicted effectors (EFFs) exhibit structural similarities with other secreted proteins previously described (Rocafort *et al.*, 2022; Derbyshire and Raffaele, 2023; Seong and Krasileva, 2023). Among these 51 proteins, BLASTp searches identified sequence similarities for 20/51, while FoldSeek searches identified structural

similarities for 10/51; notably, 6/51 proteins were detected by both approaches. The full results of these analyses are provided in **Supplementary Table 10**, while selected examples are presented in **Table 1**, highlighting the functional diversity and potential evolutionary relationships between our clusters and secreted proteins from *F. graminearum* and other pathosystems in rice, barley, and maize. These preliminary analyses suggest that the 357 core secretory proteins represent an original source of unexplored and uncharacterized effectors with pivotal role during the early stages of the FHB disease.

Identification of secretory proteins targeting the plant nucleus

As already suggested by Rocher *et al.* (2022), we hypothesized that a subset of the secretory proteins could be imported into the plant nucleus. The predicted localization of the 357 secretory proteins using the current version of the LOCALIZER tool identified 56 secretory protein sequences putatively able to enter the plant nucleus (**Figure 1B** and **Supplementary Table 10**). Among these, 13 core secretory proteins were further short listed based both on their consistent expression changes during infection in different strains and on their expression profiles that matched with the effector delivery waves hypothesis (Fabre *et al.*, 2019; Rocher *et al.*, 2022) (**Table 2**). More specifically, this subset included five and eight secretory proteins belonging to the early and late delivery wave, respectively, with fold changes ranging from 0.48 to 4.55 Log₂ normalized counts between 48 and 72 hours post inoculation (hpi) (Rocher *et al.*, 2022).

To confirm the nuclear localization of the 13 selected secretory proteins *in planta*, GFP-fused constructs without signal peptide were designed for further *Agrobacterium*-mediated transient expression in *N. benthamiana* leaf cells using an NLS-mRFP marker as a control. Confocal microscopy observations from at least two independent experiments revealed that eight out of 13 proteins effectively and consistently accumulated in the nucleus (**Figure 2A** and **Supplementary Table 11**). The remaining five proteins, which either showed weak expression, unreproducible accumulation or did not localize to the nucleus, such as EFF02 and EFF24 (**Figure 2A**) served as valuable negative controls for this experiment, demonstrating the specificity of our nuclear localization assay. Among the eight nuclear localized proteins, distinct sub-nuclear localizations were observed. For instance, EFF18-GFP fusion protein showed a strong nucleoplasm accumulation and was excluded from the nucleolus while EFF01-GFP strongly

accumulated into the nucleolus. To further demonstrate that secretory proteins effectively target the nucleus of the host cells, transient expression in wheat protoplasts was repeated for a subset of two proteins that showed strong nuclear accumulation in *N. benthamiana*, but with distinct subnuclear localization (EFF11-GFP and EFF18-GFP). Both secretory proteins-GFP fusion proved to also accumulate in wheat nuclei. Similar sub-nuclear localization was observed for EFF11-GFP fusion proteins in *N. benthamiana* and wheat, while the fluorescence signal of EFF18-GFP fusion protein was heterogeneous and appeared in small domains throughout the wheat nucleus as well as potentially within the nuclear membrane, as observed in the co-localization experiment with NLS-mRFP (**Figure 2B** and **Supplementary Table 11**).

Molecular characterization of the nuclear-targeted secretory proteins

To get further insights into the eight selected secretory proteins, which localize to the plant nucleus, we then explored their molecular structure using a set of *in silico* tools. First, conserved domains were defined using the Conserved Domain Database (CDD) (Lu *et al.*, 2020). Despite the small size of the core secretory proteins, most of them contained one or more protein domains referenced in the pfam database, with EFF18 and EFF22 sharing a common pfam domain (peptidase_M14; pfam00246) (**Figure 3A**). Many of these domains were annotated as peptidases and hydrolases, suggesting a function in protein degradation and catabolism, while others have possible functions in the redox metabolism (**Supplementary Table 7**). Amino acid identity of the eight nuclear-targeted proteins was searched among a set of 22 fungal species whose genomes are fully sequenced, including several *Fusarium* species up to *Ustilago maydis*, a more distant species. Seven proteins were found in 21 to 22 fungal species and the mean percentage of identity of the eight proteins was of $75,7 \pm 23\%$ (mean without *Fgr*) (**Figure 3B** and **Supplementary Table 12**). This suggest that the eight core secretory proteins are well-conserved within the Ascomycota division when compared to the well conserved RPA protein (82.5 ± 18.5) or a tyrosine kinase protein (52.8 ± 29.0) selected for its high accumulation of non-synonymous mutations. Interestingly, while EFF18 and EFF22 shared a common pfam domain, they do not belong to the same FoldSeek cluster (**Supplementary Table 7** and **8**), indicating that secondary and tertiary structural predictions provide complementary information. This prompted us to study the tertiary structures of the eight secretory proteins using ColabFold (**Figure 3C**). Most proteins displayed a very compact globular structure made of helixes and beta sheets, while EFF04 is not well predicted

with a high Predicted Template Modeling (pTM) average value of 0.61. Indeed, low percentages of disorder (absence of protein structure) were predicted in the eight proteins (mean: $16 \pm 11\%$, **Supplementary Table 13**) using DEPICTOR2 (Basu *et al.*, 2023). Again, among the eight core secretory proteins, EFF04 is the exception with 29% of disorder (**Figure 3D**) which corroborate its low pTM values. Finally, despite their high level of identity across a large panel of fungi species, seven out of the eight proteins belonged to the fast-evolving fraction of the *Fusarium* genome (**Figure 3E**). Interestingly, independent RNA-seq datasets confirmed the dynamic expression of these eight nuclear-targeted effectors across wheat genotypes (Recital and Chinese Spring) and in multiple *F. graminearum* strains (MDC_Fg1, MDC_Fg13, MDC_FgU1), with consistent profiles between 48hpi and 96 hpi (**Supplementary Figure 1**). Taken together, the eight secretory proteins form highly compact tertiary structures and as core secretory proteins, they were conserved among a large panel of fungal species and harboured similar strain-independent regulations upon FHB process in different wheat cultivars.

Identification of wheat proteins interacting with EFF18

To identify potential interactors of the secretory proteins that may act as susceptibility genes, we employed the yeast two-hybrid (Y2H) assay, a classical method for detecting direct protein-protein interactions. We selected EFF18 as a candidate to perform a Y2H screen, as EFF18 was consistently expressed in host plant with efficient transient expression assay that undoubtedly confirmed its nuclear localization in wheat protoplasts. EFF18 is well conserved among 22 pathogen species, contains low disorder, high pTM (**Figure 3C-D**) and its expression varies significantly (**Table 2**) between 48 and 72 hpi. To optimize our chance to identify relevant protein interactors, a specific cDNA library was generated using the wheat Recital cultivar infected with the MDC_Fg1 strain, as in Rocher *et al.* (2022).

A first Y2H screen was carried out using the stringent LexA system yielding 20 positive clones, among which only one cDNA sequence (LOC123138250), annotated as a 5-oxoprolinase-like, was identified as multiple clones with a high confidence B score. A second screen, carried out with the more sensitive Gal4 system, identified 100 positive clones, four of them having high confidence scores (A or B). As in the LexA screen, LOC123138250 was identified, as well as three more cDNAs: LOC123105923, encoding a beta-amylase 2 chloroplastic-like protein involved in polysaccharide catabolic processes, LOC543119 annotated as thylakoid formation 1 involved in

the photosystem II assembly in chloroplasts and LOC123116977, an ER membrane protein complex subunit 1-like acting in the glutathione metabolic process (**Supplementary Table 14**). The four proteins included conserved domains referenced in pfam (**Figure 4A**). Consistent with EFF18's classification as a member of the core secretome targeting conserved function in the host, we found that all four targets were conserved across a large panel of plant species representing the green lineage, exhibiting an average sequence identity of $67 \pm 10.8\%$ (**Supplementary Table 15**). This value positioned the four targets as well-conserved proteins when compared to the *MUTS HOMOLOG 1* (MSH1), a mismatch repair protein involved in genome stability which is well-conserved in the green lineage (Bai and Guo, 2023) and to the AGAMOUS-LIKE 62 (AGL62) protein, a member of type I MADS-Box transcription factors described to have a high rate of gene duplication and reduced purifying selection (Nam *et al.*, 2004) with a percentage of identity of $61.1 \pm 10.8\%$ and $36.9 \pm 5\%$ respectively (**Supplementary Table 15**). The phylogenetic trees illustrate this conservation with evolution rates expressed as the sum of the branch lengths for monocot species, similar to MSH1 but lower than AGL62 (**Figure 4B**). In good agreement with the fact that the Y2H library was constructed from spikes tissues, analysis of previous transcriptomic data gained in wheat spikes between 48 and 96 hpi (Rocher *et al.*, 2024) indicated that the four proteins are expressed in spikes with a dynamic pattern of expression during infection. LOC543119 and LOC123105923 are induced in response to Fusarium infection, with mRNA levels increasing as infection progresses. Instead, LOC123116977 transcript levels decrease in response to infection, while expression of the LOC123138250 is not affected by *F. graminearum* infection (**Figure 4C** and **supplementary Table 16**). Interactions between EFF18 and its four putative interactors were then confirmed by performing an Y2H interaction assay carried out by reversing the bait and the prey (**Figure 4D**). Finally, to determine whether the four targets were addressed to the nucleus as for EFF18, a transient expression assay was performed revealing that all four proteins were effectively addressed to and accumulated in the nucleus when co-expressed with EFF18 in *N. benthamiana* (**Figure 4E**), suggesting that they could interact physically within the nucleus. To assess their localization independently of EFF18, we also performed transient expression assays of the four targets in the absence of EFF18. We did not observe striking differences in their localization patterns compared to co-expression, except for LOC123138250-mRFP, which localized outside the nucleus when expressed alone but became nucleoplasmic in the presence of EFF18 (**Supplementary Figure 2**). Taking together, these results suggest that

EFF18 and its four interacting targets are conserved across a large panel of species, are expressed in spikes after anthesis during pivotal stages of the FHB progress and that both EFF18 and its targets can be concurrently addressed to the plant nucleus.

Characterization of the interactions between EFF18 and the beta-amylase target

ColabFold_Multimer tool was used to predict the 3D structures of the complexes between EFF18 and its four interacting targets. The Predicted Aligned Error (PAE) score that defines the confidence of the residue positions within the predicted structures showed that intra- and intermolecular interactions were predicted (**Figure 5A**). However, for all four predicted complexes, evaluation of the quality of interactions between residues by the interface predicted Template Modeling (ipTM) score, gave low scores, below 0.7, that is commonly used as a threshold to predict significant protein-protein interactions. This is likely explained by the low predicted Template Modeling (pTM) score ranging from 0.59 to 0.74 (**Figure 5A**). For this reason and to avoid considering the protein complex as a whole, local interactions were predicted by using AlphaFold-Multimer Local Interaction Score (AFM-LIS) (Kim *et al.*, 2024). AFM-LIS predicted the interaction between EFF18 and LOC123105923 (beta-amylase 2), but failed to detect any other interaction. To strengthen the evidence for a direct interaction between EFF18 and LOC123105923 in a plant cellular context, we performed BiFC assays in *N. benthamiana* leaves. Co-expression of nYFP-EFF18 and cYFP-LOC123105923 reconstituted YFP fluorescence specifically in the nucleus, where the signal co-localized with the nuclear marker NLS-mRFP, confirming that the interaction occurs within the plant nucleus (**Figure 4F** and **Supplementary Figure 3**). The interacting residues at a distance of less than 5 Ångströms from each other were then predicted with ChimeraX (**Figure 5B, 5C** and **Supplementary Table 17**). This analysis identifies one main region in EFF18 between position 304 to 350AA (45 AA), including a large alpha helix (313-330 AA), and four smaller interacting regions in the LOC123105923 between 226 to 422AA (196AA) (**Figure 5D**). This prediction was used to design deletion derivatives of EFF18, in which 45 AA or its C-terminal region was removed (**Figure 5E**). Using these new constructs in an Y2H interacting assay, we demonstrated that deleting the 45AA region was sufficient to impair the interaction between EFF18 and the LOC123105923 (**Figure 5F**), without affecting the predicted structure of EFF18 (**Supplementary Figure 4**). To further explore their co-variation during the FHB progress, we re-investigated independent RNA-seq datasets from four wheat genotypes infected with three

F. graminearum strains (MDC_Fg1, MDC_Fg13, MDC_FgU1). This analysis confirmed that EFF18 and the beta-amylase target (LOC123105923) are concomitantly induced during infection, with both showing increased expression between 48 and 96 hpi (**Supplementary Figure 5A and 5B**). To sum-up, the ColabFold_Multimer tool predicted complexes with low ipTM scores (<0.7) indicating weak protein-protein interactions. Prompting the use of AFM-LIS to confirm interaction with LOC123105923, further validated the interaction by identifying key interacting residues and regions using ChimeraX and functional assay in Y2H.

Discussion

In this study, we examined putative core secretory proteins expressed by several strains of *F. graminearum* of varying aggressiveness in different wheat cultivars during FHB infection and predicted to be targeted to the plant nucleus. These effector-like proteins were found to be significantly regulated during the FHB infection dynamics, suggesting a pivotal role in the infection success. Among the 357 putative secretory proteins described in our previous study (Rocher *et al.*, 2022), 56 were predicted to be localized in the nucleus. In this study, we shortlisted 13 nuclear secretory proteins, all of which harbored a predicted nuclear localization signal (NLS). Transient expression of these proteins in *N. benthamiana* demonstrated the effective nuclear localization for two third with specific subnuclear accumulations, and two were further successfully validated in wheat cell nuclei. By targeting a specific subnuclear structure, these effector-like proteins may interfere with various critical host processes, exerting broad-spectrum effects on the primary functions of the plant cells, thereby helping the pathogen in establishing, proliferating and spreading in infected host tissues. Several nuclear-targeting effectors have already been described in plant pathogens, such as Mlp124478 from *Melampsora larici-populina* known to disrupt host gene expression by directly interacting with plant transcription factors (Petre *et al.*, 2016) and VdSCP7 from *Verticillium dahliae* that targets the plant nucleus and alters plant immunity (Zhang *et al.*, 2017). In *F. graminearum*, very few nuclear-targeting effectors are known. FgNLS1 has been recently described for its ability to reach the cell nuclei and to suppress plant immunity at early stages of the infection (Hao *et al.*, 2023). Here, this study suggests that *F. graminearum* likely deploys a complex arsenal of effector-like molecules into wheat cell nuclei, enabling multiple nuclear targets to manipulate complementary host functions effectively. Some of our 56 secreted proteins predicted to be targeted to the nucleus have significant similarities with

1 already described effectors, such as IIRGP1/OSP4, a well-known effector found in *F.*
 2 *graminearum* (Jiang *et al.*, 2020). We also noticed that EFF04 shows significant similarity to the
 3 Pathogenesis-Related (PR)-1-like 4 protein (FgPR-IL-4) from *F. graminearum*. Notably, deletion
 4 of FgPR-IL-4 affects infection outcomes in both susceptible and resistant wheat cultivars (Lu and
 5 Edwards, 2018). Furthermore, beyond the strong sequence conservation and expression observed
 6 among *F. graminearum* strains with contrasting aggressiveness, this study also revealed a broad
 7 conservation of this set of secreted proteins across different fungal species. Notably, significant
 8 homologies were identified for 26 of the 56 predicted nucleus-localized secreted proteins, as well
 9 as for 7 of the 98 nuclear-targeting secretory proteins, which were found to be well conserved
 10 across 22 fungal species. These findings suggest that such secretory proteins may act as core
 11 components of basal infection mechanisms shared by multiple pathogens across diverse host
 12 plants, thereby supporting the idea that they could serve as promising candidates for the
 13 development of broad-spectrum resistance in crops.

14 To further understand the functional relevance of the nuclear-targeted effector-like proteins
 15 identified in this study, we analyzed the conserved protein domains predicted by the Conserved
 16 Domain Database (CDD). Several of these domains were associated with hydrolase and peptidase
 17 activities, such as those found in EFF18 and EFF22, which harbor the Peptidase_M14 domain.
 18 These enzymatic functions suggest a potential role in host protein degradation, possibly targeting
 19 key nuclear components such as histones, transcription factors, or other DNA-associated proteins
 20 to disrupt gene regulation and promote susceptibility. Other identified domains were linked to
 21 oxidoreductase or redox-related functions, which may be involved in modulating the host's
 22 oxidative stress responses, a known component of plant immune signaling. These domain
 23 predictions support the idea that *F. graminearum* secretes effectors with biochemically active
 24 domains that allow direct manipulation of plant nuclear processes, either by degrading regulatory
 25 proteins or by altering the host cellular environment to facilitate infection. Such conserved drivers
 26 of pathogenicity among various infectious pathogens supports the idea that common functions,
 27 likely essential for the infectious process, are maintained across species to facilitate infection in
 28 diverse host plants. Examples of conserved nuclear effectors playing a significant role in the
 29 infection process by manipulating primary host defense responses have already been reported. For
 30 example, XoPD, a type III secretion effector from *Xanthomonas euvesicatoria*, plays an important
 31 role in altering host gene expression, a common strategy among different pathogens (Kim *et al.*,

2013). By targeting host transcription factors such as SIERF4, this effector is able to suppress ethylene production, thereby further reducing defense responses. Similarly, the Transcription activator-like effectors (TALEs) are important effectors found in several pathogenic bacteria, such as *Ralstonia solanacearum* and *Xanthomonas* spp, able to act as specific plant transcription factors via a programmable DNA-binding domain (Hutin *et al.*, 2015). TALEs from different bacterial strains of the same pathogen have been showed to target the same susceptibility genes and thus have conserved interactors (Boch *et al.*, 2014). Taken together, the nuclear localization and broad conservation of these proteins across diverse pathogens underscore their potential role as nuclear core effector proteins able to promote generic determinants of pathogenicity across a wide range of crops.

Because nuclear effectors are supposed to interact directly or indirectly with host susceptibility factors by targeting specific players of the host cellular processes, we chose EFF18 as a case study to identify new drivers of FHB susceptibility in wheat. EFF18 was also chosen because of its large accumulation in the nucleus and of its increasing expression patterns at the early stages of the FHB infection (Rocher *et al.*, 2022). We then screened for novel interacting partners using a Y2H assay performed on an original and relevant cDNA library obtained from wheat spikes infected with *F. graminearum*. Using this strategy, we identified several plant proteins, which may represent new susceptibility genes able to control FHB development in wheat spikelets. LOC123105923 is a predicted beta-amylase protein involved in the hydrolysis of starch to maltose. In Sweet potato tubers, it plays an important role in carbohydrate metabolism by converting stored starch into sugar, which provides energy for many metabolic processes, including growth and development (Vajravijayan *et al.*, 2018). LOC543119 is involved in the formation of thylakoids in chloroplasts and is essential for photosynthesis. Its role in photosynthesis and further in plant health and defense has already been reported (Monroe and Storm, 2018). LOC123116977 is an endoplasmic reticulum (ER) membrane protein and is important for various cellular processes including stress response and defense (Yamada *et al.*, 2011) and, LOC123138250 is a 5-oxoprolinase that has also been observed to be involved in stress response (Lei *et al.*, 2024). While initially not predicted in the nucleus, we find them to localize to the nucleus in presence of the EFF18 in transient expression assays, suggesting that they may have additional nuclear functions than the ones predicted or demonstrated so far. Only the EFF18/LOC123105923 interaction was successfully confirmed by ColabFold predictions

motivating further attention to these interacting candidates. Interestingly, a plant beta-amylase already proved to accumulate in a susceptible triticale line after infection but not in resistant ones. β -amylases are enzymes involved in starch degradation, releasing maltose and thereby regulating the carbohydrate status of plant tissues. This process is important for energy availability and is linked to growth, stress responses, and developmental pathways in cereals (Thalmann *et al.*, 2019). By targeting a β -amylase, EFF18 could enhance starch degradation, increasing soluble sugars in infected tissues. This nutrient release may benefit the pathogen, enabling more robust fungal growth and reproduction. During FHB infection in a susceptible cereal line, elevated β -amylase levels correlated with higher disease severity, suggesting that changes in β -amylase are associated with weakened plant defenses (Perlikowski *et al.*, 2016), emphasizing its potential as a susceptibility factor. This beta-amylase was highly conserved across the green lineage, echoing with the high sequence conservation of its bait, EFF18, and emphasizing that relative pathogens could target conserved processes across crops. As such, targeted suppression or alteration of β -amylase activity could serve as a promising strategy to enhance resistance against FHB. The conservation of *F. graminearum* nuclear targets, such as β -amylase found broadly in plants, suggests that effectors acting on these proteins could influence susceptibility across diverse hosts. For instance, the PT1 gene (PHI:124535) from *Verticillium dahliae*, expressed during infection of *Gossypium hirsutum* (Chen *et al.*, 2024) shares 66% identity with FGRAMPH1_01G18353 (BLASTp e-value = 0), highlighting potential for broad-spectrum resistance.

To conclude, parsing multiple candidate *F. graminearum* genes supposed to control FHB in wheat, this study provides new insights about the sophisticated strategy used by this pathogenic fungus, emphasizing its ability to deploy a complex arsenal of effector-like proteins towards multiple sub-nuclear wheat targets at early stages of infection to manipulate complementary host functions. Being part of evolutionarily conserved elements across diverse pathogens, these fungal proteins could be at the crossroad of shared mechanisms of pathogenicity that could facilitate infection in a broad range of host plants through the control of generic susceptibility pathways. While further investigations are required, a more in-depth characterization of such fungal core secretory proteins and their plant targets could unlock new opportunities for identifying novel resistance sources, enabling the development of robust, broad-spectrum strategies to control multiple pathogens across various crops.

Acknowledgments

C.T. and L.B. would like to thank Dr Freddy Boutrot and Dr Aline Probst for critical review of the manuscript. The authors thank Richard Blanc for its valuable help during plant experiments and *Fusarium graminearum* inoculations. All microscopic images were acquired on the CLIC microscopy facility (CLermont Imagerie Confocale). All images were stored in an OMERO server hosted at the Mesocentre Clermont Auvergne.

Author contributions

L.B., C.S. and C.T. supervised the study. S.A., E.V., L.L., G.P., M.J., F.R. and M.B. performed experiments. S.A., L.F., C.T. and M.B. performed the bio-informatics work. S.A., L.B. and C.T. wrote the manuscript.

Conflict of interest

No conflict of interest declared

Funding

This work was funded by the Pack Ambition Recherche project FHB-SECURE from the Region Auvergne-Rhône-Alpes. This work was also supported by Centre National de la Recherche Scientifique (CNRS), Institut National de la Santé et de la Recherche Médicale (INSERM), Université Clermont Auvergne (UCA). The authors acknowledge the support received from the Agence Nationale de la Recherche of the French government through the program “Investissements d’Avenir” (16IDEX0001 CAP 2025 CIR 1, International Research Center on Sustainable AgroEcosystems).

Data availability

All data are provided in Supplementary files.

References

- 1 **Alouane T, Rimbart H, Bormann J, González-montiel GA, Loesgen S, Schäfer W, Freitag M,**
2 **Langin T, Bonhomme L.** 2021. Comparative genomics of eight *Fusarium graminearum* strains
3 with contrasting aggressiveness reveals an expanded open pangenome and extended effector
4 content signatures. *International Journal of Molecular Science* **22**(12), 6257.
- 5 **Alouane T, Rimbart H, Fabre F, Cambon F, Langin T, Bonhomme L.** 2018. Genome Sequence
6 of *Fusarium graminearum* Strain MDC_Fg1, Isolated from Bread Wheat Grown in France.
7 *Microbiology Resource Announcements* **7**(19), e01260-18.
- 8 **Bai MZ, Guo YY.** 2023. Bioinformatics Analysis of MSH1 Genes of Green Plants: Multiple
9 Parallel Length Expansions, Intron Gains and Losses, Partial Gene Duplications, and Alternative
10 Splicing. *International Journal of Molecular Science* **24**(17), 13620
- 11 **Basu S, Kihara D, Kurgan L.** 2023. Computational prediction of disordered binding regions.
12 *Computational and Structural Biotechnology Journal* **21**, 1487-1497.
- 13 **Boch J, Bonas U, Lahaye T.** 2014. TAL effectors - pathogen strategies and plant resistance
14 engineering. *New Phytologist* **204**, 823–832.
- 15 **Cai J, Jiang Y, Ritchie ES, Macho AP, Yu F, Wu D.** 2023. Manipulation of plant metabolism by
16 pathogen effectors: More than just food. *FEMS Microbiology Reviews* **47**(2), fuad007.
- 17 **Capella-Gutierrez S, Kauff F, Gabaldón TA.** 2014. Phylogenomics approach for selecting robust
18 sets of phylogenetic markers. *Nucleic Acids Research* **42**, e54.
- 19 Chen A, Morrison S., Gregson, A., Le, D. P., Urquhart, A. S., Smith, L. J., Aitken, E. A. B. and
20 Gardiner, D. M. (2024) *Fluorescently Tagged Verticillium dahliae to Understand the Infection*
21 *Process on Cotton (Gossypium hirsutum) and Weed Plant Species*. *Pathogens*, 13 (6). p. 442.
- 22 **Dean R, Van Kan JAL, Pretorius ZA, et al.** 2012. The Top 10 fungal pathogens in molecular
23 plant pathology. *Molecular Plant Pathology* **13**(4), 414-430
- 24 **Derbyshire MC, Raffaele S.** 2023. Surface frustration re-patterning underlies the structural
25 landscape and evolvability of fungal orphan candidate effectors. *Nature Communications* **14**(1),
26 5244.
- 27 **Deslandes L, Rivas S.** 2011. The plant cell nucleus: A true arena for the fight between plants and
28 pathogens. *Plant Signaling and Behavior* **6**(1), 42-48.
- 29 **Ekwoadu TI, Akinola SA, Mwanza M.** 2021. *Fusarium* mycotoxins, their metabolites (Free,
30 emerging, and masked), food safety concerns, and health impacts. *International Journal of*
31 *Environmental Research and Public Health* **18**(22), 11741.
- 32 **Fabre F, Rocher F, Alouane T, Langin T, Bonhomme L.** 2020. Searching for FHB Resistances
33 in Bread Wheat: Susceptibility at the Crossroad. *Frontiers in Plant Science* **11**, 731.

- 1 **Fabre F, Urbach S, Roche S, Langin T, Bonhomme L.** 2021. Proteomics-Based Data Integration
2 of Wheat Cultivars Facing *Fusarium graminearum* Strains Revealed a Core-Responsive Pattern
3 Controlling Fusarium Head Blight. *Frontiers in Plant Science* **12**, 644810.
- 4 **Gietz DR, Woods RA.** 2002. Transformation of yeast by lithium acetate/single-stranded carrier
5 DNA/polyethylene glycol method. *Methods in Enzymology* **350**, 87-96
- 6 **Goddard TD, Huang CC, Meng EC, Pettersen EF, Couch GS, Morris JH, Ferrin TE.** 2018.
7 UCSF ChimeraX: Meeting modern challenges in visualization and analysis. *Protein Science* **27**,
8 14-25.
- 9 **Hao G, Naumann TA, Chen H, Bai G, McCormick S, Kim HS, Tian B, Trick HN, Naldrett**
10 **MJ, Proctor R.** 2023. *Fusarium graminearum* Effector FgNls1 Targets Plant Nuclei to Induce
11 Wheat Head Blight. *Molecular Plant Microbe Interaction* **36**, 478–488.
- 12 **Harris W, Kim S, Völz R, Lee YH.** 2023. Nuclear effectors of plant pathogens: Distinct strategies
13 to be one step ahead. *Molecular Plant Pathology* **24**(6), 637-650.
- 14 **Hutin M, Sabot F, Ghesquière A, Koebnik R, Szurek B.** 2015. A knowledge-based molecular
15 screen uncovers a broad-spectrum OsSWEET14 resistance allele to bacterial blight from wild rice.
16 *Plant Journal* **84**, 694–703.
- 17 **Jiang C, Hei R, Yang Y, et al.** 2020. An orphan protein of *Fusarium graminearum* modulates host
18 immunity by mediating proteasomal degradation of TaSnRK1α. *Nature Communications* **11**(1),
19 4382.
- 20 **Jiang Y, Yang B, Harris NS, Deyholos MK.** 2007. Comparative proteomic analysis of NaCl
21 stress-responsive proteins in *Arabidopsis* roots. *Journal of Experimental Botany* **58**, 3591–3607.
- 22 **Jin M, Hu S, Wu Q, et al.** 2024. An effector protein of *Fusarium graminearum* targets chloroplasts
23 and suppresses cyclic photosynthetic electron flow. *Plant Physiology* **196**, 2422–2436.
- 24 **Katoh K, Standley DM.** 2013. MAFFT multiple sequence alignment software version 7:
25 Improvements in performance and usability. *Molecular Biology and Evolution* **30**, 772–780.
- 26 **Kim A, Hu Y, Comjean A, Rodiger J, Mohr S, Perrimon N.** 2024. Enhanced Protein-Protein
27 Interaction Discovery via AlphaFold-Multimer. *BioRxiv* doi.org/10.1101/2024.02.19.580970
- 28 **Kim JG, Stork W, Mudgett MB.** 2013. *Xanthomonas* type III effector XopD desumoylates
29 tomato transcription factor SlERF4 to suppress ethylene responses and promote pathogen growth.
30 *Cell Host and Microbe* **13**, 143–154.
- 31 **Lee S, Kim J, Kim MS, Min CW, Kim ST, Choi SB, Lee JH, Choi D.** 2023. The *Phytophthora*
32 nucleolar effector Pi23226 targets host ribosome biogenesis to induce necrotrophic cell death.
33 *Plant Communications* **4**(5), 100606.

- 1 **Lei S, Rossi S, Yang Z, Yu J, Huang B.** 2024. Metabolic regulation of 5-oxoproline for enhanced
2 heat tolerance in perennial ryegrass. *Stress Biology* **4**, 46.
- 3 **Letunic I, Bork P.** 2024. Interactive Tree of Life (iTOL) v6: Recent updates to the phylogenetic
4 tree display and annotation tool. *Nucleic Acids Research* **52**, W78–W82.
- 5 **Lu S, Edwards MC.** 2018. Molecular Characterization and Functional Analysis of PR-1-Like
6 Proteins Identified from the Wheat Head Blight Fungus *Fusarium graminearum*. *Phytopathology*.
7 **108**(4):510-520.
- 8 **Lu S, Wang J, Chitsaz F, et al.** 2020. CDD/SPARCLE: The conserved domain database in 2020.
9 *Nucleic Acids Reserch* **48**, D265–D268.
- 10 **Luo G, Li B, Gao C.** 2022. Protoplast Isolation and Transfection in Wheat. *Methods in Molecular*
11 *Biology* **2464**, 131-141.
- 12 **Mermet S, Voisin M, Mordier J, et al.** 2023. Evolutionarily conserved protein motifs drive
13 interactions between the plant nucleoskeleton and nuclear pores. *The Plant Cell* **35**, 4284–4303.
- 14 **Mirdita M, Schütze K, Moriwaki Y, Heo L, Ovchinnikov S, Steinegger M.** 2022. ColabFold:
15 making protein folding accessible to all. *Nature Methods* **19**, 679–682.
- 16 **Mistry J, Chuguransky S, Williams L, Qureshi M, Salazar GA, Sonnhammer ELL, Tosatto**
17 **SCE, Paladin L, Raj S, Richardson LJ, Finn RD, Bateman A.** 2021. Pfam: The protein families
18 database in 2021. *Nucleic Acids Research* **49**, 412-419.
- 19 **Monroe JD, Storm AR.** 2018. Review: The *Arabidopsis* β -amylase (BAM) gene family: Diversity
20 of form and function. *Plant Science* **276**, 163-170.
- 21 **Nakagawa T, Suzuki T, Murata S, et al.** 2007. Improved gateway binary vectors: High-
22 performance vectors for creation of fusion constructs in transgenic analysis of plants. *Bioscience,*
23 *Biotechnology, and Biochemistry* **71**, 2095–2100.
- 24 **Nam J, Kim J, Lee S, An G, Ma H, Nei M.** 2004. Type I MADS-box genes have experienced
25 faster birth-and-death evolution than type II MADS-box genes in angiosperms. *Proceedings of the*
26 *National Academy of Sciences USA* **101**(7), 1910-1915.
- 27 **Nielsen H, Tsirigos KD, Brunak S, von Heijne G.** 2019. A Brief History of Protein Sorting
28 Prediction. *Protein Journal* **38**, 200-216.
- 29 **Perlikowski D, Wiśniewska H, Kaczmarek J, Góral T, Ochodzki P, Kwiatek M, Majka M,**
30 **Augustyniak A, Kosmala A.** 2016. Alterations in kernel proteome after infection with *Fusarium*
31 *culmorum* in two triticale cultivars with contrasting resistance to fusarium head blight. *Frontiers*
32 *Plant Science* **7**, 1217.

- Petersen BL, Möller SR, Mravec J, Jørgensen B, Christensen M, Liu Y, Wandall HH, Bennett EP, Yang Z.** 2019. Improved CRISPR/Cas9 gene editing by fluorescence activated cell sorting of green fluorescence protein tagged protoplasts. *BMC Biotechnology* **17**,19:36.
- Petre B, Lorrain C, Saunders DG, Win J, Sklenar J, Duplessis S, Kamoun S.** 2016. Rust fungal effectors mimic host transit peptides to translocate into chloroplasts. *Cell Microbiology* **18**, 453-465.
- Rocafort M, Bowen JK, Hassing B, Cox MP, McGreal B, de la Rosa S, Plummer KM, Bradshaw RE, Mesarich CH.** 2022. The *Venturia inaequalis* effector repertoire is dominated by expanded families with predicted structural similarity, but unrelated sequence, to avirulence proteins from other plant-pathogenic fungi. *BMC Biology* **20**, 246.
- Rocher F, Alouane T, Philippe G, Martin ML, Label P, Langin T, Bonhomme L.** 2022. *Fusarium graminearum* Infection Strategy in Wheat Involves a Highly Conserved Genetic Program That Controls the Expression of a Core Effectome. *International Journal of Molecular Science* **23**(3), 1914.
- Rocher F, Dou S, Philippe G, Martin ML, Label P, Langin T, Bonhomme L.** 2024. Integrative systems biology of wheat susceptibility to *Fusarium graminearum* uncovers a conserved gene regulatory network and identifies master regulators targeted by fungal core effectors. *BMC Biology* **22**, 53.
- Rossignol P, Collier S, Bush M, Shaw P, Doonan JH.** 2007. *Arabidopsis* POT1A interacts with TERT-V(18), an N-terminal splicing variant of telomerase. *Journal of Cell Science* **120**, 3678–3687.
- Sasi S, Krishnan S, Kodackattumannil P, Shamisi AAl, Aldarmaki M, Lekshmi G, Kottackal M, Amiri KMA.** 2023. DNA-free high-quality RNA extraction from 39 difficult-to-extract plant species (representing seasonal tissues and tissue types) of 32 families, and its validation for downstream molecular applications. *Plant Methods* **19**, 84.
- Seong K, Krasileva K V.** 2023. Prediction of effector protein structures from fungal phytopathogens enables evolutionary analyses. *Nature Microbiology* **8**, 174–187.
- Sparkes IA, Runions J, Kearns A, Hawes C.** 2006. Rapid, transient expression of fluorescent fusion proteins in tobacco plants and generation of stably transformed plants. *Nature Protocols* **1**, 2019–2025.
- Sperschneider J, Catanzariti AM, Deboer K, Petre B, Gardiner DM, Singh KB, Dodds PN, Taylor JM.** 2017. LOCALIZER: Subcellular localization prediction of both plant and effector proteins in the plant cell. *Scientific Reports* **7**, 44598.
- Sperschneider J, Dodds PN.** 2022. EffectorP 3.0: Prediction of Apoplasmic and Cytoplasmic Effectors in Fungi and Oomycetes. *Molecular Plant Microbe Interaction* **35**, 146–156.

- Steinegger M, Söding J.** 2017. MMseqs2 enables sensitive protein sequence searching for the analysis of massive data sets. *Nature Biotechnology* **35**, 1026–1028.
- Steiner B, Buerstmayr M, Michel S, Schweiger W, Lemmens M, Buerstmayr H.** 2017. Breeding strategies and advances in line selection for *Fusarium* head blight resistance in wheat. *Tropical Plant Pathology* **42**, 165–174.
- Thalmann M, Coiro M, Meier T, Wicker T, Zeeman SC, Santelia D.** 2019. The evolution of functional complexity within the β -amylase gene family in land plants. *BMC Evolutionary Biology* **19**(1), 66.
- Todd JNA, Carreón-Anguiano KG, Islas-Flores I, Canto-Canché B.** 2022. Microbial Effectors: Key Determinants in Plant Health and Disease. *Microorganisms* **10**(10), 1980.
- Vajravijayan S, Pletnev S, Mani N, Pletneva N, Nandhagopal N, Gunasekaran K.** 2018. Structural insights on starch hydrolysis by plant β -amylase and its evolutionary relationship with bacterial enzymes. *International Journal of Biological Macromolecules* **113**, 329–337.
- van Kempen M, Kim SS, Tumescheit C, Mirdita M, Lee J, Gilchrist CLM, Söding J, Steinegger M.** 2024. Fast and accurate protein structure search with Foldseek. *Nature Biotechnology* **42**, 243–246.
- Velásquez-Zapata V, Elmore JM, Banerjee S, Dorman KS, Wise RP.** 2021. Next-generation yeast-two-hybrid analysis with Y2H-SCORES identifies novel interactors of the MLA immune receptor. *PLoS Computational Biology* **17**(4), e1008890.
- Voigt CA, Schäfer W, Salomon S.** 2005. A secreted lipase of *Fusarium graminearum* is a virulence factor required for infection of cereals. *Plant Journal* **42**, 364–375.
- Xu R, Shandilya UK, Yiannikouris A, Karrow NA.** 2023. Traditional and emerging *Fusarium* mycotoxins disrupt homeostasis of bovine mammary cells by altering cell permeability and innate immune function. *Animal Nutrition* **12**, 388–397.
- Yamada K, Hara-Nishimura I, Nishimura M.** 2011. Unique defense strategy by the endoplasmic reticulum body in plants. *Plant Cell and Physiology* **52**, 2039–49.
- Zhang S, Li C, Si J, Han Z, Chen D.** 2022. Action Mechanisms of Effectors in Plant-Pathogen Interaction. *International Journal of Molecular Science* **23**(12), 6758.
- Zhang L, Ni H, Du X, Wang S, Ma XW, Nürnberger T, Guo HS, Hua C.** 2017. The *Verticillium*-specific protein VdSCP7 localizes to the plant nucleus and modulates immunity to fungal infections. *New Phytologist* **215**, 368–381.

Tables

Table 1: PHI-Base and GO analysis of 31 secreted proteins predicted to be targeted to the nucleus.

Starting from the 133 secreted proteins sharing sequence and structural similarities as well as conserved domains described in Figure 1D, BLASTp and FoldSeek search were performed against the PHI-Base database. 10 hits with LDDT ≥ 0.5 were recorded identifying orthologs or paralog proteins described in several pathosystems.

Gene	GO MF	Phi_Base	Function	LDDT	Gene	Pathogen	host	Ref.
FGRAMPH1_01G05441	GO:0016899	PHI:11989	putative D-arabinono-1,4-lactone oxidase	0.5913	MoALO1 (MGG_02689)	<i>M. oryzae</i>	<i>O. sativa</i>	Wu et al 2022
				0.5913		<i>M. oryzae</i>	<i>H. vulgare</i>	
FGRAMPH1_01G05657	GO:0008061	PHI:8753	chitinase	0.8297		<i>M. oryzae</i>	<i>O. sativa</i>	Yang et al 2019
				0.8297	MoChia1 (MGG_08054)	<i>M. oryzae</i>	<i>O. sativa</i>	
				0.8297		<i>M. oryzae</i>	<i>O. sativa</i>	
		PHI:8806		0.8297		<i>M. oryzae</i>	<i>O. sativa</i>	Han et al 2019
				0.8297	MoChi1 (MGG_08054)	<i>M. oryzae</i>	<i>O. sativa</i>	
				0.8297		<i>M. oryzae</i>	<i>H. vulgare</i>	
FGRAMPH1_01G24877	NA	PHI:7144	SCP-like extracellular protein	0.6229	FvSCP1	<i>F. verticillioides</i>	<i>Z. mays</i>	Zhang et al 2017
		Pathosystems in barley, rice and maize uncharacterized protein PHI:5711		1.00	ScOrtholog_MIF2	<i>F. graminearum</i>	<i>T. aestivum</i>	Yu et al 2015
				0.8492	FGSG_02883			

Table 2. Selected set of 13 secretory proteins predicted by LOCALIZER1.0.4 to be targeted to the plant nucleus.

Molecular features of 13 selected proteins expected to be targeted to the nucleus by LOCALIZER1.04 and characterized using EFFECTORP3.0, SIGNALP5.0, their expression profiles and Log2 fold change (FC) during wheat spike infection from Rocher et al 2022.

Accession ID	Code	SIZE (aa)	Effector Prediction (EFFECTORP)	Nuclear Localization Prediction (LOCALIZER)	Expression pat during FHB	Log(FC) between 48hpi and 72hpi
FGRAMPH1_01T01843	EFF01	383	Non-effector	Nucleus	Late	0.54
FGRAMPH1_01T05171	EFF02	617	Non-effector	Nucleus	Late	1.17
FGRAMPH1_01T11559	EFF04	247	Cytoplasmic effector	Nucleus	Late	0.49

FGRAMPH1_01T11579	EFF06	856	Non-effector	Nucleus	Late	2.25
FGRAMPH1_01T12575	EFF10	1009	Non-effector	Nucleus	Early	0.54
FGRAMPH1_01T13187	EFF11	528	Non-effector	Nucleus	Early	0.82
FGRAMPH1_01T13647	EFF13	663	Non-effector	Nucleus, Apoplast	Early	1.26
FGRAMPH1_01T13999	EFF17	1350	Non-effector	Nucleus, chloroplast	Early	0.76
FGRAMPH1_01T15319	EFF18	439	Non-effector	Nucleus	Late	1.25
FGRAMPH1_01T18783	EFF20	569	Non-effector	Nucleus	Early	2.42
FGRAMPH1_01T21315	EFF22	441	Cytoplasmic effector	Nucleus	Late	0.86
FGRAMPH1_01T21741	EFF24	876	Non-effector	Nucleus	Late	0.50
FGRAMPH1_01T25857	EFF29	505	Non-effector	Nucleus, Apoplast	Late	3.94

Table 3. Interaction scores for 4 complexes between targets and EFF18 in Y2H and with AFM-LIS.

From the PAE analysis (probability of interaction between amino acids based on the distance in Ångströms between 2 amino acids), AFM-LIS allows predicting PPIs at the scale of each amino acid by jointly using a Local Interaction Area (LIA) and a Local Interaction Score (LIS). AFM-LIS with best Local Interaction Area (LIA) and best Local Interaction Score (LIS) above the minimal values (respectively $LIA \geq 3432$ and $LIS \geq 0.203$).

Y2H interactors	Y2H interaction score	Best ipTM	Best pTM	Best pLDDT	Best LIS	Best LIA
LOC123116977	B	0.38	0.71	82.42	0.015	10
LOC543119	A	0.20	0.63	77.29	0	0
LOC123105923	A	0.69	0.75	80.80	0.215	238.2
LOC123138250	B	0.19	0.69	84.00	0	0

Figure legends

Figure 1. Characteristics of the 357 core secretory proteins.

Classification of the 357 core secretory proteins according to A) their size in amino acids (aa), B) EFFECTORP-fungi 3.0 and LOCALIZER1.0.4 predictions and C) protein sequence similarities shared within the 357 secretory proteins (EFFs). D) Venn diagrams representing the unique (left) and common (right) protein features cross-referencing (i) shared sequence similarities identified by MMSEQS2, (ii) shared conserved domains referenced in pfam predicted by a Conserved Domain Database (CDD) analysis and (iii) shared structural similarities defined by FoldSeek clustering.

Figure 2. Characterization of 13 secretory proteins by transient expression in *Nicotiana benthamiana* and *Triticum aestivum*.

A) Transient co-expression of effector candidates fused to GFP (EFF-GFP) with a nuclear localization signal-tagged mRFP (NLS-mRFP) in *N. benthamiana* leaves. Localization was assessed 3 days post-infiltration using spinning disk confocal microscopy with a 40X objective. B) Transient expression of the same constructs in isolated wheat (*Triticum aestivum*) protoplasts, imaged 3 days post-transfection using a 40X oil immersion confocal objective. NLS-mRFP was used as a nuclear localization marker to confirm nuclear targeting of EFF proteins. EFF02-GFP and EFF24-GFP, which do not accumulate in the nucleus, served as negative controls. Comparative localization patterns between GFP and NLS-mRFP fluorescence signals supported nuclear accumulation for a subset of tested effectors.

Figure 3. Characteristics of 8 secreted proteins from *F. graminearum* targeted to the plant nucleus.

A) Conserved secondary structures of the 8 secreted proteins as predicted by the Conserved Domain Database (CDD) analysis. B) Box plots with Tukey whiskers representing the percentage of the 8 secretory protein identity (% identity) in 22 species of fungi as well as for RPA and a tyrosine kinase protein (kinase) chosen for their low and high conservation, respectively. Mean value (72.5%) of percentage of identity of the 8 secreted proteins is indicated as a red line. C) 3D structures predicted by ColabFold and refined by ChimeraX to highlight beta-sheets (green) and alpha helixes (blue) secondary structures. pTM values ranging from 0 to 1, with higher scores indicating better structural predictions are indicated below each structure. D) Box plots with Tukey whiskers representing the percentage of intrinsically disordered regions (% disorder) of the 8 selected proteins. Mean value (16%) of disorder of the 8 secretory proteins is indicated as a red line. E) Positions of the 8 genes encoding the secretory proteins according to the fast and slow evolving regions along the 4 chromosomes of *F. graminearum* genome. Statistical significance of the distribution of the 8 secreted protein against the complete set of 357 proteins was assessed by Fisher's Exact Test, p-Value = 0.29.

Figure 4. Characteristics of EFF18 target genes identified by Y2H screen.

A) Protein domains identified in the 4 target proteins as predicted by the Conserved Domain Database (CDD) analysis. B) Unrooted phylogenetic trees where monocotyledons (green) and dicotyledons (blue) are indicated. Branch lengths were standardized for all trees and expressed as substitution/site. Percentage of identity (%identity) and evolution rates computed as the sum of the branch lengths for the monocotyledons species are given below each tree. Values for AGL62 and MSH1 served as indicators for high and low evolution rates. C) Expression profiles of the 4 target proteins in the 'Recital' wheat cultivar. Data are expressed as Log₂(FC) between infected (values obtained from plants facing 3 different *F. graminearum* strains in 3 replicates each) and mock (3 replicates) conditions at 48, 72 and 96 hours after infection (Hpi). D) Y2H interaction assay between EFF18 (prey) and its four interacting proteins (baits). Y2H assay was performed using permissive (YNB/-Leu/-Trp, top panel) or test (YNB/-Leu/-Trp/-Ade/-His; bottom panel) media. Empty vector pGADT7 and pGBKT7 were used respectively as prey and bait negative controls. E) Co-localization assay of the four target proteins expressed as mRFP fusions and EFF18-GFP in transient expression assay and F) Bimolecular Fluorescence Complementation (BiFC) assay between EFF18 and its target LOC123105923 performed in *N. benthamiana* and observed three days after infiltration using a Zeiss Cell Observer Spinning Disk microscope under 40X objective. Scale bar: 10µm.

Figure 5. Interactions between EFF18 and its four targets predicted by ColabFold_multimer.

A) Predicted Aligned Error (PAE) score obtained from a ColabFold_multimer prediction and representing the probability of amino acids interacting with each other. PAE plots are represented as 2D matrixes with protein A as the targets and B as EFF18. PAE values range from 0 to 30 Ångströms (y-axis), where 0 indicates high (blue) and 30 low (red) confidence in interactions, along the protein (x-axis in amino acids). Intra and intermolecular interactions can be observed for A top left, B bottom right and A-B top right, B-A bottom respectively. pTM and ipTM of the complexes are indicated below the corresponding PAE plots. B) Protein complex with interacting residues highlighted in green (EFF18, 32AA, top) and red (LOC123105923, 36AA, bottom). C) Definition of the interacting residues between EFF18 and LOC123105923 using ChimeraX and redrawn at scale highlighting EFF18 (green) and beta amylase (red) residues in interaction. Pfam

domains identified by the Conserved Domain Database (CDD) are indicated as in Figure 4. D) EFF18 deletion derivatives used in Y2H assay. E) Y2H assay between EFF18, EFF18 deletion derivatives and LOC123105923. The prey (in pGADT7 vector) together with each of the bait plasmids (in pGBKT7 vector) were transformed into the yeast strains Y187 and AH109 respectively. Y2H assay was performed using permissive (YNB/-Leu/-Trp, top panel) or test (YNB/-Leu/-Trp/-Ade/-His, bottom panel) media. Empty vector pGADT7 and pGBKT7 were used respectively as prey and bait negative controls.





Figure 1
559x559 mm (x DPI)

A



B




Figure 2
559x559 mm (x DPI)




Figure 3
559x559 mm (x DPI)




Figure 4
559x559 mm (x DPI)




Figure 5
559x559 mm (x DPI)