

Two new species of *Cytospora* (Diaporthales, Cytosporaceae) causing canker disease of *Malus domestica* and *M. sieversii* in Xinjiang, China

Guifang Cai^{1,2*}, Ying Zhao^{3*}, Yawei Zhai⁴, Meilin Yan⁵, Rong Ma^{1,6}, Daoyuan Zhang⁶

¹ College of Forestry and Landscape Architecture, Xinjiang Agricultural University, Urumqi, 830052, China

² Forestry and Grassland Technology Extension Center of Changji Prefecture, Changji 831100, China

³ Forestry and Grassland Resources Monitoring Center of Xinjiang Production and Construction Corps, Urumqi, 830002, China

⁴ China Energy Engineering Group Xin Jiang Electric Power Design Institute CO., LTD., Urumqi, 830050, China

⁵ Forestry and Grassland Bureau of Hinggan League, Hinggan, 137599, China

⁶ CAS Key Laboratory of Biogeography and Bioresource in Arid Land, Xinjiang Institute of Ecology and Geography, Urumqi, China

Corresponding authors: Rong Ma (xjaumr@xjau.edu.cn); Daoyuan Zhang (zhangdy@ms.xjb.ac.cn)

Abstract

Apple tree canker is a serious disease caused by species of *Cytospora*. Xinjiang Uygur Autonomous Region is one of the most important apple-producing areas in China. However, losses due to apple *Cytospora* canker have seriously damaged the apple industry and affected the economic development of the apple growers in this region. In this study, we used morphological characteristics combined with multilocus phylogenetic analyses of the ITS, *act*, *rpb2*, *tef1* and *tub2* loci to identify isolates from apple (*Malus domestica*) and wild apple (*M. sieversii*). As a result, *C. hippophaopsis* **sp. nov.** from *M. sieversii* and *C. shawanensis* **sp. nov.** from *M. domestica* were discovered and proposed herein. Pathogenicity tests were further conducted on 13 varieties of apple and wild apple, which confirmed *C. hippophaopsis* and *C. shawanensis* as canker pathogens. Meanwhile, *C. hippophaopsis* is generally more aggressive than *C. shawanensis* on the tested varieties of apple and wild apple.

Key words: Apple, Ascomycota, phylogeny, plant disease, taxonomy, wild apple

Introduction

Wild apple (*Malus sieversii*) is a tertiary relict plant distributed only in the Tianshan mountain of central Asia. It is located in Emin and Yumin Counties of Tacheng area, Xinyuan, Gongliu and Huocheng Counties of Ili area in Xinjiang Uygur Autonomous (XJUA) Region, China. It is rich in intraspecific variation and has strong resistance, high yield, dwarf and other characters, and is often used in rootstock grafting, cross breeding, and is also important for introduction, domestication and germplasm resources (Richards et al. 2009; Volk et al. 2013; Wang et al. 2018). It plays an irreplaceable role in the study of germplasm conservation and genetic development (Yan et al. 2008; Richards et al. 2009; Volk et al. 2013). This species has been listed in the index of China's endangered secondary pro-

Academic editor: Ning Jiang

Received: 7 July 2024

Accepted: 31 August 2024

Published: 16 October 2024

Citation: Cai G, Zhao Y, Zhai Y, Yan M, Ma R, Zhang D (2024) Two new species of *Cytospora* (Diaporthales, Cytosporaceae) causing canker disease of *Malus domestica* and *M. sieversii* in Xinjiang, China. MycoKeys 109: 305–318. <https://doi.org/10.3897/mycokeys.109.131456>

Copyright: © Guifang Cai et al.

This is an open access article distributed under terms of the Creative Commons Attribution License (Attribution 4.0 International – CC BY 4.0).

* These authors contributed equally to this work.

tected plants (Fu 1991). Apple (*M. domestica*) is one of the four major fruits in the world and one of the 11 major agricultural products identified by the Ministry of Agriculture of China. XJUA Region is one of eight high-yield apple production bases in China, accounting for more than 3.5% of production. The cultivation has a long history, due to adequate sunshine, high effective accumulated temperature, large temperature difference between day and night (Li et al. 2015; Snyder and Ni 2017). Apple tree canker has been reported as a common disease in Tekes County of Ili area in XJUA Region and the death rate of apple trees reached 30% to 40%, which caused great loss to the apple industry (Yang 2009).

More than 20 species of *Cytospora* have been reported on *Malus* spp. worldwide (Zhuang 2005; Ma et al. 2018; Azizi et al. 2020; Wang et al. 2020; Bozorov et al. 2024), of these *C. leucostoma*, *C. mali*, *C. parasitica* and *C. schulzeri* have been reported to infect *M. domestica* in XJUA Region (Zhuang 2005; Ma et al. 2018; Liu et al. 2020). Meanwhile, three species of *Cytospora* were recorded on *M. sieversii*, viz. *C. leucostoma* and *C. mali* distributed in Gongliu, Aksu, Urumqi, Turpan and Shanshan (Zhuang 2005), and *C. parasitica* distributed in Tacheng and Ili areas (Ma et al. 2018).

The genus *Cytospora* (Cytosporaceae, Diaporthales) comprises many important phytopathogens that cause dieback and canker disease on twigs, branches and stems of various woody species and can cause large areas of dieback on a wide range of plants, resulting into severe commercial and ecological damage and significant losses worldwide (Adams et al. 2005; Senanayake et al. 2017; Norphanphoun et al. 2018; Fan et al. 2020; Jiang et al. 2020; Li et al. 2024).

Cytospora was initially introduced with *C. betulina*, *C. epimyces*, *C. resinae* and *C. ribis* in 1818, and *C. chrysosperma* was added to this genus and subsequently selected as the type species (Ehrenberg 1818; Fries 1823; Donk 1964). Species of this genus are characterized by having allantoid hyaline spores in both sexual and asexual states (Zhu et al. 2019; Fan et al. 2020; Pan et al. 2020, 2021; Lin et al. 2023a, 2023b). Species identification in *Cytospora* was previously based on their hosts and morphological characters (Donk 1964; Ehrenberg 1818; Fries 1823). Molecular phylogeny based on sequence data has been applied to separate species of this genus recently (Adams et al. 2005; Fan et al. 2015; Wang et al. 2015; Norphanphoun et al. 2017; Lawrence et al. 2018).

Canker diseases caused by *Cytospora* species are leading to serious economic losses in apple plantations of XJUA Region in China. The aims of this study were to clarify the taxonomy statues of the newly collected species from the host genus *Malus* in this region, and to test their pathogenicity.

Materials and methods

Sample collection, isolation and morphology

Malus canker disease investigations were conducted during 2015 and 2022 in XJUA Region. Trees with dead stems, branches and twigs were checked, and fruiting bodies were discovered on the tree barks (Fig. 1). Samples were packed in paper bags and turned to laboratory for further research.

Ascomata and conidiomata formed on barks of *Malus domestica* and *M. sieversii* were sectioned using a sterile blade, and mucoid spore masses were removed to the surface of the potato dextrose agar (PDA; potato, 200 g;



Figure 1. Symptoms of apple tree cankers in Xinjiang, China.

glucose, 20 g; agar, 20 g; distilled water, to complete 1000 mL) media using a sterile needle. Then plates were incubated at 25 °C in darkness until spores germinated. Pieces of mycelium were cut and removed to a new PDA plate under a stereomicroscope to obtain the pure cultures.

The new species of *Cytospora* were observed mainly based on the fruiting bodies naturally formed on the tree barks. The ascomata and conidiomata were sectioned using a sterile blade and photographed under the Leica stereomicroscope (M205) (Leica Microsystems, Wetzlar, Germany). The asci, ascospores, conidiophores, conidiogenous cells and conidia were measured and photographed by a Nikon Eclipse 80i microscope (Nikon Corporation, Tokyo, Japan). The colony characters were observed and recorded on PDA plates at 25 °C under darkness.

DNA extraction, PCR amplification and phylogenetic analyses

The total DNA of strains of *Cytospora* collected in the present study was extracted from cultures growing on the PDA plates overlaid with cellophane based on the CTAB method (Doyle 1990). Five primer pairs ITS1/ITS4, ACT512F/ACT783R, fRPB2-5f/fRPB2-7cR, 983F/2218R and Bt2a/Bt2b were used to amplify the internal transcribed spacer region rDNA (ITS), the partial actin (*act*) region, RNA polymerase II second largest subunit (*rpb2*), translation elongation factor 1- α (*tef1*) and the partial beta-tubulin (*tub2*) gene loci, respectively (White et al. 1990; Glass and Donaldson 1995; Carbone and Kohn 1999; Liu et al. 1999; Rehner 2001). The polymerase chain reaction (PCR) conditions were as performed as follows: an initial denaturation step of 5 min at 94 °C, followed by 35 cycles of 30 s at 94 °C, 50 s at 52 °C (ITS), 54 °C (*tef1* and *tub2*), 55 °C (*rpb2*) or 58 °C (*act*), and 1 min at 72 °C, and a final elongation step of 7 min at 72 °C. All amplified PCR products were estimated visually with 1.4% agarose gels stained with ethidium bromide and then PCR positive products were sent to Sangon Biotech (Shanghai) Co., Ltd., (Beijing, China) for sequencing.

Sequences obtained in the present study were preliminary identified by the BLAST search to confirm their classification. The referenced sequences of

Cytospora were collected from Wang et al. (2024) and downloaded. *Diaporthe vaccinii* (CBS 160.32) was selected as the outgroup taxon. The five individual loci ITS, *act*, *rpb2*, *tef1* and *tub2* were aligned using MAFFT v. 6.0 and edited manually using MEGA v. 6.0 (Kato and Standley 2013; Tamura et al. 2013). Then five loci were combined and analyzed based on maximum likelihood (ML) and Bayesian inference in the CIPRES Science Gateway platform (Miller et al. 2010). The GTR substitution model were employed, and 1000 non-parametric bootstrap replicates were set for ML phylogenetic analysis. Four simultaneous Markov Chain runs for 1000000 generations were set during Bayesian analysis. The resulting trees were visualized in FigTree v. 1.4.0 and edited by Adobe Illustrator 2020.

Pathogenicity tests

To determine the pathogenicity of *C. hippophaopsis* and *C. shawanensis* newly proposed in the present study to 13 apple varieties (*Malus domestica*) and wild apple (*M. sieversii*). Healthy branches were collected from three-year-old healthy trees of Red Meat Apple (RMA), Jonagold (JNG), Golden Delicious (GD), Manpanzi (MPZ), Sitagan (STG), Oil fruit (OF), Red Star (RS), Fuji (FU), Erzizi (EZ), Qiulimeng (QM), Apolte (AP), Hanfu (HF), New Century (NC) and wild apple *M. sieversii* (MS) in Ili Kazakh Autonomous Prefecture in XUAR in July 2019. The branches were washed with water and disinfected with a 75% (vol/vol) ethanol solution, then branches were cut into 30-cm long segments. Both ends of the branch segments were sealed with paraffin wax to reduce desiccation. A soldering iron was used to create a wound 1 mm in diameter a third of the way along each branch segment and leaf. A 3-mm-diameter mycelial plug which cultured for three days of *C. hippophaopsis* and *C. shawanensis* species was placed onto the wound of branches. There were three replicates per treatment for branch tests. The materials of branch that inoculated with sterile PDA plugs were used as controls. All tested materials were incubated in humid chambers at room temperature (26 ± 2 °C) in 12 h light and 12 h dark. The mycelial plug was removed after 24 hours. The virulence of *C. hippophaopsis* and *C. shawanensis* was assessed by measuring the lesion length (L) and width (W) of the wounded twig segments every 2 days for 30 days to determine the lesion area (mm²). The lesion area (S) of each replicate was calculated using the following formula: $S = \pi \times L/2 \times W/2$. Re-isolations were conducted from the tested branches, and the reisolates were identified based on colony characteristics and molecular data.

The data were analyzed and figured using R (Version 4.3.3, R core team, Vienna). Two-way analysis of variance (ANOVA) was performed based on the lesion area data by taking *Cytospora* species and varieties of apple tree and wild apple as fixed factors. Post hoc test of Tukey's least significant difference was carried out by $p < 0.05$. The dominant packages used were "emmeans", "tidyverse", and "ggplot2".

Results

Phylogeny

The combined sequence dataset of ITS, *act*, *rpb2*, *tef1* and *tub2* consisted of 202 strains with *Diaporthe vaccinii* (CBS 160.32) as the outgroup taxon. In the final alignment, there are 556 characters in ITS, 324 characters in *act*,

740 characters in *rpb2*, 735 characters in *tef1* and 839 characters *tub2*. The final ML optimization likelihood value of the best RAxML tree was -60551.35, and the matrix had 2092 distinct alignment patterns, with 40.58% undetermined characters or gaps. Estimated base frequencies were as follows: A = 0.244681, C = 0.288135, G = 0.237104, T = 0.230081; substitution rates AC = 1.269712, AG = 2.848941, AT = 1.301835, CG = 0.952261, CT = 4.964970, GT = 1.0; gamma distribution shape parameter α = 0.377319. The topologies resulting from ML and BI analyses of the concatenated dataset were congruent (Fig. 1). Two isolates named XJAU 1378 and XJAU 1379 from *Malus sieversii* clustered into a distinct clade in the phylogram. Meanwhile, XJAU 866 and XJAU 867 from *M. domestica* formed another clade different from any known species (Fig. 1).

Taxonomy

Cytospora hippophaopsis R. Ma & G.F. Cai, sp. nov.

MycoBank No: 825187

Fig. 3

Etymology. Referring to its morphological similarity to *C. hippophaës*.

Descriptions. Ascstromata erumpent through the surface of bark. Disc black and circular with 6–16 ostioles per disc, ostiole black and circular, 54.5–103 μm (av. = 78.8 μm , n = 30) diam. Perithecium globular, 180.5–395 μm (av. = 285.1 μm , n = 30) diam. Asci clavate to elongate obovoid, 59.5–87 $\times$ 8–12.5 μm (av. = 73.6 $\times$ 9.9 μm , n = 30), eight-spored. Ascospores biserial, elongate-allantoid, hyaline thin-walled and smooth-walled, 13–22.5 $\times$ 3.5–5.5 μm (av. = 18.4 $\times$ 4.4 μm , n = 50).

Cultural characteristics. Colonies on PDA initially white, covering the petri dish in 7 d, becoming brown after 14 d. Ascomata formed after 16 d, white at the early stage, becoming black later.

Material examined. CHINA • Xinjiang Uygur Autonomous, Ili Autonomous Prefecture, Gongliu County, Kurdin Ning, Xiaomohe, 43°11'44.15"N, 82°43'41.66"E, 1288 m asl, on dead and dying branches of *Malus sieversii*, Rong Ma, 4 Aug 2016, holotype XJAU-1378, living cultures XJAU 1378 = CGMCC 3.18997 (ITS: PP965505, *act*: PP957863, *rpb2*: PP957870, *tef1*: PP957877, *tub2*: PP957884); *ibid.* XJAU 1379 (ITS: PP965506, *act*: PP957864, *rpb2*: PP957871, *tef1*: PP957878, *tub2*: PP957885).

Notes. Two isolates (XJAU 1378 and XJAU 1379) from the present study formed a distinct clade in the genus *Cytospora* (Fig. 2), which represents a new species named *C. hippophaopsis*. *Cytospora hippophaopsis* is morphologically similar to *C. hippophaës* from *Hippophaë rhamnoides* but differs in larger asci (59.5–87 $\times$ 8–12.5 μm vs. 38.6–44 $\times$ 5.9–7.8 μm) and ascospores (13–22.5 $\times$ 3.5–5.5 μm vs. 11.8–15 $\times$ 3–4.1 μm) (Fan et al. 2015).

Cytospora shawanensis R. Ma & G.F. Cai, sp. nov.

MycoBank No: 825189

Fig. 4

Etymology. Named after the collection site, Shawan.




Figure 2. Phylogenetic tree of *Cytospora* of ML analysis on basis of combined ITS, *act*, *rpb2*, *tef1* and *tub2* loci. Numbers above the branches indicate ML bootstraps (left, ML BS $\geq$ 50%). The tree is rooted with *Diaporthe vaccinii* (CBS 160.32). New species from the present study are marked in blue.




Figure 2. Continued.




Figure 2. Continued.

Descriptions. Stromata pycnidial, ostiolate, immersed in the host bark, scattered, producing black area on bark, circular to ovoid, with multiple locules. Conceptacle absent. Ectostromatic disc inconspicuous, producing one ostiole per disc when mature. Ostiole in the centre of the disc, black, conspicuous, 60–190 µm diam. Locules numerous, arranged circularly or elliptically with independent walls, 460–1420 µm diam. Conidiophores hyaline, unbranched or branched at the bases. Conidiogenous cells enteroblastic, phialidic, smooth-walled, tapering towards apex. Conidia hyaline, unicellular, smooth-walled, $3.5\text{--}5.5 \times 1\text{--}1.5$ µm (av. = 4.3×1.2 µm, n = 50).

Cultural characteristics. Colonies on PDA initially white, covering the petri dish in 3 d, becoming grayish green after 14 d. Conidiomata formed after 5 d, randomly distributed in the colony.

Material examined. CHINA • Xinjiang Uygur Autonomous, Tacheng City, Shawan County, Daquan Village, $44^{\circ}20'1.24''\text{N}$, $85^{\circ}37'34.40''\text{E}$, 528 m asl, on dead and dying branches of *Malus domestica*, Rong Ma, 28 Jul 2015, holotype XJAU-866, living cultures XJAU 866 = CGMCC 3.18996 (ITS: PP965507, *act*: PP957865, *rpb2*: PP957872, *tef1*: PP957879, *tub2*: PP957886); *ibid.* XJAU 867 (ITS: PP965508, *act*: PP957866, *rpb2*: PP957873, *tef1*: PP957880, *tub2*: PP957887).

Notes. *Cytospora shawanensis* is phylogenetically close to *C. olivacea* in the phylogram (Fig. 2). These two species share similar conidial morphology but differs in the sequence data (2/502 bp in ITS, 20/249 bp in *act*, 41/726 bp in *rpb2* and 39/375 bp in *tub2*). In addition, host association is helpful for species identification (*C. shawanensis* on *Malus domestica* vs. *C. olivacea* on *Sorbus tianschanica*) (Pan et al. 2020).




Figure 3. Morphology of *Cytospora hippophaopsis* (XJAU 1378) **A, B** habitat of ascostromata on the host bark **C** transverse section of the ascostromata **D** longitudinal section through the ascostromata **E–G** asci and ascospores. Scale bars: 300 µm (**B–D**); 10 µm (**E–G**).




Figure 4. Morphology of *Cytospora shawanensis* (XJAU 866) **A** habitat of conidiomata on the host branch **B** longitudinal section of the conidioma **C** transverse section through the conidioma **D** conidiogenous cells with attached conidia **E** conidia. Scale bars: 1000 µm (**A**); 200 µm (**B**, **C**); 10 µm (**D**, **E**).

Pathogenicity tests

After 30 days of incubation on apple and wild apple branches, re-isolates were obtained from the tested tissues and identified. The *Cytospora* species from the tested tissues were the same as those used for incubation, both phylogenetically and morphologically, whereas no *Cytospora* species were obtained from negative controls. As shown in the Fig. 5, *Cytospora hippophaopsis* and *C. shawanensis* create obviously larger lesion areas on apple (*Malus domestica*) and wild apple (*M. sieversii*) than that by negative control. Hence, *C. hippophaopsis* and *C. shawanensis* are both considered as *Malus* canker pathogens. Generally, *C. hippophaopsis* is more aggressive than *C. shawanensis* for the obviously larger lesion areas on Erzizi, Hanfu, Manpanz, New Century, Oil Fruit, Sitagan and Red Star. Meanwhile, these two pathogens share similar virulence to Apolte, Fuji, Golden Delicious, Jonagold, Red Star of apple and wild apple.

Discussion

Two new species of *Cytospora* are proposed from *Malus domestica* and *M. sieversii* in Xinjiang, China, namely *C. hippophaopsis* and *C. shawanensis* based on morphological and phylogenetic studies. *C. hippophaopsis* is reported from cankers of *Malus sieversii*, and *C. shawanensis* is discovered on the host *M. domestica*. Further pathogenicity tests confirmed their virulence to the branches of 13 varieties of *M. domestica* and *M. sieversii*. Hence, *Cytospora hippophaopsis* and *C. shawanensis* are two new apple canker pathogens well worth controlling.




Figure 5. Pathogenicity of different *Cytospora hippophaopsis* and *C. shawanensis* on 13 varieties of apple (*Malus domestica*) and wild apple (*M. sieversii*). Values are presented as means $\pm$ SE of three replicates. Different lowercase letters in bracket indicate significant differences among apple varieties and wild apple ($p < 0.05$). Apple varieties: AP, Apolte; EZ, Erzizi; FU, Fuji; GD, Golden Delicious; HF, Hanfu; JNG, Jonagold; MPZ, Manpanzi; NC, New Century; OF, Oil Fruit; QM, Qiulimeng; STG, Sitagan; RMA, Red Meat Apple; RS, Red Star. Wild apple: MS, *M. sieversii*.

Cytospora hippophaopsis from *Malus sieversii* is generally more aggressive than *C. shawanensis* from *M. domestica* especially on Manpanzi (Fig. 5). In addition, on the varieties Erzizi, Hanfu, Manpanz, New Century, Oil Fruit, Sitagan and Red Star, *C. hippophaopsis* is also significantly more aggressive than *C. shawanensis*. While on the natural host, *M. sieversii*, *C. shawanensis* appears a bit more aggressive (not significant) than *C. hippophaopsis*. It is inferred that Manpanzi might be the original host of *C. hippophaopsis*.

In recent years, *M. sieversii* is facing an unprecedented crisis of survival, mainly in terms of its shrinking area, reduction of biological species, single community structure, and destruction of population renewal (Volk et al. 2013). According to previous studies in the wild, it was found that the problem of pests and diseases of *M. sieversii* is very prominent (Li and Zhang 2018). This investigation found that canker disease has occurred severely on *M. sieversii* in Ili and Tacheng areas. Previous studies on the pathogenic fungal species of *M. sieversii* have lacked the records of morphological characteristics and molecular data, but this study fills this gap. To further provide more effective information for the identification of canker disease pathogens, a large number of plant specimens should be collected from various hosts and from different regions in the future. The morphological and phylogenetic analyses reported in this study provide a theoretical basis for identifying *Cytospora* and for diagnosing canker diseases and lay the foundation for further pathogenicity studies and the protection of *M. sieversii* resources.

Additional information

Conflict of interest

The authors have declared that no competing interests exist.

Ethical statement

No ethical statement was reported.

Funding

This research was funded by the Third Xinjiang Scientific Expedition Program (Grant No. 2021xjkk05005), the Study on species and pathogenic differentiation of *Cytospora* on wild apple on the north slope of Tianshan Mountains (University Research Project of the Xinjiang Uygur Autonomous Region, XJEDU2018Y023) and the National Key Research and Development Plan of China (grant number, 2016YFC05011501).

Author contributions

Conceptualization, R.M. and D.Z.; methodology, R.M. and D.Z.; software, G.C. and Y.Z.; validation, M.Y.; formal analysis, G.C. and Y.Z.; investigation, R.M. and D.Z.; resources, R.M. and D.Z.; writing—original draft preparation, G.C., Y.Z. and M.Y.; writing—review and editing, R.M. and D.Z.; visualization, G.C. and Y.Z.; supervision, R.M. and D.Z.; project administration, R.M. and D.Z.; funding acquisition, R.M. and D.Z. All authors have read and agreed to the published version of the manuscript.

Author ORCIDs

Rong Ma  <https://orcid.org/0000-0002-9865-3047>

Data availability

All of the data that support the findings of this study are available in the main text.

References

- Adams GC, Wingfield MJ, Common R, Roux J (2005) Phylogenetic relationships and morphology of *Cytospora* species and related teleomorphs (Ascomycota, Diaporthales, Valsaceae) from *Eucalyptus*. *Studies in Mycology* 52: 1–144.
- Azizi R, Ghosta Y, Ahmadpour A (2020) Morphological and molecular characterization of *Cytospora* species involved in apple decline in Iran. *Mycologia Iranica* 7(2): 205–218.
- Bozorov TA, Toshmatov ZO, Kahar G, Muhammad SM, Liu X, Zhang D, Turakulov KS (2024) Uncovering the antifungal activities of wild apple-associated bacteria against two canker-causing fungi, *Cytospora mali* and *C. parasitica*. *Scientific Reports* 14(1): 6307. <https://doi.org/10.1038/s41598-024-56969-4>
- Carbone I, Kohn LM (1999) A method for designing primer sets for speciation studies in filamentous ascomycetes. *Mycologia* 3(3): 553–556. <https://doi.org/10.1080/00275514.1999.12061051>
- Donk MA (1964) Nomina conservanda proposita 1. Proposals in fungi. *Deuteromycetes. Regnum Vegetabile* 34: 7–15.
- Doyle JJ (1990) Isolation of plant DNA from fresh tissue. *Focus* 12: 13–15.
- Ehrenberg CG (1818) *Sylvae Mycologicae Berolinenses*. Formis Theophili Bruschcke, Berlin, Germany.
- Fan XL, Hyde KD, Yang Q, Liang YM, Ma R, Tian CM (2015) *Cytospora* species associated with canker disease of three anti-desertification plants in northwestern China. *Phytotaxa* 1974(4): 227–244. <https://doi.org/10.11646/phytotaxa.197.4.1>
- Fan XL, Bezerra JDP, Tian CM, Crous PW (2020) *Cytospora* (Diaporthales) in China. *Persoonia* 45(1): 1–45. <https://doi.org/10.3767/persoonia.2020.45.01>
- Fries EM (1823) *Systema Mycologicum* 2(2): 275–620.

- Fu LG (1991) China Plant Red Data Book. Beijing: Science Press.
- Glass NL, Donaldson GC (1995) Development of primer sets designed for use with the PCR to amplify conserved genes from filamentous ascomycetes. *Applied and Environmental Microbiology* 61(4): 1323–1330. <https://doi.org/10.1128/aem.61.4.1323-1330.1995>
- Jiang N, Yang Q, Fan XL, Tian CM (2020) Identification of six *Cytospora* species on Chinese chestnut in China. *MycKeys* 62: 1–25. <https://doi.org/10.3897/mycokeys.62.47425>
- Katoh K, Standley DM (2013) MAFFT multiple sequence alignment software version 7: Improvements in performance and usability. *Molecular Biology and Evolution* 30(4): 772–780. <https://doi.org/10.1093/molbev/mst010>
- Lawrence DP, Holland LA, Nouri MT, Travadon R, Abramians A, Michailides TJ, Trouillas FP (2018) Molecular phylogeny of *Cytospora* species associated with canker diseases of fruit and nut crops in California, with the descriptions of ten new species and one new combination. *IMA Fungus* 9(2): 333–369. <https://doi.org/10.5598/ima fungus.2018.09.02.07>
- Li YX, Zhang HX (2018) Effects of dead plants on population genetic diversity of *Malus sieversii*. *Arid Zone Research* 35(1): 165–170.
- Li H, Wan Y, Wang M, Han M, Huo X (2015) History, status and prospects of the apple industry in China. *Journal of American Pomological Society* 69(4): 174–185.
- Li J, Li J, Jiang N (2024) Morphology and phylogeny of *Cytospora* (Cytosporaceae, Diaporthales) species associated with plant cankers in Tibet, China. *MycKeys* 2024(104): 51–70. <https://doi.org/10.3897/mycokeys.104.113567>
- Lin L, Pan M, Gao H, Tian C, Fan X (2023a) The potential fungal pathogens of *Euonymus japonicus* in Beijing, China. *Journal of Fungi* 9(2): 271. <https://doi.org/10.3390/jof9020271>
- Lin L, Pan M, Bezerra JD, Tian C, Fan X (2023b) Re-evaluation of the fungal diversity and pathogenicity of *Cytospora* species from *Populus* in China. *Plant Disease* 107(1): 83–96. <https://doi.org/10.1094/PDIS-02-22-0260-RE>
- Liu YJ, Whelen S, Hall BD (1999) Phylogenetic relationships among Ascomycetes: Evidence from an RNA polymerase II subunit. *Molecular Biology and Evolution* 16(12): 1799–1808. <https://doi.org/10.1093/oxfordjournals.molbev.a026092>
- Liu X, Li X, Bozorov TA, Ma R, Ma J, Zhang Y, Zhang D (2020) Characterization and pathogenicity of six *Cytospora* strains causing stem canker of wild apple in the Tianshan Forest, China. *Forest Pathology* 50(3): e12587. <https://doi.org/10.1111/efp.12587>
- Ma R, Liu YM, Yin YX, Tian CM (2018) A canker disease of apple caused by *Cytospora parasitica* recorded in China. *Forest Pathology* 48(3): e12416. <https://doi.org/10.1111/efp.12416>
- Miller MA, Pfeiffer W, Schwartz T (2010) Creating the CIPRES Science Gateway for inference of large phylogenetic trees. *Gateway Computing Environments Workshop, GCE*: 1–8. <https://doi.org/10.1109/GCE.2010.5676129>
- Norphanphoun C, Doilom M, Daranagama DA, Phookamsak R, Wen TC, Bulgakov TS, Hyde KD (2017) Revisiting the genus *Cytospora* and allied species. *Mycosphere* 8(1): 51–97. <https://doi.org/10.5943/mycosphere/8/1/7>
- Norphanphoun C, Raspé O, Jeewon R, Wen TC, Hyde KD (2018) Morphological and phylogenetic characterisation of novel *Cytospora* species associated with mangroves. *MycKeys* 38: 93–120. <https://doi.org/10.3897/mycokeys.38.28011>
- Pan M, Zhu H, Bonthond G, Tian CM, Fan XL (2020) High diversity of *Cytospora* associated with canker and dieback of Rosaceae in China, with 10 new species described. *Frontiers in Plant Science* 11: 690. <https://doi.org/10.3389/fpls.2020.00690>

- Pan M, Zhu H, Tian CM, Huang M, Fan XL (2021) Assessment of *Cytospora* isolates from conifer cankers in China, with the descriptions of four new *Cytospora* species. *Frontiers in Plant Science* 12: 636460. <https://doi.org/10.3389/fpls.2021.636460>
- Rehner SA (2001) Primers for elongation factor 1- α (EF1- α). <http://ocid.nacse.org/research/deephyphae/EF1primer.pdf>
- Richards CM, Volk GM, Reilley AA, Henk AD, Lockwood DR, Reeves PA, Forsline PL (2009) Genetic diversity and population structure in *Malus sieversii*, a wild progenitor species of domesticated apple. *Tree Genetics & Genomes* 5(2): 339–347. <https://doi.org/10.1007/s11295-008-0190-9>
- Senanayake IC, Crous PW, Groenewald JZ, Maharachchikumbura SSN, Jeewon R, Phillips AJL, Bhat JD, Perera RH, Li QR, Li WJ, Tangthirasun N, Norphanphoun C, Karunaratna SC, Camporesi E, Manawasighe IS, Al-Sadi AM, Hyde KD (2017) Families of Diaporthales based on morphological and phylogenetic evidence. *Studies in Mycology* 86(1): 217–296. <https://doi.org/10.1016/j.simyco.2017.07.003>
- Snyder F, Ni L (2017) Chinese apples and the emerging world food trade order: Food safety, international trade, and regulatory collaboration between China and the European Union. *The Chinese Journal of Comparative Law* 5(2): 253–307. <https://doi.org/10.1093/cjcl/cxx014>
- Tamura K, Stecher G, Peterson D, Filipski A, Kumar S (2013) MEGA6: Molecular evolutionary genetics analysis version 6.0. *Molecular Biology and Evolution* 30(12): 2725–2729. <https://doi.org/10.1093/molbev/mst197>
- Volk GM, Henk AD, Richards CM, Forsline PL, Chao CT (2013) *Malus sieversii*: A diverse Central Asian apple species in the USDA-ARS national plant germplasm system. *HortScience* 48(12): 1440–1444. <https://doi.org/10.21273/HORTSCI.48.12.1440>
- Wang YL, Lu Q, Decock C, Li YX, Zhang XY (2015) *Cytospora* species from *Populus* and *Salix* in China with *C. davidiana* sp. nov. *Fungal Biology* 119(5): 420–432. <https://doi.org/10.1016/j.funbio.2015.01.005>
- Wang N, Jiang S, Zhang Z, Fang H, Xu H, Wang Y, Chen X (2018) *Malus sieversii*: The origin, flavonoid synthesis mechanism, and breeding of red-skinned and red-fleshed apples. *Horticulture Research* 5(1): 70. <https://doi.org/10.1038/s41438-018-0084-4>
- Wang X, Shi CM, Gleason ML, Huang L (2020) Fungal species associated with apple *Valsa* canker in East Asia. *Phytopathology Research* 2(1): 1–14. <https://doi.org/10.1186/s42483-020-00076-5>
- Wang S, Jiang N, Ma R (2024) Morphology and phylogeny reveal three new species of *Cytospora* associated with tree cankers in China. *Journal of Fungi* 10(2): 139. <https://doi.org/10.3390/jof10020139>
- White TJ, Bruns T, Lee S, Taylor J (1990) Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: Innis MA, Gelfand DH, Sninsky JJ, White TJ (Eds) *PCR protocols: a guide to methods and applications*. Academic Press, San Diego, 315–322. <https://doi.org/10.1016/B978-0-12-372180-8.50042-1>
- Yan G, Long H, Song W, Chen R (2008) Genetic polymorphism of *Malus sieversii* populations in Xinjiang, China. *Genetic Resources and Crop Evolution* 55(1): 171–181. <https://doi.org/10.1007/s10722-007-9226-5>
- Yang FQ (2009) Integrated control of apple tree canker disease. *Xinjiang Agricultural Science and Technology* 2: 49.
- Zhu HY, Pan M, Bonthond G, Tian CM, Fan XL (2019) Diaporthalean fungi associated with canker and dieback of trees from Mount Dongling in Beijing, China. *MycKeys* 59: 67–94. <https://doi.org/10.3897/mycokeys.59.38055>
- Zhuang WY (2005) *Fungi of northwestern China*. Mycotaxon Ltd, Ithaca.