

Review

Bioactive small-molecule constituents of Lao plants

Yulin Ren^{1*}, Bethany G. Elkington^{2,3}, Joshua M. Henkin^{2,3}, Kongmany Sydara⁴, A. Douglas Kinghorn¹ and Djaja D. Soejarto^{2,3}

¹Division of Medicinal Chemistry and Pharmacognosy, College of Pharmacy, The Ohio State University, Columbus, OH 43210, United States.

²Department of Pharmaceutical Sciences, College of Pharmacy, University of Illinois at Chicago, Chicago, IL 60612, United States.

³Science and Education, Field Museum, Chicago, IL 60605, United States.

⁴Institute of Traditional Medicine, Ministry of Health, Vientiane, Lao People's Democratic Republic.

Received 9 May, 2021; Accepted 2 December, 2021

Laos has a rich plant diversity, and medicinal plants are used extensively in Lao traditional medicine for the treatment of a variety of human diseases. However, only a relatively small number of these plants have been investigated for their major components with potential antitumor, anti-infective, and other types of bioactivities. These species include *Asparagus cochinchinensis*, *Diospyros quaesita*, *Gongronema napalense*, *Marsypopetalum modestum*, *Nauclea orientalis*, *Rourea minor*, *Stemona pierrei*, and *Stemona tuberosa*. Thus far, the bioactive compounds isolated from these Lao plants include alkaloids, glycerol esters, phenolic compounds such as lignans and stilbenoids, steroids, and triterpenoids. Of these, the norlignan, niasol (1b), the triterpenes, pyracrenic acid [3 β -O-*trans*-caffeoylbetulinic acid (3)] and betulinic acid (3b), and the dimeric thiopyridine, dipyrithione (5), were found to show both cancer cell cytotoxicity and anti-infective activity. The present review focuses on examples of promising lead compounds isolated from Lao plants, with their possible development as potential therapeutic agents being discussed. It is hoped that this contribution will provide useful information on higher plants growing in Laos to help stimulate future discoveries of potential agents for the treatment of cancer, infections, and other diseases.

Key words: Lao plants, chemical compounds, human diseases.

INTRODUCTION

The Lao People's Democratic Republic, a landlocked country known widely as Laos, is a small country located in the east-central section of the Indo-Chinese peninsula, flanked by Thailand in the west, Myanmar and the Chinese province of Yunnan in the north, Vietnam in the east, and Cambodia in the south. The tropical flora of Laos is rich, and according to a recent paper (Zhu, 2017),

there are 5,005 species of seed plants distributed in 1,373 genera and 188 families in Laos, of which many plants are used as medicinal herbs in the healthcare system of Laos (Sydara et al., 2005; Libman et al., 2006). On the ethnomedical use of Lao medicinal plants, sympatric intercultural traditional knowledge has always made an impact, as indicated by a similarity investigation

*Corresponding author. E-mail: ren.41@osu.edu.

conducted among three sympatric ethnic groups, the Brou, Saek, and Kry, living in the Annamite Mountains in Laos. This study showed that medicinal plants are used commonly among these groups, but the use of these plants to treat disorders seems different (de Boer et al., 2012).

The traditional use of medicinal plants in Laos has proven useful for the discovery of new bioactive compounds, and ethnomedical collections can be supportive of plant-based drug discovery programs (Gyllenhaal et al., 2012). A field survey to identify medicinal plants growing in southern Laos showed that many Lao medicinal plants are used for treating certain infections, and their frequency of use decreased in the sequence fever, stomachache, arthritis, malaria, and leukorrhea (Elkington et al., 2014a). With the aid of field interviews with healers in Laos, several plant-derived lead compounds showing activities related to cancer, acquired immunodeficiency syndrome (AIDS) caused by infection from human immunodeficiency viruses (HIV), malaria, and/or tuberculosis (TB) infectious diseases have been identified (Soejarto et al., 2012). In addition, the Hmong ethnic group in Laos uses medicinal plants extensively for the treatment of various conditions that include gastrointestinal and gynecological conditions and sexually transmitted diseases, skin infections, kidney and bladder problems, physical trauma, and as aphrodisiacs (Dubost et al., 2019). Thus, there is a strong rationale to search for therapeutic agents from Lao plants for the potential treatment of cancer, infectious diseases, and other conditions (Soejarto et al., 2002).

Historically, plants offer valuable sources of new natural product-derived drugs, and many effective plant-derived agents have been developed for the treatment of cancer and infectious diseases over the past few decades (Dehelean et al., 2021; Newman and Cragg, 2020; Talib et al., 2020). Therefore, as a consequence of the continuing search for antineoplastic and anti-infective agents from Lao plants, substances targeting unusual mechanisms to afford possible new approaches to improve human health may be discovered (Soejarto et al., 2006; Kinghorn et al., 2016). In addition, a reservoir represented by a large number of higher plants growing in Laos offers an opportunity for such a search (Henkin et al., 2017), of which eight species have been investigated for identification of their bioactive components. Subsequently, investigations on these plants are summarized, inclusive of further studies on their bioactive principles, when collected from neighboring countries, and a perspective for future investigations on Lao plants.

ASPARAGACEAE (*ASPARAGUS COCHINCHINENSIS* [LOUR.] MERR.)

A. cochinchinensis is a perennial trailing plant (Table 1) native to Cambodia, China, Japan, Korea, Laos, the


Philippines, and Vietnam (Newman et al., 2007). Based on information provided by a healer from Salavan Province of Laos, the roots of this plant are used in folk medicine to treat chronic fever when combined with other plants (Soejarto et al., 2012), and thus they were collected locally, from which several compounds were isolated. These include a 9'-norlignan, 3-hydroxy-4-methoxy-4-dehydroxynyasol (**1**), and its analogues, 3-methoxynyasol (**1a**), nyasol (**1b**), and 1,3-bis-di-*p*-hydroxyphenyl-4-penten-1-one (**1c**) (Figure 1), along with a new spirostanol saponin, asparacoside (**2**), and its analogues asparacosins A (**2a**) and B (**2b**) (Figure 1). These compounds were evaluated for cytotoxicity against Col-2 colon, KB oral epidermoid, LNCaP prostate, and Lu-1 lung human cancer cells, and **1**, **1a**, and **2** were deemed active, but **1b**, **1c**, **2a**, and **2b** were not (Zhang et al., 2004). These results demonstrated that the 3,4-*ortho*-hydroxy and -methoxy groups are important for the mediation of cancer cell line cytotoxicity by **1**, but neither transposing the substituents between the C-3 and C-4 positions nor introducing a carbonyl group at the C-9' position seems to contribute to the resultant cytotoxic potency.

Even though nyasol (*cis*-hinokiresinol, **1b**) did not show any growth inhibitory activity against human cancer cells, it was found to inhibit both farnesyl protein transferase (FPTase) and phosphatase of regenerating liver 3 (PRL-3). FPTase is involved in the membrane association of Ras to modulate tumor formation, while PRL-3 is overexpressed in human cancer cells to contribute to the acquisition of metastasis. Thus, nyasol shows some antitumor potential by targeting FPTase and PRL-3 to inhibit tumor formation and metastasis (Song et al., 2008). In addition, nyasol (**1b**) was found to inhibit selectively the basic fibroblast growth factor (bFGF) and vascular endothelial growth factor (VEGF)-induced endothelial cell proliferation, migration, and tube formation. The vessel growth was inhibited when slow release pellets of Hydron N containing 200 ng mouse recombinant VEGF were implanted into the corneal pockets of five-week-old BALB/cA mice, and then mice were treated intraperitoneally (i.p.) with nyasol (100 mg/kg, daily) for seven days. Angiogenesis is observed commonly when tumors occur, for which inhibition may contribute potentially to cancer treatment. Thus, nyasol could be an antitumor lead compound, targeting angiogenesis inhibition (Jeong et al., 2003).

Investigation of cytotoxicity of the spirosteroid **2** and its analogues isolated from the Lao plant, *A. cochinchinensis* (Zhang et al., 2004), indicates that the glycoside moiety connected at the C-3 position is necessary for the modulation of its cancer cell cytotoxicity, while the C-3–C-5 enone group plus the C-12 hydroxy group also contribute to this activity. Steroids occur widely in both plants and animals, among which cardiac glycosides have attracted wide interest for their potential antitumor activity. Of these, digoxin has long been used for the

Table 1. Selected Lao plants and their derivatives with anti-infective and cytotoxic activities.

Lao plants	Derivative and bioactivities	References
<i>Asparagus cochinchinensis</i>	1 , 1a-1c , 2 , 2a , and 2b , of which 1 , 1a and 2 showed cytotoxicity toward Col-2, KB, LNCaP, and Lu-1 cells (IC_{50} 4.0–41 μ M), while 1b and 1c showed anti-HIV activity, with IC_{50} being 75 and 46 μ M, respectively.	Zhang et al. (2004)
<i>Diospyros quaesita</i>	3 showed cytotoxicity against KB cells (IC_{50} 4.0 μ M), anti-TB (MIC 323 μ M), antimalarial (IC_{50} <1 μ M), anti-DENV NS5 RdRp (IC_{50} 0.85 μ M), and <i>in vivo</i> anti-inflammatory (25 mg/kg) activities.	Ma et al. (2008), Tanachatchairatan et al. (2008), Otsuka et al. (1981), and Bourjot et al. (2012)
<i>Gongronema napalense</i>	4 showed antimalarial activity (IC_{50} 1–2 μ M).	Libman et al., 2008
<i>Marsypopetalum modestum</i>	5 induced HeLa cell apoptosis (IC_{50} 0.91 μ M) and showed cytotoxicity toward K562, KB, MDA-MB-231, and U937 cells (IC_{50} 4.0–9.9 μ M), anti-TB (MIC <0.15 μ M), and <i>in vivo</i> anti-inflammatory (2.5 mg/kg) activities.	Elkington et al. (2014b), Fan et al. (2007, 2013), and Huang et al. (2011)
<i>Nauclea orientalis</i>	6 , 7 , and 8 showed antimalarial activity (IC_{50} 4–13 μ M), and 7 also showed anti-TB activity (MIC 110 μ M).	He et al. (2005) and Tanachatchairatana et al. (2008)
<i>Rourea minor</i>	9 , 10 , and 10a showed antimalarial activity (IC_{50} 2–13 μ M).	He et al. (2006)
<i>Stemona pierrei</i>	12 showed weak activity against Hep-G2, KB, MCF-7, and SK-Lu-1 cells (IC_{50} 77–119 μ M).	Quang et al. (2014)
<i>Stemona tuberosa</i>	11 , 11a , 11b , 12a , and 13 , of which 11b showed cytotoxicity toward KB cells (IC_{50} 17 μ M).	Khamko et al. (2013) and Quang et al. (2014)

**Figure 1.** Structures of the norlignan 3-hydroxy-4-methoxy-4-dehydroxyynyasol (**1**) and its analogues (**1a-1c**) and the steroid asparacoside (**2**) and its analogues (**2a** and **2b**) isolated from the roots of *Asparagus cochinchinensis* collected in Laos. Zhang et al. (2004).

treatment of congestive heart failure, and it was also found to show some potential antitumor activity by targeting Na^+/K^+ -ATPase (Ren et al., 2019). Recently,

another cardiac glycoside, (+)-strebloside, was identified as a promising antitumor lead compound, targeting Na^+/K^+ -ATPase, nuclear factor κ light-chain-enhancer of

activated B cells (NF- κ B), and p53 and the extracellular signal-regulated kinases (ERK) pathways (Chen et al., 2017). Comparison of the cytotoxicity of **2**, **2a**, **2b**, and the cardiac glycosides digoxin and (+)-strebloside implies that the C-22 spiro-bicyclic system of **2** seems less important than the C-17 lactone unit of cardiac glycosides in the mediation of their cytotoxic potency. However, further modification of the C-22 spiro-bicyclic system of **2** may produce some novel non- Na^+/K^+ -ATPase-targeted antitumor steroids. Thus, **2** could be regarded as a lead compound, and the search for more cytotoxic analogues of **2** from plants collected in Laos may be supportive in this regard.

Moreover, the norlignans **1**, **1a**, and **1b**, along with their analogue, 4'-O-methylnyasol, isolated from *Anemarrhena asphodeloides*, were found to show inhibitory effects on the degranulation of rat basophilic leukemia-2H3 cells. The inhibition of β -hexosaminidase release of **1** (IC_{50} 2.9 μM) was more potent than that of ketotifen (IC_{50} 10 μM), a second-generation H_1 -antihistamine used to treat allergic conjunctivitis, and thus **1** could be promising as an antiallergic lead (Bak et al., 2016). Alteration of the substituents at the C-3 and C-4 positions of **1** resulted in a significant decrease of its antiallergic potency, indicating that this norlignan could be modified synthetically in future investigations.

Also, some of these norlignans were found to exhibit antiviral activity. For example, **1b** and **1c** exhibited moderate anti-HIV activity, but **1** and **1a** did not (Zhang et al., 2004), indicating that a hydroxy group substituted at the C-4 and C-4' positions is required for this biological property. Introducing a methoxy group at the C-3 position or substitution of a hydroxy group at the C-3 position followed by methylation of the C-4 hydroxy group resulted in the elimination of the anti-HIV activity, while changing the $\Delta^{7,8'}$ double bond to a carbonyl group did not affect the activity greatly. In addition, (–)-nyasol and (–)-4'-O-methylnyasol were found to inhibit effectively respiratory syncytial virus (RSV), which frequently infects infants. Both compounds were more potently active in this regard than ribavirin, a standard antiviral drug with a broad-spectrum of activity. Thus, these norlignans show some promise as anti-HIV or anti-RSV agents, and their activity could be improved through synthetic optimization (Bae, 2007).

Nyasol (**1b**) was identified originally as a derivative of nyasoside, a new norlignan glycoside isolated from the rhizomes of *Hypoxis nyasica* Bak. (Hypoxidaceae) collected near Zomba in Malawi (Marini-Bettolo et al., 1985). In turn, (+)-nyasol was isolated from the tubers of *A. cochinchinensis* collected in Hong Kong (Tsui and Brown, 1996). The absolute configurations of (+)- and (–)-nyasol have been determined by synthetic methods and by analysis of their vibrational circular dichroism spectra (Lassen et al., 2005). In an earlier investigation, nyasol (**1b**) was found to inhibit the lipopolysaccharide (LPS)-induced release of nitric oxide (NO) in RAW264.7 murine

monocytic cells, with no discernible cytotoxicity observed against the host cells. The inhibitory potency of **1b** was equivalent to that of 1- N_6 -(1-iminoethyl) lysine, a potent inhibitor of nitric oxide synthase (NOS), indicating some anti-inflammatory potentiality of **1b** (Song et al., 2008). Following this, (–)-nyasol isolated from the rhizomes of *A. asphodeloides* obtained in Korea was found to inhibit production of cyclooxygenase (COX)-2-mediated prostaglandin E2 (PGE2) and inducible NOS (iNOS)-induced nitric oxide (NO) in LPS-treated RAW 264.7 cells. It also suppressed the generation of 5-lipoxygenase (5-LOX)-mediated leukotriene in RBL-1 rat basophilic leukemia cells treated with A-23187 (ionophore) (Lim et al., 2009) and the expression of inflammatory cytokines such as interleukin-1 β (IL-1 β) and interferon- β (IFN- β) (Lee et al., 2014). Furthermore, nyasol (**1b**) inhibited the transcriptional activity of NF- κ B, the activation of Akt or PKB (protein kinase B), and ERK in LPS-stimulated RAW 264.7 cells (Lee et al., 2014).

Further investigations have shown that (–)-nyasol inhibited the production of NO and PGE2 and the expression of iNOS and COX-2 in LPS-activated BV-2 murine microglial cells, through suppressing LPS-induced I- κ B α degradation and p38 mitogen-activated protein kinase (MAPK) activation. Activation of the microglia contributes to host damage by excessive release of various proinflammatory cytokines such as IL-1 β and tumor necrosis factor (TNF)- α and is associated with the occurrence of several neurodegenerative diseases, including Alzheimer's disease, Parkinson's disease, and stroke. Thus, inhibition of these proinflammatory molecules could be useful for the treatment of these disorders. In this regard, (–)-nyasol may have therapeutic potential for the treatment of neuroinflammatory diseases (Lee et al., 2013).

Importantly, (–)-nyasol showed anti-inflammation activity *in vivo*. It inhibited potently λ -carrageenan (CGN)-induced paw edema when male ICR mice were treated intraperitoneally (i.p.) with (–)-nyasol (24–120 mg/kg, a single dose), and the right hind paw edema of mice was induced by injection of 0.05 mg of CGN (Lim et al., 2009). Nyasol inhibited 12-O-tetradecanoylphorbol 13-acetate (TPA)-induced mouse ear edema when ICR mice were treated (i.p.) with nyasol (10 mg/kg, a single dose), and the right ear of mice was applied topically by TPA (1.0 μg in 20 μL acetone) (Lee et al., 2014).

It is worthy of note that nyasol (**1b**) may possess antidiabetic properties. It was found recently to inhibit potently α -glucosidase, with its IC_{50} value being 3 nM. Such potency is much greater (up to 5700-fold) than that found for acarbose (IC_{50} 171 μM), a standard antidiabetic drug targeting α -glucosidase, an intestinal enzyme to digest large carbohydrates into glycoses (Tantapakul et al., 2020).

Similar to the norlignans, the spirosteroid asparacosin A (**2a**) was also found to show both anti-inflammatory and antinociceptive activities. In an anti-inflammation rat

model, the paw edema of Sprague Dawley rats (150-220 g) was inhibited significantly when the animals were treated orally with asparacosin A (20 mg/kg, single dose), and then (after 1 h) the right hind paws of rats were injected plethysmometrically by carrageenan (0.1 mL of 1% suspension in 2% gum acacia saline solution). In the same model, ear edema was suppressed when rats were treated orally with asparacosin A (**2a**, 20 mg/kg, single dose), and then (after 1 h) the inner and outer surfaces of the right ear of rats were given topically 20 μ L of xylene. Moreover, in the writhing antinociceptive mouse model, the writhes observed were reduced significantly when Swiss mice (20–30 g) were treated orally with **2a** (10 mg/kg, single dose) and then (after 0.5 h) injected (i.p.) by 1% acetic acid. Mechanistic investigations indicated that the anti-inflammatory activity could be mediated by **2a** through inhibition of COX-2 enzyme and inflammatory mediators, TNF- α , IL-1 β , and PGE2. Therefore, asparacosin A may be regarded as a lead for the development of anti-inflammatory and antinociceptive agents (Karim et al., 2019).

The roots of *A. cochinchinensis* also have been used as a Chinese traditional medicine, “Tianmendong”, to treat cancer and infectious diseases (Zhang and Shen, 2007), and they were found recently to show anti-asthmatic activity (Choi et al., 2018). Phytochemical investigation of the roots of *A. cochinchinensis* obtained from China resulted in the isolation of two 8,9-dihydroxylated analogues of niasol (**1b**), iso-agatharesinol and iso-agatharesinoside (Li et al., 2012). In addition, two steroids, dioscin and methyl protodioscin, were isolated from the roots of *A. cochinchinensis* obtained from Korea, with their analogue, aspacochioside A, being identified as a major component (2.126 g from 18.6 kg of the dried root sample) of the roots of *A. cochinchinensis* collected in China (Lee et al., 2015; Shi et al., 2004). Comparison of these components with those reported from Lao *A. cochinchinensis* indicates that the major components of this plant do not vary due to the change of the collection locations in Asia. As reviewed recently, dioscin shows multiple pharmacological activities, including antitumor, anti-infection, and tissue-protective properties (Yang et al., 2019). Both dioscin and methyl protodioscin were found to suppress the expression of the MUC5AC mucin gene in NCI-H292 pulmonary mucoepidermoid carcinoma cells. Airway mucus plays a crucial role in defense against invading pathogens, but the abnormality of mucins could cause altered airway physiology to lead to severe airway pathology. The inhibitory effect of both dioscin and methyl protodioscin on the expression of MUC5AC mucin suggesting that these steroids could show some potential for the treatment of inflammatory pulmonary diseases (Lee et al., 2015).

Similar to dioscin, methyl protodioscin has antitumor potential. It induced HeLa human cervical cancer cell apoptosis, which was mediated by the accumulation of intracellular reactive oxygen species (ROS) and by the

activation of death receptor and mitochondrial pathways (Ma et al., 2019). It suppressed MG-63 human osteosarcoma cell growth through induction of cell apoptosis by, at least in part, caspase-dependent and MAPK signaling pathways (Tseng et al., 2017). Treatment with methyl protodioscin led to cell cycle arrest at the G2/M phase and resulted in inhibition of SAS and SCC9 human oral squamous cell carcinoma cell growth, which was mediated probably by induction of apoptosis and autophagy through modulation of the p38 MAPK and JNK1/2 pathways (Hsieh et al., 2017).

Interestingly, methyl protodioscin also inhibited the A7r5 rat thoracic aorta fibroblast cell growth. Neointima formation was suppressed significantly after an extravascular administration of methyl protodioscin when the balloon catheter was introduced into the common carotid artery of three-four-month-old male rats, with the carotid artery then being coated with 30% (w/v) pluronic-F127 gel with methyl protodioscin (3.0 or 6.0 μ M) for two weeks. Also, the expression of a disintegrin and metalloprotease 15 (ADAM15) in vascular smooth muscle cells (VSMCs) and injured mouse arteries was found to be reduced markedly. Abnormal proliferation and migration of VSMCs is involved in the development of restenosis, and ADAM15 expression in VSMCs plays a critical role in pathological neovascularization and atherosclerosis. Methyl protodioscin inhibited VSMC growth and migration and prevented balloon angioplasty-induced restenosis by downregulating the expression of ADAM15 and its downstream signaling pathways, and thus it may possess some therapeutic potential for the treatment of restenosis (Chung et al., 2016).

Therefore, the norlignan niasol (**1b**) and asparacosin A (**2a**) and their analogues may be regarded as the principal active components of *A. cochinchinensis* that are responsible for the medicinal use of this herb in Lao folk medicine. Further modification of these lead compounds could yield some therapeutic agents to treat cancer and infectious diseases.

EBENACEAE (*DIOSPYROS QUAESITA* THW.)

D. quaesita is a small to medium-sized tree (Table 1), and its leaves and stems are used to treat cases of alcohol and drug dependence and headache in Laos (Soejarto et al., 2012). Thus, the leaves, twigs, and branches of *D. quaesita* were collected in Sainyabuli Province of Laos, from which a caffeoyl ester of betulinic acid, namely, pyracrenic acid [3β -O-*trans*-caffeoylbetulinic acid (**3**)] (Figure 2), was isolated. Compound **3** was found to show cytotoxicity toward KB human oral epidermoid cancer cells (Ma et al., 2008). It also exhibited activity against the human A549 and SK-Lu-1 lung, AGS gastric, COLO 205 and HCT-15 colon, HeLa cervical, HepG2 liver, MCF-7 breast, and SK-OV-3 ovarian cancer and SK-MEL-2 melanoma cell lines (Table 2) (Kim et al., 2010,




Figure 2. Structures of the triterpenoid pyracrenic acid (3β-*O*-*trans*-caffeoyl betulinic acid, **3**) and betulinic acid (**3b**) isolated from the Lao plant, *Diospyros quaesita*, and of selected derivatives **3a** and **3c–3g**. Lomchid et al. (2017), Ma et al. (2008), and Shi et al. (2014).

Table 2. Cytotoxicity of compounds **3** and **3a–3g**.

Compound	A549 ^{a,1,2}	AGS ^{b,3}	COLO205 ^{c,3}	HepG2 ^{d,3}	KB ^{e,4}
3	4.4	7.4	16	20	4.0
3a	-	-	-	-	3.0
3b	>11	-	-	-	>88
3c	>10	-	-	-	-
3d	>10	-	-	-	-
3e	-	>100	>100	>100	-
3f	-	>100	>100	>100	-
3g	-	15	15	37	-

IC₅₀ values (μM) toward the human ^aA549 non-small cell lung, ^bAGS gastric, ^cCOLO 205 colon, ^dHepG2 liver, and ^eKB oral epidermoid cancer cell lines. ¹(Kim et al., 2010), ²(Shi et al., 2014), ³(Pan et al., 2008), and ⁴(Ma et al., 2008). - (Not reported).

2017; Pan et al., 2008; Quang et al., 2020; Wang et al., 2018).

Investigation of the cytotoxicity of **3** and its analogues or derivatives indicated that the introduction of a 3-*O*-caffeoyl group enhances the cytotoxic potency of betulinic acid (**3b**) against KB cells, and acetylation of the caffeoyl unit [e.g., 3β-*O*-*trans*-diacetylcaffeoyl betulinic acid (**3a**)] led to the activity being improved further (Ma et al., 2008). In addition, this type of activity was enhanced when the C-28 carboxylic acid substituent was reduced to a primary hydroxy group (Lomchid et al., 2017). As a result of treating A549 cells with **3**, **3b**, betulin (**3c**), lupeol (**3d**), 3β-*O*-*trans*-caffeoyl betulin (**3e**), and 3β-*O*-*trans*-*p*-coumaroyl betulin (**3f**), the cytotoxic potency was increased in the sequence, **3d**, **3c**, **3f**, **3b**, and **3e**. Hence, oxidation of the C-28 methyl group of lupeol (**3d**) resulted in its activity being increased, and the presence of a 6',7'-dihydroxy group was found to be preferable for this type of activity of **3** (Shi et al., 2014). In addition,

compound **3b** was found to be more potently active than **3** when tested against A549, HCT-15, SK-MEL-2, and SK-OV-3 cells (Kim et al., 2017), indicating that introducing a 3-*O*-caffeoyl group may decrease the activity of **3b** against these cell lines tested. Thus, the effect on the cytotoxic potency of the 3-*O*-caffeoyl group of **3** may vary when investigated against different cancer cell lines.

When evaluated against AGS, COLO 205, and HepG2 cancer cells, **3** showed activities that were comparable with those of 3β-*O*-*trans*-*p*-coumaroyl betulinic acid (**3g**), and both **3** and **3g** were more potently cytotoxic than 3β-*O*-*trans*-feruloyl betulinic acid for AGS and COLO cells (Pan et al., 2008). These results indicated that removal of the C-6' hydroxy group did not affect the activity of **3**, but methylation of this hydroxy group reduced its cytotoxicity. Thus, an analogue of compound **3**, namely, 3β-*O*-*trans*-feruloyl betulinic acid, was found to be inactive when tested against MDA-MB-231 human breast cancer cells

(Ren et al., 2018). In addition, several derivatives of **3** were found to be more potently active than the corresponding analogues of **3e** when tested against AGS, COLO 205, and HepG2 cells, implying that the C-28 carboxylic acid moiety is preferred over a C-28 hydroxy group in enhancing the activity of these compounds against the cancer cell lines used (Pan et al., 2008).

Additionally, compound **3** showed antimalarial activity against the chloroquine-sensitive clone D6 and the chloroquine-resistant clone W2 of *Plasmodium falciparum*, and it exhibited an equivalent potency to 3 β -*O*-*trans*-diacetylcaffeoylbetulinic acid (**3a**) and was more active than betulinic acid (**3b**) (Ma et al., 2008). These results demonstrated that introducing a 3 β -*O*-*trans*-caffeoyl group can enhance the antimalarial effects of betulinic acid (**3b**) *in vitro*, but acetylation of **3** did not change the activity significantly. Also, **3** was found to be active when tested *in vitro* against *Mycobacterium tuberculosis* (TB) H37Ra, using the standard drugs, kanamycin sulfate, isoniazid, and rifampicin as positive controls (Tanachatchairatana et al. 2008).

Importantly, compound **3** was found to inhibit the granulation tissue formation in male Wistar rats, as induced by subcutaneous implantation of cotton pellets when these test animals were treated orally with **3** (25 or 50 mg/kg, daily) for seven days (Otsuka et al., 1981). This anti-inflammatory activity has been supported by the inhibitory effects observed for **3** on the activation of NF- κ B, a key transcription factor in chronic inflammatory diseases (Andre et al., 2013), as well its antioxidative activities, as demonstrated by its 1,1-diphenyl-2-picrylhydrazyl (DPPH) radical-scavenging and copper ion-induced oxidation suppression effects (Cho et al., 2013). In addition, **3** was found to exhibit potent inhibitory activity against the dengue virus (DENV) non-structural protein 5 (NS5) RNA-dependent RNA polymerase (RdRp), with an IC₅₀ of 0.85 μ M (3'-deoxy-GTP, IC₅₀ 0.02 μ M) (Bourjot et al., 2012). DENV NS5 RdRp is responsible for viral replication and transcription and has become an attractive target for the development of anti-dengue drugs (Lim et al., 2015; Shimizu et al., 2019), and thus, compound **3** shows some promise of having antidengue activity.

Isolated also from *D. quaesita* collected in Laos was betulinic acid (BA, **3b**), a pentacyclic lupane-type triterpenoid occurring widely in plants (Ma et al., 2008). Compound **3b** has shown wide range of pharmacological activities, including antidiabetic, antihyperlipidemic, anti-infective, and antitumor properties, as reviewed recently (Ríos and Máñez, 2018; Amiri et al., 2020). More recently, determination of the mechanisms of antitumor and anti-infective activities of **3b**, as well as improvement of its pharmaceutical potential through the development of different formulations, has attracted wide interest. Compound **3b** prohibited the proliferation and migration of BGC-823 and MNK45 human gastric cancer cells

(Chen et al., 2020b), and it reduced significantly A172, U87, and U87R (temozolomide resistant) human glioblastoma cell growth through activation of the protein kinase RNA-like endoplasmic reticulum kinase (PERK)/C/EBP homologous protein (CHOP) apoptotic pathway (Lo et al., 2020). In addition, **3b** inhibited invasion and migration of highly aggressive MDA MB-231 and BT-549 human breast cancer cells by targeting glucose-regulated protein 78 (GRP78) to suppress aerobic glycolysis in cells. It also retarded breast cancer lung colonization through GRP78/ β -catenin/c-Myc signaling (Zheng et al., 2019).

To enhance the possible therapeutic applications of betulinic acid (**3b**), several different kinds of nanoscale delivery systems have been developed (Saneja et al., 2018), and the antitumor potential of **3b** has been improved through these delivery systems. For example, nanoliposomes of **3b** showed more potent inhibitory activity against the proliferation and glucose transport of HCT116 human colon cancer cells than **3b** itself (Wang et al., 2020a). This type of activity was mediated by targeting hexokinase (HK), phosphofructokinase-1 (PFK-1), phosphoenolpyruvate (PEP), and pyruvate kinase M2 (PKM2). Thus, the liposomal **3b** could be used as an effective adjuvant colon cancer therapy, by targeting tumor metabolic pathways (Wang et al., 2020b). A carrier-free disulfide-modified glutathione (GSH)-responsive **3b** has been prepared, and a photosensitive prodrug, BA-S-S/Ce6 nanoparticles (NPs), has been constructed. Under irradiation, BA-S-S/Ce6 NPs exhibited more potent cytotoxicity than Ce6 against 4T1 musculus breast cancer cells, and it improved the antitumor efficacy of Ce6 (Cheng et al., 2021).

Several conjugated compounds, with **3b** being complexed with cabazitaxel, podophyllotoxin, and thiocolchicine, were synthesized, and the corresponding nanoassemblies were formed by the self-assembly of these synthetic products. Both the monomers and nanoparticles of the conjugates showed similarly potent activities against A2780 human ovarian cancer cells, but they were less active than those of the standard antitumor agents used. However, introduction of a properly designed self-immolative linker could facilitate the release of these drugs to improve their biological activity (Colombo et al., 2020).

Additionally, **3b** showed anti-infective activity (Ríos and Máñez, 2018; Amiri et al., 2020). Symptoms in rats with collagen-induced arthritis (CIA) were alleviated by treatment with **3b**, and the histopathological features in the joints of the test animals were improved, with the secretion of inflammatory cytokines, including IL-1 β , IL-6, and TNF- α being inhibited. These activities were mediated by inhibition of transcription of VEGF and transforming growth factor β (TGF- β), and activation of the NF- κ B pathway (Kun-Liu et al., 2020). Recently, a new betulinic acid-nucleoside hybrid was found to show potent anti-HIV activity, with an IC₅₀ value of 7.8 nM




Figure 3. Structure of the steroid glycoside gongroneside A (4) isolated from the Lao plant, *Gongronema napalense*. Libman et al. (2008).

(Wang et al, 2020c). Some inhibitory effects against DENV infection were also found for this triterpene, which inhibited a post-entry stage of the DENV replication cycle, RNA synthesis, and protein production (Loe et al., 2020).

Compound **3b** was also found to inhibit significantly the receptor activator of nuclear factor κ B ligand (RANKL)-induced osteoclastogenesis (Jeong et al., 2020) and α -glucosidase (Du et al., 2021). In EA.hy926 human vascular endothelial cells, **3b** enhanced intracellular Ca^{2+} levels to increase endothelial nitric oxide synthase (eNOS) expression through phosphorylation of 5' adenosine monophosphate-activated protein kinase (AMPK) and activation of histone deacetylase 5 (HDAC5) and ERK5. NO produced by eNOS is crucial in endothelial function, and thus, **3b** possesses some potential to support the treatment of cardiovascular diseases through preventing vascular endothelial dysfunction (Lee et al., 2020). In Mlg mouse lung fibroblast and primary pulmonary fibroblast (PPF) cells, Wntless-related integration site (Wnt) 3a-induced myofibroblast activation and Wnt/ β -catenin signaling were inhibited by **3b**. Idiopathic pulmonary fibrosis (IPF) is a fatal fibrotic lung disease, and aberrant activation of the Wnt/ β -catenin signaling cascade plays the key role in the pathogenesis of IPF. Thus, **3b** shows some antipulmonary fibrosis activity (Li et al., 2021).

Accordingly, 3 β -*O*-*trans*-caffeoylbetulinic acid (**3**) and betulinic acid (**3b**) could be the principal active components of the leaves and stems of *D. quaesita*, which are used to treat cases of alcohol and drug dependence and headache in Laos.

ASCLEPIADACEAE (*GONGRONEMA NAPALENSE* [WALL.] DECNE.)

G. napalense is a vine (Table 1) native to Bangladesh, China, India, Laos, Myanmar, Nepal, and Vietnam (Newman et al., 2007) and used to treat leucorrhea,

blennorrhoea, and traumatic injury in Laos (Soejarto et al., 2012). In a search of antimalarial agents from Lao plants, a sterol hexaglycoside, gongroneside A (4) (Figure 3), was characterized an active component of the entire vines of *G. napalense* collected in Champasak Province of Laos, and it exhibited potent antimalarial activity *in vitro* (Libman et al., 2008).

ANNONACEAE (*MARSYPOPETALUM MODESTUM* [PIERRE] B. XUE & R.M.K. SAUNDERS)

In an ethnobotanical search for tuberculosis (TB) treatments using Lao plants, healers in Bokeo, Bolikhamxai, Champasak, Luang Prabang, and Vientiane Provinces were interviewed. Based on these interviews, more than 300 plants were collected and tested for anti-TB activity against virulent *Mycobacterium tuberculosis* H37Rv, and an ethanol extract of *M. modestum* (small tree), was found to be the most potently active. A follow-up bioassay-guided isolation led to the identification of dipyrithione (**5**) (Figure 4) as a major active component, which demonstrated a minimum inhibitory concentration (MIC) of $<0.15 \mu\text{M}$ against virulent mycobacteria (Elkington et al., 2014b).

Dipyrithione (**5**) also exhibited growth inhibitory effects on *M. tuberculosis* BCG (Kawabe et al., 1967) and several fast-growing species of *Mycobacterium*, including *Mycobacterium fortuitum*, *Mycobacterium smegmatis*, and *Mycobacterium phlei* (O'Donnell et al., 2009), as well several Gram-positive and -negative bacteria (Kawabe et al., 1967). While dipyrithione (**5**) suppressed the growth of methicillin-resistant *Staphylococcus aureus*, *S. aureus* EMRSA-15, and *S. aureus* XU212, it inhibited the strain growth of a multidrug-resistant (MDR) variant of *S. aureus*, *S. aureus* SA-1199B, which overexpresses the NorA efflux transporter (O'Donnell et al., 2009).

Interestingly, two monomeric analogues of dipyrithione (**5**), 2-methyldithiopyridine *N*-oxide (**5a**) and 2-




Figure 4. Structures of the thiopyridine dipyrithione (**5**) isolated from the Lao plant, *Marsypopetalum modestum*, and its analogues (**5a–5e**) (Elkington et al. 2014b, O'Donnell et al. 2009, Krejčová et al. 2014, Liu et al. 2008, Srivastava et al. 2015, Tailler et al. 2012).

methylthiomethyldithiopyridine *N*-oxide (**5b**), were found to exhibit antibacterial activity against a small panel of bacterial strains, with both **5a** and **5b** being demonstrated as potently active against *Mycobacterium bovis* BCG and *M. tuberculosis* H37Rv. Of these, **5b** also inhibited completely the incorporation of ^{14}C -labeled acetate into soluble fatty acids, indicating that this substance is a potential fatty acid biosynthesis inhibitor (O'Donnell et al., 2009). Following this, the maximum tolerated dose (MTD) and anti-TB efficacy of **5b** were evaluated in animal models. Acute adverse effects were observed at the doses, 100 and 300 mg/kg, when C57BL/6 female mice were treated orally with **5b** (a single dose of 30, 100, or 300 mg/kg) and checked post-administration at 4 and 6 h and then twice daily for a week. However, this compound did not show any anti-infective activity when eight- to ten-week-old female specific-pathogen-free C57BL/6-gamma interferon gene-disrupted (GKO) mice were infected by *M. tuberculosis* Erdman (one day) and treated orally with **5b** (30 mg/kg) for nine days (O'Donnell et al., 2009). Thus, modification of **5b** to improve its anti-TB properties would be required in future investigations.

Inflammation is a complex biological response to harmful stimuli, including infections, injuries, and toxins, and overexpression of the C-X-C motif chemokine ligand 10 (CXCL10), a small cytokine known as IFN- γ -induced protein 10 (IP-10) or IP-10/CXCL10, regulates almost all of its stages. Dipyrithione (**5**) was found to suppress IP-10/CXCL10 expression (Han et al., 2010), but it did not show any observed effects on the activity of COX-1 and COX-2 (Krejčová et al., 2014). However, two monomeric analogues of **5**, 2-methylthiomethyldithiopyridine *N*-oxide (**5b**) and 2-methylthiomethyldithiopyridine (**5c**) (Figure 4), were found to inhibit COX-1 and COX-2, while **5**, 2-methyldithiopyridine *N*-oxide (**5a**), and some other analogues, did not. This indicated that a methylthiomethyldithio group is required to mediate COX-1 and COX-2 inhibitory activities, which are involved in the conversion of arachidonic acid to prostaglandins, and thus **5b** and **5c** both show some potential for the development as new anti-inflammatory agents (Krejčová et al., 2014).

Compound **5** was also found to inhibit LPS-induced expression of iNOS and NO production, and it also

reduced the LPS-induced increase of COX-2 protein levels in RAW264.7 murine macrophage-like cells. NO produced by iNOS and prostaglandins induced by COX-2 are two of the most prominent inflammatory mediators, and thus dipyrithione (**5**) appears to possess promising anti-inflammatory effects (Liu et al., 2008). Mouse survival rates were increased when male ICR mice were injected with LPS and treated (i.p.) (30 minutes later) with **5** (1, 2.5 or 5.0 mg/kg) or dexamethasone (2 mg/kg) (Liu et al., 2008). In a mouse acute lung injury model induced by oleic acid, the lung injury was attenuated significantly by dipyrithione (**5**) when six–seven-week-old male ICR mice were dosed by caudal intravenous (i.v.) administration of 0.3 mL/kg of oleic acid and treated (i.p., single dose) with **5** (1.25, 2.5, or 5 mg/kg) or dexamethasone sodium phosphate (5 mg/kg), before or after oleic acid induction (30 minutes). Also, dipyrithione treatment alleviated significantly microvascular leakage and inhibited oleic acid-induced increase of intercellular adhesion molecule-1 (ICAM-1) and vascular cell adhesion molecule (VCAM-1) expression in the mouse lung (Huang et al., 2011).

Pyrithione (2-mercaptopyridine-1-oxide, **5d**) (Figure 4) has been used as a bactericide and a fungicide for many years, from which **5** may be formed by self-oxidation of the free SH group of pyrithione. In RAW264.7 murine macrophage-like cells, **5** exhibited more potent activity than **5d** in inhibiting LPS-induced iNOS and COX-2 expression, and than aspirin in inhibiting LPS-induced NO release. It showed anti-inflammatory effects in an endotoxin shock mouse model, which was similar to that observed for dexamethasone. This supports further the possible use of **5** as an anti-inflammatory lead compound (Liu et al., 2008). Mechanistically, dipyrithione (**5**) attenuates the activation of Janus kinase (JAK)/signal transducers and activators of transcription-1 (STAT1) signaling pathways to suppress IP-10/CXCL10 expression (Han et al., 2010). It also targets STAT1 but not the MAPK and NF- κB pathways to mediate its anti-inflammatory property in response to LPS, so suppression of iNOS expression through reducing STAT1 phosphorylation by **5** could result in the LPS-induced augmentation of COX-2 protein levels being reduced (Liu et al., 2008). Interestingly, dipyrithione (**5**) was also found

recently to inhibit potently the growth of *Trichophyton rubrum*, a fungal species leading to dermatophytosis. The dermatophyte infection was inhibited significantly when female guinea pigs (300 g) were inoculated topically by *T. rubrum* (for ten days) and treated topically with dipyrithione (0.2 mg/cm², daily) for ten days (Song et al., 2018).

In other investigations, dipyrithione (**5**) has exhibited potential antitumor activity. For example, it suppressed HeLa human cervical cancer cell growth through induction of cell apoptosis and inhibited the proliferation of human KB nasopharyngeal and MDA-MB-231 breast cancer and U937 and K562 leukemia cells (Table 3) (Fan et al., 2007; Fan et al., 2013). *In vivo* investigations indicated that hepatoma and sarcoma growths were inhibited significantly, when six-week-old male ICR mice were inoculated with human S180 sarcoma or H22 hepatoma cells and treated (i.p.) daily with dipyrithione (0.25 or 2.5 mg/kg) for ten days. No obvious changes were observed in the body weights of the treated mice (Fan et al., 2007; Fan et al., 2013). Mechanistically, dipyrithione (**5**) induces HeLa cell apoptosis through induction of the expression of p21, cleavage of caspase-3 and poly (ADP-ribose) polymerase (PARP), and release of cytochrome c, with the p38 and ERK1/2 signaling pathways being involved (Fan et al., 2007). It induces G1 arrest in K562, KB, MDA-MB-231, and U937 cancer cells by modulation of p53, p21, and levels of cyclins D1 and E1, and it induces apoptosis in these cells through activation of caspases-3 and -9 and of PARP (Fan et al., 2013).

The monomeric analogues of dipyrithione (**5**), 2-methyldithiopyridine *N*-oxide (**5a**) and 2-methylthiomethyldithiopyridine *N*-oxide (**5b**), were also found to show cytotoxicity against human MCF-7 breast, A549 lung, and HT-29 colon cancer cells, with IC₅₀ values being in the range 0.2–1.8 μM (O'Donnell et al., 2009). It is worth noting that a complex of two pyrrithione ligands chelated to Zn²⁺, namely, zinc pyrrithione (ZnPT, **5e**) (Figure 4), is attracting increasing interest for its antitumor activity. For example, ZnPT inhibited SKOV3 human ovarian cancer cell growth (IC₅₀ 0.65 μM), and its activity was increased (IC₅₀ 0.35 μM) when cells were treated with ZnPT plus zinc chloride (20 μM) (Chen et al., 2020a). ZnPT showed cytotoxicity against human HSC2 (IC₅₀ 2 μM), MDA1986 (IC₅₀ 1.3 μM), and SCC4 (IC₅₀ 2 μM) oral squamous cell cancer cells (Table 3) (Srivastava et al., 2015), as well as HL-60 human leukemia cells (IC₅₀ 0.1–1 μM) (Tailler et al., 2012). The oral tumor growth was inhibited when NOD/SCID/Crl mice were inoculated by SCC4 human oral squamous cell cancer cells and treated (i.p.) with ZnPT (1 mg/kg, weekly) for six weeks after the tumors reached 200 mm³ (Srivastava et al., 2015). Its potent antileukemic efficacy was observed when athymic nude mice were inoculated by HL-60 cells and treated (i.p.) with ZnPT (5 mg/kg, twice a week) for 40 days (Tailler et al., 2012).

Mechanistic investigations have indicated that ZnPT induces zinc accumulation and oxidative stress in HepG2 human liver cancer cells, which led to cell apoptosis through increasing the Bax/Bcl-2 ratio, causing mitochondrial dysfunction, cytochrome c release, and increased caspase-9/-3 activity (Mo et al., 2018). In PC3 human prostate cancer cells, ZnPT inhibited metabolism and induced ERK- and protein kinase C (PKC)-dependent necrosis by decreasing cellular adenosine 5'-triphosphate (ATP) levels that cause the phosphorylation of AMPK and acetyl CoA carboxylase (ACC) (Carraway and Dobner, 2012). In addition, it mediated cytotoxicity against human oral squamous cell cancer cells, postulated as acting through the phosphoinositide 3-kinase (PI3K)/Akt/mammalian target of rapamycin (mTOR) and Wnt/β-catenin signaling pathways (Srivastava et al., 2015).

Toxicological investigation showed that ZnPT induced caspase-independent cell death, heat shock response gene expression encoding heat shock proteins (HSPs), impaired genomic integrity, and depletion of cellular ATP levels in cultured primary human skin HEKn-APF keratinocytes. In both HEKn-APF keratinocytes and HEMA-LP melanocytes, ZnPT caused DNA damage, indicating that this topical antimicrobial may target and compromise keratinocytes and melanocytes in intact human skin (Lamore et al., 2010). At a concentration of 1 μM, ZnPT exposure did not cause any genotoxic threat and/or a proliferation index change in normal lymphocytes obtained from healthy human volunteers, but it increased lactate dehydrogenase, oxidative stress, and intracellular zinc levels. Thus, this complex may exhibit toxic effects against human cellular systems via increasing oxidative stress (Güner and İlhan, 2020). Therefore, the toxicity of ZnPT could be a matter of concern for its possible development as an anticancer agent, although this may be able to be resolved by synthetic modification of the molecule.

Alkaloids are an important group of natural products showing potential anticancer activity, of which derivatives of camptothecin and bisindole alkaloids such as vincristine are used clinically, targeting topoisomerase (topo)-I and tubulin polymerization, respectively (Henkin et al., 2018; Agarwal et al., 2020). A dimeric sulfur-containing pyridine derivative, dipyrithione (**5**), was synthesized by oxidation of pyrrithione (**5d**) with hydrogen peroxide (Barton and Samadi, 1992), of which **5d** was produced when 2-bromopyridine reacted with perbenzoic acid followed by treatment with sodium sulfide (Shaw et al., 1950). As a natural product derivative, dipyrithione (**5**) has been found to occur in the New Zealand basidiomycete (mushroom), *Cortinarius* species (Cortinariaceae) (Nicholas et al., 2001), and the bulbs of *Allium stipitatum* Regel (Amaryllidaceae) (Krejčová et al., 2014; Kubec et al., 2011; O'Donnell et al., 2009). However, it was identified for the first time in the family Annonaceae, from the Lao plant, *M. modestum*

Table 3. Cytotoxicity of compounds **5**, **5a**, **5b**, and **5e**

compound	A549 ^{a,1}	HeLa ^{b,2}	HSC2 ^{c,3}	HT-29 ^{d,1}	K562 ^{e,4}	KB ^{f,4}	MCF-7 ^{g,1}
5	-	4.0–10	-	-	4–10	4–10	-
5a	0.22	-	-	1.8	-	-	0.35
5b	0.78	-	-	-	-	-	0.39
5e	-	-	2.0	-	-	-	-

IC₅₀ values (μM) toward the human ^aA549 non-small cell lung cancer, ^bHeLa cervical cancer, ^cHSC2 oral squamous cell cancer, ^dHT-29 colon cancer, ^eK562 leukemia, ^fKB oral epidermoid cancer, ^gMCF-7 breast cancer cell lines. ¹(O'Donnell et al., 2009), ²(Fan et al., 2007), ³(Srivastava et al., 2015), ⁴(Fan et al., 2013). - (Not reported).

(Elkington et al., 2014b). Its potent activity (MIC <0.15 μM) against virulent mycobacteria (Elkington et al., 2014b) and other promising bioactive properties indicates that this compound could be an important lead compound for the development of effective agents to treat either cancer or various infectious diseases.

RUBIACEAE (*NAUCLEA ORIENTALIS* (L.) L.)

N. orientalis is a small tree (Table 1) native to Laos and tropical Asia (Newman et al., 2007), of which the dried stem wood and stem bark are used to treat cases of fatigue (Soejarto et al., 2012). The stems of *N. orientalis* were collected in Bolikhamxai Province of Laos, and both the methanol extract and chloroform partition were found to be active when tested against the chloroquine-sensitive clone D6 and the chloroquine-resistant clone W2 of *P. falciparum*. A follow-up bioassay-guided fractionation of the antimalarial-active chloroform partition resulted in the isolation of two new and one known indole-like alkaloid glucosides, namely, naucleaorine (**6**), epimethoxynaucleaorine (**6a**), and strictosidine lactam (**6b**), along with several known compounds, including oleanolic acid (**7**), 3α,23-dihydroxyurs-12-en-28-oic acid (**8**), and their analogues (Figure 5) (He et al., 2005). The alkaloids **6** and **6a** were found to be active in this *in vitro* antimalarial assay used, with IC₅₀ values being in the range 6–13 μM, but **6b** was inactive. This demonstrated that a conjugated system along the C-3, C-4, C-14, C-15, C-20, and C-21 positions plus a methoxy group substituted at the C-19 position are required for the activity, while the configuration at the C-19 position seems not to contribute significantly to this effect. Two triterpenoids, oleanolic acid (**7**) and 3α,23-dihydroxyurs-12-en-28-oic acid (**8**) were also active (IC₅₀ 4–13 μM) against the chloroquine-sensitive and -resistant malaria parasites, and **7** was more potently active than ursolic acid (**8a**), while other related analogues did not show any activity (He et al., 2005). These results imply that both a C-23 hydroxy group and a C-28 carboxylic acid substituent are required for antimalarial activity, but introduction of a hydroxy group at the C-19 position followed by methyl esterification of the C-28 carboxylic

acid results in this type of activity being abolished. Malaria is a mosquito-borne infectious disease caused by the *Plasmodium* species, and, to date, it continues to present a great challenge to public health in Laos, owing to increasing cases of MDR malaria as observed during its therapy. Traditional medicines have been used for a long time to treat malaria in Laos, and many plants included in Lao herbal prescriptions show some potential for the treatment of this infectious disease (Elliott et al., 2020). Thus, the discovery of potential antimalaria alkaloids **6** and **6a** and triterpenoids **7** and **8** from *N. orientalis* collected in Laos indicates that additional interesting compounds could be identified from a search for antimalarial leads from Lao plants, and that further synthetic modification of these leads may produce potent agents to treat malaria effectively.

Both **7** and **8a** were also found to show anti-TB activity (Tanachatchairatana et al., 2008). When comparing the anti-TB potency of these triterpenoids and their analogues or derivatives, including 3β-*O-trans*-caffeoylbetulonic acid (**3**), isolated from *D. quaesita* collected in Laos (Ma et al., 2008), **3** and **7** were found to exhibit activity, while ursolic acid (**8a**) was more potently active than **3** and **7** (Tanachatchairatana et al., 2008). Interestingly, 3β-*O-trans-p*-coumaroylbetulonic acid (**3g**), 3β-*O-trans-p*-coumaroyloleanolic acid (**7a**), and 3β-*O-trans-p*-coumaroylursolic acid (**8b**) were found to show more potent antimycobacterial activity than their parent compounds, **3b**, **7**, and **8a**, as well their acetylated derivatives, 3β-*O-trans-p*-acetylcoumaroylbetulonic acid, 3β-*O-trans-p*-acetylcoumaroyloleanolic acid (**7b**), and 3β-*O-trans-p*-acetylcoumaroylursolic acid (**8c**). Also, these coumarates were more potently active overall than their caffeate analogues, including **3**, **3a**, 3β-*O-trans*-caffeoyloleanolic acid (**7c**), 3β-*O-trans*-diacetycaffeoyloleanolic acid (**7d**), 3β-*O-trans*-caffeoylursolic acid (**8d**), and 3β-*O-trans*-diacetycaffeoylursolic acid (**8e**) (Table 4). Thus, introduction of a 3β-*O-trans-p*-coumaroyl group in **3b**, **7**, and **8a** appears to increase the resultant anti-TB potency within this triterpenoid class (Tanachatchairatana et al., 2008).

When tested against KB human oral epidermoid cancer cells, compound **6** showed weak activity, but compound

Table 4. Antimycobacterial activity of compounds **3** and **7** and the derivatives of **3**, **7**, and **8**

compound	TB ^a	compound	TB ^a	compound	TB ^a	compound	TB ^a
3	200	7a	6.25	8a	12.5	8e	100
3a	200	7b	12.5	8b	6.25		
3b	50	7c	200	8c	25		
7	50	7d	12.5	8d	200		

^aMIC₅₀ values (μg/mL) toward *Mycobacterium tuberculosis* H37Ra bacteria (Tanachatchairatana et al., 2008).




Figure 5. Structures of the alkaloid naucleaorine (**6**) and its analogues (**6a** and **6b**) and the triterpenoids oleanolic acid (**7**) and 3α,23-dihydroxyurs-12-en-28-oic acid (**8**) isolated from the Lao plant, *Nauclea orientalis*, and selected derivatives, **7a–7d** of **7** and **8a–8e** of **8** (He et al. 2005, Tanachatchairatana et al. 2008).

6a did not, indicating that the configuration at the C-19 position of **6** is important for this compound to exhibit inhibitory activity on KB cell growth. Similar to **6a**, both **7** and **8** did not show any discernible growth inhibitory activity (IC₅₀ >10 μM) against KB cells (He et al., 2005). However, the parent 23-dehydroxylated analogue of **8**,

ursolic acid (**8a**), has been reported extensively for its potential antitumor activity. For example, **8a** exhibited cytotoxicity against MDA-MB-231 human breast cancer cells (IC₅₀ 5.9 μM) and inhibited NF-κB activity potently (IC₅₀ 31 nM), for which substitutions at both the C-3 and C-28 positions seem important (Ren et al., 2018).




Figure 6. Structures of the lignan rourinoside (**9**) and the glycerol ester rouremin (**10**) and its analogue (**10a**) isolated from *Rourea minor* collected in Laos (He et al., 2006).

Mechanistically, ursolic acid (**8a**) targets NF- κ B, COX-2, matrix metalloproteinase 9 (MM-9), cyclin D11, and other proteins to mediate its antitumor effects, and it also inhibits angiogenesis and metastasis. Thus far, a phase I cancer clinical study has been completed for this compound, with a phase II clinical trial being recommended (Qian et al., 2015).

Even though **7** was inactive ($IC_{50} > 10 \mu M$) against KB cells (He et al., 2005), several natural or semisynthetic oleanolic acid-like compounds have been reported for their potential antitumor activity (Ren and Kinghorn, 2019), including pristimerin and bardoxolone methyl (methyl 2-cyano-3,12-dioxooleana-1,9(11)-dien-28-oate, CDDO-Me). Pristimerin is a quinone methide triterpenoid isolated from species of the Celastraceae and Hippocrateaceae plant families and exhibits promising antitumor activity (Li et al., 2019), while CDDO-Me, a synthetic derivative of **7**, inhibited potently the production of nitric oxide induced by IFN- γ in mouse macrophages and redirects the activation profile of tumor-associated macrophages from tumor-promoting to tumor-inhibition (Honda et al., 2000; Ball et al., 2016). It was also found to show antitumor efficacy by improving the immune response in cancer (Ren and Kinghorn, 2019), indicating a possible immunotherapeutic role in the treatment of cancer for CDDO-Me. In addition, CDDO-Me is a potent nuclear factor erythroid 2-related factor 2 (Nrf2) activator and a NF- κ B inhibitor and has been investigated in a phase I cancer clinical trial (Ren and Kinghorn, 2019; Wang et al., 2014). Recently, a phase III CARDINAL trial in patients with Alport syndrome has been completed for CDDO-Me (Chertow et al., 2021).

As summarized recently, both **7** and **8a** show potent pharmacological properties. Compound **7** has attracted

increasing attention for its promising anti-inflammatory, anti-metabolic syndrome, antitumor and hepatoprotective propensities, as well its antiviral activity against hepatitis, herpes, HIV, and influenza (Feng et al., 2020; Khwaza et al., 2018). Similarly, **8a** showed antimicrobial, antitumor, and antiviral activities and pharmacological effects on the cardiovascular and urinary systems (Khwaza et al., 2020; Ren and Kinghorn, 2019; Sun et al., 2020). Recently, both **7** and **8a** were proposed as potential anti-SARS-CoV-2 agents, based on their binding to the key pathogenic M^{pro} protein (Pawelczyk and Zaprutko, 2020), and thus some promising anti-COVID-19 agents could be discovered from further investigations on higher plants that occur in Laos.

The dried stem wood of *N. orientalis* is used to treat cases of fatigue in Laos, with alkaloid **6** and triterpenoid **8** and their analogues being characterized as active components of this species (Soejarto et al., 2012). Also, several triterpene analogues of **8** were identified as the major components of the hard wood of *N. orientalis* collected in Japan (Fujita et al., 1967), while a number of indole alkaloid derivatives of **6** were characterized from the leaves and stems of *N. orientalis* collected in China and in Thailand (Kanchanapoom et al., 2021; Liu et al., 2018). Thus, triterpenoids and indole alkaloids could be the major bioactive constituents of *N. orientalis* growing in Laos and in other countries in Asia, including China, Japan, and Thailand.

CONNARACEAE [*ROUREA MINOR* (GAERTN.) AUBL.]

R. minor is a liana (Table 1) native to tropical Africa and tropical Asia (Newman et al., 2007), of which the dried




Figure 7. Structures of the phenanthrene stemophenanthrene A (**11**) and its analogues (**11a** and **11b**) and the stilbenoids stemofuran Y (**12a**) and isopinosylvin A (**13**) isolated from *Stemona tuberosa*, as well stemofuran X (**12**) isolated from *Stemona pierrei*, collected in Laos (Khamko et al., 2013; Quang et al., 2014).

stems are used to treat dengue fever (Soejarto et al., 2012). Thus, the dried stems of *R. minor* were collected in Bolikhamsai Province of Laos and investigated further for identification of their bioactive components. A new lignan glucoside, rourinoside (**9**), along with several glycerides, including rouremin (**10**) and 1-(26-hydroxyhexacosanoyl)glycerol (**10a**) (Figure 6), were isolated from this plant sample. All of these compounds were found to show activity against both the chloroquine-sensitive clone D6 and the chloroquine-resistant clone W2 of *P. falciparum*, with IC_{50} values occurring in the range 2–13 μ M, and thus these compounds exhibit potential antimalarial activity (He et al., 2006).

STEMONACEAE (*STEMONA PIERREI* GAGN. AND *STEMONA TUBEROSA* LOUR.)

Stemona is the largest genus of the family Stemonaceae, and several of its species have been used as antihelminthic agents and to treat coughs in Laos, as well in China, Thailand, and Vietnam (Quang et al., 2014). Of these, *S. pierrei* is a vine (Table 1) native to Laos, Thailand, and Vietnam, while *S. tuberosa* is a climbing plant (Table 1) native to Bangladesh, Cambodia, China, India, Indonesia, Laos, Myanmar, New Guinea, the Philippines, Sri Lanka, Thailand, and Vietnam (Newman et al., 2007). In a search for potential antitumor agents from *Stemona* plants, three new phenanthrenes, namely, stemophenanthrenes A (**11**), B (**11a**) and C (**11b**) (Figure 7), were isolated from the roots of *S. tuberosa* collected in Attapeu Province of Laos.

These compounds exhibited inhibitory activity against human HepG2 liver, KB oral epidermal, MCF7 breast, and SK-Lu-1 lung cancer cells (Khamko et al., 2013), and

the laboratory work conducted indicated that a 5,6,7-*ortho* substituent is required in this regard. In addition, a new phenylbenzofuran-type stilbenoid, stemofuran X (**12**), isolated from the roots of *S. pierrei* collected in Savannakhet Province of Laos, and two new normal stilbenoids, stemofuran Y (**12a**) and isopinosylvin A (**13**), identified from the roots of *S. tuberosa* collected from Attapeu Province of Laos (Figure 7), were also tested for their cytotoxicity against HepG2, KB, MCF7, and SK-Lu-1 cancer cells. However, none of these stilbenoids was found to be active against any of these four cell lines ($IC_{50} > 10 \mu$ M) (Khamko et al., 2013; Quang et al., 2014).

Stilbenoids are an important group of natural products distributed widely among edible plants, of which *trans*-resveratrol (3,5,4'-trihydroxy-*trans*-stilbene) occurs in various berries, grapes, red wine, and many other human foods, and shows a broad range of biological activities, including anti-aging, anti-infective, antioxidant, and anticarcinogenic effects. It also possesses cardio-, gastrointestinal, hepatic-, and nephroprotective activities, and protects organs and tissues against ultraviolet radiation (UVR)-induced skin cancer and carcinogen-induced damage. Thus, *trans*-resveratrol has been investigated extensively for its cancer chemopreventive and synergistic properties, and it may reduce the side effects to enhance the outcomes of cancer chemotherapy, when used in combination with approved cancer chemotherapeutic agents (Jang et al., 1997; Xiao et al., 2019). Therefore, the discovery of new stilbenoids from Lao plants could be useful for the development of further potential cancer chemoprotective agents.

The roots of *S. tuberosa* have long been used as a traditional Chinese medicine ("Baibu") as an insecticidal and antitussive agent, and several stilbenoids have been isolated from this plant source when collected in China (Li

et al., 2020a; Lin et al., 2008). Of these compounds, stilbostemin U was found to show antimicrobial activity against *Bacillus pumilus* (MIC 41–83 μ M) (Lin et al., 2008). In addition, several alkaloids have been isolated from this same species when sourced in China (Hu et al., 2020; Lin et al., 2006).

Alkaloids derived from the monocotyledonous family Stemonaceae, namely, the stemona alkaloids, are a large group of natural products, with more than 200 compounds of this type having been identified thus far. These alkaloids have been found to show multiple promising bioactivities, including acetylcholinesterase inhibitory, anti-inflammatory, antitussive, and insecticidal activities, with stemofoline showing the most potent activity in the reversal of multidrug resistance of some anticancer agents (Greger 2019). As a member of this group, bisdehydrostemoninine, a stemona alkaloid isolated in a large amount from *S. tuberosa* collected in China, showed significant antitussive activity when Dunkin-Hartley guinea pigs (300–350 g) were treated (i.p., single dose) with bisdehydrostemoninine (50 mg/kg) followed (after 30 min.) by exposure to 0.5 M citric acid aerosol at a flow rate of 0.5 mL/min for eight min (Lin et al., 2006). Another major stemona alkaloid from *S. tuberosa*, neotuberostemonine (Hu et al., 2020), was found to suppress hypoxia-induced activation and differentiation of primary mouse lung fibroblasts (PLFs). The bleomycin-induced pulmonary fibrosis was attenuated effectively when ICR male mice were injected intratracheally with bleomycin (3.5 U/kg), and, after seven days, treated orally with neotuberostemonine (30 mg/kg, daily) for 7 or 14 days. This *in vivo* activity was found to be mediated by inhibition of HIF-1 α and its downstream profibrotic factors (Lv et al., 2018). Thus, stemona alkaloids could contribute to the observed antitussive effect of *S. tuberosa* (Lin et al., 2006). However, no stemona alkaloids have been thus far isolated from the Lao *Stemona* species, and thus further phytochemical work on the plant family Stemonaceae in Laos would seem valuable.

DISCUSSION AND PERSPECTIVE

Laos is a landlocked country with a prolific and varied tropical flora, and medicinal plants remain the backbone of primary health care in this country. However, only limited phytochemical investigations have been conducted thus far on eight of these plants, namely, *A. cochinchinensis*, *D. quaesita*, *G. napalense*, *M. modestum*, *N. orientalis*, *R. minor*, *S. pierrei*, and *S. tuberosa*, of which none is endemic to the country. All of these plant species were collected in the southern part of Laos, and previous investigations have focused mainly on the identification of anti-infective and cytotoxic constituents guided by cell-based bioassays. Thus, a further search for bioactive agents from the northern area of Laos, involving evaluation of the lead compounds for

additional biological activities, and elucidation of the cellular mechanisms of action of the active compounds present, could be important for the development of one or more useful therapeutic agents.

Cancer is a serious disease and a threat to human health, and the discovery of new effective anticancer agents is urgently needed (Newman and Cragg, 2020; Kinghorn et al., 2016). Several species collected in Laos were found to be cytotoxic toward human cancer cells (Henkin et al., 2017), from which several compounds showing cancer cell cytotoxicity have been characterized (Soejarto et al., 2006; Zhang et al., 2016). Thus, a continuing search for anticancer agents from Lao plants could be of promise. In addition, infections from bacteria, tuberculosis (TB), malaria, and/or viruses are problematic in Laos, and members of local communities have used traditional herbal medicines as their first-line therapy to treat such diseases for many years. Thus far, several Lao plants and their active components against *M. tuberculosis* H37Rv (TB), HIV, and malaria have been identified (Elkington et al., 2009; Soejarto et al., 2012; Zhang et al., 2016), and this preliminary work could form the foundation for the continuous search for agents from Lao plants for the treatment of these disorders.

Plant-derived phenolic compounds are a large group of natural products, including lignans and stilbenoids, of which some show potential antitumor activity (Soobrattee et al., 2006). For example, etoposide, a semisynthetic epipodophyllotoxin glycoside derived from a naturally occurring aryltetralin lignan lactone, podophyllotoxin, is well established as a therapy clinically to treat different types of cancer, targeting DNA topo II. Recently, a structurally similar analogue, phyllanthusmin D, has been identified as a promising antitumor lead, which mediates its activity by induction of cancer cell apoptosis through activation of caspase-3 but not by inhibition of topo II (Ren et al., 2014; Young et al., 2018; Ren et al., 2019). A norlignan, nyasol (**1b**), isolated from a Lao plant, was found to show NF- κ B-targeted anti-infective activity, and it also directs FPTase, PRL-3, VEGF to exhibit antitumor potential (Jeong et al., 2003; Lee et al., 2014; Lim et al., 2009; Song et al., 2008). Thus, further modification of this molecule may be helpful for the discovery of novel anticancer and anti-infective agents, and preparation of drug conjugates from **1b** and other drugs used clinically may be promising in this regard.

The triterpenoids, 3 β -O-(*trans*-caffeoyl)betulinic acid (**3**) and oleanolic acid (**7**), and the thiopyridine, dipyrithione (**5**), isolated from Lao plants, were all found to show antitumor, anti-infective activities, with **3** and **5** known to target NF- κ B (Liu et al., 2008; Andre et al., 2013). NF- κ B functions widely in inflammation and human pathobiology (Zhang et al., 2017), and inhibition of this protein could be supportive to the treatment of autoimmune and lymphoproliferative disorders (Qin et al., 2005; Concetti and Wilson, 2018). Accordingly, compounds **3** and **5** may serve as leads for the design of NF- κ B-targeted therapeutic agents, with potential uses in the treatment of

cancer and infectious diseases.

Currently, COVID-19 has resulted from severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), which is spreading throughout the world to threaten human health seriously. The infection of SARS-CoV-2 is commonly followed by a deep inflammation known as a “cytokine storm” caused by activation of pro-inflammatory genes, such as NF- κ B, STAT-3, IL-6, IL-8 and IL-1ss which leads to lung tissue destruction. While the microRNA network has been proposed as a target for the COVID-19-induced inflammatory response (Gasparello et al., 2021), the NF- κ B pathway and Nrf2 have been regarded as the potential targets of COVID-19 (Carcattera and Caruso, 2021; Cuadrado et al., 2020; Zinovkin and Grebenchikov, 2020). Nrf2 is a transcription factor in charge of cellular redox balance and the expression of genes involved in immunity and inflammation, and several Nrf2 activators have been proposed for their inhibitory activity against COVID-19 (Cuadrado et al., 2020; Zinovkin and Grebenchikov, 2020). Of these, an herbal medicine, PB125, an ethanol extract of a mixture of rosemary (*Rosmarinus officinalis*), ashwagandha (*Withania somnifera*), and luteolin, was found to activate Nrf2 potently and to downregulate angiotensin-converting enzyme 2 (ACE2, a surface receptor of SARS-CoV-2) and transmembrane protease serine 2 (TMPRSS2) mRNA expression in HepG2 human liver cancer cells (McCord et al., 2020). Therefore, by targeting NF- κ B and Nrf2, some anti-COVID-19 agents may be discovered from repurposing the current anticancer compounds, and SARS-CoV-2 may be introduced to infect selectively human cancer cells, through engineering modification of NF- κ B and Nrf2.

Medicinal plants have been long used for the treatment of cancer and infections in Laos, and their derivatives, nyasol (**1b**), 3 β -O-*trans*-caffeoylbetulinic acid (**3**), and dipyrithione (**5**) have been found to show the NF- κ B-targeted antitumor and anti-infective activities. In addition, both betulinic acid (**3b**) and ursolic acid (**8a**) were found to mediate their pharmacological activities via activation of the Nrf2 signaling pathway (Li et al., 2020b), and oleanolic acid (**7**) and **8a** have been considered for their potential anti-COVID-19 activity (Pawelczyk and Zaprutko, 2020). Thus, searching anti-COVID-19 agents from Lao plants seems reasonable, and **3b**, **7**, and **8a** may be modified by conjugating them with **5e**, which contains N, S, and Zn that could support such an activity, for their development as an effective therapy for COVID-19.

Thus far, several different types of natural products, including alkaloids, glycerol esters, phenolic compounds (neolignans and norlignans and stilbenoids), steroids, and triterpenoids have been identified as potential antitumor and/or anti-infective agents from Lao plants. The compounds targeting NF- κ B and/or Nrf2, including **1b**, **3**, **3b**, **5**, and **7** could be supportive of the design and discovery of novel therapeutic agents. In this regard, a continuing search for bioactive agents from Lao plants

may be regarded as not only necessary but also important for the improvement of human health.

CONFLICT OF INTERESTS

The authors have not declared any conflict of interests.

ACKNOWLEDGEMENTS

Some of the laboratory work described in this review article was supported by grants U01TW001015 (DDS), funded by the Fogarty International Center, and P01 CA125066 (ADK), funded by the National Cancer Institute, NIH, Bethesda, MD, USA. The authors thank Professor Bounhong Southavong and Dr. Bounleuane Douangdeuane (Institute of Traditional Medicine, Ministry of Health of Laos) for their gracious and generous collaboration in the implementation of these research endeavors.

REFERENCES

- Agarwal G, Blanco Carcache PJ, Addo EM, Kinghorn AD (2020). Current status and contemporary approaches to the discovery of antitumor agents from higher plants. *Biotechnology Advances* 38:107337.
- Amiri S, Dastghaib S, Ahmadi M, Mehrbod P, Khadem F, Behrouj H, Aghanoori MR, Machaj F, Ghamsari M, Rosik J, Hudecki A, Afkhami A, Hashemi M, Los MJ, Mokarram P, Madrakian T, Ghavami S (2020). Betulin and its derivatives as novel compounds with different pharmacological effects. *Biotechnology Advances* 38:107409.
- Andre CM, Larsen L, Burgess EJ, Jensen DJ, Cooney JM, Evers D, Zhang J, Perry NB, Laing WA (2013). Unusual immuno-modulatory triterpene-caffeates in the skins of russeted varieties of apples and pears. *Journal of Agricultural and Food Chemistry* 61(11):2773-2779.
- Bae G, Yu JR, Lee J, Chang J, Seo EK (2007). Identification of nyasol and structurally related compounds as the active principles from *Anemarrhena asphodeloides* against respiratory syncytial virus (RSV). *Chemistry and Biodiversity* 4(9):2231-2235.
- Bak JP, Cho YM, Kim I, Park DW, Kwon JE, Jeong YJ, Kwak JH, Kang SC (2016). Inhibitory effects of norlignans isolated from *Anemarrhena asphodeloides* on degranulation of rat basophilic leukemia-2H3 cells. *Biomedicine and Pharmacotherapy* 84:1061-1066.
- Ball MS, Shipman EP, Kim H, Liby KT, Pioli PA (2016). CDDO-Me redirects activation of breast tumor associated macrophages. *PLoS One* 11(2):e0149600.
- Barton DHR, Samadi M (1992). The invention of radical reactions. Part XXV. A convenient method for the synthesis of the acyl derivatives of *N*-hydroxypyridine-2-thione. *Tetrahedron* 48(35):7083-7090.
- Bourjot M, Leyssen P, Eydoux C, Guillemot JC, Canard B, Rasoanaivo P, Guérin F, Litaudon M (2012). Flacourtosides A-F, phenolic glycosides isolated from *Flacourtia ramontchi*. *Journal of Natural Products* 75(4):752-758.
- Carcattera M, Caruso C (2021). Alveolar epithelial cell type II as main target of SARS-CoV-2 virus and COVID-19 development via NF- κ B pathway deregulation: A physio-pathological theory. *Medical Hypotheses* 146:110412.
- Carraway RE, Dobner PR (2012). Zinc pyrithione induces ERK- and PKC-dependent necrosis distinct from TPEN-induced apoptosis in prostate cancer cells. *Biochimica et Biophysica Acta* 1823(2):544-557.
- Chen M, Ding Y, Ke Y, Zeng Y, Liu N, Zhong Y, Hua X, Li Z, Xiong Y, Wu C, Yu H (2020a). Anti-tumour activity of zinc ionophore pyrithione in human ovarian cancer cells through inhibition of proliferation and migration and promotion of lysosome-mitochondrial apoptosis. *Artificial Cells, Nanomedicine, and Biotechnology* 48(1):824-833.

- Chen WL, Ren Y, Ren J, Erxleben C, Johnson ME, Gentile S, Kinghorn AD, Swanson SM, Burdette JE (2017). (+)-Strebloside-induced cytotoxicity in ovarian cancer cells is mediated through cardiac glycoside signaling networks. *Journal of Natural Products* 80(3):659-669.
- Chen X, Yuan XN, Zhang Z, Gong PJ, Yin WN, Jiang Q, Xu J, Xu XL, Gao Y, Chen WL, Chen FF, Tian YH, Wei L, Zhang JW (2020b). Betulinic acid inhibits cell proliferation and migration in gastric cancer by targeting the NF- κ B/VASP pathway. *European Journal of Pharmacology* 889:173493.
- Cheng J, Li X, Wang S, Han Y, Zhao H, Yang X (2021). Carrier-free triterpene prodrugs with glutathione response and biosafety for synergistically enhanced photochemotherapy. *ACS Applied Materials and Interfaces* 13(1):245-256.
- Chertow GM, Appel GB, Andreoli S, Bangalore S, Block GA, Chapman AB, Chin MP, Gibson KL, Goldsberry A, Iijima K, Inker LA, Knebelmann B, Mariani LH, Meyer CJ, Nozu K, O'Grady M, Silva AL, Stenvinkel P, Torra R, Warady BA, Pergola PE (2021). Study design and baseline characteristics of the CARDINAL Trial: A phase 3 study of bardoxolone methyl in patients with Alport syndrome. *American Journal of Nephrology* 52(3):180-189.
- Cho JY, Kim CM, Lee HJ, Lee SH, Cho JA, Kim WS, Park KH, Moon JH (2013). Caffeoyl triterpenes from pear (*Pyrus pyrifolia* Nakai) fruit peels and their antioxidative activities against oxidation of rat blood plasma. *Journal of Agricultural and Food Chemistry* 61(19):4563-4569.
- Choi JY, Kim JE, Park JJ, Lee MR, Song BR, Park JW, Kang MJ, Lee HS, Son HJ, Hong JT, Hwang DY (2018). The anti-inflammatory effects of fermented herbal roots of *Asparagus cochinchinensis* in an ovalbumin-induced asthma model. *Journal of Clinical Medicine* 7(10):377.
- Chung YL, Pan CH, Wang CCN, Hsu KC, Sheu MJ, Chen HF, Wu CH (2016). Methyl protodioscin, a steroidal saponin, inhibits neointima formation *in vitro* and *in vivo*. *Journal of Natural Products* 79(6):1635-1644.
- Colombo E, Polito L, Biocotino M, Marzullo P, Hyeraci M, Dalla Via L, Passarella D (2020). New class of betulinic acid-based nanoassemblies of cabazitaxel, podophyllotoxin, and thiocolchicine. *ACS Medicinal Chemistry Letters* 11(5):895-898.
- Concetti J, Wilson CL (2018). NFKB1 and cancer: Friend or foe? *Cells* 7(9):133.
- Cuadrado A, Pajares M, Benito C, Jiménez-Villegas J, Escoll M, Fernández-Ginés R, Yague AJG, Lastra D, Manda G, Rojo AI, Dinkova-Kostova AT (2020). Can activation of NRF2 be a strategy against COVID-19? *Trends in Pharmacological Sciences* 41(9):598-610.
- de Boer HJ, Lamxay V, Björk L (2012). Comparing medicinal plant knowledge using similarity indices: A case of the Brou, Saek and Kry in Lao PDR. *Journal of Ethnopharmacology* 141(1):481-500.
- Dehelean CA, Marcovici I, Soica C, Mioc M, Coricovac D, Iurciuc S, Cretu OM, Pinzaru I (2021). Plant-derived anticancer compounds as new perspectives in drug discovery and alternative therapy. *Molecules* 26(4):1109.
- Du H, Li H, Wu P, Wang C, Xue J, Wei X (2021). α -Glucosidase inhibitory pentacyclic triterpenoids from the leaves of *Cleistocalyx conspersipunctatus*. *Phytochemistry Letters* 41:109-113.
- Dubost JM, Phakeovilay C, Her C, Bochaton A, Elliott E, Deharo E, Xayvue M, Bouamanivong S, Bourdy G (2019). Hmong herbal medicine and herbalists in Lao PDR: Pharmacopeia and knowledge transmission. *Journal of Ethnobiology and Ethnomedicine* 15(1):27.
- Elkington BG, Phiapalath P, Sydara K, Somsamouth V, Goodsmith NI, Soejarto DD (2014a). Assessment of the importance of medicinal plants among communities around Khat Ngong of southern Laos. *Journal of Environmental Biology* 35(4):607-615.
- Elkington BG, Southavong B, Sydara K, Souliya O, Vanthanouvong M, Nettavong K, Thammachack B, Pak DH, Riley MC, Franzblau SG, Soejarto DD (2009). Biological evaluation of plants of Laos used in the treatment of tuberculosis in Lao traditional medicine. *Pharmaceutical Biology* 47(1):26-33.
- Elkington BG, Sydara K, Newsome A, Hwang CH, Lankin DC, Simmler C, Napolitano JG, Ree R, Graham JG, Gyllenhaal C, Bouamanivong S, Souliya O, Pauli GF, Franzblau SG, Soejarto DD (2014b). New finding of an anti-TB compound in the genus *Marsypopetalum* (Annonaceae) from a traditional herbal remedy of Laos. *Journal of Ethnopharmacology* 151(2):903-911.
- Elliott E, Chassagne F, Aubouy A, Deharo E, Souvanasy O, Sythamala P, Sydara K, Lamxay V, Manithip C, Torres JA, Bourdy G (2020). Forest fevers: Traditional treatment of malaria in the southern lowlands of Laos. *Journal of Ethnopharmacology* 249:112187.
- Fan Y, Chen H, Qiao B, Luo L, Ma H, Li H, Jiang J, Niu D, Yin Z (2007). Opposing effects of ERK and p38 MAP kinases on HeLa cell apoptosis induced by dipyrithione. *Molecules and Cells* 23(1):30-38.
- Fan Y, Liu C, Huang Y, Zhang J, Cai L, Wang S, Zhang Y, Duan X, Yin Z (2013). Dipyrithione induces cell-cycle arrest and apoptosis in four cancer cell lines *in vitro* and inhibits tumor growth in a mouse model. *BMC Pharmacology and Toxicology* 14:54.
- Feng A, Yang S, Sun Y, Zhang L, Bo F, Li L (2020). Development and evaluation of oleanolic acid dosage forms and its derivatives. *BioMed Research International* 1308749.
- Fujita E, Fujita T, Suzuki T (1967). Constituents of *Nauclea orientalis* L. I. Noreugenin and aucleoside, a new glycoside. *Chemical and Pharmaceutical Bulletin* 15(11):1682-1686.
- Gasparrillo J, Finotti A, Gambari R (2021). Tackling the COVID-19 "cytokine storm" with microRNA mimics directly targeting the 3'UTR of pro-inflammatory mRNAs. *Medical Hypotheses* 146:110415.
- Greger H (2019). Structural classification and biological activities of *Stemona* alkaloids. *Phytochemistry Reviews* 18(2):463-493.
- Güner A, İlhan S (2020). Cytotoxic, genotoxic, oxidative, and irritant effects of zinc pyrithione *in vitro*. *Toxicological and Environmental Chemistry* 102(10):607-623.
- Gyllenhaal C, Kadushin MR, Southavong B, Sydara K, Bouamanivong S, Xayveu M, Xuan LT, Hiep NT, Hung NV, Loc PK, Dac LX, Bich TQ, Cuong NM, Ly HM, Zhang HJ, Franzblau SG, Xie H, Riley MC, Elkington BG, Nguyen HT, Waller DP, Ma CY, Tamez P, Tan GT, Pezzuto JM, Soejarto DD (2012). Ethnobotanical approach versus random approach in the search for new bioactive compounds: Support of a hypothesis. *Pharmaceutical Biology* 50(1):30-41.
- Han C, Fu J, Liu Z, Huang H, Luo L, Yin Z (2010). Dipyrithione inhibits IFN- γ -induced JAK/STAT1 signaling pathway activation and IP-10/CXCL10 expression in RAW264.7 cells. *Inflammation Research* 59(10):809-816.
- He ZD, Ma CY, Tan GT, Sydara K, Tamez P, Southavong B, Bouamanivong S, Soejarto DD, Pezzuto JM, Fong HHS, Zhang HJ (2006). Rourinose and rouremin, antimalarial constituents from *Rourea minor*. *Phytochemistry* 67(13):1378-1384.
- He ZD, Ma CY, Zhang HJ, Tan GT, Tamez P, Sydara K, Bouamanivong S, Southavong B, Soejarto DD, Pezzuto JM, Fong HHS (2005). Antimalarial constituents from *Nauclea orientalis* (L.) L. *Chemistry and Biodiversity* 2(10):1378-1386.
- Henkin JM, Ren Y, Soejarto DD, Kinghorn AD (2018). The search for anticancer agents from tropical plants. In: Kinghorn AD, Falk H, Gibbons S, Kobayashi J, Asakawa Y, Liu J-K (Eds.), *Progress in the Chemistry of Organic Natural Products*. Cham, Switzerland: Springer International Publishing pp.1-94.
- Henkin JM, Sydara K, Xayvue M, Souliya O, Kinghorn AD, Burdette JE, Chen WL, Elkington BG, Soejarto DD (2017). Revisiting the linkage between ethnomedical use and development of new medicines: A novel plant collection strategy towards the discovery of anticancer agents. *Journal of Medicinal Plant Research* 11(39):621-634.
- Honda T, Rounds BV, Bore L, Finlay HJ, Favaloro FG, Suh N, Wang Y, Sporn MB, Gribble GW (2000). Synthetic oleanane and ursane triterpenoids with modified rings A and C: A series of highly active inhibitors of nitric oxide production in mouse macrophages. *Journal of Medicinal Chemistry* 43(22):4233-4246.
- Hsieh MJ, Lin CW, Chen MK, Chien SY, Lo YS, Chuang YC, Hsi YT, Lin CC, Chen JC, Yang SF (2017). Inhibition of cathepsin S confers sensitivity to methyl protodioscin in oral cancer cells via activation of p38 MAPK/JNK signaling pathways. *Scientific Reports* 7:45039.
- Hu ZX, An Q, Tang HY, Yuan CM, Li YN, Zhang Y, Hao XJ (2020). Stemtuberolines A–G, new alkaloids from *Stemona tuberosa* and their anti-TMV activity. *Fitoterapia* 143:104572.
- Huang H, Pan Y, Ye Y, Gao M, Yin Z, Luo L (2011). Dipyrithione attenuates oleic acid-induced acute lung injury. *Pulmonary Pharmacology and Therapeutics* 24(1):74-80.

- Jang M, Cai L, Udeani GO, Slowing KV, Thomas CF, Beecher CWW, Fong HHS, Farnsworth NR, Kinghorn AD, Mehta RG, Moon RC, Pezzuto JM (1997). Cancer chemopreventive activity of resveratrol, a natural product derived from grapes. *Science* 275(5297):218-220.
- Jeong DH, Kwak SC, Lee MS, Yoon KH, Kim JY, Lee CH (2020). Betulinic acid inhibits RANKL-induced osteoclastogenesis via attenuating Akt, NF- κ B, and PLC γ 2-Ca²⁺ signaling and prevents inflammatory bone loss. *Journal of Natural Products* 83(4):1174-1182.
- Jeong SJ, Higuchi R, Ono M, Kuwano M, Kim YC, Miyamoto T (2003). *cis*-Hinokiresinol, a norlignan from *Anemarrhena asphodeloides*, inhibits angiogenic response *in vitro* and *in vivo*. *Biological and Pharmaceutical Bulletin* 26(12):1721-1724.
- Kanchanapoom T, Sahakitpichan P, Chimnoi N, Petchthong C, Thamniyom W, Nangkoed P, Ruchirawat S (2021). Monoterpene alkaloid glycosides from the leaves of *Nauclea orientalis*. *Phytochemistry Letters* 41:83-87.
- Karim N, Khan I, Khan W, Khan I, Khan A, Halim SA, Khan H, Hussain J, Al-Harrasi A (2019). Anti-nociceptive and anti-inflammatory activities of asparacosin A involve selective cyclooxygenase 2 and inflammatory cytokines inhibition: An *in-vitro*, *in-vivo*, and *in-silico* approach. *Frontiers in Immunology* 10:581.
- Kawabe K, Kanazawa T, Takimoto G, Ikeda T, Sato M, Kano S (1967). Antimicrobial activity of bis(2-pyridyl *N*-oxide) disulfide. *Chemotherapy* 15(3):224-227.
- Khamko VA, Quang DN, Dien PH (2013). Three new phenanthrenes, a new stilbenoid isolated from the roots of *Stemona tuberosa* Lour. and their cytotoxicity. *Natural Product Research* 27(24):2328-2332.
- Khwaza V, Oyediji OO, Aderibigbe BA (2018). Antiviral activities of oleanolic acid and its analogues. *Molecules* 23(9):2300.
- Khwaza V, Oyediji OO, Aderibigbe BA (2020). Ursolic acid-based derivatives as potential anti-cancer agents: An update. *International Journal of Molecular Sciences* 21(16):5920.
- Kim CS, Subedi L, Oh J, Kim SY, Choi SU, Lee KR (2017). Bioactive triterpenoids from the twigs of *Chaenomeles sinensis*. *Journal of Natural Products* 80(4):1134-1140.
- Kim KH, Choi SU, Lee KR (2010). Bioactivity-guided isolation of cytotoxic triterpenoids from the trunk of *Berberis koreana*. *Bioorganic & Medicinal Chemistry Letters* 20(6):1944-1947.
- Kinghorn AD, Carcache de Blanco EJ, Lucas DM, Rakotondraibe HL, Orjala J, Soejarto DD, Oberlies NH, Pearce CJ, Wani MC, Stockwell BR, Burdette JE, Swanson SM, Fuchs JR, Phelps MA, Xu L, Zhang X, Shen YY (2016). Discovery of anticancer agents of diverse natural origin. *Anticancer Research* 36(11):5623-5637.
- Krejčová P, Kučerová P, Stafford GI, Jäger AK, Kubec R (2014). Antiinflammatory and neurological activity of pyrithione and related sulfur-containing pyridine *N*-oxides from Persian shallot (*Allium stipitatum*). *Journal of Ethnopharmacology* 154(1):176-182.
- Kubec R, Krejčová P, Šimek P, Václavík L, Hájšlová J, Schraml J (2011). Precursors and formation of pyrithione and other pyridyl-containing sulfur compounds in drumstick onion, *Allium stipitatum*. *Journal of Agricultural and Food Chemistry* 59(10):5763-5770.
- Kun-Liu, Wang JY, Zhang L, Pan YY, Chen XY, Yuan Y (2020). Effects of betulinic acid on synovial inflammation in rats with collagen-induced arthritis. *International Journal of Immunopathology and Pharmacology* 34:2058738420945078.
- Lamore SD, Cabello CM, Wondrak GT (2010). The topical antimicrobial zinc pyrithione is a heat shock response inducer that causes DNA damage and PARP-dependent energy crisis in human skin cells. *Cell Stress and Chaperones* 15(3):309-322.
- Lassen PR, Skytte DM, Hemmingsen L, Nielsen SF, Freedman TB, Nafie LA, Christensen SB (2005). Structure and absolute configuration of nyasol and hinokiresinol via synthesis and vibrational circular dichroism spectroscopy. *Journal of Natural Products* 68(11):1603-1609.
- Lee EJ, Chung HJ, Pyee Y, Hong JY, Youn UJ, Seo EK, Lee SK (2014). Suppression of inducible nitric oxide synthase expression by nyasol and broussonin A, two phenolic compounds from *Anemarrhena asphodeloides*, through NF- κ B transcriptional regulation *in vitro* and *in vivo*. *Chemistry & Biodiversity* 11(5):749-759.
- Lee GH, Park JS, Jin SW, Pham TH, Thai TN, Kim JY, Kim CY, Choi JH, Han EH, Jeong HG (2020). Betulinic acid induces eNOS expression via the AMPK-dependent KLF2 signaling pathway. *Journal of Agricultural and Food Chemistry* 68(49):14523-14530.
- Lee HJ, Li H, Chang HR, Jung H, Lee DY, Ryu JH (2013). (-)-Nyasol, isolated from *Anemarrhena asphodeloides* suppresses neuroinflammatory response through the inhibition of I- κ B degradation in LPS-stimulated BV-2 microglial cells. *Journal of Enzyme Inhibition and Medicinal Chemistry* 28(5):954-959.
- Lee HJ, Park JS, Yoon YP, Shin YJ, Lee SK, Kim YS, Hong JH, Son KH, Lee CJ (2015). Dioscin and methyl protodioscin isolated from the roots of *Asparagus cochinchinensis* suppressed the gene expression and production of airway MUC5AC mucin induced by phorbol ester and growth factor. *Phytomedicine* 22(5):568-572.
- Li HM, He TT, Zhang M, Liu JN, Zhao X, Liu J, Fang L (2020a). Stilbenoids from the roots of *Stemona tuberosa*. *Natural Product Research* DOI:10.1080/14786419.2020.1798662.
- Li JJ, Yan YY, Sun HM, Liu Y, Su CY, Chen HB, Zhang JY (2019). Anti-cancer effects of pristimerin and the mechanisms: A critical review. *Frontiers in Pharmacology* 10:746.
- Li W, Sun K, Hu F, Chen L, Zhang X, Wang F, Yan B (2020b). Protective effects of natural compounds against oxidative stress in ischemic diseases and cancers via activating the Nrf2 signaling pathway: A mini review. *Journal of Biochemical and Molecular Toxicology* e22658.
- Li X, Liu X, Deng R, Gao S, Jiang Q, Liu R, Li H, Miao Y, Zhai Y, Zhang S, Wang Z, Ren Y, Ning W, Zhou H, Yang C (2021). Betulinic acid attenuated bleomycin-induced pulmonary fibrosis by effectively intervening Wnt/ β -catenin signaling. *Phytomedicine* 81:153428.
- Li XN, Chu C, Cheng DP, Tong SQ, Yan JZ (2012). Norlignans from *Asparagus cochinchinensis*. *Natural Product Communications* 7(10):1357-1358.
- Libman A, Bouamanivong S, Southavong B, Sydara K, Soejarto DD (2006). Medicinal plants: An important asset to health care in a region of Central Laos. *Journal of Ethnopharmacology* 106(3):303-311.
- Libman A, Zhang H, Ma C, Southavong B, Sydara K, Bouamanivong S, Tan GT, Fong HHS, Soejarto DD (2008). A first new antimalarial pregnane glycoside from *Gongronema napalense*. *Asian Journal of Traditional Medicines* 3(6):203-210.
- Lim H, Nam JW, Seo EK, Kim YS, Kim HP (2009). (-)-Nyasol (*cis*-hinokiresinol), a norneolignan from the rhizomes of *Anemarrhena asphodeloides*, is a broad spectrum inhibitor of eicosanoid and nitric oxide production. *Archives of Pharmacol Research* 32(11):1509-1514.
- Lim SP, Noble CG, Shi PY (2015). The dengue virus NS5 protein as a target for drug discovery. *Antiviral Research* 119:57-67.
- Lin LG, Yang XZ, Tang CP, Ke CQ, Zhang JB, Ye Y (2008). Antibacterial stilbenoids from the roots of *Stemona tuberosa*. *Phytochemistry* 69(2):457-463.
- Lin LG, Zhong QX, Cheng TY, Tang CP, Ke CQ, Lin G, Ye Y (2006). Stemoninines from the roots of *Stemona tuberosa*. *Journal of Natural Products* 69(7):1051-1054.
- Liu YP, Ju PK, Long JT, Lai L, Zhao WH, Zhang C, Zhang ZJ, Fu YH (2018). Cytotoxic indole alkaloids from *Nauclea orientalis*. *Natural Product Research* 32(24):2922-2927.
- Liu Z, Fan Y, Wang Y, Han C, Pan Y, Huang H, Ye Y, Luo L, Yin Z (2008). Dipyrithione inhibits lipopolysaccharide-induced iNOS and COX-2 up-regulation in macrophages and protects against endotoxic shock in mice. *FEBS Letters* 582(12):1643-1650.
- Lo WL, Hsu TI, Yang WB, Kao TJ, Wu MH, Huang YN, Yeh SH, Chuang JY (2020). Betulinic acid-mediated tuning of PERK/CHOP signaling by Sp1 inhibition as a novel therapeutic strategy for glioblastoma. *Cancers* 12(4):981.
- Loe MWC, Hao E, Chen M, Li C, Lee RCH, Zhu IXY, Teo ZY, Chin WX, Hou X, Deng J, Chu JJH (2020). Betulinic acid exhibits antiviral effects against dengue virus infection. *Antiviral Research* 184:104954.
- Lomchid P, Nasomjai P, Kanokmedhakul S, Boonmak J, Youngme S, Kanokmedhakul K (2017). Bioactive lupane and hopane triterpenes from *Lepisanthes senegalensis*. *Planta Medica* 83(3/4):334-340.
- Lv XM, Li MD, Cheng S, Liu BL, Liu K, Zhang CF, Xu XH, Zhang M (2018). Neotuberostemonine inhibits the differentiation of lung fibroblasts into myofibroblasts in mice by regulating HIF-1 α signaling. *Acta Pharmacologica Sinica* 39(9):1501-1512.

- Ma CY, Musoke SF, Tan GT, Sydara K, Bouamanivong S, Southavong B, Soejarto DD, Fong HHS, Zhang HJ (2008). Study of antimalarial activity of chemical constituents from *Diospyros quaesita*. *Chemistry and Biodiversity* 5(11):2442-2448.
- Ma YL, Zhang YS, Zhang F, Zhang YY, Thakur K, Zhang JG, Wei ZJ (2019). Methyl protodioscin from *Polygonatum sibiricum* inhibits cervical cancer through cell cycle arrest and apoptosis induction. *Food and Chemical Toxicology* 132:110655.
- Marini-Bettolo GB, Nicoletti M, Messana I, Galeffi C, Msonthi JD, Chapya WA (1985). Research on African medicinal plants-X. Glucosides of *Hypoxis nyasica* Bak. The structure of nyasoside, a new glucoside biologically related to hypoxoside. *Tetrahedron* 41(4):665-670.
- McCord JM, Hybertson BM, Cota-Gomez A, Geraci KP, Gao B (2020). Nrf2 activator PB125[®] as a potential therapeutic agent against COVID-19. *Antioxidants* 9:518.
- Mo J, Lin D, Wang J, Li P, Liu W (2018). Apoptosis in HepG2 cells induced by zinc pyrithione via mitochondrial dysfunction pathway: Involvement of zinc accumulation and oxidative stress. *Ecotoxicology and Environmental Safety* 161:515-525.
- Newman DJ, Cragg GM (2020). Natural products as sources of new drugs over the nearly four decades from 01/1981 to 09/2019. *Journal of Natural Products*, 83(3):770–803.
- Newman M, Ketphanh S, Svengsuksa B, Thomas P, Sengdala K, Lamxay V, Armstrong K (2007). A checklist of the vascular plants of Lao PDR. Royal Botanic Garden Edinburgh.
- Nicholas GM, Blunt JW, Munro MHG (2001). Cortamidine oxide, a novel disulfide metabolite from the New Zealand basidiomycete (mushroom) *Cortinarius* species. *Journal of Natural Products* 64(3):341-344.
- O'Donnell G, Poeschl R, Zimhony O, Gunaratnam M, Moreira JBC, Neidle S, Evangelopoulos D, Bhakta S, Malkinson JP, Boshoff HI, Lenaerts A, Gibbons S (2009). Bioactive pyridine-*N*-oxide disulfides from *Allium stipitatum*. *Journal of Natural Products* 72(3):360-365.
- Otsuka H, Fujioka S, Komiya T, Goto M, Hiramatsu Y, Fujimura H (1981). Studies on antiinflammatory agents. V. A new antiinflammatory constituent of *Pyracantha crenulata* Roem. *Chemical & Pharmaceutical Bulletin* 29(11):3099-3104.
- Pan MH, Chen CM, Lee SW, Chen ZT (2008). Cytotoxic triterpenoids from the root bark of *Helicteres angustifolia*. *Chemistry and Biodiversity* 5(4):565-574.
- Pawelczyk A, Zaprutko L (2020). Anti-COVID drugs: Repurposing existing drugs or search for new complex entities, strategies and perspectives. *Future Medicinal Chemistry* 12(19):1743-1757.
- Qian Z, Wang X, Song Z, Zhang H, Zhou S, Zhao J, Wang H (2015). A phase I trial to evaluate the multiple-dose safety and antitumor activity of ursolic acid liposomes in subjects with advanced solid tumors. *BioMed Research International*, 809714.
- Qin H, Wilson CA, Lee SJ, Zhao X, Benveniste EN (2005). LPS induces *CD40* gene expression through the activation of NF- κ B and STAT-1 α in macrophages and microglia. *Blood* 106(9):3114–3122.
- Quang DN, Khamko VA, Trang NT, Yen LTH, Dien PH (2014). Stemofurans X-Y from the roots of *Stemona* species from Laos. *Natural Product Communications* 9(12):1741–1742.
- Quang DN, Pham CT, Le LTK, Ta QN, Dang NK, Hoang NT, Pham DH (2020). Cytotoxic constituents from *Helicteres hirsuta* collected in Vietnam. *Natural Product Research* 34(4):585–589.
- Ren Y, Anaya-Eugenio GD, Czarnecki AA, Ninh TN, Yuan C, Chai HB, Soejarto DD, Burdette JE, Carcache de Blanco EJ, Kinghorn AD (2018). Cytotoxic and NF- κ B and mitochondrial transmembrane potential inhibitory pentacyclic triterpenoids from *Syzygium corticosum* and their semi-synthetic derivatives. *Bioorganic and Medicinal Chemistry* 26(15):4452-4460.
- Ren Y, Carcache de Blanco EJ, Fuchs JR, Soejarto DD, Burdette JE, Swanson SM, Kinghorn AD (2019). Potential anticancer agents characterized from selected tropical plants. *Journal of Natural Products* 82(3):657-679.
- Ren Y, Kinghorn AD (2019). Natural product triterpenoids and their semi-synthetic derivatives with potential anticancer activity. *Planta Medica* 85(11/12):802-814.
- Ren Y, Lantvit DD, Deng Y, Kanagasabai R, Gallucci JC, Ninh TN, Chai HB, Soejarto DD, Fuchs JR, Yalowich JC, Yu J, Swanson SM, Kinghorn AD (2014). Potent cytotoxic aryl-naphthalene lignan lactones from *Phyllanthus poilanei*. *Journal of Natural Products* 77(6):1494-1504.
- Ríos JL, Máñez S (2018). New pharmacological opportunities for betulinic acid. *Planta Medica* 84(1):8-19.
- Saneja A, Arora D, Kumar R, Dubey RD, Panda AK, Gupta PN (2018). Therapeutic applications of betulinic acid nanoformulations. *Annals of the New York Academy of Sciences* 1421(1):5-18.
- Shaw E, Bernstein J, Losee K, Lott WA (1950). Analogs of aspergillilic acid. IV. Substituted 2-bromopyridine *N*-oxides and their conversion to cyclic thiohydroxamic acids. *Journal of the American Chemical Society* 72:4362-4364.
- Shi JG, Li GQ, Huang SY, Mo SY, Wang Y, Yang YC, Hu WY (2004). Furostanol oligoglycosides from *Asparagus cochinchinensis*. *Journal of Asian Natural Products Research* 6(2):99-105.
- Shi LS, Wu CH, Yang TC, Yao CW, Lin HC, Chang WL (2014). Cytotoxic effect of triterpenoids from the root bark of *Hibiscus syriacus*. *Fitoterapia* 97:184-191.
- Shimizu H, Saito A, Mikuni J, Nakayama EE, Koyama H, Honma T, Shirouzu M, Sekine SI, Shioda T (2019). Discovery of a small molecule inhibitor targeting dengue virus NS5 RNA-dependent RNA polymerase. *PLoS Neglected Tropical Diseases* 13(11):e0007894.
- Soejarto DD, Gyllenhaal C, Kadushin MR, Southavong B, Sydara K, Bouamanivong S, Xaiveu M, Zhang HJ, Franzblau SG, Tan GT, Pezzuto JM, Riley MC, Elkington BG, Waller DP (2012). An ethnobotanical survey of medicinal plants of Laos toward the discovery of bioactive compounds as potential candidates for pharmaceutical development. *Pharmaceutical Biology* 50(1):42-60.
- Soejarto DD, Pezzuto JM, Fong HHS, Tan GT, Zhang HJ, Tamez P, Aydogmus Z, Chien NQ, Franzblau SG, Gyllenhaal C, Regalado JC, Hung NV, Hoang VD, Hiep NT, Xuan LT, Hai NV, Cuong NM, Bich TQ, Loc PK, Vu BM, Southavong BH, Sydara K, Bouamanivong S, O'Neill MJ, Lewis J, Dietzman G (2002). An international collaborative program to discover new drugs from tropical biodiversity of Vietnam and Laos. *Natural Product Sciences* 8(1):1-15.
- Soejarto DD, Zhang HJ, Fong HHS, Tan GT, Ma CY, Gyllenhaal C, Riley MC, Kadushin MR, Franzblau SG, Bich TQ, Cuong NM, Hiep NT, Loc PK, Xuan LT, Hai NV, Hung NV, Chien NQ, Binh LT, Vu BM, Ly HM, Southavong B, Sydara K, Bouamanivong S, Pezzuto JM, Rose WC, Dietzman GR, Miller BE, Thuy TV (2006). "Studies on biodiversity of Vietnam and Laos" 1998–2005: Examining the impact. *Journal of Natural Products* 69(3):473-481.
- Song MC, Yang HJ, Han MW, Kim DK, Kwon BM, Lee KT, Baek NI (2008). Inhibitory activities of cis-hinokiresinol from *Trapa pseudoincisa* on FPTase, PRL-3, and NO-production. *Journal of Applied Biological Chemistry* 51(2):66-68.
- Song X, Wei YX, Lai KM, He ZD, Zhang HJ (2018). *In vivo* antifungal activity of dipyrithione against *Trichophyton rubrum* on guinea pig dermatophytosis models. *Biomedicine and Pharmacotherapy* 108:558-564.
- Soobrattee MA, Baborun T, Aruoma OI (2006). Chemopreventive actions of polyphenolic compounds in cancer. *BioFactors* 27(1-4):19-35.
- Srivastava G, Matta A, Fu G, Somasundaram RT, Datti A, Walfish PG, Ralhan R (2015). Anticancer activity of pyrrithione zinc in oral cancer cells identified in small molecule screens and xenograft model: Implications for oral cancer therapy. *Molecular Oncology* 9(8):1720-1735.
- Sun Q, He M, Zhang M, Zeng S, Chen L, Zhou L, Xu H (2020). Ursolic acid: A systematic review of its pharmacology, toxicity and rethink on its pharmacokinetics based on PK-PD model. *Fitoterapia* 147:104735.
- Sydara K, Gneunphonsavath S, Wahlström R, Freudenthal S, Houamboun K, Tomson G, Falkenberg T (2005). Use of traditional medicine in Lao PDR. *Complementary Therapies in Medicine* 13(3):199-205.
- Tailler M, Senovilla L, Lainey E, Thépot S, Métivier D, Sébert M, Baud V, Billot K, Fenaux P, Galluzzi L, Boehrer S, Kroemer G, Kepp O (2012). Antineoplastic activity of ouabain and pyrithione zinc in acute myeloid leukemia. *Oncogene* 31(30):3536-3546.
- Talib WH, Alsalahat I, Daoud S, Abutayeh RF, Mahmood AI (2020). Plant-derived natural products in cancer research: Extraction,

- mechanism of action, and drug formulation. *Molecules* 25(22):5319.
- Tanachatchairatana T, Bremner JB, Chokchaisiri R, Suksamrarn A (2008). Antimycobacterial activity of cinnamate-based esters of the triterpenes betulinic, oleanolic and ursolic acids. *Chemical and Pharmaceutical Bulletin* 56(2):194-198.
- Tantapakul C, Chaiyosang B, Promgool T, Somteds A, Suthiphasilp V, Kanokmedhakul K, Laphookhieo S, Andersen RJ, Patrick BO, Kanokmedhakul S (2020). Spirosteroids and α -glucosidase inhibitory norlignans from *Asparagus racemosus* Willd. roots. *Phytochemistry* 177:112439.
- Tseng SC, Shen TS, Wu CC, Chang IL, Chen HY, Hsieh CP, Cheng CH, Chen CL (2017). Methyl protodioscin induces apoptosis in human osteosarcoma cells by caspase-dependent and MAPK signaling pathways. *Journal of Agricultural and Food Chemistry* 65(13):2670-2676.
- Tsui WY, Brown GD (1996). (+)-Nyasol from *Asparagus cochinchinensis*. *Phytochemistry* 43(6):1413-1415.
- Wang G, Wang YZ, Yu Y, Yin PH, Xu K (2020a). The antitumor activity of betulinic acid-loaded nanoliposomes against colorectal cancer *in vitro* and *in vivo* via glycolytic and glutaminolytic pathways. *Journal of Biomedical Nanotechnology* 16(2):235-251.
- Wang G, Yu Y, Wang YZ, Zhu ZM, Yin PH, Xu K (2020b). Effects and mechanisms of fatty acid metabolism-mediated glycolysis regulated by betulinic acid-loaded nanoliposomes in colorectal cancer. *Oncology Reports* 44(6):2595-2609.
- Wang L, Tang H, Chen K, Xue LL, Ye HY, Ma LF, Li ZY (2018). Two new triterpenoids from the stems of *Celastrus orbiculatus* Thunb. *Phytochemistry Letters* 27:90-93.
- Wang Q, Li Y, Zheng L, Huang X, Wang Y, Chen CH, Cheng YY, Morris-Natschke SL, Lee KH (2020c). Novel betulinic acid-nucleoside hybrids with potent anti-HIV activity. *ACS Medicinal Chemistry Letters* 11(11):2290-2293.
- Wang YY, Yang YX, Zhe H, He ZX, Zhou SF (2014). Bardoxolone methyl (CDDO-Me) as a therapeutic agent: An update on its pharmacokinetic and pharmacodynamic properties. *Drug Design, Development and Therapy* 8:2075-2088.
- Xiao Q, Zhu W, Feng W, Lee SS, Leung AW, Shen J, Gao L, Xu C (2019). A review of resveratrol as a potent chemoprotective and synergistic agent in cancer chemotherapy. *Frontiers in Pharmacology* 9:1534.
- Yang L, Ren S, Xu F, Ma Z, Liu X, Wang L (2019). Recent advances in the pharmacological activities of dioscin. *BioMed Research International* 5763602.
- Young AN, Herrera D, Huntsman AC, Korkmaz MA, Lantvit DD, Mazumder S, Kolli S, Coss CC, King S, Wang H, Swanson SM, Kinghorn AD, Zhang, X, Phelps MA, Aldrich LN, Fuchs JR, Burdette JE (2018). Phyllanthusmin derivatives induce apoptosis and reduce tumor burden in high-grade serous ovarian cancer by late-stage autophagy inhibition. *Molecular Cancer Therapeutics* 17(10):2123-2135.
- Zhang HJ, Li WF, Fong HHS, Soejarto DD (2016). Discovery of bioactive compounds by the UIC-ICBG drug discovery program in the 18 years since 1998. *Molecules* 21(11):1448.
- Zhang HJ, Sydara K, Tan GT, Ma C, Southavong B, Soejarto DD, Pezzuto JM, Fong HHS (2004). Bioactive constituents from *Asparagus cochinchinensis*. *Journal of Natural Products* 67(2):194-200.
- Zhang Q, Lenardo MJ, Baltimore D (2017). 30 Years of NF- κ B: A blossoming of relevance to human pathobiology. *Cell* 168(1-2):37-57.
- Zhang M, Shen Y, (2007). Progress in pharmacological research of *Asparagus cochinchinensis*. *Shanghai Yiyao* 28(6):266-269.
- Zheng Y, Liu P, Wang N, Wang S, Yang B, Li M, Chen J, Situ H, Xie M, Lin Y, Wang Z (2019). Betulinic acid suppresses breast cancer metastasis by targeting GRP78-mediated glycolysis and ER stress apoptotic pathway. *Oxidative Medicine and Cellular Longevity*, 8781690.
- Zhu H (2017). Floristic characteristics and affinities in Lao PDR, with a reference to the biogeography of the Indochina peninsula. *PLoS One* 12(6):e0179966.
- Zinovkin RA, Grebenchikov OA (2020). Transcription factor Nrf2 as a potential therapeutic target for prevention of cytokine storm in COVID-19 patients. *Biochemistry* 85(7):833-837.