



**INTERNATIONAL JOURNAL OF  
PHARMACEUTICAL SCIENCES**  
[ISSN: 0975-4725; CODEN(USA): IJPS00]  
Journal Homepage: <https://www.ijpsjournal.com>



## Review Article

# Unlocking the Nutritional, Medicinal, and Cultural Treasures of Ragi Millet: A Comprehensive Exploration of Botanical, Historical, and Health Perspectives

**Bhavesh Dharmani\*, Sarita Jangra, Preeti Kumari, Mansi Kumari**

*Chitkara College of Pharmacy, Chitkara University, Punjab (140401), India*

## ARTICLE INFO

Published: 02 Jul. 2025

### Keywords:

Ragi Millet, Eleusine coracana, Finger Millet, Nutrition, Health and adulteration of Millet, and Finger Millet.

### DOI:

10.5281/zenodo.15788117

## ABSTRACT

Cereal grain Ragi (Eleusine coracana) ranks as an essential food grain with historical value since its primary use occurs in Indian states. People have grown this cereal for thousands of years thus making it an essential part of their traditional food behaviours and cooking practices. Published studies form the basis of this paper to investigate deeply the composition of ragi millet nutrition and its therapeutic advantages. The document investigates and assesses the entire range of ragi that begins with its plant characteristics and includes medicinal elements as well as culinary applications with their health benefits and their connected difficulties. Ragi millet contains numerous essential nutrients because it provides both dietary fibers and complex carbohydrates together with multiple vitamins and antioxidants and mineral content and protein fractions. Ragi contains numerous compounds that offer medicine-rich outcomes by giving protection from oxidation and anti-inflammatory results and antidiabetic effects alongside heart protection benefits. Cookers use ragi as both traditional flatbreads and porridges and for making modern breakfast foods and baked goods products. The continuous ingestion of ragi helps users maintain better blood glucose control and healthier bones and helps control their weight while shielding their cardiovascular system. Ragi products need extensive quality checks and government oversight to tackle issues related to product substitution and contamination. Ragi millet serves as a versatile nutritional grain that produces vital outcomes for farming approaches together with food nutrition and healing methods. The complete potential of ragi needs further research attention coupled with sustainable measures to optimize its usage and marketing for human health purposes.

**\*Corresponding Author:** Bhavesh Dharmani

**Address:** Chitkara College of Pharmacy, Chitkara University, Punjab (140401), India

**Email** ✉: [bhavesh20002.ccp@chitkara.edu.in](mailto:bhavesh20002.ccp@chitkara.edu.in)

**Relevant conflicts of interest/financial disclosures:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.



## INTRODUCTION

Traditionally, ragi is used as an essential component through both continental expansions and across various millennia. Archaeological evidence indicates ragi emerged as one of the initial natural grains that human beings domesticated in Africa more than three thousand years ago. The spread of Ragi from its African origin ranges across the Indian subcontinent into multiple Asian territories as humans started using the grain for their primary food source. For hundreds of years ragi served as the primary fundamental crop for Karnataka alongside Andhra Pradesh and Tamil Nadu inhabitants who integrated both culinary arts and cultural ceremonies with this fundamental plant. Throughout centuries people of different cultures have maintained their consumption of ragi because this staple food provides all required nutrients in their daily diets

### Biological Sources

Ragi belongs to the scientific family group Poaceae among plant species. Ragi grows as an annual plant with its stems extending into thin lines while elongated leaves and finger-shaped spikes contain the grains inside. Ragi plants grow to heights between 1-2 meters during their normal life cycle and thrive in various agro-climatic regions from tropical zones through temperate zones. Due to its fine adaptability ragi grows across multiple continents. Ragi uses its tiny compact seeds to produce nutritionally rich food grains, which are used for human consumption

### Medicinal Properties

Ragi has conventionally been used in traditional medicine for its therapeutic properties of relieving digestion issues, controlling diabetes, and supporting bone health by virtue of it being rich in

calcium and fibres. Plant leaves possess antioxidants and yield nutrients that can improve various aspects of health. Individuals apply ragi leaves in the preparation of healthy vegetables to bring vital nutrients into the body. The vital aspect of ragi stays intact to our modern era as it served as the principal diet component and drug in nutritional practice from the beginning of humanity up to present times (Itagi et al., 2013).

### Substitute And Adulteration of Ragi: Challenges and Implications

- **Substitute:**

Ragi is used majorly as a dietary staple because it contains high nutritional value, but despite its nutritional benefits ragi grains sometimes get replaced by alternative cereals depending on what resources are available and what prices are and consumer demands for taste. Alternative grains like sorghum (jowar) and pearl millet (bajra) and foxtail millet join wheat and rice in substituting ragi. Nutritional benefits exist in alternative cereals but people who track their calories can substitute them for different food options. The substitute ingredients deliver comparable nutrition however they lack both ragi's distinctive medicinal advantages and particular medical advantages.

- **Adulteration:**

Market adulterated ragi undergoes contamination by two main methods: adding low-quality grain materials alongside other cereals and fillers and co-mingling it with pesticides and heavy metals. Ragi adulteration happens through two main methods that use low-quality grain mixing with additional grains and filler compounds together with heavy metal and pesticide accumulation in the products. Ragi products tainted with other materials damage the nutritional value of the grain



thus presenting health safety concerns to people who consume these products (Jung et al., 2009).

## Distribution of Ragi Plant Species: A Global Perspective

### Distribution Across India and the World

The plant species *Eleusine coracana* occurs naturally in many regions of India and specific areas throughout the world. Ragi farming serves as a vital agricultural strategy among various Indian states whereas Karnataka Andhra Pradesh Tamil Nadu and Telangana lead the ragi production. The

states support ragi cultivation due to their ideal agro-climatic factors that bring moderate temperatures and sufficient rainfall. Ragi farming exists in India as well as in multiple countries situated throughout Africa along with Asia and parts of the Americas. The food crop ragi serves as a staple in Africa particularly in Uganda Nigeria along with Ethiopia where different names identify it as wimbi, akanwu, and tef respectively. The countries of Nepal and Sri Lanka as well as Myanmar cultivate ragi because it holds critical value in their traditional food systems (Refer Figure 01) (Mangala, Malleshi, et al., 1999).



**Figure 01: The geographical distribution of 297 finger millet accessions with known geo-coordinates distributed in 23 countries around the world.**

### Environmental Conditions and Climates Suitable for Cultivation

Ragi functions well across a broad range of environmental circumstances and climatic conditions allowing its growth in numerous agro-ecological zones. Ragi needs warm temperature climates combined with sufficient rainfall to achieve its best growth outcomes. The cultivation of ragi takes place best in tropical to subtropical zones where the temperatures normally stay between 20°C to 35°C. Ragi shows strong adaptability to hot temperatures and survives through arid conditions thus becoming suitable for areas where many other crops fail to thrive. The

crop tolerates cultivation across lower sea level elevations right up to mountainous altitudes which enables its growth in different geographical regions. Ragi growers normally use rainfall as their irrigation source but also continue cultivation when they have access to sustainable water resources (Matsumoto et al., 1993).

### Varieties of Ragi Plant and Their Adaptability

The different ragi plant varieties maintain distinct features because they function in different geographical zones. Leading types of ragi include Karnataka Red together with CO-9 and MR-1 and GPU-28 alongside other varieties. The various ragi

plant varieties exhibit differences in their grain size characteristics as well as their color and their time to mature and their ability to resist pests and diseases. Karnataka Red stands out as one of the most sought-after varieties because its great production capability helps it resist both living and non-living stress factors (Hegde et al., 2005). The hybrid CO-9 was developed by the University of Agricultural Sciences Bangalore because it displays enhanced resistance to lodging and diseases. Selective ragi varieties by farmers adhere to regional soil environments because of their compatibility with local climate and soil conditions and cropping practices which boosts yield production while withstanding climate stresses. Ragi varieties can thrive successfully in various geographical conditions because they grow in agro-climatic areas from tropical to temperate regions (Urooj & Puttraj, 1999). Scientists carry out breeding programs combined with research to improve both productivity and sustainability in ragi cultivation for sustaining food availability and farm incomes in communal communities. Ragi plants distribute throughout India and worldwide because they successfully adapt to various environmental conditions and climate systems. Ragi demonstrates its position as a resilient versatile crop through its existence across numerous regions of southern India and throughout different countries throughout continents. Additionally different ragi varieties help farmers adapt this nutritious cereal grain to various agro-ecological regions worldwide. Scientists behind ongoing breeding programs and research activities are developing better ragi cultivars that will advance sustainable cultivation during upcoming years (HEGDE & CHANDRA, 2005).

### **Methods And Techniques for The Collection of Millet Grains and Seeds**

Several distinct methods with complementary techniques are used to collect millet grains and seeds for maximizing yield quality and sustainability. Multiple established techniques and methods used in collecting millet grains and seeds get described below:

- **Harvesting:**

The collection process begins with harvesting because workers either by hand or machines extract mature millet grains from their plants. Wild and cultivated millet species become mature between 70 and 120 days from seed planting depending on their type and the surrounding environmental elements. Harvesters collect grains fully matured after achieving their proper moisture quantity. The process of grain collection through traditional methods requires sickles or scythes to pick grains by hand but modern techniques use mechanical harvesters for industrial purposes. The harvesting process requires attention so that mature grains alone are selected because it improves product quality while avoiding damages (Subba Rao & Muralikrishna, 2002; Chethan & Malleshi, 2007).

- **Threshing:**

During threshing operations plants are separated from harvested plant matter through conventional means such as beating or trampling with animals. Questions about threshing methods show that traditional plant processors combine two techniques: plant-beating against solid surfaces as well as animal-stomping activities. The current techniques in grain separation use mechanized threshers along with combine harvesters that combine threshing systems. Threshing both reduces the overall plant material volume and simplifies the following processes of cleaning and winnowing.



- **Cleaning:**

The process of cleaning the harvested grains requires both manual sorting to remove damaged grains and mechanical methods for impurity elimination. Cleaning operations improve grains' quality and reduce contamination risks which occur during storage as well as processing. Cleaning includes manual sorting for damaged or discoloured grains just as it includes mechanical methods between sieving and screening and air aspiration for eliminating dust and husks. Clean and uniform grains survive the processing phase after proper cleaning methods create them available for consumption (Krishnan et al., 2012).

- **Winnowing:**

Winnowing stands as a conventional grain separation process which separates grains from the chaff and other lightweight materials through directed air flow. A traditional outdoor procedure separates grains from chaff particles by throwing them into the air through baskets or trays until wind dictates heavier grains to fall while chaff items remain in the air. Modern processing of the winnowing method employs mechanical fans or blowers as air stream generators to efficiently separate grains from chaff. Winnowing acts as an additional cleaning method that determines what harvested grains remain while discarding unwanted substances prior to storage or processing (Vitaglione et al., 2009).

- **Drying and Storage:**

The harvested grains necessitate appropriate drying through proper methods that lower their moisture content to stop storage spoilage. The process of drying normally involves laying grains into thin thin layers on dry clean surfaces which receive exposure to sunlight and air at the same time. Mechanical dryers function as an alternative

method to enhance the drying speed for situations that include humid or rainy weather conditions.

Cultivation of millet seeds requires systematic procedures starting with harvest and then moving to threshing followed by cleaning steps before achieving successful winnowing procedures and completing drying and storage. Proper method usage throughout the entire process allows farmers to keep millet grains fresh with high yield together with excellent quality during their storage time (Vetriventhan et al., 2020a; Dhingra et al., 2012).

### **Traditional And Modern Cultivation Practices for Millet Plants**

Engineered agricultural systems integrate conventional farmer management practices that protect ecosystems through increased output from specific agricultural areas. Standard operating procedures for farm activities are created through the combination of recorded weather data and cultural protocols by traditional farmers. Each agricultural field gets divided into unique zones which farmers operate under the conventional and modern cultivation techniques while receiving payments from the production of millet. This section applies contemporary agricultural practices to the age-old tradition of millet farming. (Sharma & Gujral, 2010):

**Traditional Cultivation Practices:** The guidelines for millet cultivation through generations have developed from indigenous cultural methods which have been transmitted from one generation to another (Vetriventhan et al., 2020b). Thrusting manual work together with minimal agricultural input methods matches the requirements of farming on small plots of land. Traditional practices may include:



- **Crop Rotation:** Farmers rotate millet crops with other crops such as pulses, legumes, or oilseeds to maintain soil fertility and reduce pest and disease pressure.
- **Mixed Cropping:** Intercropping millet with different crops helps farmers optimize land usage for higher yield stability results. Millet farmers commonly plant their crops in combination with sorghum, maize and legumes alongside each other.
- **Agroforestry:** Some traditional farming communities practice agroforestry, where millet is cultivated alongside trees or shrubs to provide shade, soil conservation, and additional income from timber or fruit production.
- **Rainwater Harvesting:** In rainfed areas, traditional farmers may employ rainwater harvesting techniques such as contour bunding, trenching, or check dams to capture and store rainwater for irrigation during dry periods (P. & Joye, 2020).

**Modern Cultivation Practices:** Narrowed modern millets farming processes utilize networking machinery for developing effective sustainable production techniques. Commercial farms first adopted these practices which then spread to smaller operating farms. Modern cultivation practices include: (Rao et al., 2022)

- **Mechanization:** The use of automated farming equipment comprising tractors seed drills and combine harvesters helps contemporary farmers minimize labor expenses while boosting their operational efficiency during plantation and crop harvesting and handling phases (Nkhata et al., 2018).
- **Irrigation:** Modern farmers establish irrigation infrastructure using drip irrigation and sprinkler systems and pumps groundwater

in regions with dry climate to maintain water access during cultivation.

- **Precision Agriculture:** Modern farmers adopt precision agriculture techniques, including soil testing, remote sensing, and GPS-guided equipment, to optimize inputs such as fertilizers, water, and pesticides, thereby minimizing costs and environmental impact.

### Factors Influencing Successful Cultivation

A successful millet cultivation depends on the integrated relationship of soil types with water requirements which operates in conjunction with temperature and climate and pest-disease management systems. Cultivators of millet plants make decisions about traditional or modern farming based on their site-specific requirements. Traditional farming maintains food production stability through systematic implementation of purpose-built agricultural modern technologies in well-defined farming areas devoted to economic growth. Farmers seeking crop protection paired with next-season reserve stocks must create fundamental cultivation practices (G. Tripathi et al., 2023).

### Detailed Analysis of Phytoconstituents in Millet Grains

**Phenolic Compounds and Flavonoids:** Millet grains contain predominantly two phenolic compounds including ferulic and p-coumaric acid phenolic acids with luteolin flavonoids among the main components. These compounds found in grains provide all their health benefits through their antioxidant properties which additionally demonstrate anti-inflammatory abilities to protect against cancer cells. The extensive occurrence of ferulic acid in millet grains allows it to perform as the main component since this substance promotes heart health and enhances immune response.



**Vitamins and Minerals:** The essential B-complex vitamins and vitamin E present in Millet grains exist at high density levels. Millet grains provide B complex vitamins and vitamin E which the brain requires to perform correctly through energy processes and defence cell function against damage. Millet grains provide appreciable levels of necessary elements with a specific concentration of iron and magnesium and significant amounts of phosphorus and potassium as well as zinc present. Your body needs minerals for two crucial purposes including conducting biological operations while building strong bones to strengthen your immune system (B. Tripathi & Platel, 2010).

**Phytosterols:** Millet crops have phytosterols as plant components which share cholesterol structure and chemistry. The cardiovascular protection and cholesterol reduction properties arise from phytosterol compounds such as stigmasterol, campesterol, and sitosterol that exist within millet grains. Eating balanced diets with millet has two health benefits because it lowers blood cholesterol while stopping cholesterol from entering the gut.

**Dietary Fiber:** The dietary fiber in millet grains consists of two categories: insoluble and soluble fiber. The soluble form of fiber creates a gel substance that provides a feeling of satiety while

helping to regulate blood glucose levels and lipid metabolism. The insoluble fiber component in foods acts to bulk up stool and conceals digestive health and bowel regularity both. The high dietary fiber content of millet grains proves essential for both gastrointestinal welfare and general human wellbeing (Ugare et al., 2014).

Different phytoconstituents present in millet grains create a comprehensive healthcare benefit system. Millet grains deliver complete health support because they contain phenolic compounds and flavonoids together with vitamins and minerals as well as phytosterols and various other phytochemical components. Through their phytoconstituent composition millets serve as essential partners between people and optimal health protection.

### Discussion About Ragi Millet's Nutritional and Therapeutic Benefits

Indian people have consumed the versatile nutritious millet ragi as their staple diet for many generations across their country. Ragi serves as a valuable dietary supplement because it has remarkable nutritional content and therapeutic benefits. This product carries the name finger millet (Viswanath et al., 2009). The following table 1 explains, the nutritional attributes and therapeutic advantages of ragi millet:

**Table 1: Discussion on the Nutritional and Medicinal Properties of Ragi Millet.**

| Sr. No. | Bioactive Compound Name | Property                                   | Use                                                                                                                                                                                                                                                                                                                                                              |
|---------|-------------------------|--------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 1       | Phenolic Compounds      | Antioxidant, Anti-inflammatory, Anticancer | The phenolic compounds ferulic acid, p-coumaric acid, and caffeic acid provide high concentrations in ragi millet especially within its bran and outer parts. Ragi millet contains antioxidant substances which protect cell structures from oxidative damage while reducing risks of cancer development together with heart problems and neurological diseases. |
| 2       | Flavonoids              | Antioxidant, Anti-inflammatory             | The flavonoids within ragi millet consist of luteolin, apigenin along with quercetin. The antioxidant properties with anti-inflammatory effects of Ragi millet emerge from its flavonoid substances that help maintain overall health benefits.                                                                                                                  |



|   |          |                                                                 |                                                                                                                                                                                                                                                                                                                                                                            |
|---|----------|-----------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 3 | Vitamins | Essential for various physiological functions                   | The nutritional component of ragi millet contains B-complex vitamins such as thiamine (vitamin B1), riboflavin (vitamin B2), niacin (vitamin B3) and folate (vitamin B9) together with vitamin E. The neurological system depends on these vitamins together with their contribution to energy metabolism and antioxidant defence functions and red blood cell production. |
| 4 | Minerals | Essential for metabolic processes, Bone health, Immune function | A number of essential minerals exists in ragi millet including iron, magnesium, phosphorus, potassium and zinc and phosphorus. Ragi millet contains limited essential minerals which help construct muscle and immune function along with bone structure and contribute to appropriate oxygen circulation in the body.                                                     |

**Culinary Uses of Ragi Grains** Ragi grains have become popular in homes for their mineral content and special taste profile. Ragi grains find widespread use in cooking because they have (Malleshi et al., 1986):

- **Ragi Flour:** The traditional method of preparing ragi flour involves grinding the grains before using it to make flatbreads and pancakes alongside steamed cakes and porridges. Commonly, individuals prefer use of ragi grains as a sauce thickener combinedly with stews and soups.
- **Beverages:** Ragi grains customarily follow a production process which starts with their sprouting and subsequent drying and grinding practice until they become ragi malt - a well-known nutritious beverage that people in South India enjoy. To make ragi malt one combines ragi flour with water or milk until it becomes ready while adding jaggery or sugar for sweetness. This high-calcium food along

with iron content gives essential nutrients which people frequently consume as breakfast and snack.

- **Snack Foods:** One can pop ragi grains into popcorn then spice them as an attractive nutritious snack. Ragi flakes together with puffed ragi serve as nutritious components for trail mixes and granola bars apart from other snack foods.
- **Fermented Foods:** Ragi flour becomes traditional fermented foods during fermentation such as ragi mudde (steamed ragi dumplings) and idlis and dosas. Ragi flour fermentation processes both complex carbohydrates and makes nutrients more accessible for digestion.

### Nutritional Benefits of Ragi

Ragi grains - highly nutritious and offer several health benefits due to their unique composition. Some of the nutritional benefits of ragi include (Pereira et al., 2002): Refer Figure 02.




Figure 02: Nutritional Benefits of Ragi.

### Associated Risks and Toxic Effects

While ragi grains offer numerous health benefits, there are some potential risks and toxic effects associated with their consumption:

- **Antinutritional Factors:** Phytates tannins and polyphenols found in Ragi inhibit absorption of iron and zinc from body and lower their availability to the human body (Platel et al., 2010). The antinutritional factors in ragi decrease through soaking techniques that lead to sprouting as well as through fermentation processes that boost nutritional content.
- **Allergic Reactions:** Ragi causes allergies in particular individuals who react adversely to its consumption. Allergic reactions to ragi may manifest through skin itching and rashing along with swelling and breathing obstruction as well as gastrointestinal distress. Pregnant women who have displayed allergies to gluten as well as other grains need to exercise care before eating ragi and should consult medical professionals if symptoms of an allergic reaction appear.
- **Cyanogenic Glycosides:** Ragi contains low quantities of the glycoside compounds cyanogenic glycosides but these compounds produce cyanide when eaten. Moderate consumptions of ragi do not result in toxic reactions because its cyanogenic glycoside levels remain at low concentrations (Platel et al., 2010).

- **Digestive Issues:** People suffer from digestive issues which include gas as well as bloating and stomach discomfort when they eat ragi grains or consume items made from ragi flour.

Ragi grains deliver many nutritional advantages to consumers because they are staple food that people eat extensively throughout the world but individuals need to know both the potential harm and toxicity conditions. People who understand how to prepare and benefit from ragi grains alongside its health risks will be able to select this nutritious food safely for their diet.

### Examination of Potential Toxic Compounds in Ragi

Like many other foods, ragi millet contains certain compounds that may have toxic effects if consumed in large quantities or inappropriately processed (Platel & Shurpalekar, 1994). Here, we examine the potential toxic compounds found in ragi:

- **Antinutritional Factors:** Ragi includes three groups of antinutritional components named tannins phytates and polyphenols. Mineral absorption problems develop over time because these compounds prevent zinc and iron from entering the body effectively. The nutritional value of ragi becomes better

through antinutritional factor reduction achieved through soaking and fermentation alongside sprouting as processing methods.

- **Pesticide Residues:** The cultivation process of ragi leads to crop contact with pesticides at the same time as chemical fertilizers. The chemicals leave behind residual amounts on grains that create health risks when excessive amounts are eaten. The best practice to avoid pesticide exposure from ragi includes selecting organic variants and sustainably cultivated ragi together with a proper cleaning step before eating (Chandrasekara & Shahidi, 2010).
- **Mycotoxins:** The susceptibility of Ragi grains to fungal growth occurs mainly when they are stored under warm and humid environments. Ragi grains exposed to fungal growth develop mycotoxins which are toxic substances that produce detrimental health effects in human beings. Aflatoxins represent among the most common mycotoxins found in ragi grain because this substance has both cancer-causing properties and harm liver tissues. Using dry storage areas with ventilation prevents mold growth which lowers the chances of mycotoxin contamination in ragi grains (Awika et al., 2003).

### Health Risks Associated with Exclusive Consumption of Ragi

Despite ragi's nutritional benefits, individuals exclusive on ragi consumption may face some health-associated risks such as: (Refer table 2):

- **Nutrient Imbalance:** Ragi has essential nutrients including both iron and dietary fiber as well as calcium yet it lacks essential

nutrients found in a balanced diet. Consuming ragi continuously over a long period creates nutritional deficiencies because it fails to provide adequate quantities of food from all four nutritional groups including fruits and vegetables and proteins and fats (Modak et al., 2007).

- **Phytate Content:** The phytates found in ragi grains inhibit both iron and zinc from getting absorbed within the human body. Eating large amounts of unprocessed ragi-based foods leads to mineral deficiency health problems due to high phytate levels (Nkurunziza et al., 2020).
- **Gastrointestinal Discomfort:** Hydrating foods made from ragi grains offer dietary fiber to enhance digestive health as well as promote digestive regularity. The consumption of excessive ragi can cause digestive issues when fluid intake is insufficient because it leads to bloating and gas production and constipation. To avoid digestive problems people should consume ragi-based foods in controlled portions together with maintaining proper diet balance (Dias et al., 2021).
- **Potential Allergic Reactions:** Ragi allergy exists in certain people together with allergic reactions that occur when they only eat ragi-based foods. Users of ragi-based foods might develop various allergic responses that cause skin inflammations and itching alongside lung swelling and breathing trouble along with digestive issues. Help from a doctor should be sought by people allergic to other grains or gluten before they try ragi because it can cause allergic reactions (Backiyalakshmi et al., 2021).

**Table 2: Ragi & it's Ill Effects (Long-Term Use)**

| Sr No. | Active Part | Application                                           | Ill Effects (Long-term Use)                                                |
|--------|-------------|-------------------------------------------------------|----------------------------------------------------------------------------|
| 1      | Whole grain | Culinary uses: Ragi is used to make roti, dosa, idli, | Ragi consumption as a sole diet leads to nutrient imbalance and deficiency |



|   |          |                                                                                                          |                                                                                                                                                                                         |
|---|----------|----------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
|   |          | porridge, and other traditional dishes.                                                                  | problems when people do not incorporate other food categories.                                                                                                                          |
| 2 | Flour    | Baking: Ragi flour is used in baking bread, cookies, cakes, and other baked goods.                       | The combination of inadequate water intake together with excessive chia intake leads to gastrointestinal distress that manifests as bloating and gas formation along with constipation. |
| 3 | Sprouts  | Salads: Ragi sprouts can be added to salads for added texture and nutrition.                             | Some people suffer from ragi allergies together with allergic symptoms which include rashes or itching on their skin when they eat ragi only.                                           |
| 4 | Extracts | Nutritional supplements: Ragi extracts may be used in nutritional supplements for added health benefits. | Ragi contains phytates that block mineral absorption thus leading to potential mineral deficiencies when used for extended periods.                                                     |

## CONCLUSION

Research on ragi or finger millet shows that this food holds major nutritional properties together with strong medicinal qualities. As a multi-purpose grain ragi helps digestion and weight maintenance and protects against anaemia and diabetes. The realm of agricultural production together with food nutrition and human health benefits from ragi in a vital way. Scientists should dedicate their research to determining how exact bioactive compounds inside ragi regulate human wellness through proper scientific testing methods. Agricultural technology implemented with processing and breeding research develops improved cultivation patterns and nutritional products originating from the same ragi source. Farming ragi leads to financially productive farmers who generate enhanced nutrition for worldwide public wellness. Through Regi cultivation food security ensures agricultural growth remains balanced across multiple environments. Ragi functions as an important food substance because consuming it provides multiple health benefits for different medical conditions and protective health outcomes. Farmers who consume ragi benefit from multiple nutritional

values leading to enhanced economic outputs and public health advancement worldwide.

## REFERENCES

1. Abeysinghe, D. T.; Alwis, D. D. D. H.; Kumara, K. A. H.; Chandrika, U. G. Nutritive Importance and Therapeutics Uses of Three Different Varieties (Murraya Koenigii, Micromelum Minutum, and Clausena Indica) of Curry Leaves: An Updated Review. *Evid Based Complement Alternat Med* 2021, 2021, 5523252. <https://doi.org/10.1155/2021/5523252>.
2. Antony Ceasar, S.; Maharajan, T.; Ajeesh Krishna, T. P.; Ramakrishnan, M.; Victor Roch, G.; Satish, L.; Ignacimuthu, S. Finger Millet [Eleusine Coracana (L.) Gaertn.] Improvement: Current Status and Future Interventions of Whole Genome Sequence. *Front Plant Sci* 2018, 9, 1054. <https://doi.org/10.3389/fpls.2018.01054>.
3. Awika, J. M.; Rooney, L. W.; Wu, X.; Prior, R. L.; Cisneros-Zevallos, L. Screening Methods to Measure Antioxidant Activity of Sorghum (*Sorghum Bicolor*) and Sorghum Products. *J Agric Food Chem* 2003, 51 (23),



- 6657–6662.  
<https://doi.org/10.1021/jf034790i>.
4. Backiyalakshmi, C.; Vetriventhan, M.; Deshpande, S.; Babu, C.; Allan, V.; Naresh, D.; Gupta, R.; Azevedo, V. C. R. Genome-Wide Assessment of Population Structure and Genetic Diversity of the Global Finger Millet Germplasm Panel Conserved at the ICRISAT Genebank. *Front Plant Sci* 2021, 12, 692463. <https://doi.org/10.3389/fpls.2021.692463>.
5. Behl, T., Kumar, K., Brisc, C., Rus, M., Nistor-Cseppento, D. C., Bustea, C., Aron, R. A. C., Pantis, C., Zengin, G., Sehgal, A., Kaur, R., Kumar, A., Arora, S., Setia, D., Chandel, D., & Bungau, S. (2021). Exploring the multifocal role of phytochemicals as immunomodulators. *Biomedicine & pharmacotherapy* = *Biomedecine & pharmacotherapie*, 133, 110959. <https://doi.org/10.1016/j.biopha.2020.110959>
6. Chandrasekara, A.; Shahidi, F. Content of Insoluble Bound Phenolics in Millets and Their Contribution to Antioxidant Capacity. *J Agric Food Chem* 2010, 58 (11), 6706–6714. <https://doi.org/10.1021/jf100868b>.
7. Chethan Nagappa G., S.; M. Finger Millet Polyphenols: Characterization and Their Nutraceutical Potential. *Am J Food Technol* 2007, 2 (7), 582–592. <https://doi.org/10.3923/ajft.2007.582.592>.
8. Dhingra, D.; Michael, M.; Rajput, H.; Patil, R. T. Dietary Fibre in Foods: A Review. *J Food Sci Technol* 2012, 49 (3), 255–266. <https://doi.org/10.1007/s13197-011-0365-5>.
9. Dias, M. C.; Pinto, D. C. G. A.; Silva, A. M. S. Plant Flavonoids: Chemical Characteristics and Biological Activity. *Molecules* 2021, 26 (17). <https://doi.org/10.3390/molecules26175377>.
10. Forte, M.; Bianchi, F.; Cotugno, M.; Marchitti, S.; Stanzione, R.; Maglione, V.; Sciarretta, S.; Valenti, V.; Carnevale, R.; Versaci, F.; Frati, G.; Volpe, M.; Rubattu, S. An Interplay between UCP2 and ROS Protects Cells from High-Salt-Induced Injury through Autophagy Stimulation. *Cell Death Dis* 2021, 12 (10), 919. <https://doi.org/10.1038/s41419-021-04188-4>.
11. Gheorghe, G., Toth, P. P., Bungau, S., Behl, T., Ilie, M., Pantea Stoian, A., Bratu, O. G., Bacalbasa, N., Rus, M., & Diaconu, C. C. (2020). Cardiovascular Risk and Statin Therapy Considerations in Women. *Diagnostics (Basel, Switzerland)*, 10(7), 483. <https://doi.org/10.3390/diagnostics10070483>
12. Hassan, Z. M.; Sebola, N. A.; Mabelebele, M. The Nutritional Use of Millet Grain for Food and Feed: A Review. *Agric Food Secur* 2021, 10 (1), 16. <https://doi.org/10.1186/s40066-020-00282-6>.
13. Hegde Namakkal S.; Chandra, T.S., P. S. ; R. Effects of the Antioxidant Properties of Millet Species on Oxidative Stress and Glycemic Status in Alloxan-Induced Rats. *Nutrition Research* 2005, 25 (12), 1109–1120. <https://doi.org/10.1016/j.nutres.2005.09.020>.
14. Hegde T.S., P. S. ; C. ESR Spectroscopic Study Reveals Higher Free Radical Quenching Potential in Kodo Millet (*Paspalum Scrobiculatum*) Compared to Other Millets. *Food Chem* 2005, 92 (1), 177–182. <https://doi.org/10.1016/j.foodchem.2004.08.002>.
15. Itagi Vasudeva; Indiramma, A. R.; Prakash, Maya, H. N.; S. Shelf Stable Multigrain Halwa Mixes: Preparation of Halwa, Their Textural and Sensory Studies. *J Food Sci Technol* 2011, 50 (5), 879–889. <https://doi.org/10.1007/s13197-011-0423-z>.
16. Jung Hyung Joo; Hong, Wan Soo; Kim, Donggeon; Hong, Yang Hee; Hong, In Sun; Chang, Un Jae, E. Y. S. Uncooked Rice of Relatively Low Gelatinization Degree

- Resulted in Lower Metabolic Glucose and Insulin Responses Compared with Cooked Rice in Female College Students. *Nutr Res* 2009, 29 (7), 457–461. <https://doi.org/10.1016/j.nutres.2009.07.002>.
17. Krishnan Usha; Malleshi, Nagappa G., R. D. Influence of Decortication, Popping and Malting on Bioaccessibility of Calcium, Iron and Zinc in Finger Millet. *LWT - Food Science and Technology* 2012, 48 (2), 169–174. <https://doi.org/10.1016/j.lwt.2012.03.003>.
  18. Nakamura, M. T.; Yudell, B. E.; Loor, J. J. Regulation of Energy Metabolism by Long-Chain Fatty Acids. *Prog Lipid Res* 2014, 53, 124–144. <https://doi.org/10.1016/j.plipres.2013.12.001>.
  19. Nkhata, S. G.; Ayua, E.; Kamau, E. H.; Shingiro, J.-B. Fermentation and Germination Improve Nutritional Value of Cereals and Legumes through Activation of Endogenous Enzymes. *Food Sci Nutr* 2018, 6 (8), 2446–2458. <https://doi.org/10.1002/fsn3.846>.
  20. Nkurunziza, L.; Watson, C. A.; Öborn, I.; Smith, H. G.; Bergkvist, G.; Bengtsson, J. Socio-Ecological Factors Determine Crop Performance in Agricultural Systems. *Sci Rep* 2020, 10 (1), 4232. <https://doi.org/10.1038/s41598-020-60927-1>.
  21. Malleshi H. S. R.; Tharanathan, Rudrapatnam N., N. G.; D. Free Sugars and Non-Starchy Polysaccharides of Finger Millet (*Eleusine Coracana*), Pearl Millet (*Pennisetum Typhoideum*), Foxtail Millet (*Setaria Italica*) and Their Malts. *Food Chem* 1986, 20 (4), 253–261. [https://doi.org/10.1016/0308-8146\(86\)90095-6](https://doi.org/10.1016/0308-8146(86)90095-6).
  22. Mangala H.P; Udayasankar, K; Tharanathan, R.N, S. L. R. Resistant Starch Derived from Processed Ragi (Finger Millet, *Eleusine Coracana*) Flour: Structural Characterization. *Food Chem* 1999, 64 (4), 475–479. [https://doi.org/10.1016/s0308-8146\(98\)00129-0](https://doi.org/10.1016/s0308-8146(98)00129-0).
  23. Mangala Nagappa G.; Tharanathan, Rudrapatnam N.; Mahadevamma, NA, S. L. ; M. Resistant Starch from Differently Processed Rice and Ragi (Finger Millet). *European Food Research and Technology* 1999, 209 (1), 32–37. <https://doi.org/10.1007/s002170050452>.
  24. Matsumoto Fumiko; Ishigaki, Akiyo; Iwashina, Hirotoshi; Hara, Yukihiko, N. I. Reduction of Blood Glucose Levels by Tea Catechin. *Biosci Biotechnol Biochem* 1993, 57 (4), 525–527. <https://doi.org/10.1271/bbb.57.525>.
  25. Modak, M.; Dixit, P.; Londhe, J.; Ghaskadbi, S.; Devasagayam, T. P. A. Indian Herbs and Herbal Drugs Used for the Treatment of Diabetes. *J Clin Biochem Nutr* 2007, 40 (3), 163–173. <https://doi.org/10.3164/jcbtn.40.163>.
  26. Mohan Ganesan; Sathya, R. M.; Tamil, Selvi Ramjothi; Ganesan, Anbazhagan; Sudha, Vasudevan, V. R. Dietary Carbohydrates, Glycaemic Load, Food Groups and Newly Detected Type 2 Diabetes among Urban Asian Indian Population in Chennai, India (Chennai Urban Rural Epidemiology Study 59). *Br J Nutr* 2009, 102 (10), 1498–1506. <https://doi.org/10.1017/s0007114509990468>.
  27. Okolo, J. C.; Adejumo, B. A. Finger Millet: A Crop with Food Security Potentials for Africans. In *Food Security and Safety: African Perspectives*; Babalola, O. O., Ed.; Springer International Publishing: Cham, 2021; pp 139–154. [https://doi.org/10.1007/978-3-030-50672-8\\_7](https://doi.org/10.1007/978-3-030-50672-8_7).
  28. P, N. P. V; Joye, I. J. Dietary Fibre from Whole Grains and Their Benefits on Metabolic Health. *Nutrients* 2020, 12 (10). <https://doi.org/10.3390/nu12103045>

29. Pereira David R.; Pins, Joel J.; Raatz, Susan K.; Gross, Myron D.; Slavin, Joanne L.; Seaquist, Elizabeth R., M. A.; J. Effect of Whole Grains on Insulin Sensitivity in Overweight Hyperinsulinemic Adults. *Am J Clin Nutr* 2002, 75 (5), 848–855. <https://doi.org/10.1093/ajcn/75.5.848>.
30. Platel Kantha S., K.; S. Resistant Starch Content of Indian Foods. *Plant Foods Hum Nutr* 1994, 45 (1), 91–95. <https://doi.org/10.1007/bf01091233>.
31. Platel Sushma W; Srinivasan, Krishnapura, K. E. Bioaccessible Mineral Content of Malted Finger Millet (Eleusine Coracana), Wheat (Triticum Aestivum), and Barley (Hordeum Vulgare). *J Agric Food Chem* 2010, 58 (13), 8100–8103. <https://doi.org/10.1021/jf100846e>.
32. Rao G., M. V. S. S. T. S. M. Evaluation of the Antioxidant Properties of Free and Bound Phenolic Acids from Native and Malted Finger Millet (Ragi, Eleusine Coracana Indaf-15). *J Agric Food Chem* 2002, 50 (4), 889–892. <https://doi.org/10.1021/jf011210d>.
33. Rao, M. J.; ul Qamar, M.; Wang, D.; Ali, Q.; Ma, L.; Han, S.; Duan, M.; Hu, L.; Wang, L. A High-Throughput Lipidomics and Transcriptomic Approach Reveals Novel Compounds from Sugarcane Linked with Promising Therapeutic Potential against COVID-19. *Front Nutr* 2022, 9. <https://doi.org/10.3389/fnut.2022.988249>.
34. Sharma, P.; Gujral, H. Antioxidant and Polyphenol Oxidase Activity of Germinated Barley and Its Milling Fractions. *Food Chem* 2010, 120, 673–678. <https://doi.org/10.1016/j.foodchem.2009.10.059>.
35. Sripriya K.; Murty, V.S.; Chandra, T.S., G.; C. ESR Spectroscopic Studies on Free Radical Quenching Action of Finger Millet (Eleusine Coracana). *Food Chem* 1996, 57 (4), 537–540. [https://doi.org/10.1016/s0308-8146\(96\)00187-2](https://doi.org/10.1016/s0308-8146(96)00187-2).
36. Sumathi S.; Malleshi, Nagappa G.; Rao, S. Venkat, A.; V. Glycemic Response to Malted, Popped and Roller Dried Wheat-Legume Based Foods in Normal Subjects. *Int J Food Sci Nutr* 1997, 48 (2), 103–107. <https://doi.org/10.3109/09637489709006968>.
37. Tripathi, G.; Harsh, P.; B++, J.; Borah, A.; Das, H.; Bansal, S.; Singh, N.; Singh, B.; Nath, D. A Review on Nutritional and Health Benefits of Millets. *Int J Plant Soil Sci* 2023, 35, 1736–1743. <https://doi.org/10.9734/IJPSS/2023/v35i193722>.
38. Tripathi Kalpana, B. P. Finger Millet (Eleusine Coracana) Flour as a Vehicle for Fortification with Zinc. *J Trace Elem Med Biol* 2009, 24 (1), 46–51. <https://doi.org/10.1016/j.jtemb.2009.09.001>.
39. Ugare Bharati V.; Naik, Rama K.; Bharati, Pushpa; Itagi, Sunanda, R. C. Glycemic Index and Significance of Barnyard Millet (Echinochloa Frumentaceae) in Type II Diabetics. *J Food Sci Technol* 2011, 51 (2), 392–395. <https://doi.org/10.1007/s13197-011-0516-8>.
40. Urooj S., A. P. Digestibility Index and Factors Affecting Rate of Starch Digestion in Vitro in Conventional Food Preparation. *Nahrung* 1999, 43 (1), 42–47. [https://doi.org/10.1002/\(sici\)1521-3803\(19990101\)43:1<42::aid-food42>3.0.co;2-q](https://doi.org/10.1002/(sici)1521-3803(19990101)43:1<42::aid-food42>3.0.co;2-q).
41. Vetriventhan, M.; Azevedo, V.; Upadhyaya, H.; Nirmalakumari, A.; Kane-Potaka, J.; Seetha, A.; Ceasar, S. A.; Muthamilarasan, M.; Bhat, B.; K., H.; Bellundagi, A.; Cheruku, D.; Backiyalakshmi, ; Santra, ; Vanniarajan, C.; Tonapi, V. Genetic and Genomic Resources, and Breeding for Accelerating Improvement of Small Millets: Current Status

- and Future Interventions. Nucleus (India) 2020, 63. <https://doi.org/10.1007/s13237-020-00322-3>.
42. Viswanath Asna; Malleshi, Nagappa G., V. U. Evaluation of Antioxidant and Antimicrobial Properties of Finger Millet Polyphenols (*Eleusine Coracana*). *Food Chem* 2009, 114 (1), 340–346. <https://doi.org/10.1016/j.foodchem.2008.09.053>.
43. Vitaglione, P.; Lumaga, R. B.; Stanzione, A.; Scalfi, L.; Fogliano, V. Beta-Glucan-Enriched Bread Reduces Energy Intake and Modifies Plasma Ghrelin and Peptide YY Concentrations in the Short Term. *Appetite* 2009, 53 (3), 338–344. <https://doi.org/10.1016/j.appet.2009.07.013>.

**HOW TO CITE:** Bhavesh Dharmani, Sarita Jangra, Preeti Kumari, Mansi Kumari, Unlocking the Nutritional, Medicinal, and Cultural Treasures of Ragi Millet: A Comprehensive Exploration of Botanical, Historical, and Health Perspectives, *Int. J. of Pharm. Sci.*, 2025, Vol 3, Issue 7, 198-212. <https://doi.org/10.5281/zenodo.15788117>

