

In Vitro Antioxidant, Antimicrobial, *Insilico* Molecular Docking Analysis and Phylogenetic Analysis of *Anoectochilus elatus*

Sathishkumar. M *, Lakha Sri T.K * and Selvi. S**

* Research Scholar, PG and Research Department of Biochemistry, Bharathidasan College of Arts and Science, Erode, Tamil Nadu, India.

** Associate Professor PG and Research Department of Biochemistry, Bharathidasan College of Arts and Science, Erode, Tamil Nadu, India.

Corresponding Author : Selvi. S

Received: Aug 18, 2025

Accepted: Sep 12, 2025

Published: Sep 13, 2025

Abstract - Medicinal plants have long been recognized for their therapeutic effects, serving as a rich source of bioactive compounds. Throughout the millennia, diverse cultures around the world have developed intricate systems of traditional medicine, utilizing a vast array of plant-based remedies to treat several ailments.

Keywords: Pathogens, Antioxidant, *Anoectochilus elatus*, *Insilico* analysis, Maturase K.

I. INTRODUCTION

The use of medicinal plants in the management of diseases has a rich and extensive history, dating back to ancient civilizations. Throughout the millennia, diverse cultures around the world have developed intricate systems of traditional medicine, utilizing a vast array of plant-based remedies to treat several ailments [1] (Salmerón-Manzano *et al.*, 2020). Recorded history provides ample evidence of the long-standing utilization of herbal plants. The Indian medical system, rooted in ancient Ayurvedic traditions, has long recognized the remarkable therapeutic efficacy of herbal plants. Ayurveda has documented the use of extensive plants and formulations to treat numerous diseases. The rich diversity of India's flora, with an estimated 250,000 plant species, has served as the foundation for a vast array of plant-based remedies [2] (Riaz *et al.*, 2023).

Traditional healing systems, such as Ayurveda, Siddha, and Unani, have been extensively studied for their applications in treating a wide range of diseases. As per the World Health Organization, approximately 70% of the world's population depends on plants for their primary health care. In the global market, more than 50 major drugs have been derived from tropical plants, underscoring the immense potential of plant-based medicine [3, 4] (Vaou *et al.*, 2021; Wu *et al.*, 2023). As human civilization progressed, the use of herbal plants continued to evolve and expand. The knowledge of the healing properties of these plants was passed down

from generation to generation, serving as a primitive form of health insurance. Over time, ancient records meticulously documented the various applications of herbal plants to treat diseases, creating a rich verbal materia medica [5] (Tyler and Tyler, 2023). With the advancement of modern science and technology, the focus shifted from the use of whole plant extracts to the isolation and synthesis of active components. Nevertheless, botanicals continue to serve as the foundation for many modern medical therapeutics, and the study of traditional plant-based medicines has become an area of growing interest and research [6] (Kim and Kang, 2023).

Saponins, a diverse class of naturally occurring compounds found in various medicinal plants, have garnered significant attention in the scientific community due to their remarkable pharmacological properties and potential therapeutic applications [7] (Bildziukevich *et al.*, 2023). One of the most intriguing aspects of saponins is their cytotoxic potential against cancer cells. Numerous studies have demonstrated that saponins can induce apoptosis in several cancer cells, making them attractive as potential anti-cancer agents. This is particularly significant, as eliminating tumor cells through apoptosis can help reduce the side effects associated with traditional cancer therapies [8] (Zhu *et al.*, 2023). In addition to their anti-cancer efficacy, saponins have also been shown to exhibit anti-inflammatory, antioxidant, and antimicrobial properties, further expanding their potential therapeutic applications. For instance, saponin-rich plant extracts have been utilized in conventional medicine for centuries to manage numerous diseases, including infections, metabolic disorders, and even cognitive impairment [9] (Zheng *et al.*, 2020).

Despite the growing interest in the therapeutic effects of saponins, further study is required to fully understand their roles, optimize their bioavailability, and develop effective delivery systems. As our understanding of these remarkable plant-derived compounds continues to expand, the future holds great interest to develop saponin-based therapeutics that can revolutionize the way we approach the treatment of human diseases [10] (Timilsena *et al.*, 2023). *Anoectochilus elatus* is a medicinal plant originating in India and China, which is well known for its therapeutic effects [11] (Sherif *et al.*, 2016). The therapeutic effects of *A. elatus* are harnessed from various parts, including leaves, stems, and roots [12, 13] (Mohapatra *et al.*, 2022; Wu *et al.*, 2020). Historically, it has been utilized to manage several ailments, including hepatic diseases, respiratory issues, and gastrointestinal illnesses. The herb has additionally been utilized to promote cardiovascular wellness and exhibit anticancer qualities [14] (Gantait *et al.*, 2021). Despite its several beneficial properties, this plant has not undergone scientific examination, necessitating a thorough research of its therapeutic properties. The existing

evidence regarding this plant is quite limited and insufficient. This work investigates the *in vitro* anti-oxidant, antimicrobial, and *in silico* analyses of *A. elatus* roots.

II. OBJECTIVES

The present work examines the *in vitro* antioxidant, antimicrobial, and *in silico* analysis of *Anoectochilus elatus* roots.

III. MATERIALS AND METHOD

1. Collection and identification of plant

The root parts of the plant *A. elatus* was collected from the top hill of Yercaud at Salem District, and authenticated from Plant Anatomy Research Centre (PARC), Chennai, Tamil Nadu. The authentication certificate number is No.PARC/2020/4319. Soon after the collection, the root was shade dried for 15 days to remove moisture content. Then coarsely powdered by mechanical blender and passed through sieve No.40. Powdered plant material was used for the extraction.

2. Extraction procedure

The shade dried coarsely powdered root of *A. elatus*(100g) was extracted separately with 500 ml of 50% water, methanol, chloroform, petroleum ether, benzene, and ethyl acetate. Initially, it was macerated at 37°C for 72 h. Then the plant material was extracted with specific solvents using Soxhlet apparatus. After extraction, the extract was filtered and stored at 4°C for the further investigations.

3. Extraction of crude saponins

Initially, the plant powder was defatted using petroleum ether and n-hexane. The defatted substance was subsequently extracted using methanol. The methanolic extract was concentrated via rotational evaporation under vacuum conditions. Then it was suspended in distilled water and agitated with n-butanol, leading to the precipitation of a crude saponin mixture with the addition of diethyl ether. A 100 g of the sample was placed into a conical flask, and 500 ml of 20% aqueous methanol is added. The solution is heated for 4 h with constant stirring at approximately 55°C. The mixture is filtered, and the residue is further extracted with 200 ml of 20% ethanol. The amalgamated extracts are concentrated to approximately 40 ml using a water bath at around 90°C. Subsequent to evaporation, the sample is subjected to drying in an oven until a constant weight is attained, and the saponin content is determined as a percentage.

4. Phytochemical analysis

The phytochemicals including, alkaloids, flavonoids, triterpenoids, carbohydrates, saponins, steroids, amino acids, tannins, gums & mucilages, and chlorogenic compounds, found in the water, methanol, chloroform, petroleum ether, benzene, and ethyl acetate extracts of the *A. elatus* roots were evaluated using the methods previously outlined [15,16] (Harbone, 1998; Kokate *et al.*, 1995). A quantitative phytochemical analysis was conducted on the root extract of *A. elatus* to quantify total carbohydrate, starch, reducing sugar, protein, amino acids, flavonoids, total phenols, steroids, tannins, saponins, and alkaloids, employing previously established methodologies [17] (Trease and Evans, 2002).

5. *In vitro* antioxidant analysis of crude saponins extracted from *A. elatus* roots

The effect of saponins from the *A. elatus* roots on the DPPH scavenging effect was assessed using earlier technique [18] Brand-Williams *et al.* (1995). The effect of saponins from the *A. elatus* roots on the hydrogen peroxide (H₂O₂) radical scavenging was studied by previous technique [19] (Sroka and Cisowski, 2003). The ABTS scavenging property of the saponins from *A. elatus* roots was scrutinized using the previously established technique [20] (Re *et al.*, 1999). The influence of saponins from *A. elatus* roots on the scavenging of NO radicals was studied using the earlier technique [21] (Chakraborty, 2009).

6. Analysis of antimicrobial activity

The well diffusion method was utilized to study the antibacterial effectiveness of the saponins extracted from *A. elatus* roots against *Staphylococcus aureus*, *Streptococcus pyogenes*, *Bacillus subtilis*, *Enterococcus faecalis*, *Micrococcus luteus*, *Escherichia coli*, *Salmonella typhi*, *Proteus vulgaris*, *Klebsiella pneumoniae*, *Vibrio parahaemolyticus*, and a fungi *Aspergillus niger* and *Candida albicans*. The present experiment utilized the appropriate agar growth media. Following the inoculation of the plates with the respective pathogens, saponins extracted from *A. elatus* roots at 25, 50, 75, and 100 mg/ml dosages were introduced into wells created with a cork borer on the agar plate surface and incubated at 37°C for 24 h. Following an incubation time, the diameter of the inhibitory zones was recorded. Amoxicillin served as a positive control in the trials.

7. *In silico* analysis to identify the protein components responsible for the therapeutic activities of *A. elatus*

A. Sequence retrieval and structure predictions:

The FASTA sequence of proteins was acquired from the GenBank database managed by the NCBI at <http://www.ncbi.nlm.nih.gov>. We determined the theoretical isoelectric point (pI), molecular weight, total counts of positive and negative residues, extinction coefficient, instability index, aliphatic index, and grand average of hydropathy (GRAVY) for physicochemical characterization utilizing the Expasy ProtParam server at <http://us.expasy.org/tools/protparam.html>. The Self Optimized Prediction Method with Alignment (SOPMA) was employed to identify the secondary structure. The SWISS MODEL was employed to forecast the tertiary structures.

B. Functional characterization

The protein was analyzed utilizing the SOSUI and TMHMM v.2.0 databases to evaluate its solubility or transmembrane characteristics. The InterPro database comprises a comprehensive compilation of protein families, domains, and functional sites. InterPro integrates the principal protein signature databases into a cohesive resource. These include: PROSITE, which employs regular expressions and profiles; PRINTS, which utilizes Position Specific Scoring Matrix-based (PSSM) fingerprints; ProDom, which implements automatic sequence clustering; and Pfam, SMART, TIGRFAMs, PIRSF, SUPERFAMILY, Gene3D, and PANTHER, all of which utilize hidden Markov models (HMMs). The superfamily and molecular function was detected utilizing the InterPro protein sequencing and categorization platform accessible at <http://www.ebi.ac.uk/interpro/>.

C. Sequence alignment

Pairwise sequence alignment was done utilizing the NCBI-BLAST program accessible at <http://blast.ncbi.nlm.nih.gov/Blast.cgi>. Multiple sequence alignment was executed utilizing the EBI-CLUSTAL OMEGA tool accessible at <http://www.ebi.ac.uk/Tools/msa/clustalo/>. Clustal Omega offers strong functionalities

for integrating sequences into and analyzing data within established alignments, utilizing the comprehensive updated information accessible in public databases like Pfam. The main aim of this study is to pinpoint regions of sequence similarity, hence facilitating the derivation of functional and evolutionary relationships among the proteins examined in this paper.

8. Phylogenetic analysis

A phylogenetic study was performed to detect the quantity of proteins exhibiting common structural and functional characteristics. All sequences in FASTA format were submitted to Clustal Omega using the default settings. A comprehensive analysis was performed on the output of fully aligned sequences, including scores, alignment, conserved residues, substitutions, and patterns of semi-conserved substitutions. A phylogenetic tree was constructed with the bootstrap Neighbour Joining (NJ) technique.

IV. RESULTS AND DISCUSSION

1. Phytochemical analyses of *A. elatus* root extracts

Medicinal plants have long been recognized for their therapeutic efficacy, serving as a rich source of phytochemicals. Preliminary phytochemical analysis plays a pivotal role in the exploration and characterization of these valuable plant constituents. Phytochemicals possess diverse biological activities and contribute to the salutary properties of plants. The identification and quantification of these secondary metabolites through preliminary screening can provide valuable insights into the pharmacological potential of a plant [22] (Kumar *et al.*, 2023).

The diverse solvent extracts of *A. elatus* roots, including water, methanol, chloroform, petroleum ether, benzene, and ethyl acetate, underwent preliminary phytochemical analyses. Among the evaluated solvents, methanol and aqueous extracts exhibited the occurrence of more number of phytochemicals relative to other solvent extracts. The methanol extract exhibited the highest positive results for the occurrence of phytochemicals, including alkaloids, flavonoids, triterpenoids, carbohydrates, saponins, steroids, amino acids, tannins, and chlorogenic compounds (Table 1).

Table 1: Preliminary phytochemical analysis of *A. elatus* root extracts

S. No	Phytochemicals	Aqueous	Methanol	Chloroform	Petroleum ether	Benzene	Ethyl acetate
1	Alkaloid	++	+++	-	+	-	-
2	Flavonoid	+++	+++	-	-	-	-
3	Triterpenoid	++	+++	+	++	+	+
4	Carbohydrates	+	+++	-	-	-	-
5	Saponin	+++	++	-	-	-	-
6	Steroids	++	+++	++	+	+	+
7	Amino acid	-	+	-	-	-	-
8	Tannin	++	+++	+	-	+	-
9	Gums & Mucilage	-	-	-	-	-	-
10	Chlorogenic compound	++	++	-	-	-	-

Table 2: Quantitative phytochemical analysis of *A. elatus* root extract

Biochemical constituents	Amount Present (mg/g tissue)
Total carbohydrate	195± 0.02
Starch	483± 0.05
Reducing sugar	21.89 ± 0.00
Protein	225 ± 0.01
Amino Acid	95 ± 0.04
Flavonoids	90 ± 0.16
Total Phenols	282 ± 0.10
Steroids	148 ± 0.19
Tannins	127± 0.15
Saponins	88 ± 0.11
Alkaloids	67 ± 0.09

The quantitative analysis results, subsequent to the qualitative analysis, revealed the presence of various secondary metabolites, including total carbohydrate, starch, reducing sugar, protein, amino acids, flavonoids, total phenols, steroids, tannins, saponins, and alkaloids in the *A. elatus* root (Table 2). The importance of preliminary phytochemical analysis lies in its ability to predict the therapeutic value of a medicinal plant. By detecting the presence of key phytochemicals, researchers can make informed decisions about the plant's potential applications in drug discovery and development. This information can guide further investigations, such as bioactivity assays, isolation of active compounds, and pharmacological evaluations [23] (Monjotin *et al.*, 2022).

Preliminary phytochemical screening is a vital tool in the field of pharmacognosy, the study of medicinal plants and their active constituents. It serves as a starting point for the systematic evaluation of medicinal plants, as it can help in the identification of lead compounds and the development of standardized herbal formulations [24] (Abdullah *et al.*, 2022). The present findings proved the presence of various secondary metabolites in the *A. elatus* root extracts.

2. Effect of saponins from the *A. elatus* roots on the *in vitro* free radical scavenging effects

The analysis of free radicals scavenging effect of plant extracts has become an increasingly important field of study due to the growing recognition of the role of oxidative stress in various disease states. Free radicals, such as ROS and RNS, are generated during normal cellular metabolism and can cause damage to cellular components, leading to a range of health issues, including atherosclerosis, aging, cancer, and cardiovascular diseases [25] (Szerlauth *et al.*, 2019). The major group of antioxidant phytochemicals present in plants have been found to play a pivotal role in adsorbing and neutralizing free radicals. The capacity of plants to scavenge free

radicals can be assessed using in vitro assays, which provide valuable insights into the antioxidant potential of these extracts [26] (Wu *et al.*, 2023).

Table 3: DPPH radical scavenging activity of saponin from *A. elatus* roots

Samples	Concentration ($\mu\text{g/ml}$)	Inhibition (%)
Saponin from <i>A. elatus</i> roots	100	95.2 $\pm$ 3.31
	80	82.5 $\pm$ 2.8
	60	68.0 $\pm$ 1.76
	40	46.9 $\pm$ 2.72
	20	24.7 $\pm$ 3.36
Rutin	100	98.8 $\pm$ 0.32
	80	87.5 $\pm$ 3.53
	60	73.1 $\pm$ 2.82
	40	57.0 $\pm$ 5.02
	20	35.6 $\pm$ 2.12

Table 4: H_2O_2 radical scavenging activity of saponin from *A. elatus* roots

Samples	Concentration ($\mu\text{g/ml}$)	Inhibition (%)
Saponin from <i>A. elatus</i> roots	100	76.3 $\pm$ 1.53
	80	55.8 $\pm$ 2.84
	60	36.1 $\pm$ 1.83
	40	22.0 $\pm$ 1.49
	20	10.9 $\pm$ 1.90
Rutin	100	86.5 $\pm$ 2.77
	80	71.7 $\pm$ 2.38
	60	57.4 $\pm$ 3.15
	40	45.2 $\pm$ 1.80
	20	30.0 $\pm$ 4.39

A series of free radical scavenging tests were done to assess the in vitro antioxidant effects of saponins from the *A. elatus* roots, with the findings illustrated in Tables 3-8. The treatment with saponins from the *A. elatus* roots at varying dosages (20-100 $\mu\text{g/ml}$) significantly reduced the free radical levels. The saponins from the *A. elatus* roots at 20 to 100 $\mu\text{g/ml}$ dosages effectively inhibited the levels of several free radicals, including DPPH, H_2O_2 , ABTS and NO, in dose-dependently. The ability of saponins from the *A. elatus* roots to neutralize several

free radicals in vitro was demonstrated by a reduction in DPPH, H₂O₂, ABTS, LPO, and NO radicals after exposure to increased quantities of the saponins from the *A. elatus* roots (Tables 3-8). These data further demonstrate that saponins from the *A. elatus* roots exhibits exceptional antioxidant properties.

Table 5: NO radical scavenging activity of saponin from *A. elatus* roots

Samples	Concentration (µg/ml)	Inhibition (%)
Saponin from <i>A. elatus</i> roots	100	69.1±3.51
	80	46.3.815±
	60	34.7±3.92
	40	20.2±1.93
	20	14.6±1.21
Rutin	100	80.7±4.85
	80	66.4±3.10
	60	57.8±4.61
	40	43.0±2.93
	20	27.5±1.90

Table 6: ABTS radical scavenging activity of saponin from *A. elatus* roots

Samples	Concentration (µg/ml)	Inhibition (%)
Saponin from <i>A. elatus</i> roots	100	91.0±2.35
	80	78.7±3.38
	60	70.5±1.80
	40	53.8±2.82
	20	37.5±1.38
Rutin	100	95.7±1.34
	80	83.9±3.04
	60	65.6±2.78
	40	51.1±4.01
	20	45.4±1.80

The DPPH assay is extensively employed technique for evaluating the free radical scavenging effect of plant extracts. This method involves the reduction of the stable DPPH radical by antioxidants in the extract. Similarly, the H₂O₂ scavenging assay measures the capacity of the plant extract to neutralize H₂O₂, which is a reactive oxygen species. The ABTS assay, on the other hand, measures the capacity of the extract to neutralize the ABTS radical cation, while the NO scavenging assay evaluates the extract's capacity to scavenge NO [27]

(Hassanpour and Doroudi, 2023). It has been already demonstrated the free radical scavenging effect of various plant extracts [28] (Kolar *et al.*, 2018).

Table 7: LPO radical scavenging activity of saponin from *A. elatus* roots

Samples	Concentration ($\mu\text{g/ml}$)	Inhibition (%)
Saponin from <i>A. elatus</i> roots	100	83.3 $\pm$ 2.88
	80	70.5 $\pm$ 2.61
	60	55.1 $\pm$ 2.71
	40	33.0 $\pm$ 4.70
	20	17.9 $\pm$ 3.06
Rutin	100	94.0 $\pm$ 2.7
	80	81.9 $\pm$ 4.32
	60	60.8 $\pm$ 3.24
	40	48.3 $\pm$ 2.30
	20	41.2 $\pm$ 3.26

The present results also proved that *A. elatus* root extracts has the exceptional capacity in scavenging the various free radicals, including DPPH, H_2O_2 , ABTS, NO and LPO radicals.


3. Antimicrobial activity of the saponins from the *A. elatus* roots

The analysis of antimicrobial effect of plants is a crucial field of study with significant implications for various applications. Plant extracts have long been known for their therapeutic properties, including antimicrobial effects. The increasing prevalence of drug-resistant microbial strains has further highlighted the importance of exploring alternative antimicrobial sources, such as those found in medicinal plants [29] (Álvarez-Martínez *et al.*, 2021). The well diffusion method (Figure 1) was employed to study the antibacterial efficacy of the saponins from the *A. elatus* roots against several pathogens, including *S. aureus*, *S. pyogenes*, *B. subtilis*, *E. faecalis*, *M. luteus*, *E. coli*, *S. typhi*, *P. vulgaris*, *K. pneumoniae*, *V. parahaemolyticus*, and an fungi *A. niger* and *C. albicans*. The treatment of different doses of saponins from the *A. elatus* roots has demonstrated a significant impact in inhibiting the growth of these pathogens. The treatment of saponins from the *A. elatus* roots markedly inhibited the growth of various pathogens, especially *S. aureus*, *S. pyogenes*, *E. choli*, and *P. vulgaris* (Figure 1). The existence of increased inhibition zones around the wells treated with saponins from the *A. elatus* roots signifies the suppression of pathogen growth.

One of the primary applications of analyzing the antimicrobial effect of plant extracts is in the development of novel antimicrobial drugs. The vast diversity of plant-derived compounds, including secondary metabolites, has the capacity to provide a rich source of novel antimicrobial compounds. These compounds can be utilized to develop new antimicrobial formulations, potentially overcoming the limitations of conventional antibiotics and addressing the growing problem of antimicrobial resistance [30] (Fik-Jaskółka *et al.*, 2024). Additionally, plant extracts with demonstrated antimicrobial activity can be employed for several

applications. The use of plant-based antimicrobial agents can help reduce the dependence on synthetic medicines, which can have undesirable effects on human health. The analysis of antimicrobial activity of medicinal plant extracts is also crucial for traditional medicine systems, as it can provide scientific validation for the use of these plants to treat infectious diseases[31] (Ashraf *et al.*, 2023). The results of this work proved that the *A. elatus* root extracts effectively inhibited the growth of various pathogens, which proves its antimicrobial properties.

Fig 1: Antimicrobial activity of the saponins from the *A. elatus* roots



4. Results of *in silico* molecular docking analysis

Molecular docking is a powerful computational method that has gained significant attention in the field of drug development. This method involves predicting the preferred orientation of a molecule (known as a ligand) when it binds to a target protein, and evaluating the strength of this interaction. In the context of plant proteins, *in silico* molecular docking analysis has become an increasingly valuable tool, offering numerous applications and insights [32] (Deshpande *et al.*, 2023). The importance of *in silico* molecular docking analysis of plant proteins lies in its ability to provide a cost-effective and time-efficient approach to understanding protein-ligand interactions. This dry lab approach can help researchers identify potential lead compounds, screen large chemical libraries, and optimize the binding affinity and selectivity of drug candidates, ultimately accelerating the drug discovery process [33] (Hiremath *et al.*, 2021).

Table 8: Primary structures of Ribulose-1,5-bisphosphate carboxylase and Maturase K computed using expasy's


S.N O	Accessio n number	Protein	Length	Mol.Wt	PI	- R	+ R	EC	II	AI	GRAVY
1	A0A89 9JZY2	Ribulose- 1,5- bisphosphat e carboxylase	429	47460.93	6.57	50	47	61685	37.87	78.44	-0.281
2	A0A14 0CZA1	Maturase K	281	33758.81	9.87	15	38	53860	38.28	100.50	0.065

Table 9: Secondary structures of Ribulose-1,5-bisphosphate carboxylase and Maturase K computed using expasy's

S. No	secondary structures	PROTEIN	
		Ribulose-1,5-bisphosphate carboxylase	Maturase K
1	Alpha helix	38.93%	43.42%
2	Extended Strand	17.40%	19.93%
3	Beta turn	6.76%	3.20%
4	Random coil	36.83%	33.45%

The primary structural prediction was conducted utilizing the ProtParam program (Tables 9 and 10). The molecular weights of several proteins were determined by calculating the markers using Expasy's ProtParam program. The molecular weight of Maturase K and Ribulose-1,5-bisphosphate carboxylase was found to be 47460.93 and 33758.81, respectively (Table 9). The Ribulose-1,5-bisphosphate carboxylase protein exhibited a pI value of 6.57, signifying acidity, whereas Maturase K demonstrated a pI value over 9.87, suggesting basicity. The proteins are noted to be densely packed and stable at their isoelectric points. The aliphatic index (AI) values for the both proteins ranged from 78.44 to 100.50 (Table 9). The range of GRAVY of *A. elatus* proteins was found to be -0.281 and 0.065. The secondary structure prediction of Maturase K and Ribulose-1,5-bisphosphate carboxylase was studied by SOPMA that exhibited that alpha helix, extended strand, beta turn and random coil, were more predominant. The alpha helix percentage for Maturase K and Ribulose-1,5-bisphosphate carboxylase was found to be 43.42% and 38.93%, respectively. The random coil percentage of both proteins were found to be 36.83% and 33.45%, respectively (Table 10). The calculated extinction coefficients facilitate the quantitative assessment of protein–protein and protein–ligand interactions (Figure 2).

Figure 2: Tertiary structures of Ribulosebiphosphate carboxylase large chain and Maturase K proteins




One of the primary applications of *in silico* molecular docking in the plant sciences is the identification of bioactive compounds from natural sources. Plant-derived compounds have long been recognized as a rich source of therapeutic candidates, and molecular docking can be used to screen these compounds against target proteins of interest. This approach can help researchers prioritize the most talented agents for additional studies, reducing the need for expensive and time-consuming studies [34] (Hazra *et al.*, 2023). Additionally, molecular docking can be used to study the structural determinants necessary for efficient ligand-receptor binding, providing valuable insights into the mechanisms underlying the biological activities of plant-derived compounds. Another application of *in silico* molecular docking in the plant sciences is the rational design of enzyme inhibitors. Many plant proteins, such as those involved in secondary metabolism, are of great interest for their potential therapeutic applications. Molecular docking can be used to identify and optimize small-molecule inhibitors that can modulate the activity of these enzymes, paving the way to develop novel plant-based drugs and agrochemicals [35] (Imran Tousif *et al.*, 2022).

A. Results of Ramachandran plot analysis

The study of medicinal plant extracts has gained more interest in recent times, as researchers strive to uncover the pharmacological capacity of these natural resources. One crucial tool in this endeavor is the Ramachandran plot, a powerful analytical technique that provides invaluable insights into the structural and functional properties of these extracts. The Ramachandran plot is a graphical representation of the backbone dihedral angles, ϕ and ψ , in protein structures. While primarily used in the area of protein structural analysis, this technique has also found applications in the analysis of medicinal plant extracts. By examining the Ramachandran plot of the phytochemicals present in these extracts, researchers can gain a deeper understanding of their structural

characteristics, which in turn, can shed light on their potential bioactivities and pharmacological properties [36] (Dutta *et al.*, 2023).

Figure 3: Ramachandran plot analysis of Ribulose biphosphate carboxylase large chain and Maturase K proteins



The secondary structures of polypeptides in proteins are detected by hydrogen bonds between the negatively charged carbonyl oxygen atoms and the positively charged amide hydrogen atoms in the molecular backbone. The angle ψ can be modified within the range of -180° to 180° , which translates to 360° of rotation for each angle. Nevertheless, numerous combinations of these angles are hardly encountered, while others are exceedingly common in proteins. The values of ψ in relation to the values of ϕ for a globular protein was depicted in Figure 3. The objective of this work is to obtain a dataset with the spatial coordinates of each atom. This data can be acquired from many protein structural data sites. We will utilize XRD data, as it provides the utmost precision. Despite its lack of perfect accuracy, crystal packing pressures often induce tiny aberrations in proteins. Molecular magnetic resonance (NMR) data for proteins in solution lack precision but are thought to be more accurate due to the protein being in its natural environment. This data may be analyzed to determine the amino acid residues, as the XRD ".PDB" data format categorizes all atoms and assigns them to specific residues virtually automatically. Subsequently, we employ a computer to compute the dihedral angles that determine the values of ψ and ϕ . A software application may examine a ".PDB" file and deliver the recorded values of ψ and ϕ angles for each specific residue. Initially, we shall plot these values for the yeast protein hexokinase. The plot of ψ against ϕ is referred to as a Ramachandran plot (Figure 3).

The growing importance of Ramachandran plot analysis in the field of medicinal plant research is evident from the recent advancements in analytical techniques and methods. The challenges posed by the complexity of

medicinal plant matrices and the variability of their phytochemical compositions have been largely overcome through the application of modern qualitative and quantitative techniques, such as liquid chromatography, countercurrent chromatography, and ultrasound-assisted extraction [37] (Bhat *et al.*, 2024). These analytical tools have enabled researchers to identify, characterize, and quantify the bioactive compounds present in medicinal plant extracts, paving the way for a more comprehensive understanding of their structural and functional properties. The structural and functional analysis of plant extracts is crucial for correlating the type and amount of phytochemicals present with their bioactivities, which in turn, can guide the selection of specific applications for these natural products. The Ramachandran plot analysis has emerged as a valuable tool in this regard, providing insights into the conformational preferences and spatial arrangements of the phytochemicals, which can be directly linked to their biological activities [38] (Islam *et al.*, 2023).

5. Phylogenetic analysis of *A. elatus*

Phylogenetic analysis, the study of the evolutionary relationships between organisms, has become an essential tool in the area of medicinal plant research. The economic and pharmaceutical importance of herbal plants has led to their increased exploitation, often resulting in the depletion of their natural habitats. To address this issue, phylogenetic analysis has emerged as a valuable technique for identifying closely related species, understanding their evolutionary history, and developing effective conservation strategies [39] (Youssef *et al.*, 2023). The current findings indicate that the plant species exhibiting a similarity of 90% or greater with the Ribulose-1,5-bisphosphate carboxylase include *Phalaenopsis aphrodite* (Orchidaceae) at 97.2%, *Brexia madagascariensis* (Celastraceae) at 96.2%, *Platanus occidentalis* (Platanaceae) at 96.5%, *Nypa fruticans* (Arecaceae) at 96.7%, *Liquidambar styraciflua* (Altingiaceae) at 96.0%, *Alnus incana* (Betulaceae) at 96.3%, *Tasmannia insipida* (Winteraceae) at 96.8%, *Oryza meridionalis* (Oryzeae) at 97.5%, *Oenothera argillicola* (Onagraceae) at 95.8%, and *Acer yangbiense* (Sapindaceae) at 95.6% (Table 10 and Figure 4). The phylogenetic analysis identified a list of plant species with a similarity of 90% or greater to the Maturase K including, *Arpophyllum giganteum* (Orchidaceae) at 83.4%, *Musa balbisiana* (Musaceae) at 70.5%, *Leuco jumaestivum* (Amaryllidaceae) at 71.1%, *Lilium henryi* (Liliaceae) at 68.1%, *Bowiea volubilis* (Hyacinthaceae) at 69.1%, *Acorus americanus* (Acoraceae) at 65.7%, *Buxus microphylla* (Buxaceae) at 66.2%, *Amorphophallus abyssinicus* (Araceae) at 66.3%, *Nicotiana tomentosiformis* (Solanaceae) at 64.5%, *Gossypium hirsutum* (Malvaceae) at 66.9% (Table 11 and Figure 4).

Table 10: Lists of plant species showing similarity of 95% and above with the Ribulosebiphosphate carboxylase large chain


S.No	Plant species containing Ribulosebiphosphate carboxylase large chain	Family name	Accession number	Identity (%)
1.	<i>Phalaenopsis aphrodite</i> subsp. <i>formosana</i> (Moth orchid)	Orchidaceae	Q3BAN4	97.2
2.	<i>Brexia madagascariensis</i>	Celastraceae	Q31738	96.2
3.	<i>Platanus occidentalis</i> (Sycamore) (American plane tree)	Platanaceae	P28441	96.5
4.	<i>Nypa fruticans</i> (Nypa palm)	Arecaceae	P28261	96.7
5.	<i>Liquidambar styraciflua</i> (Sweetgum tree) (<i>Liquidambar macrophylla</i>)	Altingiaceae	Q01873	96.0
6.	<i>Alnus incana</i> (White alder)	Betulaceae	Q06021	96.3
7.	<i>Tasmannia insipida</i> (Pepperbush) (<i>Drimys insipida</i>)	Winteraceae	P28456	96.8
8.	<i>Oryza meridionalis</i>	Oryzeae	H6CSD3	97.5
9.	<i>Oenothera argillicola</i> (Appalachian evening primrose)	Onagraceae	B0Z4K9	95.8
10.	<i>Acer yangbiense</i>	Sapindaceae	A0A5C7GPV1	95.6

Phylogenetic analysis has played an essential role in the identification and characterization of medicinal plants, particularly those with known or potential therapeutic properties. By mapping the evolutionary relationships between different plant species, researchers can identify closely related taxa that may share similar chemical profiles and pharmacological activities. This knowledge can guide the discovery of new drug candidates, as well as the development of alternative sources for existing medicinal compounds. Furthermore, phylogenetic analysis has been instrumental in the identification and preservation of endangered medicinal plant species. By understanding the evolutionary relationships and genetic diversity within a plant group, conservation efforts can be tailored to target specific taxa or populations that are at risk of extinction [40] (Bravo-Vázquez *et al.*, 2024).

Table 11: Lists of plant species showing similarity of 95% and above with the Maturase K

S. No	Plant species containing Maturase K	Family name	Accession number	Identity (%)
1.	<i>Arpophyllum giganteum</i> (Hyacinth orchid)	Orchidaceae	Q8WJR9	83.4
2.	<i>Musa balbisiana</i> (Banana)	Musaceae	A0A4V4H224	70.5
3.	<i>Leuco jumaestivum</i> (Summer snowflake)	Amaryllidaceae	Q8GVB5	71.1
4.	<i>Lilium henryi</i> (Henry's lily)	Liliaceae	Q9GIH9	68.1
5.	<i>Bowiea volubilis</i> (Climbing onion) (<i>Ophiobos tryxvolubilis</i>)	Hyacinthaceae	Q6LA17	69.1
6.	<i>Acorus americanus</i> (Sweetflag) (<i>Acorus calamus</i> var. <i>americanus</i>)	Acoraceae	A9LYG6	65.7
7.	<i>Buxus microphylla</i> (Littleleaf boxwood) (Japanese boxwood)	Buxaceae	A6MM17	66.2
8.	<i>Amorphophallus abyssinicus</i> (Black arum) (<i>Arum abyssinicum</i>)	Araceae	Q8MEH7	66.3
9.	<i>Nicotiana tomentosiformis</i> (Tobacco)	Solanaceae	Q70D40	64.5
10.	<i>Gossypium hirsutum</i> (Upland cotton) (<i>Gossypium mexicanum</i>)	Malvaceae	H9LI95	66.9

Figure 4: Phylogenetic analysis of plant species with Ribulose biphosphate carboxylase large chain and Maturase K proteins



V. CONCLUSIONS

The present work has demonstrated the phytochemical profile, antioxidant, and antimicrobial properties of saponins derived from *A. elatus* roots. The current results from the *in silico* studies have demonstrated the crucial function of Maturase K and Ribulose-1, 5-bisphosphate carboxylase proteins in the therapeutic effects of the *A. elatus* roots. Moreover, further studies are necessary to understand the phytochemical profile and precise underlying molecular mechanisms by which *A. elatus* exhibits therapeutic properties.

REFERENCES

- 1) Salmerón-Manzano E, Garrido-Cardenas JA, Manzano-Agugliaro F. Worldwide Research Trends on Medicinal Plants. *Int J Environ Res Public Health*. 2020 May 12; 17(10):3376.
- 2) Riaz M, Khalid R, Afzal M, Anjum F, Fatima H, Zia S, Rasool G, Egbuna C, Mtewa AG, Uche CZ, Aslam MA. Phyto bioactive compounds as therapeutic agents for human diseases: A review. *Food Sci Nutr*. 2023 Apr 17; 11(6):2500-2529.
- 3) Vaou N, Stavropoulou E, Voidarou C, Tsigalou C, Bezirtzoglou E. Towards Advances in Medicinal Plant Antimicrobial Activity: A Review Study on Challenges and Future Perspectives. *Microorganisms*. 2021 Sep 27; 9(10):2041.
- 4) Wu P, Wu W, Zhang S, Han J, Liu C, Yu H, Chen X, Chen X. Therapeutic potential and pharmacological significance of extracellular vesicles derived from traditional medicinal plants. *Front Pharmacol*. 2023 Dec 1; 14: 1272241.
- 5) Tyler SEB, Tyler LDK. Pathways to healing: Plants with therapeutic potential for neurodegenerative diseases. *IBRO Neurosci Rep*. 2023 Feb 10; 14: 210-234.
- 6) Kim YJ, Kang KS. The Phytochemical Constituents of Medicinal Plants for the Treatment of Chronic Inflammation. *Biomolecules*. 2023 Jul 25; 13(8):1162.
- 7) Bildziukevich U, Wimmerová M, Wimmer Z. Saponins of Selected Triterpenoids as Potential Therapeutic Agents: A Review. *Pharmaceuticals (Basel)*. 2023 Mar 2; 16(3):386.
- 8) Zhu M, Sun Y, Bai H, Wang Y, Yang B, Wang Q, Kuang H. Effects of saponins from Chinese herbal medicines on signal transduction pathways in cancer: A review. *Front Pharmacol*. 2023 Mar 29; 14: 1159985.
- 9) Zheng Y, Lee EH, Lee JH, In G, Kim J, Lee MH, Lee OH, Kang JJ. Preclinical Research on a Mixture of Red Ginseng and Licorice Extracts in the Treatment and Prevention of Obesity. *Nutrients*. 2020 Sep 9; 12(9):2744.
- 10) Timilsena YP, Phosanam A, Stockmann R. Perspectives on Saponins: Food Functionality and Applications. *Int J Mol Sci*. 2023 Aug 31; 24(17):13538.
- 11) Sherif, N.A.; Kumar, T.S.; Rao, M.V. In vitro regeneration by callus culture of *Anoectochilus elatus* Lindley, an endangered terrestrial jewel orchid. *Vitr. Cell. Dev. Biol*. 2016, 52, 72–80.
- 12) Mohapatra, P.; Ray, A.; Jena, S. Evaluation of Genetic Stability of In Vitro Raised Orchids Using Molecular-Based Markers. In *Commercial Scale Tissue Culture for Horticulture and Plantation Crops*; Springer: Singapore, 2022; Volume 17, pp. 293–316.
- 13) Wu, Y.B.; Peng, M.C.; Zhang, C.; Wu, J.G.; Ye, B.Z.; Yi, J.; Wu, J.Z.; Zheng, C.J. Quantitative determination of multi-class bioactive constituents for quality assessment of ten *Anoectochilus*, four *Goodyera* and one *Ludisia* species in China. *Chin. Herb. Med*. 2020, 12, 430–439.
- 14) Gantait, S.; Das, A.; Mitra, M.; Chen, J.T. Secondary metabolites in orchids: Biosynthesis, medicinal uses, and biotechnology. *S. Afr. J. Bot*. 2021, 139, 338–351.
- 15) Harborne J. B. *Phytochemical Methods A Guide To Modern Techniques of Plant Analysis*. 3rd. London, UK: Chapman and Hall; 1998.
- 16) Kokate CK, Khandelwal KR, Pawar AP, Gohalz SB. *Practical Pharmacognosy*. 4th ed. New Delhi, India: Vallabh Prakashan; 1995. p. 107.
- 17) Trease G, Evans SM. *Pharmacognosy*. 15th ed. London: Bailer Tindal; 2002. pp. 23–67.
- 18) Brand-Williams, W.; Cuvelier, M.; Berset, C. Use of free radical method to evaluate antioxidant activity. *LWT-Food Sci. Technol*. 1995, 28, 25–30.
- 19) Sroka Z, Cisowski W. Hydrogen peroxide scavenging, antioxidant and anti-radical activity of some phenolic acids. *Food Chem Toxicol*. 2003 Jun; 41(6):753-8.
- 20) Re R, Pellegrini N, Proteggente A, Pannala A, Yang M, Rice-Evans C. Antioxidant activity applying an improved ABTS radical cation decolorization assay. *Free Radic Biol Med*. 1999 May; 26(9-10):1231-7.
- 21) Chakraborty GS. Free radical scavenging activity of *Costus speciosus* leaves. *Indian J Pharm Educ Res*. 2009; 43: 96–8.

- 22) Kumar A, P N, Kumar M, Jose A, Tomer V, Oz E, Proestos C, Zeng M, Elobeid T, K S, Oz F. Major Phytochemicals: Recent Advances in Health Benefits and Extraction Method. *Molecules*. 2023 Jan 16; 28(2):887.
- 23) Monjotin N, Amiot MJ, Fleurentin J, Morel JM, Raynal S. Clinical Evidence of the Benefits of Phytonutrients in Human Healthcare. *Nutrients*. 2022 Apr 20; 14(9):1712.
- 24) Abdullah, Ahmad N, Tian W, Zengliu S, Zou Y, Farooq S, Huang Q, Xiao J. Recent advances in the extraction, chemical composition, therapeutic potential, and delivery of cardamom phytochemicals. *Front Nutr*. 2022 Sep 30; 9: 1024820.
- 25) Szerlauth A, Muráth S, Viski S, Szilagyi I. Radical scavenging activity of plant extracts from improved processing. *Heliyon*. 2019 Nov 14; 5(11):e02763.
- 26) Wu W, Li G, Zhou W, Wang E, Zhao X, Song X, Zhao Y. Comparison of Composition, Free-Radical-Scavenging Capacity, and Antibiosis of Fresh and Dry Leave Aqueous Extract from *Michelia shiluensis*. *Molecules*. 2023 Aug 8; 28(16):5935.
- 27) Hassanpour SH, Doroudi A. Review of the antioxidant potential of flavonoids as a subgroup of polyphenols and partial substitute for synthetic antioxidants. *Avicenna J Phytomed*. 2023 Jul-Aug; 13(4):354-376.
- 28) Kolar FR, Gogi CL, Khudavand MM, Choudhari MS, Patil SB. Phytochemical and antioxidant properties of some Cassia species. *Nat Prod Res*. 2018 Jun; 32(11):1324-1328.
- 29) Álvarez-Martínez FJ, Barrajón-Catalán E, Herranz-López M, Micol V. Antibacterial plant compounds, extracts and essential oils: An updated review on their effects and putative mechanisms of action. *Phytomedicine*. 2021 Sep; 90: 153626.
- 30) Fik-Jaskółka M, Mittova V, Motsonelidze C, Vakhania M, Vicidomini C, Roviello GN. Antimicrobial Metabolites of Caucasian Medicinal Plants as Alternatives to Antibiotics. *Antibiotics (Basel)*. 2024 May 24; 13(6):487.
- 31) Ashraf MV, Pant S, Khan MAH, Shah AA, Siddiqui S, Jeridi M, Alhamdi HWS, Ahmad S. Phytochemicals as Antimicrobials: Prospecting Himalayan Medicinal Plants as Source of Alternate Medicine to Combat Antimicrobial Resistance. *Pharmaceuticals (Basel)*. 2023 Jun 15; 16(6):881.
- 32) Deshpande SH, Muhsinah AB, Bagewadi ZK, Ankad GM, Mahnashi MH, Yaranguppi DA, Shaikh IA, Khan AA, Hegde HV, Roy S. In Silico Study on the Interactions, Molecular Docking, Dynamics and Simulation of Potential Compounds from *Withania somnifera* (L.) Dunal Root against Cancer by Targeting KAT6A. *Molecules*. 2023 Jan 22; 28(3):1117.
- 33) Hiremath S, Kumar HDV, Nandan M, Mantesh M, Shankarappa KS, Venkataravanappa V, Basha CRJ, Reddy CNL. In silico docking analysis revealed the potential of phytochemicals present in *Phyllanthus amarus* and *Andrographis paniculata*, used in Ayurveda medicine in inhibiting SARS-CoV-2. *3 Biotech*. 2021 Feb; 11(2):44.
- 34) Hazra S, Ray AS, Das S, Das Gupta A, Rahaman CH. Phytochemical Profiling, Biological Activities, and In Silico Molecular Docking Studies of *Causonis trifolia* (L.) Mabb. & J. Wen Shoot. *Plants (Basel)*. 2023 Mar 29; 12(7):1495.
- 35) Imran Tousif M, Nazir M, Saleem M, Tauseef S, Uddin R, Zengin G, Fawzi Mahomoodally M, Abbas Z. Total Bioactive Contents, Metabolic Profiling, Docking Studies, Antioxidant and Enzyme Inhibition Activities of *Convolvulus arvensis* L. and Multivariate Analysis to Unravel a Potential Herb as Natural Resource for Pharmaceutical Industry. *ChemBiodivers*. 2022 Nov; 19(11):e202200521.
- 36) Dutta AK, Gazi MS, Uddin SJ. A systemic review on medicinal plants and their bioactive constituents against avian influenza and further confirmation through in-silico analysis. *Heliyon*. 2023 Mar 10; 9(3):e14386.
- 37) Bhat S, Rather M, Gani S, Nabi A, Ganai SA, Shah MD, Sofi P, Jeelani F, Hussain A, Ashraf S, Anwar A, Iqbal I, Nisa TU, Summuna B, Banday S. Identification of plant based potential antifungal compounds against BMK-1 protein of *Bipolaris oryzae* using molecular docking approach. *Sci Rep*. 2024 Jul 8; 14(1):15665.
- 38) Islam MA, Mondal SK, Islam S, Akther Shorna MN, Biswas S, Uddin MS, Zaman S, Saleh MA. Antioxidant, Cytotoxicity, Antimicrobial Activity, and In Silico Analysis of the Methanolic Leaf and Flower Extracts of *Clitoria ternatea*. *Biochem Res Int*. 2023 Sep 22; 2023:8847876.
- 39) Youssef D, El-Bakatoushi R, Elframawy A, El-Sadek L, Badan GE. Molecular phylogenetic study of flavonoids in medicinal plants: a case study family Apiaceae. *J Plant Res*. 2023 May; 136(3):305-322.
- 40) Bravo-Vázquez LA, García-Ortega M, Medina-Feria S, Srivastava A, Paul S. Identification and expression profiling of microRNAs in leaf tissues of *Foeniculum vulgare* Mill. under salinity stress. *Plant Signal Behav*. 2024 Dec 31; 19(1):2361174.