PhytoKeys

PhytoKeys 256: 185–195 (2025)
DOI: 10.3897/phytokeys.256.149481

Research Article

Phylogenetic position, supplementary description and phytochemical analysis of *Camellia hekouensis* (Theaceae), a critically endangered tree native to Hekou, Yunnan, China

Dongwei Zhao[1], Guiliang Zhang[2], Shixiong Yang[3]

1 *Department of Forestry, College of Forestry, Central South University of Forestry and Technology, Changsha, Hunan 410004, China*
2 *Hekou Administration Branch of Dawei Mountain National Nature Reserve, Hekou, Yunnan 661399, China*
3 *CAS Key Laboratory for Plant Diversity and Biogeography of East Asia, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming, Yunnan 650201, China*
Corresponding author: Dongwei Zhao (zhaodw@csuft.edu.cn)

Abstract

Camellia harbors unique diversity along Sino-Vietnamese border. Some species of them are under threat due to human activity. *Camellia hekouensis*, a native of Hekou, Yunnan, China, was once considered extinct as the previously known "last living tree" died in 2024. Fortunately, 11 in-situ and 32 ex-situ trees have been protected and propagated by the staff of Hekou Administration Branch of Dawei Mountain National Nature Reserve in Yunnan with their great unpublicized efforts. Molecular phylogenetic analysis suggests that *C. hekouensis* is nested in the main clade CI of *Camellia* and forms a clade with *C. corallina*, *C. gracilipes* and *C. pubicosta*, which are generally distributed in Vietnam. Morphological characters of the capsule and seed of *C. hekouensis* are supplementally described. The leaves of *C. hekouensis* contain 1.18 mg/g theobromine, which disagrees with the previous chemotaxonomic claim. Though the economic and ecological values are little known for *C. hekouensis*, the species should be conserved and propagated effectively and promptly to prevent extinction.

Key words: Conservation, extremely small population, purine alkaloids, taxonomy

OPEN ACCESS

Academic editor: Avelinah Julius
Received: 12 February 2025
Accepted: 2 May 2025
Published: 29 May 2025

Citation: Zhao D, Zhang G, Yang S (2025) Phylogenetic position, supplementary description and phytochemical analysis of *Camellia hekouensis* (Theaceae), a critically endangered tree native to Hekou, Yunnan, China. PhytoKeys 256: 185–195. https://doi.org/10.3897/phytokeys.256.149481

Copyright: © Dongwei Zhao et al.
This is an open access article distributed under terms of the Creative Commons Attribution License (Attribution 4.0 International – CC BY 4.0).

Introduction

Camellia L. (Theaceae) is an economically valued genus that contains tea plants, oil camellia and camellias. The plants of *Camellia* are naturally distributed in East, Southeast and South Asia (Sealy 1958, Ming 2000). Species of the genus were counted as either 119 (Ming 2000) or 280 (Chang 1998). Since 2000, more than 100 species have been described (Zhao 2024a). As a shrub or tree bearing evergreen leaves, *Camellia* harbors great diversity in tropical China and Southeast Asia (Zhao et al. 2023), a global biodiversity hotspot attracting conservation interests (Meng and Song 2023).

Camellia hekouensis C.J.Wang & G.S.Fan was described based on a single specimen collected in Hekou, Yunnan, China, along the Sino-Vietnamese border (Wang and Fan 1988). A Latin diagnosis and description were provided in the protologue with a simplified interpretation of the morphological differences between

C. hekouensis and *C. longissima* Hung T.Chang & S.Ye Liang in Chinese, while the description of fruit and seed was absent. *Camellia hekouensis* was placed in *C.* sect. *Longissima* Hung T.Chang in the protologue, probably because the species bore a relatively long pedicel. *Camellia* sect. *Longissima* proposed in Chang (1981) includes only two species, viz. *C. gracilipes* Merr. ex Sealy and *C. longissima*. Chang (1998) subsequently confirmed that *C*. sect. *Longissima* included *C. hekouensis* and stated that both caffeine and theobromine were absent from the leaves of the species in the section. Ming (1999) recognized *C*. sect. *Longissima* as a heterotypic synonym of *C*. sect. *Longipedicellata* Hung T.Chang and transferred *C. hekouensis* to the latter with *C. gracilipes*, *C. laotica* (Gagnep.) T.L.Ming, *C. longipedicellata* (Hu) Hung T.Chang & D.Fang and *C. longissima*. Successive gatherings of *C. hekouensis* have been collected from the same locality as the type specimen (Ming 2000), and the morphology of fruit and seed of the species remains unclear (Ming and Bartholomew 2007). Gao et al. (2005: 164) listed an account and a photo of *C. hekouensis* that was introduced to the International *Camellia* Species Garden at Jinhua, Zhejiang, China; the photo is, however, actually a plant of *C.* sect. *Thea*. (L.) Griff., probably *C. sinensis* (L.) Kuntze var. *assamica* (Royle ex Hook.) Steenis, rather than *C. hekouensis*. Zhang et al. (2016: 255–256) provided ten photos of living plants of *C. hekouensis*, which included three of the fruit and seed. Unfortunately, the rulers or scales are absent in the photos, and a detailed lingual description of the fruit and seed is awaiting addition.

Fang et al. (2010) explored the phylogenetic relationship of *C*. sections *Chrysantha* Hung T.Chang, *Longipedicellata* and *Longissima* using four plastid DNA regions for *C. hekouensis* and another 27 species of *Camellia*. Their analysis suggested that *C. hekouensis* was sister to all other samples of *Camellia*, including *C. longipedicellata* and *C. longissima*, which implies that neither *C*. sect. *Longipedicellata* nor *C.* sect. *Longissima* is a monophyletic group and *C. hekouensis* occupies a unique but less resolved phylogenetic position in *Camellia*. Pang et al. (2022) suggested that *C. hekouensis* was nesting with species of *C*. sect. *Thea* based on phylogenetic analysis using four plastid DNA regions. Their samples were all collected from the International *Camellia* Species Garden at Jinhua, Zhejiang, China, and the sample of *C. hekouensis* was collected on 28 December 2006. However, the collection of *C. hekouensis* used in the analysis of Pang et al. (2022) was probably misidentified as discussed above for the species shown in Gao et al. (2005: 164), and the first author visited the garden in March 2018 but did not find any living or processed sample of *C. hekouensis* there. Thanks to the poor availability of the DNA samples, *C. hekouensis* is generally absent from previous phylogenetic analyses based on nuclear DNA (Vijayan et al. 2009, Yan et al. 2021, Cheng et al. 2022, Zan et al. 2023, Zhao et al. 2023).

A tree previously known as the "last living tree" of *C. hekouensis* was found to have died at the end of 2024 due to the destruction of its bark (Fig. 1A, B). Fortunately, the staff of Hekou Administration Branch of Dawei Mountain National Nature Reserve (hereafter the Reserve) in Yunnan have made great unpublicized efforts to propagate the plants and currently maintain dozens of living plants of the species. A complete morphological description, including those of fruit and seed, is provided here for the species. The phylogenetic position of *C. hekouensis* in the genus is discussed and the chemical contents of its leaves are analyzed to uncover its potential value for horticulture before it is too late to conserve this unique species of *Camellia*.




Figure 1. *Camellia hekouensis* **A** habit of the previously misidentified "last living tree" **B** the tree was found to have died at the end of 2024 **C** branches and leaves **D** adaxial surface of a leaf **E** flowers **F** a dissected flower 1 bracteoles 2 sepals 3 petals 4 androecium 5 pedicel, receptacle and gynoecium **G** dry ovary **H** capsule **I** a dissected capsule with a single seed. Photos: S.X. Yang (**A, B, F, G**); D.W. Zhao (**C, D**); G.L. Zhang (**E, H, I**). Scale bars: 5 cm (**D**); 2 cm (**E**, **F, H**); 1 mm (**G**); 1 cm (**I**).

Material and methods

Fieldworks were undertaken to search the living individuals of *C. hekouensis* in Hekou and neighboring counties of Yunnan. Photos of the habitat and fresh characters of the vegetative and propagative organs were taken.

Clean and young leaves were collected and dried in silica gel, then preserved at −30 °C for total DNA extraction and chemical analysis. Voucher specimens were deposited at the herbaria shown in Suppl. material 1. Morphological characters were described based on living plants, their images or herbarium specimens conserved in CSFI, KUN, SWFC and SYS (acronyms following Thiers 2025, continuously updated).

Total genomic DNA extraction and subsequently molecular phylogenetic analyses using three nuclear regions, including *RPB2* introns 11–15 and 23 and *waxy*, were generally stated in Zhao et al. (2023), except the GTRGAMMA model was used for bootstrapping in the maximum likelihood analysis and each metropolis-coupled Markov chain Monte Carlo analysis was run for ten million generations in the Bayesian inference. The GenBank accession numbers are listed in Suppl. material 1.

The contents of total polyphenols and catechins of dry samples, including catechin (C), epicatechin (EC), epicatechin gallate (ECG), epigallocatechin (EGC), epigallocatechin gallate (EGCG), and gallic acid (GA), were detected based on the National Standard of the People's Republic of China, Determination of total polyphenols and catechins content in tea (GB/T 8313–2018). The contents of four purine alkaloids, including caffeine, theobromine, theophylline, and theacrine, were analyzed using high-performance liquid chromatography (HPLC) methods stated in Meng et al. (2018). Each chemical was analyzed with two replications and the mean value was calculated as the final content.

Results

Living trees of *C. hekouensis*

Eleven trees of *C. hekouensis* have been found in the natural forests in Hekou, Yunnan, China, including three adult trees that can bear flowers and occur dispersedly and individually, accompanied by eight juvenile trees. Meanwhile, the staff of the Reserve have successfully propagated 32 juvenile trees by seeds. Detailed localities of the living trees are unrevealed here for conservation reasons.

Phylogenetic position

Nuclear DNA of *RPB2* (introns 11–15 and 23) and *waxy* regions of a single sample of *C. hekouensis* and one of *C. pubicosta* Merr. were sequenced in this study (Suppl. material 1). These, and another 37 samples reported in Zhao et al. (2023), were employed to reconstruct a phylogenetic tree of *Camellia* provided in Fig. 2. It suggests that *C. hekouensis* is nested in a monophyletic group with *C. corallina* (Gagnep.) Sealy, *C. gracilipes* and *C. pubicosta* with strong support (Bayesian posterior probability [PP] = 1, Bootstrap [BS, %] = 100). They are nested in the main clade CI of *Camellia*. *Camellia hekouensis* is sister to *C. gracilipes* and *C. pubicosta* (PP = 1, BS = 94), which suggests a closely phylogenetic relationship between them. Both *C. longissima*, the type of *C.* sect. *Longissima*, and *C. longipedicellata*, the type of *C.* sect. *Longipedicellata*, are nested in the main clade CII of *Camellia*, which have a relatively remote relationship with *C. hekouensis* (Fig. 2).




Figure 2. The Bayesian consensus tree reconstructed by the concatenated *RPB2* (introns 11–15 and 23) and *waxy* data for *Camellia hekouensis* and other representative species of *Camellia*. Bayesian posterior probabilities (PP) ≥ 0.5 and Bootstrap values (BS; %) ≥ 50 are presented above or below branches as PP/BS. Branch lengths are proportional to the expected nucleotide substitutions per site. Numbers in bold indicate the samples were sequenced here.

Chemical contents

The contents of catechins (C, EC, ECG, EGC, EGCG), gallic acid (GA), total polyphenols and four purine alkaloids of *C. hekouensis* are listed in Table 1. *Camellia hekouensis* has 5.13 mg/g catechins and 124.78 mg/g total polyphenols. Caffeine has not been detected, and theobromine occupies 1.18 mg/g in the dry leaves of *C. hekouensis*.

Description

***Camellia hekouensis* C.J.Wang & G.S.Fan, Acta Bot. Yunnan. 10(3): 365. 1988.**

Type material. *Holotype*: China • Yunnan: Hekou, 450 m, 7 November 1986, *C.J. Wang et al. 860235* (SWFC!, Fig. 3; isotype: SWFC!).

Description. Trees or shrubs up to 7 m tall, evergreen. ***Trunk*** brownish grey, ***new branchlets*** and ***terminal buds*** glabrous. ***Petioles*** 5–15 mm long, glabrous;

Table 1. Chemical contents of the leaves of *Camellia hekouensis*.

Chemical compounds	content (mg/g, ± standard deviation)
Total polyphenols	124.78 ± 3.13
Catechin (C)	2.24 ± 0.39
Epicatechin (EC)	1.84 ± 0.26
Epicatechin gallate (ECG)	0.29 ± 0.00
Epigallocatechin (EGC)	0.64 ± 0.05
Epigallocatechin gallate (EGCG)	0.12 ± 0.01
Total catechins	5.13 ± 0.69
Gallic acid (GA)	0.45 ± 0.02
Caffeine	ND
Theacrine	0.28 ± 0.03
Theobromine	1.18 ± 0.05
Theophylline	0.05 ± 0.00

Note. ND: not detected.

leaf blades elliptic, obovate to oblong, 10–22 × 3–8 cm, thinly coriaceous, abaxially yellowish green, adaxially dark green, shiny, glabrous on both surfaces, midrib and secondary veins abaxially elevated and adaxially impressed, secondary veins 10–17 on each side of midrib, base cuneate to obtuse, margin serrulate, apex acuminate to caudate. ***Flowers*** axillary, fragrant, solitary or up to 3 in a cluster, ca. 3 cm in diam. ***Pedicels*** 15–25 mm long. ***Bracteoles*** 3–4, alternate, persistent, deltate to ovate, 1–1.5 × 1.5–2 mm, glabrous on both surfaces. ***Sepals*** 5, deltate to sub-orbicular, 2.5–4 × 2–3 mm, glabrous on both surfaces, margin membranous and ciliolate. ***Petals*** 7–9 in 2 whorls, white, ovate, elliptic or obovate, 6–12 × 9–15 mm, glabrous on both surfaces, basally connate for 1–2 mm. ***Stamens*** 5–11 mm long; filaments light yellow, glabrous, basally connate for ca. 1 mm and outer filaments basally adnate to petals for 1–2 mm. ***Ovary*** ovoid, densely pubescent. ***Styles*** 3, distinct, 6–9 mm long, basally sparsely pubescent and gradually becoming glabrous apically. ***Capsule*** globose, 3–4 cm in diam., 1-loculed with 1 seed; pericarp 2–6 mm thick. ***Seeds*** fuscous, globose, 2–2.5 cm in diam., glabrous. Fig. 1.

Phenology. Flowering December, fruiting August.

Distribution and habitat. *Camellia hekouensis* is native to Hekou, Yunnan, China and occurs in the tropical evergreen forests at elevations of 290–800 m.

Additional specimens examined. Yunnan: • Hekou, Nanxi, 360–410 m, 21 December 1986, *C.J. Wang & L.S. Xie 904* (KUN 694698); • Hekou, Nanxi, 22.68°N, 103.93°E, 297 m, 27 November 2023, *D.W. Zhao et al. 562* (CSFI, equals to *S.X. Yang et al. 7352* at KUN); • same locality, 26 December 2023, *S.X. Yang et al. 7360* (KUN).

Discussion

Taxonomic significance

A relatively long pedicel is valued in the taxonomies of Chang (1981, 1998) and Ming (2000, Ming and Bartholomew 2007). Zhao et al. (2023) suggested that *C. longissima* formed a clade with taxa of *C.* sect. *Thea* and subsequently, Zhao (2024b) treated the former as a new member of the latter. Accordingly, *C.* sect. *Longissima* became a heterotypic synonym of *C.* sect. *Thea*. Neither Chang's



Figure 3. Holotype of *Camellia hekouensis* at SWFC. Photo: D.W. Zhao.

(1981, 1998) nor Ming's (2000, Ming and Bartholomew 2007) *C*. sect. *Longipedicellata* was supported by molecular phylogenetic analyses (Fang et al. 2010, Zan et al. 2023, Zhao et al. 2023). Our phylogenetic analysis suggests that *C. hekouensis* is nested in the main clade CI of *Camellia* (Fig. 2), which mainly consists of species native to Vietnam and splits with other camellias in the early evolutionary history of the genus (Zhao et al. 2023). *Camellia hekouensis* may be placed in the clade Corallina suggested in Zhao et al. (2023) provisionally. The clade contains *C. corallina*, *C. gracilipes*, *C. hekouensis* and *C. pubicosta* as shown in Fig. 2. The latter three species are morphologically closely related and can be easily distinguished from *C. corallina* by the morphology of leaves and flowers (Ming 2000). We propose a comprehensive taxonomic revision to evaluate whether the clade Corallina should be considered to be a section in *Camellia* as a whole or may be divided into two different sections.

Chang (1998) suggested that both caffeine and theobromine were absent from *C. hekouensis* and *C. longissima*. The single sample of *C. hekouensis* is caffeine-free, but contains 1.18 mg/g theobromine (Table 1), which disagrees with the claim of Chang (1998). Chang valued the variation of chemical contents, especially purine alkaloids, in the taxonomy of *Camellia* as insisting that *C. irrawadiensis* Barua was a distinct species from *C. taliensis* (W.W.Sm.) Melch. because the former was caffeine-free (Chang et al. 1996). By contrast, Ming (2000) and Zhao (2024a, 2024b) treated the former as a heterotypic synonym of the latter. Chemical contents may be nutrition-dependent and usually vary between individuals and populations within a single species of *Camellia* (Ye et al. 1997, Zhao 2024a). They are useful to act as essential traits to select the characteristic germplasms of tea plants; great caution is, however, needed when using them overwhelmingly as the taxonomically diagnostic characters in *Camellia*.

The image of a specimen on the website of Chinese Virtual Herbarium, *T.L. Ming et al. 224* (SYS 00089904, https://www.cvh.ac.cn/spms/detail.php?id=e87563d0), was identified as *C. hekouensis* by both T.L.Ming (on 10 May 1990) and Hung T.Chang (without date on the sheet). It may be, however, a plant of *Miliusa* sp. (Annonaceae). Remarkably, Ming (2000) did not cite this collection under his description of *C. hekouensis*.

Utilization and conservation

No records of specific utilization of *C. hekouensis* have been retrieved. Local people, except those who want to conserve the species, pay little attention to the plant. Though the species contains catechins and total polyphenols (Table 1), it does not suggest that the leaves can be used as a beverage source because *C. hekouensis* has a remotely phylogenetic relationship with *C*. sect. *Thea* (Fig. 2), the group of tea plants. The seeds of all species in *Camellia* can produce oil but their qualities vary widely (Wang et al. 2007, Su et al. 2014); unfortunately, the remaining three adult trees of *C. hekouensis* bear few seeds and the top priority is propagation rather than chemical analysis using the precious seeds. Local staff of the Reserve said the flowers of *C. hekouensis* are very fragrant, and these may be useful in horticulture. However, its white petals (Fig. 1E, F) might attract little attention of *Camellia* enthusiasts, who usually prefer yellow and red camellias. The woods of *Camellia* are usually dense and can be used

for handcrafts (Mou et al. 2007). The specific wood quality of *C. hekouensis* has not been studied probably because of unavailability. Its ecological values, such as co-evolution with its pollinators, herbivores and dwellers, are barely known.

However, *C. hekouensis* represents a unique germplasm of *Camellia* (Fig. 2). The tree seems to have a relatively long life cycle. Though the adult trees can bear many flowers, the fruits are very few. Artificial pollination may be helpful to increase the seed yield. Asexual reproduction by cuttings turned out a very low rate of survival in previous trials. Nevertheless, conservation and propagation should bear the top priority for *C. hekouensis*. The death of the previously misidentified "last living tree" of *C. hekouensis* (Fig. 1B) did give us cause for alarm and brought home the fragility of the species.

Acknowledgements

We thank Mr. Guangming Li of Hekou Administration Branch of Dawei Mountain National Nature Reserve in Yunnan, and Mr. Lu Yin of Agriculture and Rural Affairs and Science and Technology Bureau of Guangnan, Yunnan for their help during fieldwork, and the anonymous reviewers for their helpful comments.

Additional information

Conflict of interest

The authors have declared that no competing interests exist.

Ethical statement

No ethical statement was reported.

Funding

This work is supported by the Scientific Research Foundation of the Central South University of Forestry and Technology (2019YJ023), the National Natural Science Foundation of China (32270224), and the project "Basic research on protection, development and utilization of ancient tea trees in Panzhou City".

Author contributions

Conceptualization, investigation and methodology: DWZ. Resources and visualization: DWZ, GLZ, SXY. Writing - original draft: DWZ. Writing - review and editing: DWZ, GLZ, SXY.

Author ORCIDs

Dongwei Zhao https://orcid.org/0000-0002-7761-7127
Shixiong Yang https://orcid.org/0000-0001-8472-1275

Data availability

All of the data that support the findings of this study are available in the main text and GenBank as shown in Suppl. material 1.

References

Chang HT (1981) A Taxonomy of the Genus *Camellia*. The Editorial Staff of the Journal of Sun Yatsen University, Guangzhou, 180 pp.

Chang HT (1998) Theaceae (1) Theoideae 1. *Camellia*. In: Delectis Florae Reipublicae Popularis Sinicae Agendae Academiae Sinicae Edita (Eds) Flora Reipublicae Popularis Sinicae (Vol. 49[3]). Science Press, Beijing, 3–195.

Chang HT, Zhang RM, Ye CX (1996) Diagnosis on the systematic development of Theaceae IV Revision of the species of sect. *Thea*, *Camellia.* Acta Scientiarum Naturalium Universitatis Sunyatseni 35(3): 11–17.

Cheng L, Li MG, Han QW, Qiao Z, Hao YL, Balbuena TS, Zhao YY (2022) Phylogenomics resolves the phylogeny of Theaceae by using low-copy and multi-copy nuclear gene makers and uncovers a fast radiation event contributing to tea plants diversity. Biology 11: 1007. https://doi.org/10.3390/biology11071007

Fang W, Yang JB, Yang SX, Li DZ (2010) Phylogeny of *Camellia* sects. *Longipedicellata*, *Chrysantha* and *Longissima* (Theaceae) based on sequence data of four chloroplast DNA loci. Acta Botanica Yunnanica 32(1): 1–13. https://doi.org/10.3724/SP.J.1143.2010.00001

Gao JY, Parks CR, Du YQ (2005) Collected Species of the Genus *Camellia*: An Illustrated Outline. Zhejiang Science and Technology Press, Hangzhou.

Meng HH, Song YG (2023) Biogeographic patterns in Southeast Asia: Retrospectives and perspectives. Biodiversity Science 31(12): 23261. https://doi.org/10.17520/biods.2023261

Meng XH, Liu C, Fan R, Zhu LF, Yang SX, Zhu HT, Wang D, Yang CR, Zhang YJ (2018) Antioxidative flavan-3-ol dimers from the leaves of *Camellia fangchengensis*. Journal of Agricultural and Food Chemistry 66(1): 247–254. https://doi.org/10.1021/acs.jafc.7b04572

Ming TL (1999) A systematic synopsis of the genus *Camellia*. Acta Botanica Yunnanica 21(2): 149–159.

Ming TL (2000) Monograph of the Genus *Camellia*. Yunnan Science and Technology Press, Kunming, 352 pp.

Ming TL, Bartholomew B (2007) Theaceae. In: Wu ZY, Raven PH, Hong DY (Eds) Flora of China (Vol. 12). Hippocastanaceae through Theaceae. Science Press, Beijing / Missouri Botanical Garden Press, St. Louis, 366–478.

Mou JP, Xu F, Liang HW (2007) Study on wood anatomical molecules belonged to Theaceae genera of Guangxi. Guangxi Forestry Science 36(2): 78–81.

Pang Z, Wang YL, Mantri N, Wang Y, Hua XJ, Quan YP, Zhou X, Jiang ZD, Qi ZC, Lu HF (2022) Molecular phylogenetic relationships and taxonomy position of 161 *Camellia* species in China. Taiwania 67(4): 560–570.

Sealy JR (1958) A Revision of the Genus *Camellia*. The Royal Horticultural Society, London, 239 pp.

Su MH, Shih MC, Lin KH (2014) Chemical composition of seed oils in native Taiwanese *Camellia* species. Food Chemistry 156: 369–373. https://doi.org/10.1016/j.foodchem.2014.02.016

Thiers BM (2025) Index Herbariorum. http://sweetgum.nybg.org/science/ih/ [Accessed 2 May 2025]

Vijayan K, Zhang WJ, Tsou CH (2009) Molecular taxonomy of *Camellia* (Theaceae) inferred from nrITS sequences. American Journal of Botany 96: 1348–1360. https://doi.org/10.3732/ajb.0800205

Wang CJ, Fan GS (1988) New taxa of *Camellia* from Yunnan. Acta Botanica Yunnanica 10(3): 365–366.

Wang YY, Song XH, Li CR, Shi XG, Ye CX, Miao AQ, Li JX (2007) Fatty acids analysis of seeds oil from eight kinds of *Camellia* plants. China Oils and Fats 32(9): 78–79.

Yan YJ, Davis CC, Dimitrov D, Wang ZH, Rahbek C, Borregaard MK (2021) Phytogeographic history of the tea family inferred through high-resolution phylogeny and fossils. Systematic Biology 70(6): 1256–1271. https://doi.org/10.1093/sysbio/syab042

Ye CX, Lin YC, Zhou HY, Chen F, Li XY (1997) Isolation and analysis of purine alkaloids from *Camellia ptilophylla* Chang. Acta Scientiarum Naturalium Universitatis Sunyatseni 36(6): 30–33.

Zan T, He YT, Zhang M, Yonezawa T, Ma H, Zhao QM, Kuo WY, Zhang WJ, Huang CH (2023) Phylogenomic analyses of *Camellia* support reticulate evolution among major clades. Molecular Phylogenetics and Evolution 182: 107744. https://doi.org/10.1016/j.ympev.2023.107744

Zhang GL, Cai J, Jiang CQ (2016) Rare and Threatened Plants of Dawei Mountain National Nature Reserve in Yunnan. Yunnan Science and Technology Press, Kunming, 300 pp.

Zhao DW (2024a) Taxonomic revision of *Camellia* (Theaceae) in Thailand. PhytoKeys 239: 29–57. https://doi.org/10.3897/phytokeys.239.113878

Zhao DW (2024b) Botany and taxonomy of tea (*Camellia sinensis*, Theaceae) and its relatives. In: Chen L, Chen JD (Eds) The Tea Plant Genome. Springer, Singapore, 13–37. https://doi.org/10.1007/978-981-97-0680-8_2

Zhao DW, Hodkinson TR, Parnell JAN (2023) Phylogenetics of global *Camellia* (Theaceae) based on three nuclear regions and its implications for systematics and evolutionary history. Journal of Systematics and Evolution 61(2): 356–368. https://doi.org/10.1111/jse.12837

Supplementary material 1

Samples used in the study

Authors: Dongwei Zhao, Guiliang Zhang, Shixiong Yang

Data type: xlsx

Explanation note: Vouchers and GenBank numbers of samples used

Copyright notice: This dataset is made available under the Open Database License (http://opendatacommons.org/licenses/odbl/1.0/). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: https://doi.org/10.3897/phytokeys.256.149481.suppl1