

Are anthropogenic soils from dumpsites suitable for arable fields? Evaluation of soil fertility and biomass response to elements

Michael Asare Opare (✉ asare@fzp.czu.cz)

Czech University of Life Sciences Prague: Ceska Zemedelska Univerzita v Praze <https://orcid.org/0000-0002-0008-1459>

Research Article

Keywords: Anthropogenic soil, Bioaccessibility, Macronutrient, Micronutrient, Potentially toxic elements, Soil fertility

Posted Date: October 6th, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-2126356/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Version of Record: A version of this preprint was published at Plant and Soil on January 13th, 2023. See the published version at <https://doi.org/10.1007/s11104-023-05870-6>.

Abstract

Background and purpose

The fertility of anthropogenic soils developed from dumpsites used for arable fields is not well-studied. The study aimed to evaluate the fertility of soils from an abandoned dumpsite in Awotan, Nigeria, by measurable indicators and assess the bioaccessibility of elements of identified plant species.

Methods

The study adopted multi-analytical approaches to determine the signatures of the soils and further parameterized the bioaccessibility of elements to plants.

Results

The comparatively high content of Ca and Na in the anthropogenic soil contributed to the slightly alkaline soil reaction against the slightly acidic control. The high amount of organic matter is well-indicated by the enrichment of organic C and N in the anthropogenic soil. Waste deposition significantly contributed to the high accumulation of macronutrients (P, Ca, K, S) and micronutrients (Mn, Na, Fe) sufficient for maximum plant growth and yields, with an adequate C/N ratio supporting effective mineralization. The high cation exchange capacity of the anthropogenic soil contributed to cations binding. Indiscriminate deposition resulted in a high accumulation of potentially toxic elements (PTEs; Cu, Zn, As, Cd, Pb) above permissible limits in agricultural soils following WHO limits. Potential effects on lives are evident by the high PTEs accumulation in roots and leaves of *Chromolaena odorata*, *Saccopetalum tectonum*, *Passiflora foetida*, and *Senna siamea*. These plant species exhibited various PTEs accumulation, especially excluding Cd and Pb.

Conclusion

Although anthropogenic soils remained fertile, the bioaccessibility of PTEs by plants indicates potential threats to consumers of crops and herbs produced from such sites.

1. Introduction

Anthropogenic soils (human-altered-soils) can develop from diverse activities, including metallurgy, farming, and often deposition of waste in specified locations (Hard 2019). Wastes deposition often occurs at locations created by households or large land sizes for an entire community, especially in rural settings. With time these waste deposition sites, often called dumpsites, are abandoned and often reused as arable fields (Oyakhilome et al. 2019).

The material composition of decomposed waste in dumpsites is a function of elements (macro, micro, or risk elements) released into soils. For example, Sajjad et al. (2022) and Wan et al. (2019) reported that the presence of plastics in soils increases the content of potentially toxic elements (PTEs) and excessive loss of soil H₂O and may lead to microbial activity limitation. The accumulation of PTEs has toxicological consequences on soil-water/plant-animal interactions (e.g., Taylor et al. 2010; Thomas et al. 2015). For example, up to 4692 mg of Pb kg⁻¹ was detected in landfill (Gupta et al. 2011), with subsequent uptake by plants (Amusan et al. 2005). The deposition of waste, including the occasional burning of dumpsites and decomposition of organic matter, can also change the matrix color and other physical properties of the previous natural soil.

Waste, including kitchen leftovers (e.g., bones, wood ash, and food residues), can decompose to release vital plant nutrients (such as P, Ca, K, Mn, and Fe) into soils. Additionally, Bassey et al. (2021) recorded beneficial and pathogenic microbes belonging to *Clostridium*, *Escherichia*, *Pseudomonas*, *Enterococcus*, *Micrococcus*, *Shigella*, *Klebsiella*, and *Bacillus* in a dumpsite soil. However, indiscriminate disposal and burning or decay of wastes can result in substances that contain elevated metal/(loid)s, e.g., As, Cd, and excess Zn (Kalina et al. 2019). Therefore, the feasibility of crop and herbage production of former dumpsites is questionable.

Repurposing old dumpsites for green space is not a new thing. Abandoned dumpsites, alternatively, are often converted into arable fields and agricultural grazing lands. Although some authors evaluated the negative impacts of solid wastes disposition on soils from dumpsites, these studies related to risk elements (e.g., Pb and Cd) abundant in industrialized localities (Fakayode and Onianwa 2002; Onipede and Bolaji 2004) and reclamation of areas dumped with mine waste (Novak et al. 2008). Hence, the need for intermittent analyses of anthropogenic soils emanating from dumpsites. So far, what is not well-known in such arable fields is the fertility derived from their chemical, physical, and biological signatures and the level of toxicity of such soils and respective biomass.

For this study, soil quality is assessed based on indicators on the concept of control charts, e.g., regional and global permissible limits. Estimation of enrichment factor of elements includes a direct comparison between soils from the dumpsite and surrounding control site without waste deposition. The evaluation of indicators of soil fertility includes one of the most pertinent aspects of sustainable agricultural production (Sarhat 2015). The uptake and translocation of elements by plant species in such sites with diverse soil characteristics are required to support the accession of soil fertility. Hence, it is relevant to estimate the level of bioaccessibility of elements in the soil to plants to provide complete proof of the properties of soils in dump sites.

The current study provides comprehensive exploratory research on the fertility of one of the largest abandoned dump sites in Africa, Awotan, Nigeria (Alia et al. 2013; Amuda et al. 2014; Raman and Narayanan 2008), considering physical, chemical, and biological indicators. Many of such sites are currently used for agricultural production worldwide, e.g., Mzedi in Malawi (Kalina et al. 2022), Thailand (Boonmahathanakorn 2020), Calabar in Nigeria (Bassey et al. 2021),

and Tucson, Arizona (USA; Hard et al. 2019). However, studies that provide a detailed evaluation of the fertility of soils from dumpsites and assess their bioavailability of potentially toxic elements are less published.

Therefore, the current study demonstrated the fertility of anthropogenic soils from the dumpsite by 1) Comparing the soil physical properties of the dumpsite to surrounding control without the deposition of waste, 2) Estimating the enrichment of chemical properties (including pH, macro, trace, and risk element contents) of the anthropogenic soils from the dumpsite, and 3) Exploring the bioavailability of risk elements of selected plant species to make conclusive inferences on the suitability of such sites for arable fields.

2. Materials And Methods

2.1. Study site

The studied abandoned dumpsite (7°27'41" N 3°50'52" E; Fig. 1) is in Awotan, Ibadan in Oyo state, approximately 128 km NE of Lagos and 345 km SW of Abuja, the federal capital of Nigeria. The studied dumpsite has been in existence since 1829. The dumpsite covers an area of approximately 22 ha, with the distance of the nearest settlement only 200 m. The study area experiences two local climates; rainy (March to ends in October) and dry/harmattan (November to February) seasons. The average annual precipitation is 1311 mm, and an altitude between 200 and 300 m a. s. l. (MacDonald et al. 2005). The temperature of the study area ranges from 21 and 35 °C in the wet and dry seasons, respectively. The study site is well-characterized by lixisol, located on an igneous and metamorphic rock (Adelana et al. 2008; MacDonald et al. 2005).

2.2. Sampling design and sample collection

The study adopted to judgmental sampling technique after Maul (2018) to cover the variability of soil and plant samples on the site. Four pits (Locus A - D) were excavated 200 m apart on the edges of the dumpsite since the waste extends to 4–5 m high in the southwest section. Soil samples representing control additionally were collected from two randomized pits (non-residential area) with no waste deposition, approximately 200 m away from the dumpsite, with the same soil and parent bedrock. The waste exhibit high variability of substances, including plastic bags, partly decomposed organic, and inorganic materials.

Random bulk soil sample collection involves the upper 0–30 cm (arable layer) and 30–60 cm (subsoil layer) depths from each pit after the removal of surface litter (Figure S1). Sampling involved the collection of three soil samples from each depth, making an overall total sample number of 36. Two duplicates of the most dominant plant species, *Chromolaena odorata* L. (Asteraceae), *Saccopetalum tectonum* (Annonaceae), *Passiflora foetida* (Passifloraceae), and *Senna siamea* (Fabaceae) subsequently were collected in proximity to each sampled pit of the dumpsite. The nomenclature of the taxa used followed that adopted in APG IV- Angiosperm Phylogeny Group (<https://www.gbif.org/species/3106>). Only the leaves of *Senna siamea* were within reach during collection. Both plant and soil samples were collected and put in labeled polyethylene bags in May 2019.

2.3. Sample preparation and analysis

i. Soil

The collected soil samples were air-dried and sieved. The homogenous fraction under a 2 mm sieve was then analyzed for the pseudo-total content of elements. Procedure for soil samples digestion:

Approximately 5.0 g of each homogenized soil sample was measured into a conical flask, with 10 ml of concentrated nitric acid (HNO₃) added. The mixture was heated in the hot plate at 120 °C for approximately 20 minutes to near dryness and was allowed to cool and then filtered (using Whatman filter papers 125 mm Ø Cat No. 1001125) before being transferred into a standard flask made up to 50 ml of distilled H₂O. The digested solutions further were analyzed for the presence of total element concentrations using Atomic Absorption Spectrophotometer (AAS) Type: S4 AA System, Nc: 942340030042).

Soil reaction (pH_[H₂O]) was measured in a 1/2.5 (w/v) suspension of soil and 0.01M CaCl₂. Soil organic C (SOC) and total N were also determined with a Skalar Primacs SNC-100 analyzer produced by Skalar (Netherlands). The cation exchange capacity (CEC) was determined by the Isotherm approach (ISO 11260, 1994).

ii. Plant biomass

Below (roots) and above-ground (leaves) biomass were detached and washed with deionized ultrapure H₂O to remove ions transported from environmental dust. All the samples were additionally dried in the electric oven (SLW 53 STD, Pol-Eko, Wodzisław Śląski, Poland) at 60°C for 72 hours. Dried plant samples of each species were then ground with an electrical plant tissue pulverizer and stored in sealable bags. Homogenized powdered samples were prepared in two replicates (for each organ) and determined for total PTEs. The extraction of PTEs in plant samples (dry weight) followed the microwave-assisted acid approach (USEPA 3052, 1995), and the contents of elements were well-determined with inductively coupled plasma absorption emission spectroscopy (ICP-AES, Agilent 5900, Technologies Inc., USA). Three replicates of every sample were well-determined.

2.4. Size distribution of soils

Analysis of the soil grain distribution followed the mechanical sieve approach by FAO (2008) with size fractions of 2 ("very coarse"), 1 (coarse), 0.5 (medium), and 0.25 (fine) mm. For each representative soil sample, 20g was used for this analysis. Moreover, texture description followed the approach according to FAO (2008).

2.5. Statistical analysis

There was homogeneity of variance among data for all the analyzed samples. Dataset met the assumption of parametric statistics according to the Shapiro-Wilk W normality test. One-way ANOVA model was used to evaluate the differences in soil chemical properties, and content of elements in dry weight of biomass samples followed by the posthoc comparison using the Tukey honest significant difference (HSD) test applied to identify significant differences between chemical properties of soils with depth and content of elements among biomass. To evaluate the relationship between the concentration of elements, soil reaction (pH), and particle size distribution we used Pearson's correlation. STATISTICA 13.4 (www.statistica.io) was used to perform all statistical analyses.

2.6. Metal transport and accumulation indices

To study the behavior of PTEs in the soil-plant system, bioaccumulation (*BF*), translocation factors (*TF*), and mobility ratio (*MR*) was calculated as the ratio of PTE in vegetative aerial structures (leaves) to those in soil and root, with values > 1 indicating enrichment of the plant structures- accumulators (Mesa et al. 2017).

Calculation of *BF* follows the equation after Baker (1981),

$$i)BF = \text{Leafcontentofelement (Dwt)} \div \text{elementcontentinsoil, depth 0-60 cm depth}$$

$$ii)BF = \text{rootcontentofelement (Dwt)} \div \text{elementcontentinsoil, 0 - 60cmdepth (1)}$$

The calculation of *TF* follows the equation according to Klink et al. (2014),

$$iii)TF = \text{leafcontentofelement (Dwt)} \div \text{root (Dwt) contentofelement}$$

2

$$iv)MR = \frac{\text{contentofelementinleaf}}{\text{contentofelementinsoil, depth0 - 60cm}}$$

3

3. Results

The statistical description of soil's physical and chemical properties, the elemental composition of plant organs, and the bioavailability of PTEs are in Figures 2-7 and Tables 1-4.

3.1. Bulk soil chemical properties

There was a significant effect on pH and the contents of macro, micronutrients, and potentially toxic elements in individual loci (Figure 2-6). Background values of pH and plant nutrients (SOC, N, P, Ca, K, Na, Fe, Mn, Cu, and Zn) were predominantly lesser compared to soils from the dumpsites (Figures 2a-c, 3, 4a-c, and 5a-b) and vice versa in the case of S (Figure 4d). Additionally, there was no increasing or decreasing pattern of elements with depth in Loci A – D. However, decreasing trends of SOC, N, P, Ca, K, S, Fe, and Na were detected only with depths in control.

The soil reaction of the anthropogenic soil was slightly alkaline (Figure 2a) under the influence of Ca and Na (Figures S2a and b) compared to slight acidity in control. The results further show suitable conditions for mineralization of nutrients, according to the C/N ratio of approximately 12 in the soil from the dumpsite, due to the comparatively increased contents of SOC and total N (Figures 2b and c). Increased SOC, N, and C/N ratio support higher content of organic matter in the anthropogenic soil than control of this study.

The soil composition was evaluated principally against the limits of metallic elements, according to WHO's permissible limits for agricultural soils (Chiroma et al. 2014). Waste deposition significantly increased the content of Cu, Zn, As, Cd, and Pb in the soil from the dumpsite above the regulatory levels for agricultural soils and background values in control (Figures 5 and 6). The Ni contents in both the dumpsite and control were below limits in arable soils (Figure 6b). The results of waste deposition cause increases in CEC (125.7 - 141.7 mmol₊ kg⁻¹) in all the locus of the dumpsite in comparison with control- 9.2 – 15.7 mmol₊ kg⁻¹ (Figure 7).

3.2. Physicochemical characteristics of soil

Soils from the study site and control are primarily similar, exhibiting sandy loam characteristics (72.1 sand: 7.8 silt: 19.3 clay) according to FAO (2008) (Table 1). Sub-soil layers across the loci predominantly characterize coarse particles inter-grained by fine soils (clay and silt) serving as adhesives, which may prevent a high rate of leaching of elements. Meanwhile, a comparatively high percentage of coarse particles may support the preferential gravitational flow of elements in solution form during precipitation.

3.3. Relationship between soil chemical properties and particle size distribution

Gravels (> 2mm), coarse (< 2 - 1 mm) mm), and medium (> 1 - 0.5 mm) soil particles exhibited no significant positive correlation with the studied elements (Table S1). Except for Cu ($r = 0.60$, $p < 0.05$), there was no significantly positive correlation between silt (< 0.5 – 0.25) and the remaining studied elements. The correlation model indicates no adsorption of elements on clay particles (< 0.25 mm). There was almost no correlation between soil reaction (pH) and content of S, an indication of no or minimal leaching of elements ($r = -0.08$, $p = 0.722$, $y = 1.5484 - 0.0363 \times x$).

3.4. Potentially toxic elements in plant organs

Individual plant species display different contents of Cu, Zn, As, Cd, and Pb accumulation in their above- and below-ground biomass (Table 2). Out of the overall content of 966 mg Cu kg⁻¹ in the soil from the dumpsite, *Senna siamea* accumulated 55.4% in the leaf (d wt., dry weight), 15.3% by *Chromolaena odorata* L, and approximately 10% for the remaining species. Copper root accumulation was 33.4% in *Passiflora foetida*, 24.4% in *Saccopetalum tectonum* (24.4%), and 9.1% in *Chromolaena odorata* L.

Accumulation percentage according to the mean content of Zn in the soils was 64.5% in *Senna siamea*, followed by 55.1% in *Passiflora foetida*, 23.6% in *Saccopetalum tectonum*, with 21.6% in *Chromolaena odorata* L. Root accumulation of Zn was 86.7, 48.4, and 17.1% in *Chromolaena odorata* L, *Saccopetalum tectonum*, and *Passiflora foetida*, respectively. Except for *Saccopetalum tectonum* (22.9%), both leaf and root accumulation of As were ≤10% in all remaining plant species. Additionally, Cd accumulation in leaves was 201.5, 83.2, 38.2, and 27.7% in *Senna siamea*, *Saccopetalum tectonum*, *Chromolaena odorata* L, and *Passiflora foetida*, respectively. Root Cd accumulation, except for *Chromolaena odorata* L (101.2%), was over 200% for all the species. The accumulation efficiency of Pb was > 100% for all organs of the studied species except for *Saccopetalum tectonum* (82.8%) in leaves.

3.5. Bioaccumulation indicators

The selected plant species exhibit diverse transport systems of PTEs in different organs (Table 3). *Chromolaena odorata* L has a high uptake ability for Cd and Pb (BF >1) and an efficient translocation of Pb (BF/MR >1) and Cu (TF >1) to the leaf. Meanwhile, *Saccopetalum tectonum* has a high soil-root quotient (BF >1) for both Cd and Pb, together with effective leave transport (TF <1). Similarly, *Passiflora foetida* shows high Cd and Pb root uptake, leaf translocation of Pb, and TF >1 for Zn, while *Senna siamea* has high leaf uptake for both Cd and Pb (BF < 1).

4. Discussion

One of the misconceptions, especially in many rural settings, is that degradation/decomposition of waste can eventually fertilize soils for crop production. Current studies have shown the adverse effects of different waste deposition on soils (e.g., Chernykh et al. 2021; Gao and Minh 2022; Yeilagi et al. 2021). The current study offers valuable practical insight into the effects of indiscriminate deposition of waste to soils and the trophic levels, suggesting the critical assessment of several fertility evaluators to make conclusive inferences on the use of dumpsites for agricultural fields.

Although the grain distributions of the soils from the dumpsite and control show relatively high similarity with depth, the deposition of waste, particularly from domestic waste (e.g., food remains, furniture, and polyethylene materials) and occasional municipal loadings (e.g., plastic materials and a by-product of paints from industries) can further affect many soil physical properties. The pore size of the soils with a high fraction of coarse particles ($\geq 2-0.5$ mm) can support aeration, which contributes to microorganisms' mobility and the hydraulic properties of the soil (Hallam 2018; Mangalassery et al. 2013). The relatively high fraction of silt and clay provides a condition of element retention. Clay particles usually provide a surface area for element sorption. However, the pH (alkaline) of the anthropogenic soil affects clay dispersion by changing the net charge. In more alkaline soils, CaCO₃ formation leads to charge reduction on clay, which results in flocculation (Chorom and Rengasamy 1995).

The soil reaction as a “master soil variable” in the anthropogenic soil was slightly alkaline, primarily due to the comparatively high presence of base-forming cations associated with carbonates and bicarbonates (Ca, K, and Na). Conversely, the acidic conditions occur in control soil with parent material high in elements, such as silica, and high levels of sand with low buffering capacities- the ability to resist pH change (Jiang et al. 2018). Organic matter content evident by the comparatively high contents of SOC and N in the anthropogenic soil is often acidic, and the slightly alkaline condition produces a circum-neutral soil reaction that supports relative mineralization of nutrients for plant uptake. High organic matter and quality, often indicated by C/N values, contribute to optimal mineralization, as in the case of the anthropogenic soil of this study.

The soil's ability to hold and supply nutrients is partly related to its CEC, the number of parking spaces for nutrients on soil particles. Cation exchange capacities can also be influenced by soil pH. However, in this study, the slightly alkaline pH may produce little or no positive ions. Soils with high amounts of silt/clay and organic matter often show high CEC, which can bind more cations, e.g., Ca, K, and Na. They also have greater buffering capacity.

The decomposition of organic materials indicated by kitchen waste (e.g., animal and fish bones and food remains) contributed to high macro and micronutrients in the anthropogenic soil compared to the control. For instance, the P in the anthropogenic soil (0.16–0.23%) was higher than the level for crops and vegetables with high P (0.01–0.02%) requirements (Šrek et al. 2010). Besides the comparatively high nutrient (P, Ca, K, S, Fe, Na, and Mn) concentrations in the anthropogenic soils, these were within the values in agricultural soils without the application of mineral fertilizers (Marschner 1995; Šrek et al. 2010), which can promote maximum crop growth. Considering the withered nutrient-poor soils of the control in this study, often from the tropics, many farmers may resort to anthropogenic soil from the dumpsite for crop and herbage production.

Meanwhile, the Cu content as a nutrient for crops in this study was within the most recorded (5–20 mg kg⁻¹) without physiological impairment (Marschner 1995). The content of Cu and Zn as plant nutrients was above the permissible limit in agricultural soils, according to FAO/WHO (100 mg kg⁻¹ Cu) and EU (140

mg kg⁻¹ Cu) guidelines, and 300 mg kg⁻¹ for Zn, rendering them toxic for plant (Chiroma et al. 2014; EU guidelines 2002). The higher content of PTEs in the soil from the dumpsite is partly the result of the deposition of recycled plastic waste. For example, brominated organics and phthalate esters are extensively applied in plastic products as flame retardants and plasticizers, respectively (Hopewell et al. 2009). These additives also include plasticizers, pigments, stabilizers, and reinforcement components that contaminate soil- known to contain heavy metals (see also Luo et al. 2011; Nakashima et al. 2012; Rochman et al. 2013). Morf et al. (2007) reported the upper content range of Pb at 100–2000 mg kg⁻¹ in the plastic fraction of waste electrical and electronic equipment.

Plants can break the core rules governing the uptake and accumulation of PTEs. For example, some plants produce root exudates and attract specific microbes for PTEs dissolution for effective uptake and nutrient fixation (Visioli et al. 2015), while others possess (*Arabidopsis thaliana*) active protein transporters (Balafrej et al. 2020). *Acinetobacter*, *Bacillus*, *Gluconacetobacter*, and *Pseudomonas* are well-characterized by Zn solubility (Costerousse et al. 2017), which involves the reduction of soil pH, Zn chelation, or improvement of root growth. However, these mechanisms differ from one microorganism to another. Studies by Vogel-Mikuš et al. (2006) concluded that colonization by arbuscular mycorrhizal in *Nocceae praecox* increased the Zn shoot-to-root ratio, indicating better Zn uptake.

Several plants have different strategies for accumulating excess PTEs in their organs, including the most edible parts, e.g., root/tuber/rhizomes and leaves. In an attempt to exclude high Cu, *Passiflora foetida* retains 40% Cu in the root. Meanwhile, 40% of 966 mg Cu kg⁻¹ in the soil represents 386 mg Cu kg⁻¹, still above the permissible limit for edible crops (FAO/WHO; Table 4). It is worth noting that plants, such as *Senna siamea*, are used locally in treating malaria fever, especially the leaves (Nas et al. 2018). Meanwhile, this study shows that the leaves of *Senna siamea* can accumulate Cu, Zn, As, Cd, and Pb above permissible limits when compared to FAO/WHO and EU (Commission Regulation (EC) No 1881/2006) standards.

These plant species exhibit promising accumulation of PTEs in different organs. The accumulation abilities have resulted in their persistence in the contaminated soils; adaptation by tolerance or by accumulation and detoxification. All the studied plant species exhibited high tolerance and excluded Cd and Pb (BF > 1), which provide a basis for their inclusion in phytostabilization and other phytoremediation schemes of contaminated sites. Notably, many food crops and vegetables have the high ability to uptake and distribute PTE in their organs in urban, peri-urban, and rural areas, e.g., *Rumex alpinus*, *Brassica oleracea* var. *capitata*, and *Daucus carota* (see Jungová et al. 2022; Mensah et al. 2009). Even though Pb shows restrictive transport of to shoot due to apoplastic blockage in the root, e.g., the Casparian barrier (Dogan et al. 2018), the leaf content was higher than the allowable limit.

The redevelopment of waste landfills for agricultural purposes remained sufficiently researched in other parts of the world with mixed results. Green et al. (2014) concluded that places with historic landfills repurposed for grazing are generally safe for grazers, while animals remain exposed to high metal ingestion rates. A study in Eritrea focused on crops grown in landfill soil and observed that although the landfill soil improved the fertility of farmers' fields, it had alarming contents of heavy metals (Tesfai and Dresher 2009). It is now evident that plants can grow well on soils from dumpsites due to nutrient availability. However, these plants may still support the uptake and accumulation of PTEs in different compartments and can adversely affect potential consumers. Unfortunately, food security and quality are less assessed at many local government levels. Although several studies have reported high contents of PTEs, including Pb and Cd, in soil dumpsites (e.g., Giao and Minh 2022; Musa et al. 2019), a direct relationship with plants will indicate the level of impact on the food chain.

The study can represent a guide for the spontaneous analysis of anthropogenic soils from dumpsites and a recipe for their arable land use, as many waste disposal sites are not well-streamlined to strategically address their agricultural use.

5. Conclusions

The study revealed the characteristics of anthropogenic soils developed from the indiscriminate deposition of wastes, as these soils are often transformed into arable lands. Physical properties, including the porosity of the studied anthropogenic soils, remained relatively similar compared to surrounding control without the mark of waste deposition. The anthropogenic soils were well-characterized by slightly neutral soil reactions and high CEC, resulting in the presence of high cationic nutrients (Ca, K, and Na). The high content of SOC and N contributed to the optimal C-N ratio of the anthropogenic soils, which provides a basis for the mineralization of nutrients. The fertility of the soils from the dumpsite was coupled with a substantially high accumulation of macro (P, Ca, K, and S) and micro (Mn, Na, and Fe) nutrients, with no or minimal indication of leaching to groundwater, according to the content of elements at different depths of the soil.

Although the results of the fertility indicators of the anthropogenic soil permit cultivation, the contents of PTEs (Cu, Zn, As, Cd, and Pb) were above regulatory limits in agricultural soils, according to WHO. Availability of Cu, Zn, As, Cd, and Pb in the organs of tolerant plant species (*Chromolaena odorata*, *Saccopetalum tectonum*, *Passiflora foetida*, and *Senna siamea*) in the dumpsite indicate that these crops and herbages produced from this sites can be detrimental to the health of consumers. All plant species studied exhibited high tolerance and excluded Cd and Pb (BF > 1; below-ground organ) and can be considered in phytostabilization and other phytoremediation schemes of contaminated sites.

Declarations

Author's contribution

MOA conceived and executed the research and wrote the paper.

Competing interests

The author declares no conflict of interest.

Data availability statement

The datasets generated during and/or analyzed during the current study are available from the corresponding author.

Funding

This research received support from the Nutrisk project (European Regional Development Fund - Project No. CZ.02.1.01/0.0/0.0/ 16_019/0000845).

References

1. Adelana SMA, Olasehinde PI, Bale RB, Vrbka P, Edet AE, Goni IB (2008) An overview of the geology and hydrogeology of Nigeria. In: Adelana S, MacDonald A (eds) Applied groundwater studies in Africa, CRC Press, London, pp 171-197.
2. Alia AA, Shahida NK, Bushra J, Saeed AA (2013) Phosphate solubilizing bacteria associated with vegetables roots in different ecologies. Pakistan J Bot 5 535-544.
3. Amuda OS, Adebisi SA, Jimoda LA, Alade AO (2014) Challenges and Possible Panacea to the Municipal Solid Wastes Management in Nigeria. J Sust Dev Stud 1: 64-70.
4. Amusan AA, Ige DV, Olawale R (2005) Characteristics of soils and crops' uptake of metals in municipal waste dump sites in Nigeria. J Human Ecol 17: 167-171.
5. Baker AJM (1981) Accumulators and Excluders Strategies in Response of Plants to Heavy Metals. J Plant Nutr 3:643-654. <https://doi.org/10.1080/01904168109362867>
6. Balafrej H, Bogusz D, Triqui ZA, Guedira A, Bendaou N, Smouni A, Fahr M (2020) Zinc Hyperaccumulation in Plants: A Review. *Plants*, 9, 562. <https://doi.org/10.3390/plants9050562>
7. Bassey IU, Edet UO, Umoafia NG, Nwachi AC, Ebenge IA, Odokuma L (2021) Microbial structure and function diversity of open dumpsite compost used as fertilizer by peasant farmers. Sci Afri 11: e00699 <https://doi.org/10.1016/j.sciaf.2021.e00699>.
8. Boonmahathanakorn S (2020) Rubbish dump turned lush urban farm. Available via <https://www.iied.org/rubbish-dump-turned-lush-urban-farm>. Accessed 9 Sep 2022.
9. Chernykh N, Baeva Y, Thoma A (2021) Content of heavy metals and arsenic in soils near Sharra landfill (Tirana, Albania). E3S Web Conf 265: 03003. <https://doi.org/10.1051/e3sconf/202126503003>
10. Chiroma TM, Ebewele RO, Hymore FK (2014) Comparative assessment of heavy metal levels in soil, vegetables, and urban grey wastewater used for irrigation in Yola and Kano. Intern Ref J Eng Sci 3: 1-9.
11. Chorom M, Rengasamy TP (1995) Dispersion and zeta potential of pure clays as related to net particle charge under varying pH, electrolyte concentration and cation type Eur J Soil Sci 46: 657-665
12. Costerousse B, Schönholzer-Mauclaire L, Frossard E, Thonar C (2017) Identification of Heterotrophic Zinc Mobilization Processes among Bacterial Strains Isolated from Wheat Rhizosphere (*Triticum aestivum* L.). Appl Environ Microbiol 84: e01715-17. <https://doi.org/10.1128/AEM.01715-17>
13. European guidelines (2002) Communication from the Commission to the Council, the European Parliament, the Economic and Social Committee, and the Committee of the Regions. Towards a Thematic Strategy for Soil Protection. Available via <https://eur-lex.europa.eu/LexUriServ/LexUriServ.do?uri=COM:2002:0179:FIN:EN:PDF>. Accessed 12 Sep 2022
14. Dogan M, Karatas M, Aasim M (2018) Cadmium and lead bioaccumulation potentials of an aquatic macrophyte *Ceratophyllum demersum* L.: A laboratory study. Ecotoxicol Environ Safety 148: 431-440
15. Fakayode S, Onianwa P (2002) Heavy metal contamination of soil, and bioaccumulation in Guinea grass (*Panicum maximum*) around Ikeja Industrial Estate, Lagos, Nigeria. Environ Geo 43:145-150. <https://doi.org/10.1007/s00254-002-0633-9>
16. FAO (2008) Guide to laboratory establishment for plant nutrient analysis. FAO fertilizer and plant nutrition bulletin, Rome, Italy. Available via <http://www.fao.org/3/a-i0131e.pdf>. Accessed 22 October 2021.
17. Giao NT, Minh VQ (2022) Risk associated with the occurrence of toxic elements in the environment surrounding landfills in An Giang Province, Vietnam. Soil Water Res 17: 80-90
18. Green ID, Boughey K, Diaz A (2014) Potentially toxic metals in historic landfill sites: Implications for grazing animals. Water Air Soil Pollut 225:1-12. [10.1007/s11270-014-2110-y](https://doi.org/10.1007/s11270-014-2110-y).
19. Gupta D, Ghosh R, Mitraet AK, et al (2011) Nondestructive characterization of municipal-solid-waste- contaminated surface soil by energy-dispersive X-ray fluorescence and low-Z (atomic number) particle electron probe X-ray microanalysis. J Air Waste Manag Assoc 61:1102-14
20. Hallam J (2018) Soil hydraulic function: Earthworm-plant root interactions. Ph.D. thesis, Environment and Geography University of York.
21. Hard HR, Brusseau M, Ramirez-Andreotta M (2019) Assessing the feasibility of using a closed landfill for agricultural graze land. Environ Monit Assess 191:458.
22. Hopewell J, Dvorak R, Kosior E (2009) Plastics recycling: challenges and opportunities. Philos Trans R Soc Lond B Biol Sci 364: 2115-2126. [10.1098/rstb.2008.0311](https://doi.org/10.1098/rstb.2008.0311)
23. Jiang J, Wang Y-P, Mengxiao Yu M, Cao N, Yan J (2018) Soil organic matter is important for acid buffering and reducing aluminum leaching from acidic forest soils. Chem Geol 501: 86-94. <https://doi.org/10.1016/j.chemgeo.2018.10.009>.

24. Jungová M, Asare MO, Jurasová V et al (2022) Distribution of micro- (Fe, Zn, Cu, and Mn) and risk (Al, As, Cr, Ni, Pb, and Cd) elements in the organs of *Rumex alpinus* L. in the Alps and Krkonoše Mountains. *Plant Soil* 477: 553–575 (2022). <https://doi.org/10.1007/s11104-022-05440-2>
25. Kalina M, Tilley E (2019) Blurred lines: agricultural production on the margins of a dumpsite in Blantyre, Malawi. Available via https://www.researchgate.net/publication/343222728_'BLURRED_LINES'_AGRICULTURAL_PRODUCTION_ON_THE_MARGINS_OF_A_DUMPSITE_IN_BLANTYRE_MALAWI. Accessed 20 Sep 2022.
26. Klink A, Stankiewicz A, Wisłocka M, Polechońska L (2014) Macro- and microelement distribution in organs of *Glyceria maxima* and biomonitoring applications. *Environ Monit Assess* 186:4057– 4065 <https://doi.org/10.1007/s10661-014-3680-2>
27. Luo Y, Qin G, Zhang J (2011) D-myo-inositol-3-phosphate affects phosphatidylinositol-mediated endomembrane function in *Arabidopsis* and is essential for auxin-regulated embryogenesis. *Plant Cell* 23:1352-72
28. MacDonald AM, Cobbing J, Davies J (2005) Developing groundwater for rural water supply in Nigeria. British Geological Survey Commissioned Report CR/05/219N. Available via <https://core.ac.uk/reader/55390>. Accessed 10 Jun 2022.
29. Mangalassery S, Sjögersten S, Sparkes D, Sturrock C, Mooney SJ (2013) The effect of soil aggregate size on pore structure and its consequence on the emission of greenhouse gases. *Soil Till Res* 132:39- 46
30. Maul A (2018) Judgment Sampling. In: Frey BB (ed) *The SAGE Encyclopedia of Educational Research, Measurement, and Evaluation*, SAGE Publications, Inc, California, USA, pp 914-914.
31. Marschner H (1995) *Mineral Nutrition of Higher Plants* (2nd ed) Academic, San Diego
32. Mensah E, Kyei-Baffour N, Ofori E, Obeng G (2009) Influence of Human Activities and Land Use on Heavy Metal Concentrations in Irrigated Vegetables in Ghana and Their Health Implications. In: Yanful EK (ed) *Appropriate Technologies for Environmental Protection in the Developing World*. Springer, Dordrecht.
33. Mesa V, Navazas A, González-Gil R et al (2017). Use of Endophytic and Rhizosphere Bacteria To Improve Phytoremediation of Arsenic-Contaminated Industrial Soils by Autochthonous *Betula celtiberica*. *Appl Environ Microbiol* 83:e03411-16.
34. Morf LS, Tremp J, Gloor R, Schuppisser F, Stengele M, Taverna R (2007) Metals, non-metals and PCB in electrical and electronic waste – Actual levels in Switzerland. *Waste Manage* 27:1306-1316
35. Musa JJ, Bala JD, Mustapha HI et al (2019) Determination of the elemental composition of soil samples from selected dumpsites in Nasarawa, Kogi and Niger States, Nigeria. *J Solid Waste Tech Manage* 45. <https://doi.org/10.5276/JSWTM/2019.457>
36. Nakashima K, Takasaki H, Mizoi J, Shinozaki K, Yamaguchi-Shinozaki K (2011) NAC transcription factors in plant abiotic stress responses. *Biochim et Biophys Acta (BBA) - Gene Reg Mec* 1819: 97-103
37. Nas FS, Oyeyi TI, Ali M (2018) Antibacterial efficacy and phytochemical screening of *Senna siamea* leaves extracts on some pathogenic bacteria. *J Microbiol Exp* 6:159-163. DOI: 10.15406/jmen.2018.06.00208
38. Nigeria Population Commission and National Bureau of Statistics (2016) National Population Estimates. Available via <https://nigerianstat.gov.ng>. Accessed 23 July 2022
39. Novak J, Hák J, Steklou J, Jana Ceska J, Dvorakova E (2008) Reclaimed spoil heap from an open-cast mine: analysis of meadow communities. Available via https://www.researchgate.net/publication/295714957_RECLAIMED_SPOIL_HEAP_FR_OM_AN_OPEN-CAST_MINE_ANALYSIS_OF_MEADOW_COMMUNITIES. Accessed 1 Sep 2022.
40. Onipede AIM, Bolaji BO (2004) Management and disposal of industrial waste in Nigeria. *Nig J Mech Eng* 2: 49-58.
41. Oyakhilome GI, Ajiwe VI, Akinola OK (2019) The Microbial Status and Physico-Chemical Pollutants Studies of Rivers Owan and Evbiobe in Edo State, Nigeria. *Sci J Anal Chem* 7: 57-64. DOI: 10.11648/j.sjac.20190702.13
42. Raman N, Narayanan DS (2008) Impact of solid waste effect on groundwater and soil quality nearer to Pallavaram solid waste landfill site in Chennai. *Rasayan J Chem* 4: 828-836.
43. Rochman C, Browne M, Halpern B et al (2013) Classify plastic waste as hazardous. *Nature* 494: 169–171. <https://doi.org/10.1038/494169a>
44. Sajjad M, Huang Q, Khan S, Khan MA et al (2022) Microplastics in the soil environment: A critical review. *Environ Technol Innov* 27:102408 <https://doi.org/10.1016/j.eti.2022.102408>.
45. Sarhat AR (2015) Assessment of some indicators of soil fertility in a number of agricultural lands under wheat production system in Kalar. Available via https://www.researchgate.net/publication/327824490_Assessment_of_some_indicators_of_soil_fertility_in_a_number_of_agricultural_lands_under_wheat_production_system_in_Kalar. Accessed 31 Aug 2022.
46. Srek P, Hejcmána M, Kunzová E (2010) Multivariate analysis of relationship between potato (*Solanum tuberosum* L.) yield, amount of applied elements, their concentrations in tubers, and uptake in a long-term fertilizer experiment. *Field Crops Research* 118:183-193
47. Taylor MP, Mackay AK, Hudson-Edwards KA, Holz E (2010) Soil Cd, Cu, Pb, and Zn contaminants around Mount Isa city, Queensland, Australia: Potential sources and risks to human health. *Appl Geochem* 25: 841–855. <https://doi.org/10.1016/j.apgeochem.2010.03.003>.
48. Tesfai M, Drescher S (2009) Assessment of benefits and risks of landfill materials for agriculture in Eritrea. *Waste manage* 29: 851–858.
49. Thomas IHL, Bautista F, Ruíz FRC, Delgado MC, Owen PQ, Aguilar D, Goguitchaichvili A (2015) Concentration of toxic elements in topsoil of the metropolitan area of Mexico City: Spatial analysis using ordinary kriging and indicator kriging. *Revista Internacional de contaminación Ambiental* 31: 47-62.
50. Visioli, G., D'Egidio, S., & Sanangelantoni, A. M. (2015). The bacterial *rhizobiome* of hyperaccumulators: future perspectives based on omics analysis and advanced microscopy. *Frontiers in Plant Sciences*, 5, 752. <https://doi.org/10.3389/fpls.2014.00752>

51. Vogel-Mikuš K, Pongrac P, Kump P, Nečemer, Regvar M (2006) Colonization of a Zn, Cd, and Pb hyperaccumulator *Thlaspi praecox* Wulfen with indigenous arbuscular mycorrhizal fungal mixture induces changes in heavy metal and nutrient uptake. Environ Pollut 139: 362–371.
52. Wan Y, Wu C, Xue Q, Hui X (2019) Effects of plastic contamination on water evaporation and desiccation Sci Total Environ 654: 576-582
53. WHO (1996) Permissible Limits of Heavy Metals in Soil and Plants. World Health Organization, Geneva, Switzerland.
54. Yeilagi S, Rezapour S, Asadzadeh F (2021) Degradation of soil quality by the waste leachate in a Mediterranean semi-arid ecosystem. Sci Rep 11: 11390. <https://doi.org/10.1038/s41598-021-90699-1>

Tables

Table 1. Physical characteristics, including particle size distribution of soils from the dumpsite and neighboring control (cont.) soils

Sampled location	Depth [cm]	Texture ^a	Consistency	Grain size distribution [%]				
				≥ 2 mm	< 2 – 1mm	< 1 – 0.5	< 0.5 – 0.25	< 0.25
				Gravel	Coarse sand	Medium sand	Fine sand	Fines
L. A	0 – 30	Sandy loam	Coarse slightly sticky + slightly plastic	19.4 ± 0.9 ^a	11.5 ± 0.6 ^{ab}	41.9 ± 2.3 ^a	8.1 ± 0.4 ^a	18.2 ± 1.1 ^a
	30 – 60	Sandy loam	Slightly coarse, sticky + slightly plastic	17.6 ± 0.3 ^a	11 ± 0.2 ^{ab}	45 ± 1.9 ^{ab}	6 ± 0.1 ^a	19.9 ± 0.7 ^a
L. B	0 – 30	Sandy loam	Coarse slightly sticky + slightly plastic	21.1 ± 1.1 ^{ab}	12.1 ± 2.2 ^{ab}	43.8 ± 2.6 ^{ab}	5.6 ± 0.2 ^b	16.9 ± 1 ^{ab}
	30 – 60	Sandy clay loam	Coarse, sticky + plastic	18.7 ± 0.7 ^a	10 ± 0.4 ^a	40.9 ± 1.7 ^a	10 ± 0.09 ^b	20 ± 0.7 ^a
L. C	0 – 30	Sandy loam	Coarse slightly sticky + slightly plastic	21.4 ± 1.5 ^{ab}	10.9 ± 0.1 ^a	41.6 ± 3.2 ^a	6 ± 0.3 ^b	19.9 ± 0.3 ^a
	30 – 60	Sandy loam	Slightly coarse, sticky + slightly plastic	20.7 ± 0.9 ^{ab}	10.9 ± 0.6 ^a	40.6 ± 2.4 ^{ab}	8.1 ± 0.5 ^c	19 ± 0.6 ^a
L. D	0 – 30	Sandy loam	Coarse slightly sticky + slightly plastic	19.7 ± 0.8 ^a	12.5 ± 2 ^b	44.3 ± 2.8 ^{ab}	5.2 ± 0.02 ^b	18 ± 0.3 ^a
	30 – 60	Sandy clay loam	Coarse, sticky + plastic	18.1 ± 1.3 ^a	12 ± 2.3 ^{ab}	40.2 ± 3 ^a	12.3 ± 0.6 ^d	17.2 ± 0.9 ^{ab}
Cont. 1	0 – 30	Sandy loam	Coarse, sticky + plastic	18.1 ± 0.7 ^a	10.3 ± 0.04 ^a	43 ± 0.8 ^{ab}	8.6 ± 0.8 ^a	20 ± 2 ^a
	30 – 60	Sandy loam	Coarse, sticky + plastic	18 ± 0.2 ^a	10 ± 0.2 ^a	42.6 ± 1.4 ^{ab}	9.1 ± 0.2 ^c	19.7 ± 0.9 ^a
Cont. 2	0 – 30	Sandy loam	Coarse, sticky + plastic	18.7 ± 0.1 ^a	11 ± 0.6 ^{ab}	41.4 ± 1 ^a	9.7 ± 0.2 ^c	19.9 ± 1.3 ^a
	30 – 60	Sandy loam	Slightly coarse, sticky + slightly plastic	16.8 ± 0.4 ^b	10.9 ± 0.2 ^a	43.7 ± 2.8 ^{ab}	5.3 ± 0.01 ^b	23 ± 1.5 ^b
<i>p</i> -value				0.0271	0.001	0.001	0.001	0.001

A represents texture according to FAO (2008)

Table 2. The content of risk elements accumulated in plant organs [leaf and root] in the soils [mean values of 0 - 60 cm depths] from the dumpsite

Location	Species	Cu [mg kg ⁻¹]			Zn [mg kg ⁻¹]			As [mg kg ⁻¹]			Cd [mg kg ⁻¹]			Pb [mg kg ⁻¹]	
		Soil	leaf	root	Soil	leaf	root	Soil	leaf	root	Soil	leaf	root	Soil	leaf
L. A	<i>Chromolaena odorata L</i>	969.3	171	94.5	713.2	140	806.5	253.9	21	4.35	4.75	3.05	6.5	102.2	103
L. B	<i>Chromolaena odorata L</i>	964	135	81.85	802.2	141	466.5	246.7	17.5	6.75	4.88	2.56	6.84	86.17	122
L. C	<i>Chromolaena odorata L</i>	991	146.5	81.5	829.8	193.5	470.5	384.9	22.4	8.1	12.3	2.52	7.73	116.9	119
L. D	<i>Chromolaena odorata L</i>	939.5	138.5	91.85	725	187.5	917	294.4	20	7.45	5.47	2.37	6.8	94.8	148.5
	Mean	966	147.8	87.4	767.6	165.5	665.1	295	20.21	6.66	6.85	2.62	6.97	100	123.1
L. A	<i>Saccopetalum tectonum</i>	969.3	74	219.5	713.2	213	323	253.9	51	32	4.75	7.3	102	102.2	85.5
L. B	<i>Saccopetalum tectonum</i>	964	77.5	253.5	802.2	167.5	370	246.7	41.5	27.8	4.88	1.1	114	86.17	70
L. C	<i>Saccopetalum tectonum</i>	991	94	243	829.8	144	421	384.9	71.4	33	12.3	3.6	112	116.9	80.7
L. D	<i>Saccopetalum tectonum</i>	939.5	119	230	725	199	372.5	294.4	106	29.5	5.47	10.7	104	94.8	95
	Mean	966	91.1	236.5	767.6	180.9	371.6	295	67.5	30.6	6.85	5.7	107.8	100	82.8
L. A	<i>Passiflora foetida</i>	969.3	102.9	309	713.2	474.5	128.5	253.9	21.85	23	4.75	1.62	67	102.2	125.5
L. B	<i>Passiflora foetida</i>	964	107	331	802.2	403	115.5	246.7	12.25	27.6	4.88	2.23	69	86.17	144.5
L. C	<i>Passiflora foetida</i>	991	92.9	337.5	829.8	371.5	144.5	384.9	12.75	24.1	12.3	1.99	75	116.9	124.5
L. D	<i>Passiflora foetida</i>	939.5	121.5	311.5	725	442	136	294.4	10.45	21.2	5.47	1.72	91	94.8	160.5
	Mean	966	106.08	322.3	767.6	422.75	131.1	295	14.3	24.0	6.85	1.9	75.5	100	138.8
L. A	<i>Senna siamea</i>	969.3	530	-	713.2	463.5	-	253.9	9.95	-	4.75	10.8	-	102.2	169
L. B	<i>Senna siamea</i>	964	517.5	-	802.2	482.5	-	246.7	14.6	-	4.88	17.2	-	86.17	119
L. C	<i>Senna siamea</i>	991	505	-	829.8	515	-	384.9	23.75	-	12.3	11.8	-	116.9	161
L. D	<i>Senna siamea</i>	939.5	587.5	-	725	519.5	-	294.4	24.85	-	5.47	15.7	-	94.8	137.8
	Mean	966	535		767.6	495.1		295	18.3		6.85	13.8		100	146.7

Table 3. Bioavailability indices, including bioaccumulation [BF], translocation [TF] factors, and mobility ratio [MR] of potentially toxic [PTEs] of selected plant species

PTEs	<i>Chromolaena odorata L</i>				<i>Saccopetalum tectonum</i>				<i>Passiflora foetida</i>				<i>Senna siamea</i>	
	BF[l/s]	BF[r/s]	TF	MR	BF[l/s]	BF[r/s]	TF	MR	BF[l/s]	BF[r/s]	TF	MR	BF[l/s]	
Cu	0.15	0.09	1.68	<i>0.15</i>	0.09	0.24	0.39	<i>0.09</i>	0.11	0.33	0.33	<i>0.11</i>	0.55	
Zn	0.22	0.87	0.25	<i>0.22</i>	0.24	0.48	0.49	<i>0.24</i>	0.55	0.17	3.22	<i>0.55</i>	0.51	
As	0.07	0.02	3.03	<i>0.07</i>	0.23	0.1	2.21	<i>0.23</i>	0.05	0.08	0.6	<i>0.05</i>	0.06	
Cd	0.38	1.02	0.38	<i>0.38</i>	0.83	15.74	0.05	<i>0.83</i>	0.28	11.02	0.03	<i>0.28</i>	2.01	
Pb	1.23	1.82	0.68	1.23	0.83	1.83	0.45	<i>0.83</i>	1.39	4.78	0.29	1.39	1.47	

Bioavailability indices were estimated from the mean contents of elements in soil (0-60 cm) and plant organs. Abbreviation: l= leaf; s= soil, and r= root. BF; TF; MR > 1 indicates the plant could potentially be a hyperaccumulator [Baker (1981) and Mesa et al. (2017)]

Table 4. Mean content of potentially toxic elements in organs of selected plants species and their respective permissible limits

PTE	<i>Chromolaena odorata</i>		<i>Saccopetalum tectonum</i>		<i>Passiflora foetida</i>		<i>Senna siamea</i>	WHO/FAO permissible limit in plants [mg kg ⁻¹]	EU leaf vegetables/fresh herbs [mg kg ⁻¹]	EU root vegetables [mg kg ⁻¹]
	leaf	root	leaf	root	leaf	root	root			
Cu	147.8	87.4	91.1	236.5	106.08	322.3	535	10	0.3	
Zn	165.5	665.1	180.9	371.6	422.75	131.1	495.1	0.6	< 0.5	
As	20.21	6.66	67.5	30.6	14.3	24.0	18.3	0.5	-	
Cd	2.62	6.97	5.7	107.8	1.9	75.5	13.8	0.02	0.2	0.1
Pb	123.1	182.3	82.8	182.8	138.8	478	146.7	2	0.3	

Commission Regulation (EC) No 1881/2006 of 19 December 2006 set maximum levels for certain contaminants in foodstuffs (Text with EEA relevance).

<https://eur-lex.europa.eu/legal-content/EN/TXT/?uri=celex:32006R1881>

WHO (1996)

Figures




Figure 1

Map of Nigeria showing the location of the studied dumpsite in Awutan and associated waste deposition




Figure 2

Effect of waste deposition on (a) pH and mean content of (b) organic (org) C, (c) total (Tot) N, and (d) C/N ratio from different sampled locations. The F-and p-values were obtained by One-way ANOVA. Using Tukey *posthoc*HSD test, mean values with the same letters were significantly not different. The error bars indicate the standard error of the mean. The xx indicates the trend of movement (increase and decrease in content)




Figure 3

Effect of waste deposition on the mean content of total (a) P, (b) Ca, (c) K, and (d) S from different sampled locations. The F-and p-values were obtained by One-way ANOVA. Using Tukey *posthoc*HSD test, mean values with the same letters were significantly not different. The error bars indicate the standard error of the mean. The xx indicates the trend of movement (increase and decrease in content)




Figure 4

Effect of waste deposition on the mean content of total (a) Na, (b) Fe, (c) Mn, and (d) Si from different sampled locations. The F-and p-values were obtained by One-way ANOVA. Using Tukey *posthoc*HSD test, mean values with the same letters were significantly not different. The error bars indicate the standard error of the mean. The xx indicates trend of movement (increase and decrease in content)




Figure 5

Effect of waste deposition on the content of (a) Cu, (b) Zn, (c) As, and Cd from different sampled locations. The F-and p-values were obtained by One-way ANOVA. Using Tukey *posthoc* HSD test, mean values with the same letters were significantly not different. The error bars indicate the standard error of the mean. Abbreviation: x- permissible limit in agricultural soils according to WHO (Chiroma et al., 2014), xx indicates a trend of increases.




Figure 6

Effect of waste deposition on the content of (a) Pb and (b) Ni from different sampled locations. The F-and p-values were obtained by One-way ANOVA. Using Tukey *posthoc* HSD test, mean values with the same letters were significantly not different. The error bars indicate the standard error of the mean. Abbreviation: x- permissible limit in agricultural soils according to WHO (Chiroma et al., 2014), xx indicates a trend of increases.




Figure 7

Effect of waste deposition on the content of Cation exchange capacity (CEC) from different sampled locations. The F-and p-values were obtained by One-way ANOVA. Using Tukey *posthoc*HSD test, mean values with the same letters were significantly not different.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SupplementaryMaterial.docx](#)