

Morphological Response of *Lycopersicum esculentum* Mill (Tomato) to Salt (NaCl and Na₂SO₄) Stress

Jacob Olusola Oyetunji*; Samuel Kennedy Banja** and Elizabeth Folake*

Department of Botany University of Ibadan
Plant Physiology Unit, Department of Botany, University of Ibadan, Oyo State, Nigeria

Department of Biological Sciences, Taraba state University, Jalingo, Taraba State
kennedysamuel161@yahoo.com

Abstract:

Soil salinity is regarded as the major factor affecting crop productivity. To evaluate the effect of NaCl and Na₂SO₄ at different concentrations on the morphological features of Tomato plant, a variety of tomato was grown in green house condition subjected to varying concentrations of NaCl and Na₂SO₄. All experimental units were laid out in Completely Randomized Design (CRD) with three replications. The result obtained from the study showed a significant decline in plants height and number of leaves in stressed plants. Tomato plants subjected to high salinity (120 and 160mM Na₂SO₄) died off after 2weeks of treatment while plants treated with 160 mM of NaCl had the least biomass (0.82g). Fresh weight of *Lycopersicum esculentum* declined with increase in NaCl and Na₂SO₄ where Na₂SO₄ had least biomass and recorded crop mortality at 120 mM and 160mM concentration. This study, revealed that despite the ability of tomato plant to tolerate salinity stress at low concentration, a significantly decline in growth and biomass at higher salt concentration can occur. Both NaCl and Na₂SO₄ inhibited growth, metabolic activity, and ionic uptake. However, Na₂SO₄ affected plant growth more adversely than NaCl. Furthermore, tomato plants can die off at prolong subjection to salinity especially Na₂SO₄ stress.

Keywords —Salinity, NaCl, Na₂SO₄, *Lycopersicon esculentum*, growth, morphology.

I. INTRODUCTION

Soil Salinity is considered a major factor threatening crop production in arid and semi – arid regions, where soil salt content is naturally high and precipitation can be insufficient for leaching [1,2]. Salt accumulation in agrarian soil has been aggravated by the requirement of irrigation for crop production in arid and semi-arid environments. Salinity problems increases as salt in the soil increases due to irrigation [3]. It is estimated that at least 20% of all irrigated lands are salt-affected [4]. Even though only around 17% of the land used for cultivation is irrigated, irrigated agriculture accounts for more than 30% of all agricultural production. [5].

Salinity decreases the productivity of agricultural land and jeopardises the sustainability of agriculture [6]. Complex morphological, physiological, and metabolic changes occur in plants as a result of high salinity [7,8]. The higher ratios of toxic salts in the leaf can lead to dehydration and turgor loss, and death of leaf cells and tissues [9].

Tomato (*Lycopersicon esculentum* Mill) is edible, often red fruit/berry of the night shade family Solanaceae [10] Renato Vicario, 2014). Tomato is an important agricultural commodity worldwide. More than 80% of tomatoes are consumed in the form of processed products such as tomato juice, paste, ketchup, sauce, and salsa or as a salad or as cooked. Lycopene is responsible for the characteristic deep-red colour of ripe tomato fruits and tomato products [11]. Tomato is rich in vitamins [12], minerals and lycopene, an excellent antioxidant [13] that helps to reduce the risk of prostate and breast cancer [14].

A comprehensive review of research carried out up to 1999 on the responses of tomato to salinity has been presented by Cuartero and Munoz (1999) [15]. As indicated in previous investigations, the tomato plant is moderately sensitive to salinity [16] (Maas and Hoffman, 1977). The effect of salinity

on the germination, vegetative growth, and yield of Tomato has been studied by a number of researchers [17-19; 15, 20]. Although extensive work has been carried out on the effects of salinity on tomato (*Lycopersicon esculentum*) using NaCl as a source of salinity. However, little or no researched has been carried out using Na₂SO₄ in some cultivated tomato plants. This work, therefore, aim to analyze the effect of NaCl and Na₂SO₄ at different concentrations on the morphological features of Tomato plant.

II. Materials and Methods

The Experiments was conducted in a greenhouse at the Department of Botany, Faculty of Science, University of Ibadan, located at Lat. 003° 53' 47.1" and Long. 07° 26' 35.0" N. Tomato seeds were planted in experimental pots of 15cm diameter and 17 cm depth; each pot was filled with 5.0 kg of soil.

NaCl and Na₂SO₄ with varied equimolar concentration (0 mM, 40 mM, 80 mM, 120 mM, and 160 mM) were applied twice a week except for the control (0 mM) that was only watered occasionally with distilled water. After approximately nine weeks of treatment, plant were harvested and analyzed for various growth characters.

2.1 Plant Morphology and Biomass

Plant morphological characters such as plant height and number of leaves were monitored at weekly interval; Root length, Shoot and root fresh weight were obtained at end of the experiment. Plant height was determined using a meter rule while the number of leaves were counted and recorded on weekly basis.

Plant shoots and roots were recorded at the time of harvest. For fresh shoot weight, plants were washed with distilled water and prior to weighing: excess water was removed with paper towels. For fresh root weight, roots were removed from soil with running tap water: excess moisture was absorbed in

paper towels prior to weighing. Shoot and root dry weights were determined after drying them at 80°C till constant weight in air dried oven.

2.2 Determination of Relative Growth Rate

Relative growth rate (RGR) which is the change in relative growth over time was determined using the formula propose by Wareing and Philips (1981) [21]

$$RGR = \frac{\log n_2 - \log n_1}{t_2 - t_1}$$

Where n_2 -final plant height (cm) at t_2 – final time (day) and n_1 –initial plant height (cm) at t_1 – initial time (day).

2.3 Experimental Design and Statistical Analysis

All experiments were laid out in Completely Randomized Design (CRD) with three replications. The data collected were analyzed using Analysis of Variance with Statistical Analysis System (SAS). The means were compared by Duncan Multiple Range Test (DMRT) at 95% level of pr3.

3.0 RESULTS

3.1 Effect of varying concentrations of NaCl and Na₂SO₄ on plant Morphology.

The effects of salinity on the plant height and number of leaves are summarized in Tables 1 and 2.

The results in Table 1 showed that Tomato height was significantly affected by different salt treatments. Plant height decreased significantly with increase in the concentrations of the salts applied. One week after treatment, there was no significant difference in the tomato treated with NaCl and control (0mM) but significant difference was observed in both tomatoes treated with Na₂SO₄. Control pots had the tallest plants (17.87cm). However, tomato treated with 120mM and 160mM Na₂SO₄ had significantly shortest

(12.37cm and 12.83cm) plants respectively. Two weeks after treatment, control (0 mM) had the tallest plant height (25.10cm) followed by plants treated with 40mMNaCl (22.63cm) while the shortest (13.20 and 13.77cm) was observed in plants treated with 120mM and 160mMNa₂SO₄. Third week after treatment, control (0 mM) had the tallest (32.63 cm). No significant difference was observed in both plants treated with NaCl concentrations and 40 mM Na₂SO₄ compare to control (0 mM) although significant difference was observed in plants treated with 80 mM, 120 mM and 160 mM Na₂SO₄ compare to control (0 mM). Fourth week after treatment, height of the tomatoes were significantly affected by both salt treatments. Application of 120mM and 160mM of Na₂SO₄ resulted in stunted growth and death of the plants (4.27cm and 4.67cm). No significant difference was observed in plants treated with 160mM NaCl with those treated with 40 mM and 80 mM of Na₂SO₄. Fifth week after treatment, control (0 mM) had the tallest (49.67 cm) while the shortest (26.00 cm) was observed in plants treated with 80 mM Na₂SO₄. There is no significant difference of pot treated with 160 mM of NaCl and 40 mM Na₂SO₄. Application of 120 mM and 160 mM of Na₂SO₄ resulted in total elimination of treated tomatoes. So, it is clear that highest salinity level 120 mM and 160 mM exerted the maximum drastic effects on plant height as compared to the other salinity levels.

Table 2 shows the effect of different concentrations of NaCl and Na₂SO₄ applications on number of leaves of tomato.

After the first week of NaCl and Na₂SO₄ application, there was no significant difference in the number of leaves in all salt treatments. However, on the second week, The plants treated with 160mM Na₂SO₄ had the fewest leaves (13.00), but the control (0mM) had the highest average number (26.67) of leaves. Both plants treated with various NaCl and 40mM Na₂SO₄ concentrations showed no significant difference by the third week following treatment. Although the control pot (0 mM) had the most leaves on average (40.00), 120 and 160 mM of Na₂SO₄ killed several

duplicate plants, leaving the plants treated with those concentrations with the fewest leaves (3.67 and 2.33). In comparison to the control (0 mM), both plants treated with varying amounts of NaCl and Na₂SO₄ showed significant differences at the fourth week post-treatment. The plant treated with 120mM Na₂SO₄ had the least amount of leaves (2.67) whereas a control (0mM) plant had the most (50.00). On the fifth week following treatment, the control (0 mM) had the most leaves (63.67), while Na₂SO₄ at 80 mM had the fewest leaves (26.00). Salt concentrations of NaCl and Na₂SO₄ were not significantly different. 120 mM and 160 mM of Na₂SO₄ caused complete defoliation and plant death in plants.

Figure 1 shows the effect of NaCl and Na₂SO₄ salinity on relative growth rate of tomato plants. The relative growth rates (RGR) in the two salts declined significantly compare to control (0mM). The relative growth rate of tomato was highest in plants grown at 0mM followed by plants grown at 40mMNaCl. The lowest growth rate was in plants grown at 160mMNaCl and 80mM Na₂SO₄ respectively (Figure 1). Further increase in salinity caused a decreased in RGR to a low value at higher salinities compared to control.

Effect of varying concentration of NaCl and Na₂SO₄ on biomass of tomato plant is presented in (Table 3). Reduction in fresh and dry weights of root and shoot was observed with the increase in salts stress, as the concentrations of NaCl and Na₂SO₄ increased, dry and fresh weight decreased also. The control showed the highest (5.13g) shoot dry mass, followed by 40mMNaCl (2.54g), while 160 mM NaCl (0.82g) had the lowest value. Salt stress also exerted a drastic effect on root growth and development by reducing its mass. All the two salt concentrations caused reduction in root dry weight compared to the control. The plants grown under control (0 mM) exhibited the maximum value (6.73g) for total dry weight as compared to remaining treatments, which indicated that salinity is responsible for the reduction.

Fresh mass of shoots and roots was decreased significantly ($p>0.05$) by all salt treatment (Table 3). The highest shoot fresh weight was obtained from control treatment (24.51g) and lowest shoot fresh weight (6.96 g) was obtained from 160 mM NaCl. Table 3, showed that shoot fresh weight decreased as salt concentration increased in tomato plant. It was also found that as the salt concentration increased, the root fresh weight decreases (Table 3). The highest root fresh weight was obtained in control (11.01g) and the lowest (1.07g) in 160 mM NaCl.

As the concentration of NaCl and Na₂SO₄ increased, dry and fresh weight decreased. The decreased was more pronounced due to Na₂SO₄ treatments compare to NaCl of the same concentrations.


Table 1.Effect of varying concentrations of NaCl and Na₂SO₄ on height (cm) of *Lycopersicon esculentum* Mill.

Salts	Concentrations (mM)	1 ST	2 ND	3 rd	4 TH	5 TH
WEEKS AFTER TREATMENT						
	0	17.87 _a	25.10 ^a	32.63 ^a	41.00 ^a	49.67 ^a
NaCl	40	17.33 _a	22.63 ^a	28.33 ^a	37.00 _b	42.17 ^a
	80	13.70 _a	18.33 ^a	21.17 ^a	28.17 _b	38.00 ^a
	120	15.27 _a	18.83 ^a	23.50 ^a	28.67 _b	34.67 ^b
	160	13.87 _a	17.33 ^a	20.00 ^a	24.67 ^b	27.43 ^b
Na ₂ SO ₄	40	14.87 _b	17.43 ^b	21.33 ^b	25.17 ^b	33.17 ^b
	80	13.20 _c	14.93 ^b	17.83 ^b	22.50 ^b	26.00 ^c
	120	12.37 _c	13. ^c	13.83 ^b	4.27 ^c	0.00 ^d
	160	12.83 _c	13.77 ^c	9.17 ^c	4.67 ^c	0.00 ^d

Each value is a mean of three replicates. Values in the same column with the same letter(s) were not significantly different at $P>0.05$ using Duncan's Multiple Range Test (DMRT).

Table 2. Effect of varying concentrations of NaCl and Na₂SO₄ on Number of Leaves *Lycopersicon esculentum* Mill

Salts	Concentrations (mM)	1 ST	2 ND
	0	17.00 ^a	26.67 ^a
NaCl	40	18.00 ^a	26.67 ^a
	80	14.67 ^a	22.66 ^a
	120	18.33 ^a	29.00 ^a
	160	13.33 ^a	18.33 ^a
Na ₂ SO ₄	40	11.33 ^b	16.67 ^b
	80	11.33 ^b	16.33 ^b
	120	11.00 ^b	13.67 ^b
	160	10.00 ^b	13.00 ^b



NaCl and Na₂SO₄ on Relative GrowthRate of *Lycopersicon esculentum* Mill.

Each value is a mean of three replicates. Values in the same column with the same letter(s) were not significantly different at $P>0.05$ using Duncan's Multiple Range Test (DMRT).

Table 3.Effect of varying concentrations of NaCl and Na₂SO₄ on Biomass components of *Lycopersicon esculentum* Mill.

Salts	Concentration (mM)	RFW	SFW	TFW
NaCl	0	11.01 ^a	24.51 ^a	35.52 ^a
	40	8.55 ^b	18.66 ^b	27.21 ^b
	80	6.55 ^c	15.42 ^{bc}	21.97 ^{bc}
	120	6.32 ^c	10.06 ^{cd}	16.38 ^{cd}
	160	1.07 ^d	6.96 ^d	8.03 ^d
Na ₂ SO ₄	40	27.16 ^a	18.02 ^a	25.01 ^a
	80	5.74 ^a	13.34 ^b	19.08 ^a

Each value is a mean of three replicates. Values in the same column with the same letter(s) were not significantly different at P>0.05 using Duncan's Multiple Range Test (DMRT).

RFW: Root fresh weight SFW:Shoot fresh weight
TFW: Total fresh weight RDW:Root dried weight
SDW: Shoot dried weight TDW:Total dried weight

4. Discussion

Salinity causes several and specific structural changes that affects plant water balance [22, 23]. These structural changes include fewer and smaller leaves, less number of stomata per unit leaf area, thickening of leaf cuticle and wax deposition on leaf surface [24,25].

The continuous increase in the number of leaves in this study agrees with finding of other workers who suggested

Salinity inhibits plant growth for two reasons: first, water deficit and second due to salt-specific or ion-

excess effects [26]. In this work, reductions in plants height and number of leaves in stressed plants were observed, plants subjected to high salinity (40 and 160 mM Na₂SO₄) died off after 2 weeks of treatment. Corroborating the results obtained by Neves et al., 2004 [27], in Umbu plants. These negative effects of salt stress may be due to reduction of both cell division and cell enlargement [28, 29]. Otherwise, inhibition of shoot growth has been considered a whole plant adaptation to salt stress [30,31]. The suppression of growth under salt stress may also be due to direct effects on ion toxicity especially Na⁺ or indirect effects of saline ions that cause soil/plant osmotic imbalance [1, 32]. These results supports those obtained by other researchers [33,34,35, 36].

Although plant height is genetically controlled, environmental factors also have strong influence on the expression of genes as it is very clear from the current work that all salt treatments significantly influenced the plant. Salinity has been reported to negatively limit germination, plant vigor and yield [37]. Salt tolerance has usually been assessed as the percentage biomass production in saline versus control conditions over a prolonged period of time [38]. It is reported that salt stress affects the plant growth and development by influencing fresh and dry weights of roots, shoots along with shoot length development [39]. Salinity causes stunted growth in glycophytes which results reduction in shoot and root fresh weight [40, 41]. The decrease in dry weight of tomato in response to salt stress might be due to several factors like adverse effects of salinity on photosynthesis, reduction in turgor pressure of expanding tissues and salinity response of root to down regulate shoot growth through a long distance signal [40]. These results are in accordance with previous findings that decrease in shoot and root fresh weight might be due to decreased uptake and accumulation of nutrients in the plant body [42].

5. Conclusions

From the results of this study, it could be concluded that salinity stress significantly reduced tomato growth and its biomass. Both NaCl and Na₂SO₄

inhibit growth, metabolic activity, and ion uptake. However, Na₂SO₄ affected plant growth more adversely than NaCl. In addition, prolonging the duration of time to which plant is exposed resulted in more negative effect.

Author Contributions: the conceptualization, supervision and editing of this study was done by Oyetunji O. J., while Methodology, provision of resources, investigation, and manuscript was done by Samuel K B and Folake Elizabeth.

Funding: "This research received no external funding"

Conflicts of Interest: "The authors declare no conflict of interest."

References

- ¹Zhao, J., Shi, K and Wei, F. 2007: research and application of remote sensing techniques in Chinese agricultural statistics. Paper presented at the forth international conference on agricultural statistics, October 22-24.
- ²Isayenkov, S V and Maathuis, F. J. M .2019. Plant Salinity Stress: Many Unanswered Questions Remain. *Front. Plant Sci.* 10:80.
- ³Abdellatef, A.A. 2010.Changes of antioxidative enzymes in salinity tolerance among different wheat cultivars. *Cereal Res Comm.* 38: 43-55.
- ⁴Pitman, M.G. and Lauchli, A. 2002.Global impact of Salinity and agricultural ecosystem. In *Salinity: Environmental-Plants Molecules*, Eds. *Dordrecht Kluwer*, 3-20.
- ⁵Hillel, D. 2000.Salinity Management for sustainable irrigation. The World Bank, Washing, D. C.
- ⁶Mckee K.L, Mendelssohn, I.A and Mateme, M.D 2004. Acute salt marsh dieback in the Mississippi River deltaic plain: a drought-induced phenomenon? *Global Ecol Biogeog.*13:65-73
- ⁷Hilal, M., Zenoff, A.M., Ponessa, G., Moreno, H. and Massa, E.M. 1998. Saline stress alters the temporal patterns of xylem differentiation and alternative oxidase expression in developing Soybean roots. *Plant Physiol.*, 117:695-701.
- ⁸ Ozturk A, Ukara A, Ipek A, Gurbuz B (2004). Effects of salt stress and water deficit on plant growth and essential of Lemon balm (*Melissa officinalis* L.). *Pak. J. Bot.* 36(4): 787-792.
- ⁹Marschner H .1995. Mineral nutrition of higher plants. (2nded). *Academic Press, London*, New York, pp. 861.
- ¹⁰Renato Vicario. 2014. Slow Food upstate
- ¹¹Helyes, L., Lugasi, A., Pogonyi, A. and Pek, Z. 2009. Effect of variety and grafting on lycopene content of tomato (*Lycopersicon lycopersicum* L. Karsten) fruit. *Acta Alimentari.*, 38:27-34.
- ¹²John, D, Suthin, RT, Usha, RS and Udhayakumar, R. 2010. Role of defense enzymes activity in tomato as induced by *Trichoderma virens* against Fusarium wilt caused by *Fusarium oxysporum* F. Sp *Lycopersici*. *J. Biopesticide*, 3, 158-162.
- ¹³Osemwegie, O. O. Ogbenekaro, A O. and Owolo, I. O. 2010.Effect of Pulverized Ganoderma spp. On Sclerotium rolfsii Sacc and Post-harvest tomatoes fruit preservation *J. appl. Sci Rs.* 6. 1794 -1800.
- ¹⁴Giovannucci, E. 1999.Tomatoes, tomato-based products lycopene, and cancer; review of the epidemiologic literature *J. Natl cancer inst.* 17: 91, 317 -331.

- ¹⁵Cuartero, J. and Fernández-Muñoz, M. 1999. Tomato and salinity. *Scientia Horticulturae*. 78:83–125
- ¹⁶Maas, E.V. and Hoffman, G.J. 1997. Crop salt tolerance current assessment. *Journal of Irrigation and Drainage Division*. 103: 115-134.
- ¹⁷Tal, M. and Shannon, M.C. 1983. Salt tolerance in the wild relatives of cultivated tomato: Responses of *Lycopersicon esculentum*, *L. cheesmanii*, *L. peruvianum*, *Solanum pennellii* and F1 hybrids to high salinity. *Aust. J. Plant Physiol.*, 10:109-117.
- ¹⁸Cruz, V. and Cuartero, J. 1990. Effects of salinity at different developmental stages of six genotypes of tomato (*Lycopersicon* Sp). Proceedings of the 11th Eucarpia meeting on Tomato Genetics and Breeding, March 6-8, 1990, Malaga, Spain. Pp:81-86.
- ¹⁹Groot, S. P. and Karssen, C. M. 1992. "Dormancy and germination of Abscic and deficient tomato seeds studied with sitiens Mutant". *Plant Physiol.* 99: 952-958.
- ²⁰Dasgan, H.Y., Aktas, K., Abak and Calcmak, I. 2002. Determination of screening techniques to salinity tolerance in tomatoes and investigation genotype responses. *Plant Sci.* 63: 695-703.
- ²¹Wareing, P.F. and Phillips, I.D. 1981. The control of growth and differentiation in plants, 3rd Edn, Pergamon Press, New York.
- ²²Robinson, S. P., W.J. Downton and J.A. Millhouse. 1983. Photosynthesis and ion content of leaves and isolated chloroplasts of salt stressed spinach. *Plant Physiol.* 73:238-242.
- ²³Rahdari, P. and Hoseini, S. M. 2011. Salinity Stress: A Review. *Tech J Engin & App Sci.*, 1 (3): 63-66.
- ²⁴Yeo, A.R. and Flowers T.J. 1984. Mechanisms of salinity resistance in rice and their role as physiological criteria in plant breeding. p. 151-170. In: Salinity Tolerance in Plants- Strategies for Crop Improvement. R.C. Staples and G.A. Toenniessen (eds.). John Wiley and Sons, New York, USA.
- ²⁵Khan MA, Irwin A, Allan MS (2000). The effect of salinity on the growth, water status and ion content of a leaf succulent perennial halophyte, *Suaeda fruticosa* L.. *Arid Environment Journal*. 45: 73-84.
- ²⁶Munns, R., James, R.A. and Lauchli A. 2006. Approaches to Increasing the Salt Tolerance of Wheat and other Cereals. *Botany Journal of Experimental*, 57: 1025-1043.
- ²⁷Neves, O.S., Carvalho, J.G. and Rodrigues, C.R. 2004. Crescimento e nutrição mineral de mudas de umbuzeiro (*Spondias tuberosa* Arr. Cam.) submetidas a níveis de salinidade em solo, e ao nutriente. *Ciência e Agrotec.* 28: 997–1006.
- ²⁸Yasseen, B.T., Jurjee, J.A. and Sofajy, S.A. 1987. Changes in some growth processes induced by NaCl in individual leaves of two barley cultivars. *Ind J Plant Physiol.* 30:1-6.
- ²⁹Sairam RK, Tyagi A (2004). Physiology and molecular biology of salinity stress tolerance in plants. *Current Sci. J.* 86(3): 407-421.
- ³⁰Mulholland, B.J., Taylor, I.B., Jackson, A.C. and Thompson, A.J. 2003. Can ABA mediate responses of salinity stressed tomato? *Environ. Exp Bot.* 50:17-28.
- ³¹Qaderi, M.M., Kurepin, L.V. and Reid, D.M. 2006. Growth and physiological responses

of canola (*Brassica napus*) to three components of global climate change: temperature, carbon dioxide and drought. *Physiol Plant*. 128:710-721.

CIMMYT hexaploid wheat germplasm using photosynthetic capacity and water relations as selection criteria. *Photosynthetica* 41: 273-280.

- ³²Hajiboland, R, Aliasgharzadeh, A., Laiegh, S.F and Poschenrieder, C. 2010. Colonization with arbuscular mycorrhizal fungi improves salinity tolerance of tomato (*Solanum lycopersicum* L.) plants. *Plant Soil*. 331:313-327.
- ³³Chookhampaeng, S., Pattanagul, W., and Theerakulpsiut, P. 2008. Effects of Salinity on Growth, Activity of Antioxidant Enzymes and Sucrose Content in Tomato (*Lycopersicon esculentum* Mill.) at the Reproductive Stage. *Science Asia* 34:69-75.
- ³⁴Shores, h M., Spivak, M and Bernstein, N. 2011. Involvement of calcium- mediated effects on ROS metabolism in the regulation of growth improvement under salinity. *Free Rad Biol MED*. 51: 1221-1234.
- ³⁵Abdel Latef, A.A, Chaoxing, E.I .2011. Effect of Arbuscular mycorrhizal fungi on growth, mineral nutrition, antioxidant enzymes activity and fruit yield of tomato grown under salinity stress. *SciElotic*. 127:228-233.
- ³⁶Khalid, A. and Cai, W . 2011. The effects of mannitol and salinity stress on growth and biochemical accumulations in lemon balm. *Acta Ecological Sinica*. 31:112-120.
- ³⁷Munns, R. and Tester, M. 2008. Mechanisms of salinity tolerance. *Annual Review of Plant Biology*. 59:651-681
- ³⁸Munns, R. 2002. Comparative physiology of salt and water stress. *Plant Cell Environ*. 25(2): 239- 250.
- ³⁹Ashraf, M. and Shahbaz, M. 2003. Assessment of genotypic variation in salt tolerance of early
- ⁴⁰Parida, A. K. and A.B. Das. 2005. Salt tolerance and salinity effects on plants a review. *Ecotoxicology and Environmental Safety* 60: 324-49.
- ⁴¹Hajer, A. S., A. A. Malibari, H. S. Al-Zahrani and O. S. Almaghrabi. 2006. Response of three tomato cultivars to sea water salinity 1. Effect of salinity on theseedling growth. *Afr. J. Biotechnol*. 5:855-861.
- ⁴²Dadkhah, A.R. and Griffiths, H. 2006. The effect of Salinity on growth, inorganic ions and dry matter partitioning in sugar beet cultivars. *J. Agricult. Sci. Technol.*, 8;199-210.