Phytochemicals and their role in disease prevention

Sagar Pal[† 1], Vinit Pal[† 1], Vaibhav Kumar[† 1], Puneet Kumar[*,1]

[1]Bhagwant Institute of Pharmacy, Bhagwantpuram,17th Milestone, Muzaffarnagar-251315, Uttar Pradesh, India

[†]Email ID: - sagarpal99170@gmail.com

[*]Corresponding Author Email ID: - puneetatpharmacy@gmail.com

Abstract

Phytochemicals are naturally occurring compounds in plants that contribute to their color, flavor, and defense mechanisms. These bioactive substances are linked to a reduced risk of chronic diseases such as cancer, heart disease, and diabetes. Common classes of phytochemicals include flavonoids, carotenoids, and polyphenols, many of which are abundant in everyday foods. Their health-promoting effects are largely attributed to their antioxidant properties, anti-inflammatory actions, and ability to influence gene expression. Numerous studies have demonstrated the protective potential of phytochemicals against various conditions, including cardiovascular diseases, neurological disorders, and several forms of cancer. Recent research has brought new insights into the role of phytochemicals in disease prevention, highlighting advances in understanding their bioactivity, the development of functional foods, and the growing field of nutraceuticals. These findings underscore the promising role of phytochemicals in promoting long-term health and preventing disease. Phytochemicals often exhibit synergistic effects, meaning their combined action can produce greater health benefits than when consumed individually. This synergy can lead to enhanced nutrient absorption, increased antioxidant activity, and more targeted disease prevention. However, studying the role of phytochemicals in disease prevention presents several challenges, including issues related to bioavailability, variability in individual responses, complex compound interactions, standardization, and optimal dosing. Despite these limitations, ongoing research into phytochemicals has the potential to revolutionize public health. By promoting plant-based diets and developing functional foods designed to prevent chronic illnesses, phytochemical science can contribute significantly to global wellness. A deeper understanding of their mechanisms will enable the development of targeted strategies aimed at reducing disease

incidence, enhancing overall health, and supporting sustainable, preventative healthcare practices worldwide.

Keywords: Phytochemicals, Polyphenols, Disease Prevention, Alkaloids, Terpenoids

Introduction

Phytochemicals are naturally occurring chemical compounds produced by plants, primarily as secondary metabolites. These compounds play a crucial role in plant growth, reproduction, and defense mechanisms. They are classified into several groups, including phenolics, terpenoids, alkaloids, flavonoids, and saponins.[1] The production of phytochemicals is closely linked to the plant's metabolic processes. While primary metabolites—such as sugars and amino acids—are vital for essential functions like photosynthesis and respiration, secondary metabolites like phytochemicals are synthesized to help plants adapt to environmental stresses, resist pathogens, deter herbivores, and attract pollinators. For instance, anthocyanins contribute to the bright colors of flowers and fruits, aiding in pollination and seed dispersal.[2]

The functions and biosynthesis of phytochemicals have been extensively documented through historical studies and review papers. For example, research has highlighted their use in traditional medicine, where compounds with anti-inflammatory properties, such as salicin from willow bark, were utilized.[3] Additionally, the potential applications of secondary metabolites in the food, pharmaceutical, and nutraceutical industries have been explored. The health benefits of phytochemicals, particularly their ability to prevent chronic conditions such as diabetes, obesity, cancer, and cardiovascular diseases, have garnered significant attention. Their antibacterial, anti-inflammatory, and antioxidant properties make them valuable in functional foods and nutraceuticals.[4] Furthermore, studies have underscored their importance in both modern dietary practices and traditional medicine.

By examining their modes of action, including antibacterial, anti-inflammatory, and antioxidant properties, this review seeks to explore the potential role of phytochemicals in disease prevention. This involves understanding how these plant-derived substances help reduce the risk of chronic conditions such as diabetes, cardiovascular diseases, and cancer[56].

Classification of Major Groups of Phytochemicals

The chemical structure and biological activity of phytochemicals allow them to be broadly classified into several primary categories. Some notable examples include:

1. **Polyphenols**: This group includes flavonoids, phenolic acids, and tannins, which are found in foods like tea, wine, fruits, and vegetables. Known for their potent antioxidant properties, they play a significant role in protecting cells from oxidative stress (e.g., catechins in tea, resveratrol in wine).[7]

2. **Carotenoids**: These pigments provide the red, orange, and yellow colors in plants. Beta-carotene, found in carrots, and lycopene, present in tomatoes, are two prominent examples. They are important for their antioxidant effects and contribute to eye health and cancer prevention.[8]

3. **Alkaloids**: These nitrogen-containing compounds include substances like morphine, nicotine, and caffeine. They are known for their diverse biological activities and are commonly found in plants such as poppies (morphine), tobacco (nicotine), and coffee (caffeine)[9,10].

4. **Terpenoids**: These compounds, which include essential oils like camphor and menthol, are responsible for the distinctive scents of many plants. For example, camphor is found in the camphor tree, while menthol is present in peppermint. These terpenoids are commonly found in herbs, spices, and aromatic plants like basil, rosemary, and lavender, contributing to their fragrance and potential therapeutic effects.[11,12]
5. **Saponins**: Found in legumes and certain herbs, saponins are glycosides with soap-like properties. For example, they are present in beans, chickpeas, and quinoa. These compounds are believed to help lower cholesterol levels and may offer other health benefits, such as supporting immune function.[13,14]
6. **Glucosinolates**: Renowned for their cancer-preventive properties, glucosinolates are found in cruciferous vegetables such as broccoli, cabbage, and kale. These compounds, including sulforaphane, are thought to support detoxification processes and protect against carcinogenesis[15,16].




Figure 1. Phytochemicals from the plants

Phytochemicals may help prevent diseases through the following general mechanisms:

1. **Antioxidant Properties**: Phytochemicals like flavonoids and carotenoids help reduce oxidative stress by scavenging free radicals. This action lowers the risk of chronic diseases such as cancer and heart disease. For example, the antioxidant properties of beta-carotene in carrots and lycopene in tomatoes contribute to their protective effects.[17–19]
2. **Anti-inflammatory Effects**: Compounds such as polyphenols can reduce the production of pro-inflammatory cytokines, like TNF-α and IL-6, thereby inhibiting inflammatory pathways. This mechanism plays a role in managing conditions such as inflammatory bowel disease and arthritis. For instance, the polyphenol resveratrol found in grapes has been shown to have anti-inflammatory effects.[20–22]
3. **Modulation of Cell Signaling Pathways**: Phytochemicals regulate key cell signaling pathways, such as NF-κB and MAPK, which are involved in inflammation and cell survival. By modulating these pathways, phytochemicals can prevent the progression of diseases like cancer. For example, curcumin in turmeric is known to affect these pathways.[23–26]

4. **Impact on Gene Expression**: Some phytochemicals influence gene expression by affecting epigenetic mechanisms, including DNA methylation and histone modification. This regulation can suppress the genes associated with disease progression. For example, compounds like sulforaphane from broccoli have been linked to gene expression changes that help prevent cancer.[16,27,28]

Specific Disease Areas and Relevant Phytochemicals: Phytochemicals play a crucial role in the management and prevention of various diseases. Below is an overview of specific disease areas and the phytochemicals associated with them:

1. **Cardiovascular Diseases**: Flavonoids, found in tea and berries, are renowned for their anti-inflammatory and antioxidant properties. These compounds help reduce the risk of atherosclerosis, improve endothelial function, and decrease oxidative stress, thereby supporting heart health.[29–31]
2. **Cancer**: Lycopene, a carotenoid present in tomatoes, has strong antioxidant properties and has been linked to a reduced risk of prostate cancer. Additionally, cruciferous vegetables like broccoli contain isothiocyanates, which are known for their ability to inhibit cancer cell growth and activate detoxifying enzymes.[32–34]
3. **Neurodegenerative Disorders**: Curcumin, the active compound in turmeric, is known for its antioxidant and anti-inflammatory properties. Research is exploring its potential to slow the progression of neurodegenerative diseases such as Alzheimer's.[35–37]
4. **Diabetes**: Resveratrol, found in grapes, has been shown to improve insulin sensitivity, reduce oxidative stress, and influence glucose metabolism. These effects suggest its potential as a therapeutic agent for managing diabetes.[38,39]

Bioavailability and metabolism of phytochemicals

The chemical structure of phytochemicals, their dietary matrix, and interactions with digestive enzymes all play a significant role in their bioavailability and metabolism. Key points include:

1. **Bioavailability**: Phytochemicals in complex forms, such as glycosides, require enzymatic hydrolysis for absorption. Factors like solubility, molecular size, and the presence of

transporters in the intestinal lining influence their bioavailability. For example, after enzymatic breakdown, polyphenols are absorbed as aglycones.

2. **Metabolism**: Once absorbed, phytochemicals undergo extensive metabolism in the gut microbiota and liver. This metabolic process can alter their structure and potentially affect their biological activity. For instance, flavonoids are metabolized into smaller phenolic acids, which may provide unique health benefits.
3. **Factors Affecting Bioavailability**: The absorption of certain phytochemicals can be influenced by the dietary matrix, which includes proteins and lipids, either enhancing or hindering their bioavailability. Additionally, the metabolism of these substances is significantly affected by individual variations in gut microbiota composition, which can vary from person to person.[1,40]

Potential synergistic effects of multiple phytochemicals:

When multiple phytochemicals work together to enhance their therapeutic effects beyond the sum of their individual actions, this is known as synergistic effects. This phenomenon is particularly significant in plant-based nutrition and medicine. Key points include:

1. **Improved Bioavailability:** Some phytochemicals can enhance the absorption and effectiveness of other compounds. For example, piperine in black pepper is known to increase the bioavailability of curcumin in turmeric, making it more effective.

2. **Complementary Mechanisms:** When different phytochemicals target distinct biological pathways, they can produce a more comprehensive therapeutic effect. For instance, combining antioxidants like carotenoids and flavonoids can provide broader protection against oxidative stress.

3. **Decreased Negative Effects:** Synergistic combinations may reduce the required dosage of individual substances, thereby minimizing potential side effects. This is often observed in herbal formulations that combine multiple plant-based compounds.

4. **Enhanced Efficacy**: Research has demonstrated that mixtures of phytochemicals can exhibit stronger antimicrobial, anti-inflammatory, or anticancer activities than individual compounds alone.[41–44]

Current research trends:

The current focus of phytochemical research is on understanding the mechanisms of action and health benefits of these compounds. Here's a summary:

1. **Epidemiological Research**: These studies explore the link between diets rich in phytochemicals and a reduced risk of chronic diseases. For example, plant-based diets abundant in fruits, vegetables, and whole grains have been associated with a lower risk of cardiovascular disease and certain cancers.

2. **In Vitro and Animal Studies**: Laboratory and animal studies examine the specific actions of phytochemicals. Research on carotenoids and flavonoids, for instance, has demonstrated their anti-inflammatory and antioxidant properties, suggesting potential therapeutic applications.

3. **Clinical Trials**: Human clinical trials are increasingly focusing on the effectiveness of phytochemical-based therapies. Studies on compounds like resveratrol and curcumin are investigating their potential to treat diseases such as cancer, diabetes, and neurological disorders.[31,45]

Challenges in phytochemical research:

Phytochemical research faces several challenges, including:

1. **Isolating Active Compounds from Whole Foods**: Extracting specific phytochemicals from complex food matrices can be challenging. Bioactive compounds often need to be isolated and identified using advanced techniques such as spectroscopy and chromatography.

2. **Determining Optimal Doses**: The ideal dosage for therapeutic benefits can be difficult to determine due to variations in bioavailability and metabolism. Factors such as individual health status and interactions within the food matrix further complicate dosage estimation.

3. **Considering Individual Genetic Variations**: Genetic differences can influence the absorption, metabolism, and efficacy of phytochemicals. As a result, personalized nutrition approaches are being explored to tailor phytochemical intake to individual genetic profiles.

Conclusion

Phytochemicals, as naturally occurring plant compounds, hold immense promise in the prevention and management of chronic diseases due to their potent antioxidant, anti-inflammatory, and gene-regulating properties. Their diverse mechanisms—ranging from neutralizing free radicals to modulating cell signaling pathways—underscore their therapeutic potential in conditions such as cancer, cardiovascular disease, diabetes, and neurodegenerative disorders. Despite challenges related to bioavailability, dosage optimization, and individual variability, ongoing research continues to shed light on their multifaceted health benefits. Emphasizing the consumption of phytochemical-rich foods and further exploring their synergistic interactions can significantly contribute to the development of effective, plant-based preventative healthcare strategies. As science advances, phytochemicals are poised to play a pivotal role in shaping future approaches to global wellness and sustainable health.

References

(1) Choudhari, A. S.; Mandave, P. C.; Deshpande, M.; Ranjekar, P.; Prakash, O. Phytochemicals in Cancer Treatment: From Preclinical Studies to Clinical Practice. *Front. Pharmacol.* **2019**, *10*, 1614. https://doi.org/10.3389/fphar.2019.01614.

(2) Epifano, F.; Genovese, S.; Menghini, L.; Curini, M. Chemistry and Pharmacology of Oxyprenylated Secondary Plant Metabolites. *Phytochemistry* **2007**, *68* (7), 939–953. https://doi.org/10.1016/j.phytochem.2007.01.019.

(3) De Cássia Da Silveira E Sá, R.; Andrade, L. N.; De Sousa, D. P. A Review on Anti-Inflammatory Activity of Monoterpenes. *Molecules* **2013**, *18* (1), 1227–1254. https://doi.org/10.3390/molecules18011227.

(4) Gigant, B.; Cormier, A.; Dorléans, A.; Ravelli, R. B. G.; Knossow, M. Microtubule-Destabilizing Agents: Structural and Mechanistic Insights from the Interaction of Colchicine and Vinblastine with Tubulin. *Top. Curr. Chem.* **2009**, *286*, 259–278. https://doi.org/10.1007/128_2008_11.

(5) Liu, T.; Zhang, L.; Joo, D.; Sun, S.-C. NF-KB Signaling in Inflammation. *Signal Transduct. Target. Ther.* **2017**, *2* (1), 17023. https://doi.org/10.1038/sigtrans.2017.23.

(6) Lin, Z.-Y.; Wu, C.-C.; Chuang, Y.-H.; Chuang, W.-L. Anti-Cancer Mechanisms of Clinically Acceptable Colchicine Concentrations on Hepatocellular Carcinoma. *Life Sci.* **2013**, *93* (8), 323–328. https://doi.org/10.1016/j.lfs.2013.07.002.

(7) Magalhães, P. J.; Dostalek, P.; Cruz, J. M.; Guido, L. F.; Barros, A. A. The Impact of a Xanthohumol-Enriched Hop Product on the Behavior of Xanthohumol and Isoxanthohumol in Pale and Dark Beers: A Pilot Scale Approach. *J. Inst. Brew.* **2008**, *114* (3), 246–256. https://doi.org/10.1002/j.2050-0416.2008.tb00335.x.

(8) Rohmer, M. Methylerythritol Phosphate Pathway. *Compr. Nat. Prod. III* **2020**, 560–590. https://doi.org/10.1016/B978-008045382-8.00702-4.

(9) Johari, S. A.; Mohtar, M.; Syed Mohamad, S. A.; Mohammat, M. F.; Sahdan, R.; Mohamed, A.; Mohamad Ridhwan, M. J. In Vitro Evaluations and In Vivo Toxicity and Efficacy Studies of MFM501 against MRSA. *Biomed Res. Int.* **2017**, *2017*, 8032865. https://doi.org/10.1155/2017/8032865.

(10) Yuan, H.; Ma, Q.; Ye, L.; Piao, G. The Traditional Medicine and Modern Medicine from Natural Products. *Molecules* **2016**, *21* (5). https://doi.org/10.3390/molecules21050559.

(11) Porras, G.; Chassagne, F.; Lyles, J. T.; Marquez, L.; Dettweiler, M.; Salam, A. M.; Samarakoon, T.; Shabih, S.; Farrokhi, D. R.; Quave, C. L. Ethnobotany and the Role of Plant Natural Products in Antibiotic Drug Discovery. *Chem. Rev.* **2021**, *121* (6), 3495–3560. https://doi.org/10.1021/acs.chemrev.0c00922.

(12) Friedrich, S.; Serbian, I.; Hoenke, S.; Kancana, R.; René, W. CHEMISTRY Synthesis and Cytotoxic Evaluation of Malachite Green Derived Oleanolic and Ursolic Acid Piperazineamides. *Med. Chem. Res.* **2020**, 926–933. https://doi.org/10.1007/s00044-020-02536-1.

(13) Pandey, P.; Mehta, R.; Upadhyay, R. Physico-Chemical and Preliminary Phytochemical Screening of Psoralea Corylifolia. *Sch. Res. Libr. Arch. Appl. Sci. Res.* **2013**, *5* (2), 261–265.

(14) Shamsudin, N. F.; Ahmed, Q. U.; Mahmood, S.; Ali Shah, S. A.; Khatib, A.; Mukhtar, S.; Alsharif, M. A.; Parveen, H.; Zakaria, Z. A. Antibacterial Effects of Flavonoids and Their

Structure-Activity Relationship Study: A Comparative Interpretation. *Molecules (Basel, Switzerland)*. Drug Design and Synthesis Research Group, Department of Pharmaceutical Chemistry, Kulliyyah of Pharmacy, International Islamic University Malaysia, Kuantan 25200, Pahang D. M., Malaysia. 2022, p 1149. https://doi.org/10.3390/molecules27041149.

(15) Yonekura-Sakakibara, K.; Saito, K. Functional Genomics for Plant Natural Product Biosynthesis. *Nat. Prod. Rep.* **2009**, *26* (11), 1466–1487. https://doi.org/10.1039/b817077k.

(16) Clark, S. M.; Vaitheeswaran, V.; Ambrose, S. J.; Purves, R. W.; Page, J. E. Transcriptome Analysis of Bitter Acid Biosynthesis and Precursor Pathways in Hop (Humulus Lupulus). *BMC Plant Biol.* **2013**, *13* (1). https://doi.org/10.1186/1471-2229-13-12.

(17) Chen, Y.-F.; Wu, S.-N.; Gao, J.-M.; Liao, Z.-Y.; Tseng, Y.-T.; Fülöp, F.; Chang, F.-R.; Lo, Y.-C. The Antioxidant, Anti-Inflammatory, and Neuroprotective Properties of the Synthetic Chalcone Derivative AN07. *Molecules* **2020**, *25* (12). https://doi.org/10.3390/molecules25122907.

(18) Checker, R.; Sandur, S. K.; Sharma, D.; Patwardhan, R. S.; Jayakumar, S.; Kohli, V.; Sethi, G.; Aggarwal, B. B.; Sainis, K. B. Potent Anti-Inflammatory Activity of Ursolic Acid, a Triterpenoid Antioxidant, Is Mediated through Suppression of NF-KB, AP-1 and NF-AT. *PLoS One* **2012**, *7* (2). https://doi.org/10.1371/journal.pone.0031318.

(19) Littarru, G. P.; Tiano, L. Bioenergetic and Antioxidant Properties of Coenzyme Q10: Recent Developments. *Mol. Biotechnol.* **2007**, *37* (1), 31–37. https://doi.org/10.1007/s12033-007-0052-y.

(20) Sharma, S.; Gupta, S.; Khajuria, V.; Bhagat, A.; Ahmed, Z.; Shah, B. A. Analogues of Boswellic Acids as Inhibitors of Pro-Inflammatory Cytokines TNF-α and IL-6. *Bioorganic Med. Chem. Lett.* **2016**, *26* (2), 695–698. https://doi.org/10.1016/j.bmcl.2015.11.035.

(21) Cattoor, K. O.; Bracke, M.; Deforce, D.; De Keukeleire, D.; Heyerick, A. Transport of Hop Bitter Acids across Intestinal Caco-2 Cell Monolayers. *J. Agric. Food Chem.* **2010**,

58 (7), 4132–4140. https://doi.org/10.1021/jf904079h.

(22) Ghate, M.; Kusanur, R. A.; Kulkarni, M. V. Synthesis and in Vivo Analgesic and Anti-Inflammatory Activity of Some Bi Heterocyclic Coumarin Derivatives. *Eur. J. Med. Chem.* **2005**, *40* (9), 882–887. https://doi.org/10.1016/j.ejmech.2005.03.025.

(23) Gallo, C.; Dallaglio, K.; Bassani, B.; Rossi, T.; Rossello, A.; Noonan, D. M.; D'Uva, G.; Bruno, A.; Albini, A. Hop Derived Flavonoid Xanthohumol Inhibits Endothelial Cell Functions via AMPK Activation. *Oncotarget* **2016**, *7* (37), 59917–59931. https://doi.org/10.18632/oncotarget.10990.

(24) Sun, Y.; Liu, Y.; Ma, X.; Hu, H. The Influence of Cell Cycle Regulation on Chemotherapy. *Int. J. Mol. Sci.* **2021**, *22* (13). https://doi.org/10.3390/ijms22136923.

(25) Chakraborty, S.; Kumar, A.; Faheem, M. M.; Katoch, A.; Kumar, A.; Jamwal, V. L.; Nayak, D.; Golani, A.; Rasool, R. U.; Ahmad, S. M.; Jose, J.; Kumar, R.; Gandhi, S. G.; Dinesh Kumar, L.; Goswami, A. Vimentin Activation in Early Apoptotic Cancer Cells Errands Survival Pathways during DNA Damage Inducer CPT Treatment in Colon Carcinoma Model. *Cell Death Dis.* **2019**, *10* (6), 467. https://doi.org/10.1038/s41419-019-1690-2.

(26) Deng, N.; Qiao, M.; Li, Y.; Liang, F.; Li, J.; Liu, Y. Anticancer Effects of Licochalcones: A Review of the Mechanisms. *Front. Pharmacol.* **2023**, *14* (January), 1–10. https://doi.org/10.3389/fphar.2023.1074506.

(27) You, H. J.; Choi, C. Y.; Kim, J. Y.; Park, S. J.; Hahm, K. S.; Jeong, H. G. Ursolic Acid Enhances Nitric Oxide and Tumor Necrosis Factor-Alpha Production via Nuclear Factor-KappaB Activation in the Resting Macrophages. *FEBS Lett.* **2001**, *509* (2), 156–160. https://doi.org/10.1016/s0014-5793(01)03161-1.

(28) Madar, I. H. Identification of Marker Genes in Alzheimer's Disease Using a Machine-Learning Model. *Bioinformation* **2021**, *17* (2), 363–368. https://doi.org/10.6026/97320630017363.

(29) Batovska, D.; Todorova, I. Trends in Utilization of the Pharmacological Potential of Chalcones. *Curr. Clin. Pharmacol.* **2010**, *5* (1), 1–29.

https://doi.org/10.2174/157488410790410579.

(30) Mlala, S.; Oyedeji, A. O.; Gondwe, M.; Oyedeji, O. O. Ursolic Acid and Its Derivatives as Bioactive Agents. *Molecules* **2019**, *24* (15), 1–25. https://doi.org/10.3390/molecules24152751.

(31) Li, C.-C.; Wang, T.-L.; Zhang, Z.-Q.; Yang, W.-Q.; Wang, Y.-F.; Chai, X.; Wang, C.-H.; Li, Z. Phytochemical and Pharmacological Studies on the Genus Psoralea: A Mini Review. *Evidence-Based Complement. Altern. Med.* **2016**, *2016* (1), 8108643. https://doi.org/https://doi.org/10.1155/2016/8108643.

(32) Kumar, A.; Singh, B.; Mahajan, G.; Sharma, P. R.; Bharate, S. B.; Mintoo, M. J.; Mondhe, D. M. A Novel Colchicine-Based Microtubule Inhibitor Exhibits Potent Antitumor Activity by Inducing Mitochondrial Mediated Apoptosis in MIA PaCa-2 Pancreatic Cancer Cells. *Tumour Biol. J. Int. Soc. Oncodevelopmental Biol. Med.* **2016**, *37* (10), 13121–13136. https://doi.org/10.1007/s13277-016-5160-5.

(33) Blanquer-Rosselló, M. M.; Oliver, J.; Valle, A.; Roca, P. Effect of Xanthohumol and 8-Prenylnaringenin on MCF-7 Breast Cancer Cells Oxidative Stress and Mitochondrial Complexes Expression. *J. Cell. Biochem.* **2013**, *114* (12), 2785–2794. https://doi.org/10.1002/jcb.24627.

(34) Boyle, P. Triple-Negative Breast Cancer: Epidemiological Considerations and Recommendations. *Ann. Oncol. Off. J. Eur. Soc. Med. Oncol.* **2012**, *23 Suppl 6*, vi7-12. https://doi.org/10.1093/annonc/mds187.

(35) Mellado, M.; Salas, C. O.; Uriarte, E.; Viña, D.; Jara-Gutiérrez, C.; Matos, M. J.; Cuellar, M. Design, Synthesis and Docking Calculations of Prenylated Chalcones as Selective Monoamine Oxidase B Inhibitors with Antioxidant Activity. *ChemistrySelect* **2019**, *4* (26), 7698–7703. https://doi.org/10.1002/slct.201901282.

(36) Sasazawa, Y.; Kanagaki, S.; Tashiro, E.; Nogawa, T.; Muroi, M.; Kondoh, Y.; Osada, H.; Imoto, M. Xanthohumol Impairs Autophagosome Maturation through Direct Inhibition of Valosin-Containing Protein. *ACS Chem. Biol.* **2012**, *7* (5), 892–900. https://doi.org/10.1021/cb200492h.

(37) Barber, K.; Mendonca, P.; Evans, J. A.; Soliman, K. F. A. Antioxidant and Anti-Inflammatory Mechanisms of Cardamonin through Nrf2 Activation and NF-KB Suppression in LPS-Activated BV-2 Microglial Cells. *Int. J. Mol. Sci.* **2023**, *24* (13). https://doi.org/10.3390/ijms241310872.

(38) Dzobo, K. The Role of Natural Products as Sources of Therapeutic Agents for Innovative Drug Discovery. *Comprehensive Pharmacology*. 2022, pp 408–422. https://doi.org/10.1016/B978-0-12-820472-6.00041-4.

(39) Sun, H.; Wang, D.; Song, X.; Zhang, Y.; Ding, W.; Peng, X.; Zhang, X.; Li, Y.; Ma, Y.; Wang, R.; Yu, P. Natural Prenylchalconaringenins and Prenylnaringenins as Antidiabetic Agents: α-Glucosidase and α-Amylase Inhibition and in Vivo Antihyperglycemic and Antihyperlipidemic Effects. *J. Agric. Food Chem.* **2017**, *65* (8), 1574–1581. https://doi.org/10.1021/acs.jafc.6b05445.

(40) Borris, R. P. Natural Products Research: Perspectives from a Major Pharmaceutical Company. *J. Ethnopharmacol.* **1996**, *51* (1–3), 29–38. https://doi.org/10.1016/0378-8741(95)01347-4.

(41) Wen, L.; Zhou, T.; Jiang, Y.; Gong, L.; Yang, B. Identification of Prenylated Phenolics in Mulberry Leaf and Their Neuroprotective Activity. *Phytomedicine* **2021**, *90*, 153641. https://doi.org/10.1016/j.phymed.2021.153641.

(42) Badshah, S. L.; Faisal, S.; Muhammad, A.; Poulson, B. G.; Emwas, A. H.; Jaremko, M. Antiviral Activities of Flavonoids. *Biomed. Pharmacother.* **2021**, *140*, 111596. https://doi.org/https://doi.org/10.1016/j.biopha.2021.111596.

(43) Jia, W.; Gao, W.-Y.; Yan, Y.-Q.; Wang, J.; Xu, Z.-H.; Zheng, W.-J.; Xiao, P.-G. The Rediscovery of Ancient Chinese Herbal Formulas. *Phytother. Res.* **2004**, *18* (8), 681–686. https://doi.org/10.1002/ptr.1506.

(44) Combination, T.; Nf-, B.; Szaefer, H.; Cykowiak, M.; Krajka-ku, V.; Kleszcz, R.; Kuci, M.; Piotrowska-kempisty, H.; Plewi, A. Pathways in HepG2 Cells In Vitro and Tumor Burden In Vivo. **2021**.

(45) Murali Krishna, P.; Knv, R.; S, S.; Banji, D. A Review on Phytochemical, Ethnomedical

and Pharmacological Studies on Genus Sophora, Fabaceae. *Brazilian J. Pharmacogn.* **2011**, *22*. https://doi.org/10.1590/S0102-695X2012005000043.