

Qualitative phytochemical analysis of fruits of *Shorea robusta* C.F. Gaertn.

Priya Darshani¹, Neelakshi Sharma², Manas Jyoti Kapil³, Rahul Uniyal⁴, Debasmita Mahanta⁵, Nibedita Jena⁶ and Sanjeet Kumar^{6*}

¹Department of Pharmacognosy, Institute of Teaching and Research in Ayurveda, Jamnagar, Gujarat, India

²Department of Pharmacognosy, Royal School of Pharmacy, The Assam Royal Global University, Guwahati, Assam, India

³Department of Pharmaceutics, Royal School of Pharmacy, The Assam Royal Global University, Guwahati, Assam, India

⁴Department of Agroforestry, VCSG UHF College of Forestry, Ranichauri, Tehri Garhwal, Uttarakhand, India

⁵Department of H & T, Government of Odisha, India

⁶Ambika prasad Research Foundation, Odisha, India

*Email-id: sanjeetaprf@gmail.com



Licensed under a Creative Commons Attribution 4.0 International License

Abstract

This study aimed to report the bioactive compounds available in the fruits of *Shorea robusta* C.F. Gaertn. A qualitative phytochemical analysis was conducted using different extracts (aqueous, methanol, and ethanol). The results revealed the presence of various bioactive compounds, including tannins, saponins, flavonoids, phenolic compounds, reducing sugars, and steroids. These compounds have been linked to several medicinal uses, such as treating intestine infections, anti-cancer properties, antimutagenic properties, antidiabetic properties, and anti-inflammatory activities. The study highlights the potential of *S. robusta* fruits as a rich source of phytochemicals with therapeutic applications.

Key words: Anticancer, antidiabetic, antimutagenic, therapeutic applications

Introduction

Shorea robusta, commonly known as Sal (Marandi *et al.*, 2016), is a tree species native to the Indian subcontinent and Southeast Asia (Rahman *et al.*, 2010). The tree is highly valued for its timber, fruits, leaves, and bark. It has been used in traditional medicine for centuries (Yadav and Antil, 2023). The traditional uses of *S. robusta* include treating various ailments such as fever, cough, and skin diseases (Bala *et al.*, 2020). Despite its widespread use in traditional medicine, there is a lack of scientific evidence to support the medicinal properties of *S. robusta* (Singh *et al.*, 2024). The phytochemical composition of the plant, particularly its fruits, remains largely unexplored. Phytochemical analysis is a critical tool in the discovery of novel medicinal compounds from plants (Egbuna *et al.*, 2020). Therefore, it is essential to investigate the phytochemical composition of *S. robusta* fruits to understand their potential medicinal applications. The increasing demand for natural products with medicinal properties has led to a surge in research on secondary metabolites from various plant species (David *et al.*, 2015). However, the phytochemical analysis of *S. robusta* fruits has received little attention. This study aims to bridge this knowledge gap by conducting a qualitative phytochemical analysis of *S. robusta* fruits. The need for this study arises from the potential of *S. robusta* fruits as a rich source of phytochemicals with therapeutic applications. The identification of bioactive compounds in *S. robusta* fruits could provide insight into their potential medicinal uses, such as treating various diseases and disorders. Furthermore, this study could contribute to the development of new drugs and therapies from natural sources, addressing the growing concern of antibiotic resistance and the need for alternative treatments.

Materials and methods

The fruits of *S. robusta* were collected from the Rourkela Forest Division of Sundargarh district of Odisha, India (Figure 2; Kumar *et al.*, 2021). The plant species was identified by the authors (Figure 1). The Soxhlet extraction method was adopted using different solvents for phytochemical analysis (Kumar *et al.*, 2013; Devi *et al.*, 2023; Marandi *et al.*, 2024).

Qualitative phytochemical analysis

Detection of nine secondary metabolites were conducted using standard methods (Thakur *et al.*, 2025).



Figure 1: Different plant parts of *S. robusta*

Test for Tannin

About 1 ml of the fruit extract was taken. Added 3-5 drops of 10% lead acetate solution to it. The formation of gelatinous precipitate confirmed the positive results for the presence of tannin (Kumar *et al.*, 2024).

Test for saponin

About 1 ml of the fruit extract was taken and 1 ml of distilled water was added and shaken well. The persistent froth formation confirmed the presence of saponin.

Test for flavonoids

About 1 ml of the fruit extract was taken. Added 2 ml of 2% NaOH solution and then dilute HCL to it. The colour initially turned to an intense yellow with NaOH solution and later became colourless. This colour changing transformation confirmed for the presence of flavonoids (Ankari *et al.*, 2024).




Figure 2: Dried fruit powder of *S. robusta* for experimental analysis

Test for terpenoids

About 1 ml of the filtrate was added with 6 drops of chloroform and placed in the water bath for a few minutes. Then 6 drops of concentrated sulphuric acid were added. The appearance of reddish-brown interface confirmed the presence of terpenoids.

Test for phenolic compounds

About 1 ml of the fruit extract was taken and added few drops of 5% ferric chloride solution to it. The dark bluish black appearance confirmed the presence of phenolic compounds (Singh *et al.*, 2024).

Test for reducing sugars

About 1 ml of the fruit extract was taken and 2-3 drops of Fehling's solution A and B were respectively added. Then kept in the water bath for some time. The presence of red-orange precipitate confirmed the presence of reducing sugar (Table 1).

Test for steroids

About 1 ml of the fruit extract was taken. 1 ml of chloroform and 1 ml of concentrated sulphuric acid was added into it. The appearance of 2 phases with the upper red and lower yellow with green fluorescence confirmed the presence of steroids (Acharya *et al.*, 2024).

Test for alkaloids

About 1 ml of the extract was taken and added 3 to 4 drops of Dragendorff's reagent. The formation of reddish-brown precipitate confirmed the presence of alkaloids.

Test for carbonyl compounds

About 1 ml of the fruit extract was taken added 3 to 4 drops of 2,4-dinitrophenylhydrazine (DNPH) reagent. The yellow crystal formation confirmed the presence of carbonyl compounds (Jena *et al.*, 2024).

Results and discussion

The results of the phytochemical analysis, as presented in Table 1, revealed the presence of various bioactive compounds in *S. robusta* fruits. The aqueous, methanol, and ethanol extracts contain tannins, saponins, flavonoids, phenolic compounds, reducing sugars, and steroids, with varying degrees of presence across different extracts (Figure 3-5). These compounds have been linked to several medicinal uses, including the treatment of intestine infections, anti-cancer properties (Figure 6), antimutagenic properties, antidiabetic properties, and anti-inflammatory activities, as summarized in Table 2. Overall, the study highlights the potential of *S. robusta* fruits as a rich source of phytochemicals with therapeutic applications.

Table 1: Qualitative phytochemical analysis of *S. robusta* fruits using different extracts

Bioactive compounds	Extracts		
	Aqueous	Methanol	Ethanol
Tannin	+	+	-
Saponin	+	+	+
Flavonoids	-	+	+
Terpenoids	-	-	-

Phenolic compounds	+	+	+
Reducing sugars	+	+	+
Steroids	+	+	+
Alkaloids	-	+	-
Carbonyl compounds	-	-	-

(+: Positive; -:Negative)



Figure 3: Detection of secondary metabolites of *S. robusta* using aqueous extract




Figure 4: Detection of secondary metabolites of *S. robusta* using methanolic extract




Figure 5: Detection of secondary metabolites of *S. robusta* using ethanolic extract

Table 2: Probable medicinal uses of *S. robusta* due to the presence of following bioactive compounds

Bioactive compounds	Medicinal uses	Source
Tannin	Tannin present in aqueous and methanol extracts have potentials to treat intestine infections	Pizzi, (2019)
Saponin	Presence of saponins in aqueous, methanol and ethanol have anti-cancer properties	Man <i>et al.</i> , (2010)
Phenolic compounds	Phenolic groups present in aqueous, methanol and ethanol extracts have antimutagenic properties.	Huang <i>et al.</i> , (2010)
Reducing sugars	Reducing sugars present in aqueous,	McKenzie and Lee, (2022)

	methanol and ethanol extracts have antidiabetic properties	
Steroids	Steroids present in aqueous, methanol and ethanol extracts have anti-inflammatory activities	Williams, (2018)




Figure 6: Pharmacological potential (anticancer activity) of *S. robusta* fruits

Conclusion

The present study has successfully demonstrated the presence of various bioactive compounds in *S. robusta* fruits. The study highlights the potential of *S. robusta* fruits as a rich source of phytochemicals with therapeutic applications. Future studies should focus on the isolation and characterization of the bioactive compounds present in *S. robusta* fruits. In vitro and in vivo studies should be conducted to evaluate the efficacy and safety of these compounds for various medicinal applications. Additionally, clinical trials should be conducted to confirm the therapeutic potential of *S. robusta* fruits. Furthermore, studies on the cultivation and conservation of *S. robusta* trees should be undertaken to ensure the sustainable use of this valuable plant species. Overall, the results of this study provide a foundation for further

research on the medicinal properties of *S. robusta* fruits and their potential applications in the development of new drugs and therapies.

References

- Acharya S, Das K, Soni P, Kaur L, Ray P, Jena N and Kumar S. (2024). Primary and secondary metabolites of kusum seeds. Plants and Secondary Metabolites, Volume 2. Ambika Prasad Research Foundation, India.
- Ankari AB, Mahendru N, Paniyappanayara V, Tripathi G, Jena N and Kumar S. (2024). Evaluation of medicinal values of a fern species [*Thelypteris parasitica* (L.) Tardieu] of India. Plants & Secondary Metabolites, Volume 2, Ambika Prasad Research Foundation, India.
- Bala SK, Parida S and Mahalik G. (2020). Studies on the use of plants and plant parts by the tribals of Malkangiri district, Odisha for different rituals and medicinal uses. Plant Archives. 20(2): 3308-3312.
- David B, Wolfender JL and Dias DA. (2015). The pharmaceutical industry and natural products: historical status and new trends. Phytochemistry Reviews. 14: 299-315.
- Devi RS, Satapathy KB and Kumar S. (2023). Validation of tribal claims for formulation of future drugs through evaluation of ethno-pharmacological values of *Ludwigia adscendens*. Medicinal Plants. 15(4): 691-697.
- Egbuna C, Kumar S, Ifemeje JC, Ezzat SM and Kaliyaperumal S. (2020). Phytochemicals as Lead Compounds for New Drug Discovery. Elsevier Science Publishing Co Inc, Netherlands.
- Huang WY, Cai YZ and Zhang Y. (2010). Natural phenolic compounds from medicinal herbs and dietary plants: potential use for cancer prevention. Nutrition and Cancer. 62(1): 1-20.
- Jena N, Rout S, Devi RS and Kumar S. (2024). Phytochemical and cytotoxicity analysis of fruits of *Zanthoxylum asiaticum* (L.) Appelhans, Groppo & J. Wen: A minor fruit plant of Odisha, India. E-planet. 22(1): 62-70.
- Kumar A, Jadhav JY, Dehar PS, Gupta PSP, Mondal K, Vishwakarma K, Jena N and Kumar S. (2024). Qualitative phytochemical screening of a medicinally-ornamental wild fern species (*Pteris vittata* L.) of India. Plants & Secondary Metabolites, Volume 2. Ambika Prasad Research Foundation, India.
- Kumar S, Behera SP and Jena PK. (2013). Validation of tribal claims on *Dioscorea pentaphylla* through phytochemical screening and evaluation of antibacterial activity. Plant Science Research. 35: 55-61.
- Kumar SN, Mishra S and Kumar S. (2021). Documentation of Indigenous Traditional Knowledge (ITK) on Commonly Available Plants in Koira Range, Bonai Forest Division, Sundargarh, Odisha, India. Asian Plant Research Journal. 8(4): 83-95.

- Man S, Gao W, Zhang Y, Huang L and Liu C. (2010). Chemical study and medical application of saponins as anti-cancer agents. *Fitoterapia*. 81(7): 703-714.
- Marandi RR, Britto SJ and Soreng PK. (2016). Phytochemical profiling, antibacterial screening and antioxidant properties of the sacred tree (*Shorea robusta* Gaertn.) of Jharkhand. *International Journal of Pharmaceutical Sciences and Research*. 7(7): 2874-2888.
- Marndi S, Barik M and Kumar S. (2024). Phytochemical and cytotoxicity analysis of *Eulophia explanata* Lindl. (Tuber extracts). *African Journal of Biological Sciences African Journal of Biological Sciences*. 6(4):2751-2756; DOI: 10.48047/AFJBS.6.Si4.2024.2751-2756.
- McKenzie E and Lee SY. (2022). Sugar reduction methods and their application in confections: a review. *Food Science and Biotechnology*. 31(4): 387-398.
- Pizzi A. (2019). Tannins: prospectives and actual industrial applications. *Biomolecules*. 9(8): 344-344.
- Rahman MM, Rahman M, Guogang Z and Islam KS. (2010). A review of the present threats to tropical moist deciduous sal (*Shorea robusta*) forest ecosystem of Central Bangladesh. *Tropical Conservation Science*. 3(1): 90-102.
- Singh B, Rao K, Aarti, Yadav N, Chauhan J, Jena N and Kumar S. (2024). Phytochemical screening and thin layer chromatography (TLC) of bark of *Alstonia scholaris* (Apocynaceae). *Plants & Secondary Metabolites, Volume 1*. Ambika Prasad Research Foundation, India.
- Singh SV, Johar V and Dhyani A. (2024). A review on importance of sal tree (*Shorea robusta*) as an interminable wood. *Advanced Journal of Molecular Biology and Environmental Sciences*. 26(1): 153-156.
- Thakur K, Jena N and Kumar S. (2025). Pharmacological evaluation of *Vitex altissima* L. *Medico-Biowealth of India, Volume 19*. Ambika Prasad Research Foundation (APRF), India.
- Williams DM. (2018). Clinical pharmacology of corticosteroids. *Respiratory Care*. 63(6): 655-670.
- Yadav P and Antil RS. (2023). Utilizing the multi-faceted potential of *Shorea robusta* in 2023: a comprehensive review. *International Journal for Research in Applied Science & Engineering Technology*. 11(10): 161-168.