

Chapter

Long-Term Observation of *In Vitro*-Derived *Malus Sylvestris* (L.) Mill., the Path from the Bud to the Tree

*Martina Komárková, Helena Cvrčková, Jaroslav Dostál,
Václav Buriánek and Pavlína Máchová*

Abstract

The European wild apple (*Malus sylvestris* L.), a wild contributor to the domesticated apple, belongs to the endangered species in the Czech Republic. Thus, an efficient protocol was developed for *in vitro* plantlet regeneration using the post-dormant buds. The highest shoot induction was obtained on MS medium supplemented with 0.5 mg.l⁻¹ BAP, 5 mg.l⁻¹ GA and 0.1 mg.l⁻¹ IBA. Shoot multiplication and elongation took place on the same medium with 0.2 mg.l⁻¹ BAP and 0.1 mg.l⁻¹ IBA. Indole-3-butyric acid at 0.5 mg.l⁻¹ was most effective for rooting. The micropropagated plantlets were successfully acclimatized in greenhouse conditions and were transplanted into soil in forest. Finally, qualitative and quantitative parameters of tissue culture-derived plants were evaluated. Monitoring of *in vitro* plantings on experimental trials suggests that micropropagated wild apple trees retain the growth characteristics of generative individuals.

Keywords: acclimatization, establishment of *in vitro* culture, long-term forest trial, *Malus sylvestris* L., rooting, shoot multiplication

1. Introduction

M. sylvestris L., the only native wild apple species in Europe, belongs to the family Rosaceae. It is mainly pollinated by bees and flies, and thanks to its small and hard fruits, the trees are often called crab apples. This species occurs across Western and Central Europe, from southern Scandinavia to the Iberian Peninsula and from the Volga to the British Isles [1]. With a high up to 10 m, it grows in low-density populations and the growth habit looks like shrubs, more than trees. Under good growth conditions (high light requirements), the crab apple can live up to 80–100 years [2]. Though many different varieties of apples were developed over time, it was shown that old varieties have higher nutritional quality when compared to commercial ones [3–5].

In the Czech Republic, the wild apple belongs to the endangered species [6] because of its shrinking habitat, fragmentation of populations [7–9] and likelihood of a genetic admixture with domesticated apple (*Malus domestica* Borkh.). Recent

studies have underlined the significant contribution of wild apples to the cultivated apple genome, *M. domestica* Borkh., during the domestication history from the Central Asian progenitor *Malus sieversii* (Ldb.) M. Roem [10]. Because of hybridization, it can be very difficult to reliably identify ‘true type’ wild apples from existing hybrids based on morphological characteristics, such as the fruit width or the hairiness of leaves [2]. Easy hybridization of wild apple with the domesticated apple is due to the absence of prezygotic isolation mechanisms [11], which results in fitness reduction of the wild apple populations and can lead to their reduction or extinction. Moreover, due to human activities, such as forest clearing, industrialization, population increase and intensive agriculture, natural and artificial forest stand regeneration is almost impossible. To avoid loss of wild apple genetic resources, they must be conserved and the conservation strategy should be applied sustainably.

The solution to the problem of preservation and restoration of the gene pool of endangered wild apples could be the *in vitro* cultivation using the micropropagation method. Micropropagation is used to multiply a wide variety of plants by a number of tissues, cells or organ culture methods. It means the aseptic culture of small plant explant of tissue or organs, closed in vessel with defined culture media and under controlled conditions. It provides the large-scale production of disease-free seedlings within a short time and with limited space. In the Czech Republic, biotechnology approaches for *in vitro* conservation of different woody plant species are well established and widely used (e.g., see [12–17]). During the past years, many studies have been carried out using *in vitro* cultures of apples (for review, see [18]). A key objective of apple *in vitro* cultures is to multiply disease-free clones, suitable for rooting, acclimatization and planting. However, as in many other plant species, the medium composition, plant growth regulators requirement or specific growth conditions are cultivar-dependent [19].

Additionally, micropropagation enables the protection of endangered species, thanks to providing the plant material in a larger amount for plant breeding programs at specific sites [20]. However, not many available papers and reviews dealing with studies on tissue cultures of forest trees include subsequent *ex vitro* evaluation of their quantitative and qualitative traits. The aim of our study was to find the optimal composition of explant culture media used for initiation of culture, as well as long-term multiplication and rooting of wild apples *in vitro*. In addition, the growth parameters of *in vitro*-derived wild apple trees were evaluated.

2. Material and methods

2.1 Plant material and *in vitro* culture establishment

M. sylvestris L. trees were carefully determined, not to be mistaken for domesticated apples *M. domestica* Borkh.

Juvenile branches with dormant leaf buds of the crown of *M. sylvestris* trees from central Bohemia were collected and cut into 10–15 cm long twigs. Twigs were immediately packed into the plastic bags and stored at 4°C before being cultured *in vitro*. To decrease microbial contamination, the twigs were cut into 5 cm long segments with buds, rinsed for 30 min in running water and surface disinfected with Tween®20 (2 drops/10 ml) for 20 min, followed by soaking in KORSOLEX for 20 min, rinsed with sterile distilled water for 20 min, incubated in HgCl₂ solution (1 mg.l⁻¹) for 15 min and subsequently immersed in distilled water three times for 15 min.

The surface disinfected bud explants (0.3–0.5 cm large) were placed in glass jars containing 100 ml of culture medium: MS medium [21] supplemented with 0.5 mg l⁻¹ BAP, 0.1 mg l⁻¹ IBA, 10 mg l⁻¹ glutamine, 2 mg l⁻¹ glycine, 30 g l⁻¹ sucrose and 6 g l⁻¹ agar. The pH value was adjusted to 5.8 before autoclaving at 121°C, 150 kPa for 20 min. The aseptic cultures were incubated in a growth chamber at 24 ± 1°C with a 16-h photoperiod (30 µE m⁻² s⁻¹).

2.2 Shoot multiplication

New shoots from explant cultures were separated into stems approximately 2.0 cm long and transferred into shoot induction media. The medium used was MS medium containing 0.2 mg l⁻¹ BAP, 0.1 mg l⁻¹ IBA, 200 mg l⁻¹ glutamine, 2 mg l⁻¹ glycine, 200 mg l⁻¹ casein, 30 g l⁻¹ sucrose, 6 g l⁻¹ agar and final pH adjusted to 5.8. Explants were cultured in the growth conditions as described above and repeatedly subcultured at a constant 4-week subculture interval.

2.3 Root induction, *ex vitro* acclimatization and hardening

Healthy shoots (1.5–2 cm) were excised and cultured on ¼ MS medium supplemented with 0.5 mg l⁻¹ IBA, 10 g l⁻¹ sucrose, 6 g l⁻¹ agar and final pH adjusted to 5.8. Explants were cultured in the growth conditions as described above.

After 2–3 weeks, the rooted shoots were carefully washed with distilled water to remove agar and the rooted shoots were hardened in a culture room in conical planter Quick Pot T 35 with perforated bottom, filled with perlite (Perlite Praha spol. s.r.o., Czech Republic) and placed into the transparent plastic box fully closed. Plants were regularly watered with 1/2 strength liquid MS medium devoid of sucrose and phytohormones and diluted with distilled water in a ration 1:10. After 2–3 weeks, the plants were transferred into the planter Quick Pot T 60 containing soil (Zahradnický substrát a.s. Soběslav, Czech Republic): perlite (2:1) and kept in a bigger transparent plastic box. During 2 weeks, plantlets were adapted to lower moisture conditions by gradually tilting the upper part of the plastic box. Fully adapted plants were moved into greenhouse and subsequently transferred into the outdoor flower beds to obtain capable seedlings.

3. Results

3.1 Plant material

The wild apple trees are characterized by pressed buds, leaves and combs. Only in spring, the leaves can have inconspicuous hairs on the reverse side at the base of the thicker veins. The size of the leaves and fruits are also different, with clearly smaller size in the wild apple tree. *M. sylvestris* L. has leaves of up to 6.5 cm long, while the leaves of the domestic apple tree are 6–12 cm long. The fruits (pome) of the wild apple tree are 2–3.5 cm in diameter, while the apples of the domestic apple tree are at least 5 cm in size. The crown slices of flowers of the wild apple tree are also smaller and tend to be narrower [22].

Juvenile branches with axillary buds were collected during the early spring. In agreement with Ref. [23], buds collected during spring and summer seasons produced a significantly higher percentage of explant establishment and were less contaminated than buds collected during autumn or winter.

3.2 Shoot propagation from the buds

As for surface sterilization of buds, different methods have been described. The disinfection procedures include washing in sodium hypochlorite (NaOCl) solution [24], mercuric chloride (HgCl_2) solution [19], 75% alcohol followed by HgCl_2 solution [25] or calcium hypochlorite [$\text{Ca}(\text{OCl})_2$] solution [26]. In our experiment, we used an HgCl_2 solution (1 mg.l^{-1}), which is highly efficient for surface sterilization of buds from field-grown trees.

As in other plant species, the optimal basal medium is often cultivar-dependent in *Malus* species [27]. Additionally, Kabylbekova et al. [28] have used response surface methodology (RSM) and showed that each apple cultivar requires a different composition of mineral nutrition for its optimal growth. Similarly, the selection of the optimal plant growth regulators is also genotype-dependent, as has been shown in the study of different apple scions and rootstocks [29, 30]. Thus, different media types for bud induction and shoot development were tested. The MS medium with 0.5 mg.l^{-1} BAP, 0.1 mg.l^{-1} IBA, 10 mg.l^{-1} glutamine and 2 mg.l^{-1} glycine was the most effective medium, based on monitoring following parameters: percentage of contamination, percentage of necrotic explants and percentage of explants with shoot initiation. In our experiment, plant growth regulators BAP and IBA were used for culture establishment and also for shoot multiplication, consistently with the studies from Refs. [31] or [28]. However, application of meta-topolin [32] or TDZ [33] in the growth medium was also described in apples.

3.3 Shoot proliferation

Newly developed shoots from bud explants were transferred into shoot induction media (**Figure 1**). Based on determination of multiplication parameters, that is, multiplication index (the number of newly formed shoots per initial shoot tip) and length of lateral shoots, MS medium consisted of 0.2 mg.l^{-1} BAP, 0.1 mg.l^{-1} IBA, 200 mg.l^{-1} glutamine, 2 mg.l^{-1} glycine and 200 mg.l^{-1} casein was chosen. Same as in our experiment, Sota et al. [34] have used MS medium for shoot multiplication of wild apples. However, the BAP concentration was quite higher than in our experiment (1 mg.l^{-1} BAP), and instead of IBA, α -Naphthaleneacetic acid (NAA) was used. Additionally, the application of other growth nutrient media has also been reported [27, 35].

3.4 *In vitro* rooting, acclimatization and hardening

Well-multiplied shoots were subjected to MS medium lacking cytokinins but supplemented with auxins. All the treatments resulted in root production. The highest rooting percentage and roots per cultured shoot were obtained on $\frac{1}{4}$ MS supplemented with 0.5 mg.l^{-1} IBA, 10 g.l^{-1} sucrose and 6 g.l^{-1} agar. Although some authors found that NAA [36] or IAA [37] is more effective for rooting, the application of IBA showed the best results in our experiment.

The rooted shoots obtained from the best treatment (**Figure 2**) were removed from the rooting medium and the plantlets were then transferred to planter with perlite and watered with liquid MS medium. After 2 weeks, plantlets were transferred into bigger planter containing a mixture of perlite and soil and gradually adapted to lower moisture conditions. Finally, fully adapted plants (**Figure 3**) were transferred into greenhouse.



Figure 1.
In vitro shoot multiplication of *M. sylvestris* L.

3.5 Qualitative and quantitative traits of wild apples at experimental trials

The study was conducted at the Oldřichov (425 m a.s.l., central Bohemian Highlands) in 2003 and at Polná II (550 m a.s.l., Czech-Moravian Highlands) in 2007 in the Czech Republic. At both experimental sites, row planting with 2×2 m spacing was used and the plots were fenced off for the entire monitoring period. The subject of the evaluation of quantitative traits was survival rate, height and diameter at breast height (DBH). The DBH was measured using a millimetre calliper, and the height was determined using a measuring rod and a Vertex III ultrasonic altimeter (Haglöf Sweden AB, Langsele, Sweden). As for qualitative traits, trunk shape, forkness, branching angle, branch thickness and vitality were established. The qualitative traits were determined according to the manual in **Table 1**.

In Oldřichov, grafters of wild apples were planted together with *in vitro*-derived plantlets (**Figure 4**). For statistical evaluation, data from 2010 to 2018 were used



Figure 2.
In vitro rooting of *M. sylvestris* L.



Figure 3.
Fully adapted *in vitro*-derived *M. sylvestris* L. plantlets.

Trunk shape		Forkness		Branching angle		Branch thickness		Vitality	
1	Straight	1	Stem continuous to the crown	1	Horizontal	1	Thin (up to 10% DBH)	1	Highly vital
2	Slightly curved on one side	2	Fork in the upper third of the tree's height	2	Ascending	2	Medium thick (10–25% DBH)	2	Vital
3	Strongly curved on one side	3	Fork in the second third of the tree's height	3	Overhanging	3	Thick (over 25% DBH)	3	Less vital
4	Trunk at least 2x Slightly ace-curved	4	Fork in the lower third of the tree					4	Declining tree
5	Significantly bent trunk	5	Repeated multiple forkness						

Table 1.
The list of qualitative traits and their descriptors.



Figure 4.
M. sylvestris L. trees grown at Oldřichov research plot in 2010.

(**Table 2**). The mortality of tissue culture plantlets was zero, while grafters' mortality reached 24%, mostly due to withering of the trees. ANOVA did not show any significant differences in height between *in vitro*-derived plantlets and grafters ($\alpha = 0.05$). However, in 2018, grafters were significantly higher than tissue culture plantlets, according to ANOVA, which could be caused by the growth rate of rootstock. No differences were observed between qualitative traits (**Table 2**). The trunk shape reached

Group	2010					2018											
	Height (m)					Height (m)					DBH (cm)		TrSh	Bra	BrAng	BrTh	Vit
	Count	Mean	Median	StD	Count	Mean	Median	StD	Mean	Median	StD	Median	Median	Median	Median	Median	
<i>In vitro</i>	15	3.03	3.25	0.90	15	6.89	7.35	2.95	5.24	5.80	1.61	3	3	2	2	2	1
Grafters	195	2.79	3.30	1.45	148	8.01	8.63	2.80	7.26	7.50	1.56	3	3	2	2	2	1
All	210	2.80	3.28	1.41	163	7.91	7.40	1.66	7.08	8.40	2.82	3	3	2	2	2	1

Table 2. Average growth parameters of *Malus sylvestris* L. plantlets at Oldřichov research plot. DBH - diameter of breast height, TrSh – trunk shape, Bra – Branching, BrAng – branching angle, BrTh – branch thickness, Vit - vitality and StD – standard deviation.

Clone	2008					2017					2017					2017				
	Height (m)					Height (m)					DBH (cm)					TrSh				
	Count	Mean	Median	StD	Count	Mean	Median	StD	Count	Mean	Median	StD	Count	Mean	Median	Count	Mean	Median	StD	Count
J1	9	1.14 ^{a, b}	1.25	0.34	9	5.03 ^{a, b}	4.9	1.00	9	4.16	4.05	0.93	9	4.16	4.05	9	4.16	4.05	0.93	9
J2	10	1.21 ^{a, b}	1.12	0.29	10	4.88 ^{a, b}	4.95	0.68	10	3.96	3.93	1.01	10	3.96	3.93	10	3.96	3.93	1.01	10
J4	23	0.98 ^b	0.96	0.21	23	3.86 ^{b, c}	3.7	0.78	23	3.14	3.05	1.14	23	3.14	3.05	23	3.14	3.05	1.14	23
J5	18	1.11 ^{a, b}	1.10	0.23	17	4.35 ^{a, b, c}	4.6	0.72	17	3.99	4.1	1.11	17	3.99	4.1	17	3.99	4.1	1.11	17
J6	50	1.15 ^{a, b}	1.15	0.29	50	3.82 ^c	3.75	1.07	50	3.48	3.35	1.71	50	3.48	3.35	50	3.48	3.35	1.71	50
J7	18	0.99 ^b	1.01	0.27	18	3.94 ^{b, c}	4.05	0.61	18	3.26	3.4	1.23	18	3.26	3.4	18	3.26	3.4	1.23	18
J8	12	1.13 ^{a, b}	1.23	0.39	12	5.21 ^a	4.9	0.61	12	4.43	4.28	0.61	12	4.43	4.28	12	4.43	4.28	0.61	12
J9	10	1.20 ^{a, b}	1.22	0.20	10	4.82 ^{a, b, c}	4.8	1.03	10	3.83	4.1	1.39	10	3.83	4.1	10	3.83	4.1	1.39	10
J10	14	0.89 ^b	0.87	0.33	14	4.25 ^{a, b, c}	4.15	1.09	14	3.84	4.28	1.70	14	3.84	4.28	14	3.84	4.28	1.70	14
J12	20	1.10 ^{a, b}	1.06	0.22	20	4.36 ^{a, b, c}	4.5	0.88	20	3.27	3.3	1.37	20	3.27	3.3	20	3.27	3.3	1.37	20
J13	19	1.10 ^{a, b}	1.15	0.22	19	4.37 ^{a, b, c}	4.4	0.77	19	3.53	3.55	1.18	19	3.53	3.55	19	3.53	3.55	1.18	19
J26	20	1.33 ^a	1.40	0.27	20	4.25 ^{a, b, c}	4.4	1.06	20	3.68	3.45	1.62	20	3.68	3.45	20	3.68	3.45	1.62	20
J28	13	1.14 ^{a, b}	1.22	0.20	13	3.98 ^{b, c}	3.8	0.78	13	4.51	4.7	1.63	13	4.51	4.7	13	4.51	4.7	1.63	13
J30	14	1.12 ^{a, b}	1.18	0.25	14	4.25 ^{a, b, c}	4.4	0.94	14	4.27	4.45	1.33	14	4.27	4.45	14	4.27	4.45	1.33	14
All	250	1.11	1.12	0.28	249	4.24	4.30	0.96	249	3.69	3.75	1.42	249	3.69	3.75	249	3.69	3.75	1.42	249

Table 3. Average growth parameters of *Malus sylvestris* plantlets on Pohná II research plot. DBH - diameter of breast height, TrSh - trunk shape, Bra - branching, BrAng - branching angle, BrTh - branch thickness, Vit - vitality and StD - standard deviation. Values followed by the same letter(s) (a-c) within the same column are not significantly different according to Tukey's test ($P < 0.05$).



Figure 5.
Malus sylvestris L. trees grown at Polná II research plot in 2011.

the third degree (strongly curved on one side), same as forkness (forks in the second-third of the tree's height). The second degree was observed in the branching angle (ascending branches) and in the branch thickness (medium thickness with 10–25% DBH). The vitality reached the first degree (highly vital).

The growth characteristics of *in vitro*-derived wild apple plantlets recorded from years 2008 and 2017 grown at Polná II research plot are given in **Table 3**. The mortality was very low, only one individual in clone J5 died between the monitored years. ANOVA showed significant differences in height of wild apples in both years ($\alpha = 0.05$), whereas no differences were observed in DBH. Among the qualitative traits, only small variances were noted. The trunk shape is on the third degree (strongly curved on one side) with the exception of clone J2, where the trunk was at least 2x slightly ace-curved. Forkness reached mostly the third degree. Only for clones J1 and J26, the fourth degree of forkness (forks in the lower third of the tree) was noted, and for clones J8 and J9, an intermediate stage between the third and the fourth degree was noted. All the wild apple clones had ascending branches (the second degree of branch angle). For clone J12, medium branch thickness was recorded (10–25% DBH), whereas all other clones reached the third degree of branch thickness (strong thickness with over 25% DBH). All tested clones were highly vital (**Figure 5**).

The comparison of our results with domestic or foreign studies is very problematic. There is a lack of evidence about the growth characteristics of wild apples, due to their low abundance in nature and difficult determination. Moreover, wild apples grown in our research plots are residual individuals selected in the Czech Republic, with different qualitative traits and growth potential.

4. Conclusions

The results of the presented study indicated that our methodology for the micro-propagation of *M. sylvestris* L. from dormant bud could be efficiently used for *in*

in vitro conservation of endangered wild apples. Moreover, according to long-term experimental trials, *in vitro*-derived plantlets show quality and balanced growth, comparable with grafters. The growth characteristics of wild apples and grafters on the research plots at Oldřichov and Polná II will be still evaluated in the future.

In this regard, implementation of *in vitro* conservation of endangered wild apple *M. sylvestris* L. in practice can speed up the process of protection and reproduction.

Acknowledgements

This research was supported by Institutional Support (MZE-RO0118).

Conflict of interest

The authors declare no conflict of interest.

Author details

Martina Komárková*, Helena Cvrčková, Jaroslav Dostál, Václav Buriánek
and Pavlína Máchová
Forest and Game Management Research Institute, Jíloviště, Czech Republic

*Address all correspondence to: komarkova@vulhm.cz

IntechOpen

© 2022 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. 

References

- [1] Robinson JP, Harris SA, Juniper BE. Taxonomy of the genus *Malus* Mill. (Rosaceae) with emphasis on the cultivated apple, *Malus domestica* Borkh. *Plant Systematics and Evolution*. 2001;**226**:35-58. DOI: 10.1007/s006060170072
- [2] Stephan R, Wagner I, Kleinschmit J. EUFORGEN Technical Guidelines for Genetic Conservation and Use for Wild Apple and Pear (*Malus sylvestris* and *Pyrus pyrausta*). Italy: International Plant Genetic Resources Institute; 2003
- [3] Donno D, Beccaro GL, Mellano MG, Torello Marinoni D, Cerutti AK, Canterino S, et al. Application of sensory, nutraceutical and genetic techniques to create a quality profile of ancient apple cultivars. *Journal of Food Quality*. 2012;**35**:169-181
- [4] Iacopini P, Camangi F, Stefani A, Sebastiani L. Antiradical potential of ancient Italian apple varieties of *Malus domestica* Borkh. in a peroxynitrite-induced oxidative process. *Journal of Food Composition and Analysis*. 2010;**23**:518-524. DOI: 10.1016/j.jfca.2009.05.004
- [5] Jakobek L, Barron AR. Ancient apple varieties from Croatia as a source of bioactive polyphenolic compounds. *Journal of Food Composition and Analysis*. 2016;**45**:9-15. DOI: 10.1016/j.jfca.2015.09.007
- [6] Grulich V. Red List of vascular plants of the Czech Republic. *Preslia*. 2012;**84**:631-645
- [7] Larsen AS, Asmussen CB, Coart E, Olrik DC, Kjær ED. Hybridization and genetic variation in Danish populations of European crab apple (*Malus sylvestris*). *Tree Genetics & Genomes*. 2006;**2**:86-97. DOI: 10.1007/s11295-005-0030-0
- [8] Reim S, Proft A, Heinz S, Höfer M. Diversity of the European indigenous wild apple *Malus sylvestris* (L.) Mill. in the East Ore Mountains (Osterzgebirge), Germany: I. Morphological characterization. *Genetic Resources and Crop Evolution*. 2012;**59**:1101-1114. DOI: 10.1007/s10722-011-9746-x
- [9] Wagner I, Maurer WD, Lemmen P, Schmitt HP, Wagner M, Binder M, et al. Hybridization and genetic diversity in wild apple (*Malus sylvestris* (L.) MILL.) from various regions in Germany and from Luxembourg. *Silvae Genetica*. 2014;**63**:81-93. DOI: 10.1515/sg-2014-0012
- [10] Cornille A, Giraud T, Smulders MJM, Roldán-Ruiz I, Gladieux P. The domestication and evolutionary ecology of apples. *Trends in Genetics*. 2014;**30**:57-65. DOI: 10.1016/j.tig.2013.10.002
- [11] Larsen AS, Jensen M, Kjær ED. Crossability between wild (*Malus sylvestris*) and cultivated (*M. domestica*) apples. *Silvae Genetica*. 2008;**57**:127-130. DOI: 10.1515/sg-2008-0019
- [12] Máchová P, Malá J, Cvrčková H, Dostál J, Buriánek V. In vitro reproduction of rare and endemic species of rowan tree. *Journal of Forest Science*. 2013;**2013**(59):386-390. DOI: 10.17221/46/2013-JFS
- [13] Malá J, Máchová P, Cvrčková H, Karady M, Novák O, Mikulík J, et al. The role of cytokinins during micropropagation of wych elm. *Biologia Plantarum*. 2013;**57**:174-178. DOI: 10.1007/s10535-012-0252-6

- [14] Pokorná E, Buriánek V, Máchová P, Dostál J, Komárková M. Nové poznatky o reprodukci topolu šedého v in vitro podmínkách. ZPRÁVY LESNICKÉHO VÝZKUMU. 2017;**62**:279-287
- [15] Cvrčková H, Komárková M, Trčková O, Máchová P. Micropropagation of *Tilia cordata* Mill. and verification of genetic diversity of donor trees. In: Bonga JM, Park YS, Trontin JF, editors. In: Proceedings of the 5th International Conference of the IUFRO Unit 2.09.02 on "Clonal Trees in the Bioeconomy Age: Opportunities and Challenges." September 10-15, 2018. Coimbra, Portugal
- [16] Machova P, Malá J, Cvrčková H. Micropropagation of *Betula nana*. Zprávy Lesnického Výzkumu. 2012;**57**:202-206
- [17] Cvrčková H, Malá J, Machova P. In vitro reproduction of *Jurinea cyanoides* (L.) Rchb. Zprávy Lesnického Výzkumu. 2012;**57**:27-32
- [18] Teixeira da Silva JA, Gulyás A, Magyar-Tábori K, Wang M-R, Wang Q-C, Dobránszki J. In vitro tissue culture of apple and other *Malus* species: Recent advances and applications. *Planta*. 2019;**249**:975-1006. DOI: 10.1007/s00425-019-03100-x
- [19] Dobránszky J, Jámboř-Benczúr E, Hudák I, Magyar-Tábori K. Model experiments for establishment of in vitro culture by micrografting in apple. *International Journal of Horticultural Science*. 2005;**11**:47-49. DOI: 10.31421/IJHS/11/1/557
- [20] Chokheli VA, Dmitriev PA, Rajput VD, Bakulin SD, Azarov AS, Varduni TV, et al. Recent development in micropropagation techniques for rare plant species. *Plants*. 2020;**9**:1733. DOI: 10.3390/plants9121733
- [21] Murashige T, Skoog F. A revised medium for rapid growth and bio assays with tobacco tissue cultures. *Physiologia Plantarum*. 1962;**15**:473-497
- [22] Kaplan Z, Danihelka J. Klíč ke květeně České republiky. Prague: Academia; 2019
- [23] Modgil M, Sharma DR, Bhardwaj SV. Micropropagation of apple cv. Tydemans Early Worcester. *Scientia Horticulturae*. 1999;**81**:179-188. DOI: 10.1016/S0304-4238(98)00259-3
- [24] Grant NJ, Hammatt N. Increased root and shoot production during micropropagation of cherry and apple rootstocks: effect of subculture frequency. *Tree Physiology*. 1999;**19**:899-903. DOI: 10.1093/treephys/19.13.899
- [25] Wang Q, Tang H, Quan Y, Tang YQ. Phenol induced browning and establishment of shoot-tip explants of 'Fuji' apple and 'Jinhua' pear cultured in vitro. *Journal of Horticultural Science*. 1994;**69**:833-839
- [26] Yepes LM, Aldwinckle HS. Micropropagation of thirteen *Malus* cultivars and rootstocks, and effect of antibiotics on proliferation. *Plant Growth Regulation*. 1994;**15**:55-67
- [27] Ciccotti A, Bisognin C, Battocletti I, Salvadori A, Herdemertens M, Jarausch W. Micropropagation of apple proliferation-resistant apomictic *Malus sieboldii* genotypes. *Agronomy Research*. 2008;**6**:445-458
- [28] Kabylbekova B, Kovalchuk I, Mukhitdinova Z, Turdiyev T, Kairova G, Madiyeva G, et al. Reduced major minerals and increased minor nutrients improve micropropagation in three apple cultivars. *Vitro Cell Dev Biol-Plant*. 2020;**56**:335-349. DOI: 10.1007/s11627-019-10019-1

- [29] Dobránszky J, Abdul-Kader A, Magyar-Tábori K, Jámbor-Benczúr E, Bubán T, Szalai J, et al. In vitro shoot multiplication of apple: comparative response of three rootstocks to cytokinines and auxin. *International Journal of Horticultural Science*. 2000;**6**:36-39
- [30] Dobránszky J, Abdul-Kader A, Magyar-Tábori K, Jámbor-Benczúr E, Bubán T, Szalai J, et al. Single and dual effects of different cytokinins on shoot multiplication of different apple scions. *International Journal of Horticultural Science*. 2000;**6**:76-78
- [31] Boudabous M, Mars M, Marzougui N, Ferchichi A. Micropropagation of apple (*Malus domestica* L. cultivar Douce de Djerba) through in vitro culture of axillary buds. *Acta Botanica Gallica*. 2010;**157**:513-524
- [32] Magyar-Tábori K, Dobránszky J, Jámbor-Benczúr E. High in vitro shoot proliferation in the apple cultivar Jonagold induced by benzyladenine analogues. *Acta Agronomica Hungarica*. 2002;**50**:191-195
- [33] Bacha N, Darkazanli K, Abdul-Kader A. Direct organogenesis and plantlet multiplication from leaf explants of in vitro-grown shoots of apple (*Malus domestica* Borkh.) cv.'golden delicious' and 'MM111' rootstock. *Fruit, Vegetable and Cereal Science and Biotechnology*. 2009;**3**:28-34
- [34] Sota V, Benelli C, Çuko B, Kongjika E. Effect of a double phase culture system and activated charcoal on in vitro propagation of *Malus sylvestris* (L.) Mill. *Advances in Horticultural Science*. 2022;**35**:361
- [35] Muleo R, Morini S. Physiological dissection of blue and red light regulation of apical dominance and branching in M9 apple rootstock growing in vitro. *Journal of Plant Physiology*. 2008;**165**:1838-1846
- [36] Zhang Y, Bozorov T, Li D, Zhou P, Wen X, Ding Y, et al. An efficient in vitro regeneration system from different wild apple (*Malus sieversii*) explants. *Plant Methods*. 2020;**16**:1-10
- [37] Bommineni V, Mathews H, Samuel S, Kramer M, Wagner D. A new method for rapid in vitro propagation of apple and pear. *HortScience*. 2001;**36**:1102-1106. DOI: 10.21273/HORTSCI.36.6.1102