



Development, structure, and secretion of leaf colleter in *Clusia criuva* Cambess. subsp. *criuva* (Clusiaceae)

Roberta Silva Teixeira¹ , Diego Ismael Rocha² , Jailma Rodrigues Gonçalves¹
and Valdneá Casagrande Dalvi^{1*}

Received: March 23, 2021

Accepted: February 05, 2022

ABSTRACT

Colleters produce a secretion composed of hydrophilic and/or lipophilic substances which lubricates and protects the shoot apical meristem against biotic and abiotic agents. Little attention has been given to these structures in Clusiaceae. In the present study, the structure and development of the leaf colleters of *Clusia criuva* subsp. *criuva* were described and variations in the exudate composition of the colleters at different stages of leaf development were identified. The samples were collected and processed according to techniques for light and scanning electron microscopy. Colleters are of the standard type and not vascularized, and during leaf development, changes in color, structure, and secretion abundance were observed. Asynchrony in the development was noticeable in the leaf primordia and young leaves, from colleters in early formation to those in early senescence. In early phases there was an abundance of polysaccharide, lipid, and protein secretions, whereas adult and senescent leaves revealed an accumulation of phenolic compounds and cell degradation. The secretion was released by the rupture of the cuticle. The structural changes and secretion composition during leaf development emphasize the role of colleters in protecting meristems and developing organs.

Keywords: exudate, glands, histochemistry, Malpighiales, morphoanatomy

Introduction

Throughout the evolution of terrestrial plants, protection strategies have been incorporated into the plant body that allow for greater competition and resistance to environmental adversities, thus ensuring the survival of plants in different environments (Paiva & Machado 2006; Paiva 2016; Prado & Demarco 2018). One of these strategies

involves chemical defense through compounds produced by secretory structures (Prado & Demarco 2018). Plant secretions are often associated with defense strategies (Paiva 2016). They are constituted with different primary and/or secondary metabolites. The chemical predominance of certain compounds in the secretion may suggest specificity in the activity of the secretory cells and also on the ecological role played by such secretion (Castro & Machado 2006; Ascensão 2007; Paiva 2016).

¹ Instituto Federal de Educação, Ciência e Tecnologia Goiano, 75901-970, Rio Verde, GO, Brazil

² Departamento de Agronomia, Universidade Federal de Viçosa, 36570-900, Viçosa, MG, Brazil

* Corresponding author: valdnea.dalvi@ifgoiano.edu.br

Colleters are external structures that produce a sticky secretion composed of mucilage, resins, or a mixture of both (Fahn 1979; Thomas 1991). The colleter secretion present on the surface of vegetative and reproductive organs, protects developing meristems and organs against desiccation (Thomas 1991), acts as an insect repellent (Smith 1963; Curtis & Lersten 1974) or herbivore and pathogen deterrent, and provides inhibition of fungi and bacterial growth through its antimicrobial properties (Castro & Demarco 2008; Calvo *et al.* 2010; Cardoso-Gustavson *et al.* 2014). The colleters have an early development and usually complete their differentiation and even their senescence long before the final development of the plant organ that surrounds them (Canaveze 2012).

Colleters are widely distributed among the angiosperms and were reported in about 60 families (Thomas 1991), including Clusiaceae (Malpighiales), a family known worldwide for the economic importance of its secretions (Judd *et al.* 2009). However, the structural and biological aspects of colleters are poorly investigated within the family being restricted to *Garcinia* and *Clusia*. Filiform colleters occur in the flowers of *Garcinia brasiliensis* (Leal *et al.* 2012); lachrymiform colleters (cone shaped) were found on the leaf axes of *Clusia fluminensis* and *C. lanceolata* (Silva *et al.* 2019) and on the petiole of *C. grandiflora* (Machado & Emmeric 1981). Recently, standard type colleters were found at the base of the petiole of *C. burchellii* (Teixeira *et al.* 2021).

Clusia is one of the largest genera of the Clusiaceae (300 - 400 spp.) with 68 species found in different Brazilian biomes (Lüttge 2007; BFG *et al.* 2015; Bittrich *et al.* 2015). During field expeditions in a fragment of Atlantic Forest located in the State of Espírito Santo (Brazil), the observation of copious secretions covering the apical meristems and young leaves of *Clusia criuva* subsp. *criuva* led us to hypothesize that such exudates were produced by colleters. In the present study we aim to confirm the structure identity responsible for the exudation of the secretion observed in the field. Additionally, we discussed the chemical composition of the secretion throughout the leaf development in a way to contribute to the understanding of the morphofunctionality of such leaf glands present in Clusiaceae.

Material and methods

Collection and sampling location

Samples of the (i) shoot apical meristem with leaf primordia and the base of (ii) young (first and second nodes), (iii) completely expanded (third and fourth nodes) and (iv) senescent leaves (sixth node on) were collected (Fig. 1A - D). The samples were collected from three individuals of *Clusia criuva* Cambess. subsp. *criuva* in the Parque Estadual da Pedra Azul (20°24' S, 41°01' W), in the municipality of Domingos Martins, located in the state of Espírito Santo,

Brazil (Fig. 1E). The area is an Upper Montane Rain Forest with rupestrian vegetation (Fig. 1F) which belongs to the Atlantic Forest Domain. Female (Fig. 1G) and male (Fig. 1H) individuals were collected in a transition region from forest to rupestrian area, growing on a rocky outcrop in full sun (Fig. 1I). Reproductive branches were dried and deposited in the collection of the herbarium at the Instituto Federal Goiano, campus Rio Verde (IFRV): Dalvi VC 111, Dalvi VC 112 and Dalvi VC 113. The species identification was confirmed by Dr. José Elvino do Nascimento Júnior.

Light microscopy

In the field, the samples were fixed in FAA (formalin, acetic acid, and 70 % ethanol 1:1:18 v/v) (Johansen 1940) for structural characterization, as well as in neutral-buffered formalin solution (phosphate buffer: formalin, 9:1 v/v) (Lillie 1965) for histochemical tests. After 48 h, all samples were subjected to dehydration in an ethanol series until their storage in 70% ethyl alcohol.

Subsequently, the samples were embedded in methacrylate resin (Historesin Leica; Leica Microsystems, Heidelberg, Germany), transversely and longitudinally sectioned using a rotary microtome (1508R; Logen Scientific). Sections at 5 µm thickness were stained with toluidine blue (pH 4.6) (O'Brien *et al.* 1964), mounted on slides with synthetic resin (Permunt; Fisher Scientific, NJ, USA) and anatomically characterized.

Samples fixed in neutral-buffered formaldehyde solution were included in methacrylate resin, as described above, and subjected to histochemical tests including: periodic acid and Schiff's reagent (McManus 1948); ruthenium red (Johansen 1940); potassium dichromate (Gabe 1968) and ferric chloride (Johansen 1940); Sudan III (Pearse 1985); and Coomassie brilliant blue (Fisher 1968) for detection of total polysaccharides, pectins/mucilage, phenolic compounds, total lipids, and proteins, respectively. The sections with no reagent (control) were also observed under a light microscope. Material analysis and photographic documentation were performed using an Olympus (model BX61) light microscope equipped with an image capture system, a DP-72 camera.

Scanning Electron Microscopy

For scanning electron microscopy, leaf samples of all four developmental stages were fixed in Karnovsky's fixative (0.1 M sodium phosphate buffer, pH 7.2) (Karnovsky 1965) for 48 h. After dehydration in an ethanol series, the specimens were critical point-dried using CO₂ (Autosamdri®, 815, Séries A) (Bozzola & Russel 1992), mounted on stubs with double-sided adhesive tape, and sputter-coated with gold (Denton Vacuum, Desk V). Photographs were taken using a scanning electron microscope (Jeol, JSM - 6610), equipped with Energy-Dispersive X-Ray Spectroscopy (EDS) and NSS Spectral Imaging (Thermo Scientific, USA).



Results

Variations in the morphology and secretory activity of the colleters

The colleters were located on the adaxial surface of the leaves, in the region of insertion with the stem,

and they presented different coloration according to the leaf developmental stages (Fig. 2). We observed a large accumulation of translucent secretion (Fig. 2A-C) covering the leaf primordia. The young leaves displayed three to four rows of colleters with a whitish color and a brownish apical portion (Fig. 2D). In these two stages we observed asynchrony in the development of the colleters and the beginning of senescence was noticeable by the wilting




Figure 1. Map with the geographic distribution of the Atlantic Forest in Brazil, collection site of *Clusia criuva* Cambess. subsp. *criuva* and sampled material. **A-D.** Leaves at different developmental stages. **A.** Stem apical meristem with leaf primordia (arrow). **B.** Young leaf. **C.** Adult (completely expanded) leaf. **D.** Senescent leaves. **E.** Map for the sampling area, Domingos Martins, State of Espírito Santo, Brazil. **F.** Collection site (arrow) near the Parque Estadual da Pedra Azul. **G.** Female flower. **H.** Male flower. **I.** Details of individual (arrow) growing in a rocky outcrop area.




Figure 2. Colleters in *Clusia criuva* Cambess. subsp. *criuva*. **A-C.** Abundant secretion (asterisk) covering the leaf primordia (lp). **D-F.** Colleters on young leaves. **D.** Brownish color at the apex of the colleters. **E.** Constriction at the apical portion (arrows). **F.** Cuticle rupture (arrowheads). **G.** Extravasated secretion (asterisks). **H, I.** Colleters on adult leaves. **H.** Accentuated brown color. **I.** Collapsed cell, reduced secretion (asterisks) and constriction at the apex of the colleter (arrows). **J-L.** Colleters in senescent leaves. **J.** Blackened colleters. **K.** Collapsed colleters. **L.** Cuticle rupture (arrowheads). Abbreviation: co = colleters. Bars: A, D, H = 2 mm; B, C, I = 500 μ m; E, L = 100 μ m; F = 20 μ m; G = 2 μ m; J = 2 mm; K = 200 μ m.

of cells at the apical portion (Fig. 2E). Intense production and release of secretion via cuticular rupture (Fig. 2F-G) was observed.

In the adult leaves, the brown color of the collectors was intensified and a reduction in the amount of secretion was observed both in the field (Fig. 2H) and with SEM (Fig. 2I). In the apical portion of the collector, a constriction was noted that separates the brownish apex, where wilting and cell disruptions were common, from the lower region (Fig. 2I).

Finally, collectors acquired a blackish color (Fig. 2J), coinciding with the senescence of the leaves. In this phase little or no secretion was observed and the collectors were withered (Fig. 2K). Cellular disruptions, especially in the secretory portion, were common (Fig. 2L).

Development and histology of collectors

Collectors were formed during the first stage of leaf development (Fig. 3). In early development, most cells




Figure 3. Anatomical sections of the leaf collectors of *Clusia criuva* Cambess. subsp. *criuva*. **A-F.** Collectors in early leaf. **A.** Asynchronous development from protodermal and subepidermal cells. **B.** Detail of the undifferentiated cells of the collectors, which have an evident nucleus and nucleolus. **C.** Intense vacuolization (stars). **D.** Collector with central axis and secretory epidermis. **E.** Standard type collectors, sessile with multiseriate epidermis. **F.** Non-vascularized collector. **G.** Cuticle stained with Sudan III. **H, I.** Collectors starting senescence (arrowheads). **J.** Senescent collector. Asterisks indicate secretion. Abbreviations: ca = central axis; se = secretory epidermis. Bars: A, D, I, J = 500 μ m; E, F, H = 200 μ m; B, C = 20 μ m; G = 50 μ m.

of the protoderm elongated (Fig. 3A) and successive anticlinal divisions resulted in a projection formed by the multiseriate epidermis (Fig. 3A). The newly divided cells presented dense cytoplasm, voluminous nuclei and nucleoli, and high nucleus/cytoplasm ratio (Fig. 3B). The subepidermal cells then divided and formed the central axis made up of parenchymal cells (Fig. 3A, D), and at this stage, the vacuolization process was observed throughout the cytoplasm of the parenchymal cells (Fig. 3C). Divisions were intensified, culminating in the formation of colleters (Fig. 3D).

Colleters were sessile and non-vascularized with a truncated or tapered apex (Fig. 3D-E). They presented a multicellular head formed by a central axis of parenchymal cells and a secretory epidermis (Fig. 3D-F) coated with a thick cuticle (Fig. 3G), following into the standard-type colleters. The secretory epidermis had commonly more than two layers of columnar juxtaposed cells which tended to palisade (Fig. 3E).

In adult leaves, the degradation of cells in the apical portion was accentuated (Fig. 3H). In senescent leaves, the secretory epidermis of most colleters was completely degraded with collapsed and/or ruptured cells (Fig. 3I-J). Although senescent, *Clusia criuva* subsp. *criuva* colleters did not come into abscission.

Histochemical characterization

Histochemical tests showed polysaccharides as the main components of the secretion produced by *Clusia criuva* subsp. *criuva*. Total polysaccharides (Fig. 4A-C) and pectins/mucilage (Fig. 4D-F) were observed in the exudates of the colleters at all leaf development stages. In the primordia, young, and adult leaves, accumulation of these compounds was observed in the cytoplasm of secretory cells (Fig. 4A, B, D, E), in the subcuticular space (Fig. 4B, E), and in extravasated secretion (Fig. 4A, C, F).

Proteins were observed in the exudate of the colleters from the leaf primordia (Fig. 4G) through to the senescent leaves (Fig. 4I). However, the accumulation of proteins in secretory cells was evident only in colleters from leaf primordia and young leaves (Fig. 4G). Lipid droplets were observed within the secretory cells as well as in the cells from the central axis in the young leaves. In adult and senescent leaves, lipids were detected only in secretory epidermal cells (Fig. 4H), and structural lipids impregnated in cell walls in cells of the central axis (Fig. 4H).

Phenolic compounds were registered in the colleters present in all leaf development stages. However, in leaf primordia and young leaves, these compounds were restricted to the apical portion of the colleters (Fig. 4J), whereas in adult and senescent leaves, phenolics were observed throughout the entire structure of the colleter, particularly in the secretory cells of the epidermis (Fig. 4K, L).

Discussion

The abundant secretion observed in the field in the leaf primordia and young leaves of *Clusia criuva* subsp. *criuva*, the distribution as well as the structural and chemical nature of the secretion confirm our hypothesis for the presence of colleters on the leaves of this species. The colleters of *C. criuva* subsp. *criuva* are derived from the protoderm and the ground meristem and are classified as emergencies according to Evert (2006). In *Ilex* species, González (1998) reported that a group of protodermal cells undergoes radial enlargement and anticlinal division and some subepidermal cells divide to form a protuberance. Thus, the anticlinal division gives rise to the secretory epidermis and the subepidermal cells produce the parenchymal axis of the colleter, and the same pattern has been observed here. Other studies report that the most common occurrence is that of colleters forming only by the protoderm from periclinal and anticlinal divisions, without no involvement of the ground meristem (Paiva & Machado 2006; Paiva 2009; Paiva & Martins 2011; Rocha *et al.* 2011). As pointed out by our results, the involvement of the protoderm and the ground meristem in the process for colleter was reported in *C. fluminensis*, *C. lanceolata* (Machado & Emmerich 1981; Silva *et al.* 2019) and *C. burchellii* (Teixeira *et al.* 2021), which suggests that it may be a conservative characteristic in the group, although studies on colleters in the genus are still scarce.

The standard type colleters present in *C. criuva* subsp. *criuva* is constituted by a central parenchymal axis surrounded by a layer of palisade-secreting epidermal cells, as previously reported by Lersten (1974). The standard type colleter also was found in *C. grandiflora* (Machado & Emmerich 1981) and *C. burchellii* (Teixeira *et al.* 2021). In *C. fluminensis* and *C. lanceolata*, lachrymiform cone-shaped colleters were described (Silva *et al.* 2019). However, we understand that the lachrymiform refers to the external morphology of the colleter, which is anatomically organized as the standard type, according to Lersten (1974). Studies on the morphological differences of colleters can be important to offer contributions to taxonomy studies and to establish phylogenetic relationships within the Clusiaceae. Our results clearly show cuticle ruptures in the apical portion of the colleters, which is a common way to secrete substances by colleters (Paiva & Machado 2006; Silva *et al.* 2017).

The chemical composition of the colleter exudates in *C. criuva* subsp. *criuva* is diverse as it is composed of a mixture of hydrophilic and lipophilic compounds, that is, polysaccharides, proteins, lipids, and phenolic compounds. However, variations in the composition of the secretion were observed depending on the leaf development stage. Mucilage consisting mainly of polysaccharide polymers with high molecular weight (Fahn 1988) together with pectin may act as a water absorber due to its hygroscopic properties (Hall 1981; Nobel *et al.* 1992), which may protect



meristem and young leaves of *C. criuva* subsp. *criuva* against desiccation. This corroborates the primary function of colleters in producing secretions that protect young organs during their development (Lersten & Horner 1968; Lersten

1974; Fahn 1979) and/or that promote the lubrication of the stem meristem, minimizing friction of the developing leaf tissues and preventing dehydration (Fahn 1979; Thomas 1991; Mayer *et al.* 2011; Silva *et al.* 2012; Mayer *et al.* 2013).




Figure 4. Histochemical tests of *Clusia criuva* Cambess. subsp. *criuva*. **A-C.** Reaction with PAS showing the presence of polysaccharides. **D-F.** Presence of pectin confirmed by the red ruthenium test. **G, I.** Reaction with Coomassie brilliant blue showing proteins in the cytoplasm and extravasated secretion. **H.** Lipid droplets (black arrow) and impregnation of lipids in the cells of the central axis (white arrow), evidenced by Sudan III. **J-L.** Ferric chloride test demonstrates the increase in the amounts of phenolic compounds in the colleters during leaf development. Asterisks indicate extravasated secretion. Bars: A, D, G, J = 500 μ m; B, C, E, F, I, K, L = 100 μ m; H = 20 μ m.

Proteins are involved in the defense against fungal activity and attack of pathogens (Klein *et al.* 2004; Miguel *et al.* 2006) and are commonly reported in the secretion of colleters (González & Tarragó 2009; Coelho *et al.* 2013; Dalvi *et al.* 2014; Lacchia *et al.* 2016; Pinheiro *et al.* 2019).

The phenolic compound production was intensified, in colleters, of adult and senescent leaves of *C. criuva* subsp. *criuva*. Phenolics have been reported for other species of *Clusia* both in the secretion present in the meristem and in the adult leaves (Silva *et al.* 2019). According to Coelho *et al.* (2013), phenolic compounds and lipids are the final products of the secretion of colleters in a senescence stage. The cuticle wrinkling, the formation of a constriction at the base of the colleters and cell disruption also indicate the senescence of the colleters (Pinheiro *et al.* 2019), which were also shown here for *C. criuva* subsp. *criuva* colleters even they did not come into abscission.

The color of colleters changes in the field during the development of *C. criuva* subsp. *criuva* leaf, as reported for *C. burchellii* (Teixeira *et al.* 2021) and species from other families, including Apocynaceae (Thomas & Dave 1989; Appezzato-da-Glória & Estelita 2000; Rio *et al.* 2002), Gentianaceae (Dalvi *et al.* 2013; 2014), and Rubiaceae (Lersten 1974; Miguel *et al.* 2006; 2009; Vitarelli & Santos 2009; Klein *et al.* 2010; Pinheiro *et al.* 2019). These authors generally associate changes in color with the presence of phenolic compounds, which was confirmed in colleters of *C. criuva* var. *criuva*. The accumulation of phenolic compounds in colleters is also related to a change in the function of these glands, being produced in a second stage of the secretory phase (Ribeiro *et al.* 2017), which would act to avoid predation (Castro & Demarco 2008). However, the position of colleters in this species and others would not preclude the predation of leaves by herbivores.

In summary, we report here the occurrence of standard-type colleters in *C. criuva* subsp. *criuva* as well as the morphoanatomical variations of the structure and chemical composition of the exudate throughout the leaf development of this species. Our results contribute to a better understanding of the morphofunctionality of colleters in Clusiaceae, structures that are scarcely studied in this group of plants. Comparative studies with a larger number of Clusiaceae species are also needed in order to understand the evolution of colleter for this family and their correlation with the biomes where such species occur.

Acknowledgments

We thank the Ministério da Ciência e Tecnologia/ Conselho Nacional de Desenvolvimento Científico e Tecnológico (Grant #406824/2016-9) and the IF Goiano, campus Rio Verde, for financial support; the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES), Finance Code 001, for the Master's scholarship granted to

first author; the Laboratório Multiusuário de Microscopia de Alta Resolução (LabMic/UFG) for the preparation and analysis of electron microscopy samples; Fernando Henrique Antonioli Farache for creating the map, and Patrícia Oliveira da Silva for the drawings in Figure 1.

References

- Appezzato-Da-Glória B, Estelita MEM. 2000. Development, structure and distribution of colleters in *Mandevilla illustris* and *M. velutina* (Apocynaceae). *Revista Brasileira de Botânica* 23: 113-120.
- Ascensão L. 2007. Estruturas secretoras em plantas. Uma abordagem morfo-anatômica. In: Figueiredo AC, Barroso JG, Pedro LG. (eds.) *Potencialidades e aplicações das plantas aromáticas e medicinais. Curso Teórico-Prático*. 3. ed. Lisboa: Faculdade de Ciências da Universidade de Lisboa/Centro de Biotecnologia Vegetal. p. 19-28.
- BFG - The Brazil Flora Group. 2015. Growing knowledge: an overview of seed plant diversity in Brazil. *Rodriguésia* 66: 1085-1113.
- Bittrich V, Trad RJ, Cabral FN, Nascimento-Jr JE, Souza VC. 2015. Clusiaceae in lista de espécies da flora do Brasil. *Jardim Botânico do Rio de Janeiro*. <http://floradobrasil.jbrj.gov.br/jabot/floradobrasil/FB28292>. 03 Jan. 2020.
- Bozzola JJ, Russel LD. 1992. *Electron microscopy*. Boston, Jones and Bartlett Publishers.
- Calvo TR, Demarco D, Santos FV, et al. 2010. Phenolic compounds in leaves of *Alchornea triplinervia*: anatomical localization, mutagenicity, and antibacterial activity. *Natural Product Communication* 5: 1225-1232.
- Canaveze Y. 2012. Estrutura, origem e desenvolvimento de laticíferos e coléteres em plantas de *Tabernaemontana catharinensis* a.d.c. (Rauvolfioideae, Apocynaceae) em diferentes fases do desenvolvimento vegetativo. MSc Thesis, Universidade Estadual Paulista "Júlio de Mesquita Filho", Botucatu, Brasil.
- Cardoso-Gustavson P, Campbell LM, Viveiros-Mazzoni S, Barros F. 2014. Floral colleters in Pleurothallidinae (Epidendroideae: Orchidaceae). *American Journal of Botany* 101: 587-597.
- Castro MM, Demarco D. 2008. Phenolic compounds produced by secretory structures in plants: a brief review. *Natural Product Communications* 3: 1273-1284.
- Castro MM, Machado SR. 2006. Células e tecidos secretores. In: Appezzato-da-Glória B, Carmello-Guerreiro SM (eds.) *Anatomia vegetal*. 2nd. edn. Viçosa, Universidade Federal de Viçosa. p. 179-203.
- Coelho VPM, Leite JPV, Fietto LG, Ventrella MC. 2013. Colleters in *Bathysa cuspidata* (Rubiaceae): Development, ultrastructure and chemical composition of the secretion. *Flora* 208: 579-590.
- Curtis JD, Lersten NR. 1974. Morphology, seasonal variation, and function of resin glands on buds and leaves of *Populus deltoides* (Salicaceae). *Botany* 61: 835-845.
- Dalvi VC, Cardinelli LS, Meira RMSA, Azevedo AA. 2014. Foliar colleters in *Macrocarpaea obtusifolia* (Gentianaceae): anatomy, ontogeny, and secretion. *Botany* 92: 59-67.
- Dalvi VC, Meira RMSA, Francino DMT, Silva LC, Azevedo AA. 2013. Anatomical characteristics as taxonomic tools for the species of *Curtia* and *Hockinia* (Saccifolieae-Gentianaceae Juss.). *Plant Systematics and Evolution* 300: 99-112.
- Evert RF. 2006. *Esau's Plant anatomy: meristems, cells, and tissues of the plant body: their structure, function, and development*, 3rd. edn. Hoboken, John Wiley & Sons.
- Fahn A. 1979. *Secretory Tissues in Plants*. London, Academic Press.
- Fahn A. 1988. *Plant anatomy*. Oxford, Pergamon Press.
- Fisher DB. 1968. Protein staining of ribboned epon sections for light microscopy. *Histochemistry Cell Biology* 16: 92-96.
- Gabe M. 1968. *Techniques histologiques*. Paris, Masson and Cie.
- González AM, Tarragó JR. 2009. Anatomical structure and secretion compounds of colleters in nine *Ilex* species (Aquifoliaceae) from southern South America. *Botanical Journal of the Linnean Society* 160: 197-210.



- González AM. 1998. Collecters in *Turnera* and *Piriqueta* (Turneraceae). Botanical Journal of the Linnean Society 128: 215-228.
- Hall WC. 1981. Biomass as an alternative fuel. Rockville, Government Institutes Inc.
- Johansen DA. 1940. Plant microtechnique. New York, Mc GrawHill.
- Judd WS, Campbell CS, Kellogg EA, Stevens PF, Donoghue MJ. 2009. Sistemática Vegetal - um enfoque filogenético 3rd. edn. Porto Alegre, Artmed.
- Karnovsky MJ. 1965. A formaldehyde-glutaraldehyde fixative of high osmolality for use in electron microscopy. Journal Cell Biology 27: 137-138.
- Klein DE, Gomes VM, Silva-Neto SJ, Cunha M. 2004. The structure of collectors in several species of *Simira* (Rubiaceae). Annals of Botany 94: 733-740.
- Klein DE, Oliveira MA, Cunha M. 2010. Ultrastructure of secretory and senescence phase in collectors of *Bathysa gymnocarpa* and *B. stipulata* (Rubiaceae). Revista Brasileira de Botânica 33: 425-436.
- Lacchia APS, Tölke EEAD, Demarco D, Carmello-Guerreiro SM. 2016. Presumed domatia are actually extrafloral nectaries on leaves of *Anacardium humile* (Anacardiaceae). Rodriguesia 67: 19-28.
- Leal DO, Malafaia C, Cesar R, et al. 2012. Floral structure of *Garcinia brasiliensis* in relation to flower biology and evolution. International Journal of Plant Sciences 173: 172-83.
- Lersten NR. 1974. Morphology and distribution of collectors and crystals in relation to the taxonomy and bacterial leaf nodule symbiosis of *Psychotria* (Rubiaceae). American Journal of Botany 61: 973-981.
- Lersten NR, Horner HTJR. 1968. Development, structure and function of secretory trichomes in *Psychotria bacteriophila* (Rubiaceae). American Journal of Botany 55: 1089-1099.
- Lillie RD. 1965. Histopathologic technic and practical histochemistry. 3rd. edn. New York, McGraw-Hill.
- Lüttge U. 2007. *Clusia* – A woody Neotropical genus of remarkable plasticity and diversity. Ecological studies, vol. 194. New York, Springer.
- Machado MMP, Emmerich M. 1981. Presença de coléteres em *Clusia lanceolata* Cambess. Boletim do Museu Nacional de Botânica 59: 1-7.
- Mayer JLS, Cardoso-Gustavson P, Appezzato-Da-Glória B. 2011. Collecters in monocots: New record for Orchidaceae. Flora 206: 185-190.
- Mayer JLS, Carmello-Guerreiro SM, Mazzafera P. 2013. A functional role for the collectors of coffee flowers. AoB Plants 5: 1-13.
- McManus JFA. 1948. Histological and histochemical uses of periodic acid. Stain Technol. 23: 99-108.
- Miguel EC, Gomes VM, Oliveira MA, Cunha M. 2006. Collecters in *Bathysa nicholsonii* K. Schum. (Rubiaceae): Ultrastructure, secretion protein composition, and antifungal activity. Plant Biology 8: 715-722.
- Miguel EC, Moraes DG, Cunha M. 2009. Stipular collectors in *Psychotria nuda* (Cham. & Schltdl.) Wawra (Rubiaceae): micromorphology, anatomy and crystals microanalysis. Acta Botanica Brasilica 23: 1034-1039.
- Nobel PS, Alm DM, Cavelier J. 1992. Growth respiration, maintenance respiration and structural-carbon costs for roots of three desert succulents. International Journal of Plant Sciences 153: 102-107.
- O'Brien TP, Feder N, McCully ME. 1964. Polychromatic staining of plant cell walls by toluidine blue O. Protoplasma 59: 368-373.
- Paiva EAS, Machado SR. 2006. Collecters in *Caryocar brasiliensis* (Caryocaraceae), ontogenesis, ultrastructure and secretion. Brazilian Journal of Biology 66: 301-308.
- Paiva EAS, Martins LC. 2011. Calycinal trichomes in *Ipomoea cairica* (Convolvulaceae): ontogenesis, structure and functional aspects. Australian Journal of Botany 59: 91-98.
- Paiva EAS. 2009. Occurrence, structure and functional aspects of the collectors of *Copaifera langsdorffii* Desf. (Fabaceae, Caesalpinioideae). Comptes Rendus Biologies 332: 1078-1084.
- Paiva EAS. 2016. How do secretory products cross the plant cell wall to be released? A new hypothesis involving cyclic mechanical actions of the protoplast. Annals of Botany 117: 533-540.
- Pearse AGE. 1985. Histochemistry: theoretical and applied. Vol II. Edinburgh, Livingstone.
- Pinheiro SKP, Teófilo FBS, Lima AKM, Cordoba BV, Miguel TBAR, Miguel EC. 2019. Ontogenesis and secretion mechanism of collectors in *Morinda citrifolia* L. (Rubiaceae). South African Journal of Botany 121: 26-33.
- Prado E, Demarco D. 2018. Laticifers and secretory ducts: similarities and differences. In: Hufnagel L (ed.) Ecosystem services and global ecology. London, IntechOpen. p. 103-123.
- Ribeiro JC, Ferreira MJP, Demarco D. 2017. Collecters in Asclepiadoideae (Apocynaceae): protection of meristems against desiccation and new functions assigned. International Journal of Plant Sciences 178: 465-477.
- Rio MCS, Castro MM, Kinoshita LS. 2002. Distribuição e caracterização anatômica dos coléteres foliares de *Prestonia coalita* (Vell.) Woodson (Apocynaceae). Revista Brasileira de Botânica 25: 339-349.
- Rocha JF, Pimentel RR, Machado SR. 2011. Estruturas secretoras de mucilagem em *Hibiscus pernambucensis* Arruda (Malvaceae): distribuição, caracterização morfoanatômica e histoquímica. Acta Botanica Brasilica 25: 751-763.
- Silva CJ, Barbosa LCDA, Marques AE, Barakat-Pereira MC, Pinheiro AL, Meira RMSA. 2012. Anatomical characterisation of the foliar collectors in Myrtoideae (Myrtaceae). Australian Journal of Botany 60: 707-717.
- Silva KMM, Luna BN, Joffily A, Paiva SR, Barros CF. 2019. Revealing the development of secretory structures in the leaves of *Clusia fluminensis* and *Clusia lanceolata* (Clusiaceae). Flora 256: 69-78.
- Silva MDS, Coutinho ÍAC, Araújo MN, Meira RMSA. 2017. Collecters in *Chamaecrista* (L.) Moench sect. *Chamaecrista* and sect. *Caliciopsis* (Leguminosae-Caesalpinioideae): anatomy and taxonomic implications. Acta Botanica Brasilica 31: 382-391.
- Smith RH. 1963. Preferential attack by *Dendroctonus terebrans* on *Pinus elliotti*. Journal of Economic Entomology 56: 817-819.
- Teixeira RS, Rocha DI, Dalvi V. 2021. Leaf collectors in *Clusia burchellii* Engl.: Structural and ultrastructural features of a little-known gland in Clusiaceae. Flora 280: 151834.
- Thomas V, Dave Y. 1989. Structure, origin, development and senescence of collectors in *Nerium indicum* mill. (*Nerium odoratum* Soland, Apocynaceae). Korean Journal Botany 32: 163-172.
- Thomas V. 1991. Structural, functional and phylogenetic aspects of the collector. Annals of Botany 68: 287-305.
- Vitarelli NC, Santos M. 2009. Anatomia de estípulas e coléteres de *Psychotria carthagenensis* Jacq. (Rubiaceae). Acta Botanica Brasilica 23: 923-928.

