

PREPRINT

Author-formatted, not peer-reviewed document posted on 22/03/2023

DOI: <https://doi.org/10.3897/arphapreprints.e103766>

A taxonomic synopsis of Heliotropiaceae (Boraginales) in Thailand

 Kanokorn Rueangsawang,  Pranom Chantaranothai

Disclaimer on biological nomenclature and use of preprints

The preprints are preliminary versions of works accessible electronically in advance of publication of the final version. They are not issued for purposes of botanical, mycological or zoological nomenclature and **are not effectively/validly published in the meaning of the Codes**. Therefore, nomenclatural novelties (new names) or other nomenclatural acts (designations of type, choices of priority between names, choices between orthographic variants, or choices of gender of names) **should NOT be posted in preprints**. The following provisions in the Codes of Nomenclature define their status:

International Code of Nomenclature for algae, fungi, and plants (ICNafp)

Article 30.2: "An electronic publication is not effectively published if there is evidence within or associated with the publication that its content is merely preliminary and was, or is to be, replaced by content that the publisher considers final, in which case only the version with that final content is effectively published." In order to be validly published, a nomenclatural novelty must be effectively published (Art. 32.1(a)); in order to take effect, other nomenclatural acts must be effectively published (Art. 7.10, 11.5, 53.5, 61.3, and 62.3).

International Code of Zoological Nomenclature (ICZN)

Article: 21.8.3: "Some works are accessible online in preliminary versions before the publication date of the final version. Such advance electronic access does not advance the date of publication of a work, as preliminary versions are not published (Article 9.9)".

A taxonomic synopsis of Heliotropiaceae (Boraginales) in Thailand

Kanokorn Rueangsawang¹, Pranom Chantaranothai^{2,3}

1 Department of Biology, Faculty of Science, Ramkhamhaeng University, Bangkok 10240, Thailand **2** Applied Taxonomic Research Center, Department of Biology, Faculty of Science, Khon Kaen University, Khon Kaen 40002, Thailand **3** Honorary Research Associate, Royal Botanic Gardens, Kew, Richmond, Surrey TW9 3AE, UK.

Corresponding author: Pranom Chantaranothai (pranom@kku.ac.th)

Abstract

A synopsis of Heliotropiaceae in Thailand is presented together with the taxonomic history, identification keys, distribution, phenology, herbarium specimen citations and diagnostic characters of each species. The taxonomic character of the family is of terminal style with a conical stigma structure forming a basal ring around the style. There are 12 species: five species each in *Euploca* and *Tournefortia* and two species in *Heliotropium*. Three species are newly recorded in Thailand, *Euploca ovalifolia*, *Tournefortia hookeri* and *T. montana*. Lectotypes are selected for *T. boniana* and *T. tetrandra* var. *angustifolia*.

Keywords

Euploca, *Heliotropium*, morphology, Thai flora, *Tournefortia*

Introduction

Heliotropiaceae (=Boraginaceae s.l., subfamily Heliotropioideae) comprises ca. 450 species in four genera: *Heliotropium* L. (incl. *Tournefortia* L.), *Euploca* Nutt., *Myriopus* Small. and *Ixorhea* Fenzl, (Luebert et al. 2016; Diane et al. 2016) and is widely distributed in the tropical, subtropical and temperate zones, with the centre of diversity in the Neotropics (Mabberley 2017). In the previous classification based on morphology and anatomy, Förther (1998) recognised eight genera: *Argusia* Böhm, *Ceballosia* Kunkel, *Heliotropium*, *Hilgeria* Förther, *Ixorhea*, *Nogalia* Verde, *Schleidenia* Endl. and *Tournefortia*. The family varies from herbs, shrubs, lianas or small trees. The distinct characteristics are scorpioid cymes and terminal style with a conical stigmatic head having a basal ring-shaped stigma (Luebert et al. 2016; Diane et al. 2016). Recently, the genus *Euploca* is accepted as a valid genus, including *Heliotropium* sect. *Orthostachys*, *Hilgeria* and *Schleidenia*, all these are confirmed by the molecular results (Hilger and Diane 2003). *Euploca* is separated from the other genera due to the bracteate inflorescence and fruit breaking up into four distinct nutlets (Feuillet 2016).

The contributions to the knowledge of *Tournefortia* and *Heliotropium* remain controversial. Recently, the *trnL-F* and ITS1 sequence data analysis has shown that *Heliotropium* and *Tournefortia* are not monophyletic (Diane et al. 2002; Hilger and Diane 2003; Luebert et al. 2011). *Tournefortia* sect. *Tournefortia* recognised by drupaceous fruit and distinctly lobed has been shown nested within *Heliotropium* s.s. in a sister group to a HELIOTROPIUM I. In moving *Tournefortia* to *Heliotropium* which was unclarified in Hilger and Diane (2003). However, new combinations

of *Euploca* and *Heliotropium* have been made (e.g. Craven 2005; Feuillet 2016; Frohlich et al. 2020; Antony and Javad 2020).

In SE Asian species are unclear and maintain *Tournefortia* in their old circumscription (e.g. Gagnepain and Courchet 1914; Ridley 1923; Johnston 1935; Backer and Bakhuizen van den Brink Jr 1965; Zhu et al. 1995; Riedl 1997). The first checklist by Fletcher and Kerr (1951) was reported in the *Florae Siamensis Enumeratio*, with nine genera in the former Boraginaceae. Since the phylogenetic studies have been changed, a small family of Heliotropiaceae. There are three genera in Thailand: *Euploca*, *Heliotropium* and *Tournefortia*. The present study is to provide a synopsis of Heliotropiaceae for the Flora of Thailand, with the key to the determination and nomenclature of all species.

Materials and methods

The extensive collections of herbarium specimens at AAU, ABD, BK, BKF, BM, C, CMU, CMUB, E, K, K-W, KKU, L, QBG and SING (abbreviations follow Thiers 2022 continuously updated) were studied together with living material observation in the field. Species delimitation was based on the examination of original publications including the relevant literature: Clarke 1885; Gagnepain and Courchet 1914; Ridley 1923; Johnston 1935, 1951; Backer and Bakhuizen van den Brink Jr 1965; Zhu et al. 1995; Riedl 1997; Mill 1999. The type specimens were observed online in JSTOR Global Plants (<https://plants.jstor.org/>). Most measurements were taken from dried specimens and spirit-preserved material and colour descriptions were studied from living material. Distribution, ecology and phenology were derived from specimen labels. All collections cited that have been seen by the authors are marked with ‘!’, and those seen as digital images are indicated with ‘image’.

Taxonomic treatment

Heliotropiaceae Schrad., Commentat. Soc. Regiae Sci. Gött. Recent. 4: 192. 1819, nom.cons.

Type genus: *Heliotropium* L.

Description. Annual or perennial herbs, climbing shrubs or small trees. Leaves simple, alternate, exstipulate, petiolate or sessile; lamina linear to ovate, apex acute to acuminate, margin entire or revolute. Inflorescences terminal or axillary, spike-like, scorpioid, subcapitate or subcorymbose cymes; with or without bracts. Flowers 5-merous, rarely 4-merous, bisexual, actinomorphic. Calyx 5-lobed, divided almost to the base, lobes linear to ovate, entire, persistent. Corolla white to pinkish white, pale green or greenish to yellowish, campanulate to funnel-shaped, lobes orbicular. Stamens 5, filaments adnate to the corolla tube. Pistil: ovary superior, bicarpellate, usually 4-loculate, 1 ovule per locule, nectar disk at the base surrounds the ovary; style terminal, with a conical stigma structure forming a basal ring around style. Fruit dry, separating into Fruit dry or fleshy, separating into four 1-seeded nutlets or two 2-seeded nutlets. Seeds straight or curved, with endosperm.

Three genera and 12 species are found in Thailand.

Key to the genera in Thailand

- 1 Climbing shrub; inflorescence dichotomous branched, scorpioid cymes *Tournefortia*

- Annual herb or small tree; inflorescence spike-like, scorpioid, subcapitate or subcorymbose 2
- 2 Inflorescence bracteate; fruit separating into four nutlets ***Euploca***
- Inflorescence ebracteate; fruit separating into two nutlets ***Heliotropium***

***Euploca* Nutt., Trans. Amer. Philos. Soc., ser. 2, 5: 189. 1836.**

Heliotropium sect. *Orbostachys* R.Br., Prodr. Fl. Nov. Holland.: 493. 1810. Type species:

Heliotropium foliatum R.Br.

Preslaea Mart., Nov. Gen. Sp. Pl. 2: 75. 1827, nom. illeg.

Schleidenia Endl., Gen. Pl.: 646. 1839. Type species: *Preslaea paradoxa* Mart.

Hilgeria Förther, Sendtnera 5: 132. 1998. Type species: *Hilgeria hypogaea* (Urb. & Ekman) Förther.

Heliotropium sect. *Orthostachys* subsect. *Bracteata* I.M.Johnst., Contr. Gray Herb. 81: 47. 1928.

Type species: *Heliotropium procumbens* Mill.

Type species. *Euploca convolvulacea* Nutt.

Description. Annual herbs. Leaves petiolate or sessile; lamina linear to ovate, apex acute to obtuse, margin entire or revolute, pubescent to strigose or silky silver hairs on both surfaces. Inflorescences terminal or axillary, spike-like, scorpioid or subcapitate cymes, bracteate. Flowers 5-merous. Calyx 5-lobed, lobes linear to lanceolate. Corolla white with a yellow-orange throat inside, tubular or funnel-shaped, lobes orbicular to oblong, throat pubescent inside. Stamens included, sessile or with short filaments; anthers, elliptic-oblong. Ovary divided into 4 lobes. Fruits separating into four 1-seeded nutlets.

Key to the species of *Euploca*

- 1 Inflorescence spike-like or scorpioid cymes 2
- Inflorescence subcapitate cymes 4
- 2 Leaves elliptic to obovate or oblanceolate; cymes 2-rowed scorpioid 3. ***E. ovalifolia***
- Leaves linear to narrowly elliptic; cymes unilateral or spike-like 3
- 3 Prostrate much-branched stem; leaves narrowly elliptic with strigose hairs 5. ***E. strigosa***
- Erect unbranched to few-branched stem; leaves linear with greyish tomentose hairs 4. ***E. paniculata***
- 4 Plants suberect with white pubescent hairs; leaves elliptic 1. ***E. bracteata***
- Plants prostrate with stiff bristly white hairs; leaves narrowly lanceolate 2. ***E. marifolia***

1. *Euploca bracteata* (R.Br.) M.W.Frohl. & M.W.Chase, Phytotaxa 434(1): 15. 2020.

Fig. 1A

Heliotropium bracteatum R.Br., Prodr. Fl. Nov. Holland.: 493. 1810. Type: Australia, Carpentaria Lison (Carpentaria), *Brown s.n.* (holotype P [P00610128 image!]; isotype K [K000998273!]).

Heliotropium cyrtoachyum Miq., Fl. Ned. Ind. 2: 924. 1858. Type: Indonesia, Java, *Horsfield s.n.* (holotype K [K000998248!]; isotypes L [L0281634!], K [K000998249!]).

Heliotropium bracteatum var. *leptostachyum* Benth., Fl. Austral. 4: 451. 1869. Type: Australia,

Cape York, *Daemel s.n.* (holotype K [K000998274!]; isotype BM [BM001040578!]).

Type. Based on *Heliotropium bracteatum* R.Br.

Distribution. Pakistan, India, Sri Lanka, Myanmar, China (Hainan, Kwangtung), Laos, Cambodia, Vietnam, Malaysia, Java, Australia.

Ecology. Open areas in deciduous forest, dry secondary forest; 50–300 m alt., flowering and fruiting from February to October.

Specimens examined. Thailand, Northern: Nakhon Sawan, 10 km north-west of Nakhon Sawan, 150 m alt., 21 July 1973, *Murata et al. T-16587* (BKF, L). South-western: Uthai Thani, Tapoh, 4 Jan 1962, *Larsen 9135* (C, K); Kanchanaburi, Sai Yok, 15 Aug 1967; Khao Pu Maklai, Nong Hoi, 10 July 1978, *Phengkai et al. 4247* (BKF, K); *ibid.*, 40 m alt., 8 July 1922, *Marcan 889* (BM, K); Khao Tok, 50 m alt., 13 July 1930, *Kerr 19547* (BM, BK, K, E); *ibid.*, 50 m alt., 10 July 1930, *Kerr s.n.* (K); Wangpho, 15 Oct 1967, *Chersmsirivathana 790* (BK). Central: Lop Buri, Lam Narai, 13 Nov 1975, *Smitinand 12108* (BKF); *ibid.*, 22 Aug 1975, *Boonkurd 272* (BK); Saraburi, Phraphutthabat, 18 Sept 2004, *Pooma et al. 4798* (BKF); Phu Khae, 100 m alt., 26 June 1947, *Bunpheng 147* (BKF, K); Mae Nam Sak, 40 m alt., 29 May 1923, *Kerr 7020* (BK, K); South-eastern: Sa Kaeo, Aranyaprathet, 9 Apr 1930, *Kerr 19593* (ABD, BM, BK, BKF, E, K, L); *ibid.*, 20 Oct 1828, *Put 2073* (BK, BM). Peninsular: Trang, Palian, 20 m alt., 26 May 1976, *Smitinand 12229* (BKF).

Diagnostic characters. *Euploca bracteata* is suberect, with sparsely white pubescent hairs on both leaf surfaces, subcapitate cyme inflorescence at the end of branches and sessile flower or with pedicels up to 1 mm long. This species is similar to *E. marifolia*, sharing a similar habit and inflorescence form but differs in its elliptic leaves (vs linear to lanceolate leaves in *E. marifolia*).

2. *Euploca marifolia* (J.Köenig ex Retz.) Ancy & P.Javad, Nord. J. Bot. 38(11): 2. 2020.

Fig. 1B

Heliotropium marifolium J.Köenig ex Retz., Observ. Bot. 2: 8. 1781. Type: Asia, 1781, Köenig 7052 (holotype LD [LD1748458 image!]).

Heliotropium scabrum Retz., Observ. Bot. 2: 8. 1781. Type: India, Coromandel coast, Köenig s.n. (C [C10008736!]).

Euploca scabra (Retz.) M.W.Frohl. & M.W.Chase, Phytotaxa 434: 19. 2020. Type as above.

Type. Based on *Heliotropium marifolium* J.Köenig ex Retz.

Distribution. Pakistan, India, Sri Lanka, China, Cambodia, Vietnam, Peninsular Malaysia, Java.

Ecology. In the open deciduous forest, dry secondary forest; 50–300 m alt., flowering and fruiting from January to October.

Specimens examined. Thailand, Eastern: Nakhon Ratchasima, Nong Sarai, 300 m alt., 5 Sept 1963, *Smitinand & Sleumer 8359* (K, L). South-western: Kanchanaburi, Si Sawat, 16 Nov 1917, *van Beusekom et al. 3809* (BKF, L); Sai Yok, 15 Aug 1967; Hin Lat, 50 m alt., 28 Nov 1957, *Smitinand 3875* (BKF); Khao Ai Mao, 15 Nov 1968, *Sangkhachand 1576* (BKF, K); *ibid.*, 40 m alt., 8 July 1922, *Marcan 889* (BM, K); Thong Pha Phum, 215 m alt., 6 Aug 2012, *Middleton et al. 5261* (BKF, E); Phetchaburi, Kaeng Krachan NP, 285 m alt., 11 Aug 2002, *Middleton et al. 972* (BKF, E, K). Central: Sing Buri, Mueang, 5 Jun 1880, *Put 2614* (BM, BK, K, BKF), Suphan Buri, Bang Plama, 22 Sept 1930, *Kerr s.n.* (K); Phra Nakhon Si Ayutthaya, 9 Sept 1922, *Marcan 1000*

(ABD, BM, K).

Diagnostic characters. *Euploca marifolia* is characterised by its prostrate, much-branched with stiff bristly white hairs on the stems and leaves, lanceolate leaves (6–17 × 1–4 mm), revolute margin and inflorescence with leaf-like bracts.

3. *Euploca ovalifolia* (Forssk.) Diane & Hilger, Bot. Jahrb. Syst. 125(1): 48. 2003.

Fig. 1C–E

Heliotropium ovalifolium Forssk., Fl. Aegypt.-Arab.: 38. 1775. Type: Yemen, Al-Hadiyah [Hadie], Forsskål, 299 (holotype C [C10002362!], isotype BM [BM000795522!]).

Heliotropium coromandelianum J.Koenig ex Retz., Observ. Bot. 2: 9. 1781. Type: India, Koenig s.n. (holotype C [C10008743!]; isotype BM [BM000795508!]).

Heliotropium gracile R.Br., Prodr. Fl. Nov. Holland.: 493. 1810. Type: Australia, Queensland, Carpentaria, Brown s.n. (isotype MEL [MEL604552 image!]).

Type. Based on *Heliotropium ovalifolium* Forssk.

Distribution. Tropical Africa, Madagascar, Arabian Peninsula, Pakistan, India, Laos, Cambodia, Vietnam, Australia, Solomon Islands.

Ecology. Open area, sandy soil on riverbanks, rice fields, grassland, along roadsides; ca 200 m alt., flowering and fruiting from December to May.

Specimens examined. Thailand, Northern: Lamphun, Mae Ping NP, 22 May 2019, Thammarong et al. 673 (QBG); Tak, Tha Song Yang-Mae Sa Rieng, Moei river, 22 Mar 2006, Pooma et al. 6220 (BKF, L); Bhumibol Dam, 200 m alt., 29 May 2008, Pooma et al. 7073 (BKF); Ban Maesong, 23 June 2005, Pooma et al. 5443 (BKF); Mae Sa Rieng-Mae Sot road, 15 May 2007, Pooma et al. 6786 (BKF, E).

Vernacular. Nguang chang dok khao (งวงช้างดอกขาว).

Diagnostic characters. *Euploca ovalifolia* is newly recorded for Thailand. It is recognised by elliptic to obovate or oblanceolate leaves with silky silver hairs, spike-like or scorpioid cymes inflorescence with pedicels up to 2 cm long and a white corolla with a yellowish to yellow center. This species is similar to *Heliotropium indicum* in its inflorescence form but differs in fruit structure.

4. *Euploca paniculata* (R.Br.) M.W.Frohl. & M.W.Chase, Phytotaxa 434(1): 18. 2020.

Fig. 2A–C

Heliotropium paniculatum R.Br., Prodr. Fl. Nov. Holland.: 494. 1810. Type: Australia, Queensland: Sweer's Island, 28 Nov 1802, Brown 2930 (holotype BM [BM001040573!]; isotype K [K000998298!]).

Heliotropium tenunifolium var. *paniculatum* (R.Br.) Domin, Biblioth. Bot. 89: 549. 1928.

Heliotropium zeylanicum subsp. *paniculatum* (R.Br.) Kazmi, J. Arnold Arbor. 51: 156. 1970.

Type. Based on *Heliotropium paniculatum* R.Br.

Distribution. Pakistan, India, Myanmar, Cambodia, Australia.

Ecology. Open area of bare rock in dry dipterocarp and dry open deciduous forests, roadsides or wheat field; 50–1,000 m alt., flowering and fruiting from June to December.

Specimens examined. Thailand, Northern: Chiang Mai, 300 m alt., 19 Jan 1914, Kerr 3291

(BM, K); Tak, Ban Tak, 150 m alt., 1 Sept 2016, *Pooma & Pattharahirantricin* 7944 (BKF); Nakhon Sawan, 31 km north, 100 m alt., 2 June 1959, *Smitinand* 5815 (BKF); Ban Dong Lan, Wang Jan, Sam Ngao, 28 Aug 2010, *Norsaengsri* 7030 (QBG); Lamphun, Phra Puttabat Tak Pa, 24 Aug 2010, *Norsaengsri* 7205 (QBG). North-eastern: Udon Thani, Ban Phen, 24 Dec 1964, *Chermsirivathana* 225 (BK); Sakon Nakhon, Phu Phan NP, 400 m alt., 6 Aug 2004, *Nielsen et al.* 1542 (BKF); Bueng Kan, Ban Tong, Seka, 220 m alt., 20 May 2004, *Pooma et al.* 4116 (BKF); Tham Noi waterfall, 320 m alt., 30 July 2008, *Pooma et al.* 7345 (BKF); Khon Kaen, Mueang Phon, 200 m alt., 5 July 1967, *Smitinand* 12385 (BKF). Eastern: Chaiyaphum, Pa Khok Yai Chiu, 190 m alt., 28 Aug 1966, *S.N.* 235 (BKF); Nong Bua Deng, 400 m alt., 15 Aug 1972, *Larsen et al.* 31869 (AAU, BKF, E, K); Nakhon Ratchasima, Bannot Phattana, Road 226, km 42–41, 14 Sept 2004, *Pooma et al.* 4677 (AAU, BKF, E); Phimai, 200 m alt., 26 Aug 1958, *Smitinand* 4783 (BKF); *ibid.*, 5 Aug 1968, *Pradit* 316 (BK); Bua Yai, 200 m alt., 19 May 1931, *Kerr* 20488 (ABD, BK, BKF, BM, K, L); Ban Chum Seng, 22 May 1929, *Noe* 153 (BM, E, K); Surin, Chumphon Buri, 8 June 1982, *Paisooksantivatana & Sutheesorn* y 913–82 (BK); Roi Et, Kaset Wisai, 150 m alt., 22 June 1969, *Smitinand & Nalamphun* 10719 (BKF); *ibid.*, 9 June 1982, *Paisooksantivatana & Sangkhachand* y1000–82 (BK); Ubon Ratchathani, Khua Nangnee waterfall, Pha Taem NP, 200 m alt., 22 Aug 2001, *Pooma et al.* 2281 (BKF), Phu Chong Nayoi NP, 20 Oct 2009, *Middleton et al.* 5196 (BKF, E, K); *ibid.*, 250 m alt., 20 May 1998, *Suksathan* 1043 (QBG) *ibid.*, 100 m alt., 25 Jan 1924, *Kerr* 8345 (BK, BM, K). South-western: Kanchanaburi, Khao Meng, 14 Apr 1965, *Chantanamuck* 1067 (BK); 50 m alt., 13 Sept 1931, *Kerr* 20552 (BK, BM, E, K, L); *ibid.*, 50 m alt., 13 Sept 1931, *Kerr* 20553 (E, L); Prachinburi, Huai Saai, Cha-am, 19 Aug 2003, *Puudjaa* 1222 (BKF); Wat Luke Chang, Luke Chang Village, Tha Mai Ruak Subdistrict, Tha Yang, 5 July 2007, *Maxwell* 07–410 (QBG); Prachuap Khiri Khan, Hua Hin, 31 July 1976, *Maxwell* 76–455 (AAU, BK, L); Khao Tao, Hua Hin, *Smitinand* 1459 (BKF); *ibid.*, 10 Nov 1928, *Marcan* 2450 (BM, K); *ibid.*, 20 m alt., 9 Nov 1928, *Kerr* 16164 (AAU, BK, BM, K). Central: Lop Buri, Lam Narai, 50 m alt., 13 Nov 1975, *Smitinand* 12102 (BKF); *ibid.*, 50 m alt., 13 Nov 1975, *Smitinand* 12109 (BKF); Phatthana Nikhom, Ban Diluang, km 29–30, 100 m alt., 17 May 2004, *Pooma et al.* 3989 (BKF); Saraburi, 5 Sept 1954, *Bunnag* 10 (BK). Peninsular: Chumphon, Bang Son, Pathio, 20 m alt., 10 Jan 1927, *Kerr* 11351 (BK, BM, E, K, L).

Diagnostic characters. *Euploca paniculata* is widespread, and stems woody at the base. The distinguishing features are linear leaves with greyish tomentose hairs, the spike-like inflorescence arranged in one rank, elongate up to 10 cm long and flowers with pedicels up to 2 mm long. The herbarium specimens from Thailand, have mostly been confused with *Heliotropium strigosum*, but the two species can be recognised by vegetative characteristics.

5. *Euploca strigosa* (Willd.) Diane & Hilger, Bot. Jahrb. Syst. 125(1): 49. 2003.

Fig. 2D–F

Heliotropium strigosum Willd., Sp. Pl., ed. 4, 1(2): 743. 1798. Type: Ghana [Guinea], *Isert s.n.* (holotype B [B-W 219/3253, microfiche], isotypes C [C10003972!], P [P-JU6571 image!]).
Lithospermum chinense Hook. & Arn., Bot. Beechey Voy.: 202. 1837. Type: China, Canton, *Vachell* 286 (holotype E [E00369167!]; isotype BR [BR0000006966485 image!]).

Type. Based on *Heliotropium strigosum* Willd.

Distribution. Africa, Afghanistan, Pakistan, India, Nepal, Bhutan, Myanmar, Laos, Cambodia, Vietnam, China (Hainan, Kwangtung), Malesia, Australia.

Ecology. Open area of bare rock in dry dipterocarp and dry open deciduous forests; 50–400 m alt., flowering and fruiting from April to November.

Specimens examined. Thailand, North-eastern: Sakon Nakhon, Phu Phan NP, 400 m alt., 6 Aug 2004, *Nielsen et al.* 1542 (BKF); Phrae, Ban Nun, Song, 193 m alt., 25 June 2012, *Norsaengsri & Tathana* 9480 (QBG). Eastern: Surin, Nadi, 17 May 1965, *Sakol* 224 (BK); Ubon Ratchathani, Chiet, 100 m alt., 21 May 1932, *Kerr* 21537 (BK, BM, E, K). Central: Chai Nat, Utapao, 20 Sept 1930, *Kerr* 19692 (ABD, BM, BK). South-western: Kanchanaburi, 5 km west of Thong Pha Phum Town, 16 Oct 2015, *Tanming* 873 (QBG); Thong Pha Phum, along route 323, 4 km, 240 m alt., 29 Nov 1982, *Koyama et al.* T-30469 (BKF); Prachinburi, Watananakorn, 23 July 1987, *Paisooksantivatana & Sangkhachand* 2109-87 (BK). South-eastern: Sa Kaeo, Aranyaprathet, 50 m alt., 3 Apr 1930, *Kerr* 19575 (BM, BK, K). Peninsular: Songkhla, Padang Besar, 50 m alt., 24 Dec 1927, *Kerr* 13602 (BM, BK, K).

Diagnostic characters. *Euploca strigosa* is similar to *E. paniculata*, but differs in the former is the prostrate to much-branched stem (vs erect unbranched to few-branched stem in *E. paniculata*), the narrowly elliptic leaves with strigose hairs on both surfaces, 5–10 × ca 3 mm (vs linear leaves with greyish tomentose hairs, 7–40 × 1–2 mm in *E. paniculata*).

***Heliotropium* L., Sp. Pl.: 130. 1753.**

Type species. *Heliotropium europaeum* L.

Description. Annual herbs or small trees. Leaves with petiole; lamina ovate, elliptic to obovate, apex acute, obtuse or rounded, margin entire or irregularly undulate, strigose, harsh hairs or greyish white tomentose on both surfaces. Inflorescences terminal or axillary, scorpioid cymes or subcorymbose, ebracteate. Flowers 5-merous. Calyx 5-lobed, lobes lanceolate. Corolla pale green, white to pinkish white, greenish to yellowish, with a yellow-orange throat inside, funnel-shaped, lobes broadly ovate to orbicular, throat glabrous inside. Stamens included, sessile or with short filaments; anthers ovate-lanceolate. Ovary entire. Fruits drupe, deeply bilobed, mesocarp thinly fleshy, endocarp dividing into two 2-seeded nutlets.

Key to the species of *Heliotropium*

- 1 Small trees; leaves 8–20 cm long with densely greyish tomentose hairs **1. *H. foertherianum***
- Annual herbs; leaves 3–6 cm long with strigose and harsh hairs **2. *H. indicum***

1. *Heliotropium foertherianum* Diane & Hilger, Bot. Jahrb. Syst. 125(1): 46. 2003.

Fig. 3A–C

Tournefortia argentea L.f., Suppl. Pl.: 133. 1781. Type: Sri Lanka, *Köenig s.n.* (lectotype LINN 193/6).

Messerschmidia argentea (L.f.) I.M.Johnst., J. Arnold Arbor. 16: 164. 1935.

Argusia argentea (L.f.) Heine, Fl. Nouv.-Calédonie & Dépend. 7: 109. 1976.

Type. Based on *Tournefortia argentea* L.f.

Distribution. China (Hainan), Taiwan, Japan (Ryukyu), Vietnam, Malesia, New Guinea, Australia.

Ecology. Sandy beaches, flowering and fruiting from November to February.

Specimens examined. Thailand, Peninsular: Surat Thani, Ban Bua Put, Ko Samui, 14 May 1928, *Kerr 15728* (BK, E, K, TCD); Phangnga, Similan NP, 14 Nov 1996, *Santisuk s.n.* (BKF); Krabi, Lan Ta NP, 7 Jan 1992, *Niyomdham 2854* (BKF); Ko Lan Ta Yai, 20 Mar 1998, *Chamchumroon 37* (BKF); Satun, Adang, La-ngu, 15 Jan 1928, *Kerr 14095* (BK, K).

Vernacular. Nguang chang thale (เงาะช้างทะเล).

Diagnostic characters. *Heliotropium foertherianum* can be recognised by a small tree of seashore plant, oblanceolate or obovate leaves up to 20 cm long with densely greyish tomentose hairs, the corolla as long as the calyx, fleshy fruit with fleshy mesocarp and spongy endocarp. In Thailand known as *Tournefortia argentea* or *Argusia argentea*, the currently accepted name is *Heliotropium foertherianum*.

2. *Heliotropium indicum* L., Sp. Pl.: 130. 1753.

Fig. 3D–F

Type. India utraque, Herb. Hermann 1: 9, no.70 (lectotype, designated by Mill in Cafferty and Jarvis 2004, p.802: BM [BM00061256!],).

Distribution. North America, South America, Pakistan, India, Sri Lanka, China, Myanmar, Taiwan, Japan, Laos, Cambodia, Vietnam, Sumatra, Malaysia, Java, Borneo, Australia.

Ecology. Common on sandy soil near stream or open areas, rice fields, roadsides; 50–1,000 m alt., flowering and fruiting from January to April.

Specimens examined. Thailand, Northern: Mae Hong Son, Mueang, 256 m alt., 10 Sept 2013, *Lakoet 0352* (QBG); 350 m alt., 18 June 1973, *Geesink et al. 5937* (AAU, BKF, E, K, TCD); Chiang Mai, Doi Sa Ket, 12 Dec 2007, *Warintorn 07–105* (QBG); San Kamphang, 325 m alt., 10 July 1996, *Panatkool et al. 37* (BKF); Doi Suthep, 300 m alt., *Hosseus s.n.* (K); *ibid.*, 1,000 m alt., 12 May 1915, *Winit 402* (K, E); *ibid.*, 1,000 m alt., 5 Jan 1911, *Kerr 1845* (BM, K, E); Hod, 8 Sept. 1996, *BGO Staff 28* (QBG); Ban Chang Keong, 488 m alt., 28 May 2019, *Pingyot 276* (QBG); Chiang Rai, Thoeng, 12 Feb 2011, *Norsaengsri & Tathana 7444* (QBG); San Pa Tong, 325 m alt., 26 Apr 1988, *Maxwell 88–521* (BKF, CMUB); Phayao, Ban Khun Kam Lang, Pong, 400 m alt., 19 May 2017, *Pongamorkul 6201* (QBG); Nan, Na Noi, 9 Apr 2018, *Khattiyot et al. 778* (QBG); Huai Haeng, 318 m alt., 26 Apr 2017, *Muaengyen 2011* (QBG); Lampang, Sop Prap, 28 Feb 2012, *Norsaengsri et al. 9055* (QBG); Uttaradit, Nam Pat, 22 Mar 2011, *Romklao Botanical Garden 0224/2554* (QBG); Tak, Mae Sot, 22 July 1959, *Floto 7653* (C); Tha Song Yang, 15 May 2007, *Pooma et al. 6785* (BKF, E, L); Phitsanulok, Tham Pha Phon Non-Hunting Area, 5 Mar. 2012, *Maknoi 4627* (QBG); Wat Hua Khao, 31 Jan 2012, *Maknoi 4565* (QBG). North-eastern: Loei, Na Haeo, Yaa Ngoung Chaang, 31 July 1995, *Nanakorn et al. 4077* (QBG); Phu Kradueng, 27 Aug 1988, *Tamura T-60450* (BKF); Sithan, Huai Yang, 300 m alt., 21 Feb 1958, *Ploenchit 1342* (BKF); Bueang Kan, Mueang, 27 Dec 2011, *Norsaengsri & Tathana 8581* (QBG); Maha Sarakham, Kosum Phisai, 3 May 2001, *Norsaengsri et al. 1440* (QBG); Ban Maong Ai, 130 m alt., 3 Apr 2001, *Norsaengsri et al. 1440* (QBG); Chiang Yuen, 14 Apr 2018; Khon Kaen, Chum Phae, 2 Sept 1967, *Shimizu et al. T-8685* (BKF, E, TCD). Eastern: Nakhon Ratchasima, Bua Yai, 18 Mar 2009, *Norsaengsri 5135* (QBG); Ban Chum Sang, 25 May 1929, *Nai Noe 277* (ABD, BM, BK, K); Ubon Ratchatani, Khong Chiam, 170 m alt., 25 May 2001, *Greijmans 90* (BKF); Phibun Mangsahan, 250 m alt., 9 Dec 1982, *Koyama et al. T-30698* (AAU, BKF); Det Udom, 270 m alt., 10 Dec 1982, *Koyama et al. T-30778* (BKF); Buri Ram, 7 Aug 1970, *Sutheesorn 2065* (BK); Ban Na, Suwan Naphum, 16 Aug 1982, *Sutheesorn & Saraphunphichit 5354* (BK). South-western: Kanchanaburi, Si Sawat, 150 m alt., 6 Nov 1971, *van Beusekom et al. 3955* (BKF, C, K); Thum

Pha, 13 Dec 1961, *Phengklai* 228 (BKF); Kwae Noi, 30 Apr 1946, *Wichian* 304 (K); Na Ta Hom, 23 Apr 1965, *Chantanamuck* 1094 (BK); Prachuap Khiri Khan, 18 Aug 1967, *Shimizu et al.* T-7689 (AAU, BKF, E, K). Central: Chat Nat, 5 Mar 1958, *Sørensen et al.* 1922 (E, C); Pathum Thani, 50 m alt., 12 Mar 1958, *Smitinand* 4444 (BKF); Nakhon Nayok, Khao Yai, 1,000 m alt., 5 May 1964, *R.S.* 116 (BKF); 40 m alt., 6 Aug 1970, *Phengklai et al.* 3748 (BKF); Nonthaburi, Sai Maa Tai, 14 Apr 1992, *Puudjaa* 65 (BKF); Bangkok, 5 July 1920, *Marcen* 293 (BM, K); 27 Oct 1919, *Kerr* 3840 (ABD, BM, C, K); 19 Apr 1926, *Lakshnakara* 52 (BK); Sept 1921, *Smith* 49 (BK); 1899, *Zimmermann s.n.* (K); Ang Thong, 30 Dec 1929, *Put* 2595 (AAU, ABD, BK, BM, K). South-eastern: Chachoengsao, Phanom Sarakham, 50–100 m alt., 1 Oct 1984, *Murata et al.* T-37042 (BKF); Chon Buri, Si Racha, 30 Nov 1927, *Collins* 2025 (ABD, BM, BK, K, TCD); Chanthaburi, 8 Apr 1959, *Sørensen et al.* 7214 (C). Peninsular: Chumphon, 31 Jan 1958, *Sørensen et al.* 829 (BKF, C); Ranong, Kapoe, 19 Jan 2016, *Kertsawang* 3937 (QBG); Satun, La-ngu, Ko Kabeng, 11 Apr 2003, *Phengklai et al.* 13650 (BKF); Songkhla, 21 Aug 1995, *Larsen et al.* 45836 (AAU, BKF); Satun, Terutao, 5 Mar 1966, *Hansen & Smitinand* 12542 (BKF, C).

Vernacular. Ya nguang chang (หญ้าเงาช้าง)

Diagnostic characters. *Heliotropium indicum* is a common medicinal weed species and most easily recognised by ovate to elliptic leaves with strigose and harsh hairs on both surfaces, truncate or obtuse base, irregularly undulate margin, white to purple or purplish corolla with a yellow-orange throat inside, fruit in which its ribs angular, deeply 2-lobed with apical divergent lobes.

***Tournefortia* L., Sp. Pl.: 140. 1753.**

Argusia Boehm. in Ludwig, Def. Gen., ed. Boehmer: 507. 1760. Type species: *Argusia sibirica* (L.) Dandy.

Messerschmidia L. ex Hebenstr., Nov. Comm. Acad. Sci. Imp. Petrop. 8: 315, t. 11. 1763. Type species: unknown.

Tetrandra (DC.) Miq., Fl. Ind. Bat. 2: 928. 1858. Type species: *Tetrandra wallichii* Miq. [= *Tournefortia tetrandra* Blume].

Mallotonia (Griseb.) Britt., Ann. Missouri Bot. Gard. 2: 47. 1915. Type species: *Tournefortia gnaphalodes* R.Br.

Type species. *Tournefortia hirsutissima* L. [= *Heliotropium verdcourtii* Craven]

Description. Climbing shrubs. Leaves with petiole; lamina ovate or elliptic to lanceolate, apex acute to acuminate, margin entire, tomentose or strigose on both surfaces. Inflorescences terminal or axillary, dichotomously branched, dense 2-rowed scorpioid cymes, ebracteate. Flowers 4–5-merous. Calyx 5-lobed, lobes lanceolate to ovate. Corolla pales green, white, greenish to yellowish, tube cylindrical, lobes spreading. Stamens included, with short filaments; anthers elliptic-oblong. Ovary entire. Fruits drupe and distinctly lobed, mesocarp fleshy and juicy, endocarp dividing into two 2-seeded nutlets.

Key to the species of *Tournefortia*

- 1 Flower 4-merous; lower surface of leaves pale, brownish-black when dry **5. *T. tetrandra***
- Flower 5-merous; both surfaces of leaves the same colour, light brown when dry **2**
- 2 Leaves oblong-elliptic to ovate-lanceolate lanceolate, apex acute **3**
- Leaves ovate, apex long acuminate **4**

- 3 Leaves oblong-elliptic; inflorescence dense dichotomous branched; corolla tube angular **2. *T. intonsa***
- Leaves ovate-lanceolate; inflorescence loosely dichotomous branched; corolla tube not angular **3. *T. montana***
- 4 Leaves with minutely tuberculate **4. *T. ovata***
- Leaves with golden brown tomentose hairs **1. *T. hookeri***

1. *Tournefortia hookeri* C.B. Clarke, Fl. Brit. India 4: 147. 1883.

Tournefortia viridiflora Gamble, Darjeeling List: 57. 1878. nom. illeg.

Type. Sikkim, Himalayas, *Hooker s.n.* (holotype K [K000998133!]).

Distribution. East Himalaya, Myanmar.

Ecology. Deciduous forest, flowering and fruiting in June.

Specimens examined. Thailand, Northern: Chiang Mai, Ban Pa Kar, Samoengtai, Samoeng, 1,424 m alt., 26 June 2008, *Jatupol 08–275* (QBG).

Diagnostic characters. *Tournefortia hookeri* is newly recorded for Thailand, a specimen of this distinctive species has been seen from QBG. This species is recognisable by broadly ovate leave with dense golden brown on the lower surface and loosely dichotomous cymose.

2. *Tournefortia intonsa* Kerr, Bull. Misc. Inform. Kew 1940: 185. 1940.

Fig. 4A

Type. Thailand, Doi Sutep, ca 900 m, evergreen forest, *Kerr 2285* (holotype K [K000998142!]; isotypes BM [BM001209041!], E [E00766166!]).

Distribution. Laos.

Ecology. In deciduous forest, evergreen forest, flowering and fruiting from November to February.

Specimens examined. Thailand, Northern: Mae Hong Son, Pang Mapha, 580 m alt., 26 Feb 1968, *Hansen & Smitinand 12741* (AAU, BKF, C, E, K); Chiang Mai, Doi Chiang Dao, 20 Mar 1956, *Suvarnakoses 1121* (BKF, K); *ibid.*, 700 m alt., 18 Feb 1958, *Sørensen et al. 1312* (BKF); *ibid.*, 23 Dec 1931, *Put 4570* (BK, E, K); *ibid.*, 600–800 m alt., 14 Mar 1956, *Garrett 1449* (K, L); *ibid.*, 600 m alt., 14 Mar 1956, *Garrett 1476* (K, L); *ibid.*, 1,500 m alt., 17 Feb 1958, *Smitinand 4218* (BKF); *ibid.*, 525 m alt., 26 Feb 1989, *Maxwell 89–273* (BKF, CMU, CMUB, L); *ibid.*, 550 m alt., 10 Mar 1990, *Maxwell 90–294* (CMU); Chiang Rai, Doi Tung, 750 m alt., 15 Feb 2012, *van de Bult 1241* (BKF); Mae Sai, Huai Nam Dang, 612 m alt., 13 Jan 2011, *Norsaengsri & Tathana 7537* (BKF, QBG); Mae Sai, Ban Phamee, 572 m alt., 15 Feb 2012, *Norsaengsri & Tathana 9003* (QBG); Hua Yane, Mae Chan, 25 Jan 1970, *Sutheesorn 1438* (BK); Doi Tung, 1,400 m alt., 29 Jan 1989, *Bragg 8* (CMU, L); Phrae, Mae Khaen stream, 440 m alt., 5 Jan 1972, *van Beusekom, et al. 4636* (C, BKF, K); *ibid.*, 22 Mar 1912, *Vanpruck 454* (K); Mae Yom NP, 400 m alt., 15 Dec 1993, *Maxwell 93–1509* (CMUB, L)]. North-eastern: Phetchabun, Chon Daen, 15 Jan 1969, *Vacharapong 304* (BK); Loei, Phu Suan Sai NP, 11 Mar 2008, *Maknoi 2029* (BKF, QBG); route to Sam Nuck Bab, 15 May 2008, *Maknoi & Srisaga 2302* (BKF, QBG); Tad Hueang waterfall, 1,000 m alt., 16 May 2006, *Maknoi 824* (QBG); Na Haeo, 650 m alt., 27 Apr 1995, *Nanakorn et al. 3239* (QBG); Khon Kaen, Pha Bhroung Cave, Chum Phae, 14 Feb 1963, *Chantanamuck 295* (BK). South-western: Uthai Thani, Huai Haeng, 27 Apr 1963, *Kasem*

329 (BK). South-eastern: Chanthaburi, Pong Nam Ron, 600 m alt., 12 Jan 1956, *Smitinand 3192* (BKF).

Vernacular. Ya Nguang Chang Luang (หญ้างวงช้างหลวง)

Diagnostic characters. *Tournefortia intonsa* is similar to *T. montana* in its inflorescence form but differs in its oblong or elliptic leaves (vs ovate-lanceolate leaves in *T. montana*), densely greyish tomentose hairs on the inflorescence (vs glabrous to minutely strigose hairs in *T. montana*).

3. *Tournefortia montana* Lour., Fl. Cochinch. 1: 122. 1790.

Fig. 4B

Tournefortia viridiflora (Lehm.) Wall., Fl. Ind. 2: 4–5. 1824. Type: India, *Wallich Numer. List 907* (holotype K [K001110251!]).

Tournefortia sampsonii Hance, J. Bot. 6: 330. 1868. Type: China, Sai-chii-shan, Cantonesis, *Sampson 13035* (holotype BM [BM001014454!]).

Tournefortia boniana Gagnep., Noutul. Syst. (Paris) 3: 33. 1914. Type: Vietnam, Tonkin m ridional, *Bon 1932*, (lectotype, designated here: P [P03897617 image!]; isoelectotypes: P [P03897616 image!], K [K000998141!]).

Tournefortia gaudichaudii Gagnep., Notul. Syst. (Paris) 3: 34. 1914. Type: Vietnam, Tourane, *Gaudichaud 180* (holotype K [K000998140!]).

Tournefortia brachyantha Merr. & Chun, Sunyatsenia 2: 314. 1935. Type: China, Jiangsu, Hainan, *How70424* (holotype K [K000998121!]).

Type. not indicated.

Distribution. India (Assam), Bangladesh, southern China, Myanmar, Laos, Vietnam, Cambodia,

Ecology. Open roadside, along the stream in evergreen forest, bamboo forest, slope of the hill in evergreen forest and mixed deciduous forests, 250–1,500 m alt., flowering and fruiting from February to December.

Specimens examined. Thailand, Northern: Chiang Rai, Mae Chan, 450 m alt., 17 Mar 2005, *Pooma et al. 4859* (BKF); Chiang Mai, Mae Hia Thai Literary Botanical Garden, 350 m alt., 13 June 2003, *TLBG 165* (QBG); 300 m alt., 21 Jan 1921, *Kerr 4692* (BM, BK, K); Nan [Doi Phu Kha NP 1,500 m alt., 24 Jan 2003, *Srisanga 2684* (QBG); *ibid.*, 11 Apr 2002, *Srisanga 1253* (QBG); *ibid.*, 1,200 m alt., 19 Mar 2000, *Srisanga 1340* (BKF, QBG); Ban Den Thara, Pra That, Chiang Klang District, 346 m alt., 15 Mar 2018, *Khattiyot et al. 616* (QBG); Tam Paa Toop, Ta Wang Paa, 10 km from Nan, 360 m alt., 24 Feb 1998, *Srisanga et al. 220* (QBG); Lampang, Doi Luang NP, Wahng Gaye waterfall, 625 m alt., 26 Mar 1997, *Maxwell 97-753* (BKF); Uttaradit, Phu Soi Dao NP, 1,000 m alt., 18 Mar 2002, *Chamchumroon & Puff V.C. 1438* (BKF); Tak, Doi Muser, Hui Sakulee, 700 m alt. 27 Feb 1987, *Paisooksantivatana Y2032-87* (BK); Mae Sot to Mae Sariang, 19 Jan 1995, *Pooma 1007* (BKF, CMUB). South-western: Uthai Thani, Ban Rai, 4 Mar 1977, *Sutheesorn 4017* (BK); Kanchanaburi, Sangkhla Buri to Thong Pha Phum, 180 m alt., 17 Dec 2009, *Pooma et al. 7436* (BKF); Sai Yok, Khwae Noi, 21 Dec. 1961, *Larsen 8893* (BKF, C, K); *ibid.*, 21 Dec 1961, *Simitnand 9500* (BKF); Sai Yok, 23 Dec 1961, *Larsen 8916* (C); Pompee village near Khwae Noi, 250 m alt., 26 Mar 1968, *van Beusekom & Phengklai 129* (BKF, E, K, L); Mae Nam Noi near waterfall, 2 Jan 1962, *Phengklai 398* (BKF); Huai Banlcae, 8 Nov 1971, *van Beusekom et al. 3528* (BKF); Thong Pha Phum NP, 260 m alt., 3 Dec. 2003, *Sirimongkol 69-2* (BKF). Peninsular: Krabi, Ban Nai Chong, 17 Dec 1965, *Umpai 208* (BK);

Khap Thong Thai, 19 Jan. 1966, *Hansen & Smitinand 12002* (BKF, C); Nakhon Si Thammarat, Nop Phitum, 80 m alt., 13 Feb 2005, *Williams et al. 1475* (BKF, E).

Diagnostic characters. *Tournefortia montana* is newly recorded for Thailand and can be recognised by ovate-lanceolate leaves with glabrous or minutely strigose on both surfaces and minutely tuberculate on the lower surface. The specimens from Thailand have mostly been misidentified as *T. intonsa*. In the protologue, Gagnepain (1914) described *T. boniana* based on two different collections (*Bon 1357 & Bon 1932*). *Bon 1932* P[P03897617] is chosen as lectotype because it is the best preserved, with complete inflorescences and numerous flowers, and has two duplicates as isolectotypes.

4. *Tournefortia ovata* Wall. ex G.Don, Gen. Hist. 4: 369. 1837.

Type. Myanmar, Rangoon, *Wallich Numer. List 908* (holotype K [K001110253!]).

Distribution. India (Andaman Islands, Nicobar Islands), Myanmar.

Ecology. Open or shaded, slightly disturbed area in mixed evergreen forest 400–500 m alt., flowering and fruiting from November to April.

Specimens examined. Thailand, Northern: Chiang Rai, Tham Luang-Khun Nam Nang Norn NP, Mae Sai, 560 m alt., 27 Mar 2012, *Norsaengsri & Tathana 9227* (QBG); Mae Tak, 3 Mar 1958, *Sørensen et al. 1865* (BKF); Chiang Mai, Gnai, Tintok, 500 m alt., 8 Mar 1965, *C.A. & B.S. 270* (BKF); Doi Chiang Dao, 750 m alt., 29 Mar 1995, *Maxwell 95–269* (BKF); *ibid.*, 575 m alt., 11 Mar 1989, *Maxwell 89–317* (BKF, CMU, L); *ibid.*, 550 m alt., 8 Mar 1965, *Hambanond 270* (BK); Phayao, Doi Pha Dam, Rom Yen Subdistrict, Chiang Kham, 890 m alt., 21 Feb 2013, *La-onsri et al.*, 2662 (QBG); Lampang, Mae Pukaung, 6 Mar 1925, *Winit 1282* (BK, K); Nan, Song Khwae, Nam Pan Village, 350 m alt., 12 Jan 2011, *Srithi 630* (QBG); Tak, Lan Sang NP, 2 Jan 1969, *Cheviwat & Nimanong 10* (BKF); Phrae, 500 m alt., 22 Mar 1913, *Vanpruck 454* (BKF, K); Huai Rai, 13 Mar 1961, *Chanthamuk 6* (BKF); *ibid.*, 13 Mar 1961, *Chanthamuk 9* (BKF); *ibid.*, 23 Mar 1961, *Phengklai 66* (BKF, C, K); 26 Jan 1913, *Vanpruck 357* (BKF, K); Sukhothai, Srisatchanalai NP, route to Tad Dao waterfall, 3 Feb 2015, *Maknoi 7243* (QBG). North-eastern: Phetchabun, Nam Nao NP, 18 Feb 2014, *Maknoi 6553* (QBG). Eastern: Nakhon Ratchasima, Bua Yai, 400 m alt., 29 Nov 1924, *Kerr 9487* (BM, BK, E, K, L). South-western: Kanchanaburi, Si Sawat, Dongyai, 50 m alt., 17 Aug 1971, *Phengklai et al. 3016* (BKF), *ibid.*, 600 m alt., 17 Jan 1926, *Kerr 10233* (BM, BK, K, E); Thung Yai Naresuan WS, Sangklaburi, 475 m alt., 11 Apr 1994, *Maxwell 94–489* (CMUB, L); Prachuap Khiri Khan, Pran Buri, 25 Nov 1929, *Put 2449* (BM, BK, K).

Vernacular. Liang (เหลียง).

Diagnostic characters. *Tournefortia ovata* is similar to *T. tetrandra* in its ovate or ovate-lanceolate leaf shape but differs in the 5-merous flower and dense minutely tuberculate on the lower leaf when dry (vs 4-merous flowers and minutely tuberculate on both leaf surfaces in *T. tetrandra*).

5. *Tournefortia tetrandra* Blume, Bijdr. Fl. Ned. Ind. 14: 845. 1826.

Tournefortia wallichii DC., Prodr. 9: 527. 1845. Type: Singapore, Penang, *Wallich Numer. List 911* (holotype K-W [K00111261]; isotypes K-W [K00111262, K000998138], M [M0188731 image!], GZU [GZU000106058 image!], MEL [MEL2502678 image!]).

Tournefortia tetrandra Blume var. *angustifolia* Moritzi, Syst. Verz. 52: 1845. Type: Java, *Zollinger 939* (lectotype, designated here: K[K000998147!]; isolectotype: K[K000998148!])

Tetrandra zollingeri Miq., Fl. Ind. Bat. 2: 928. 1858. Type: Java, Zollinger 395 (holotype K [K000998149!])

Heliotropium biblianum Craven, Blumea 50(2): 379. 2005. Type: Based on *Tournefortia tetrandra* Blume.

Type. Java, Blume s.n. (holotype L [L0281636!]).

Distribution. India (Nicobar Islands), Sri Lanka, Cambodia, Vietnam, Malesia, New Guinea

Ecology. Deciduous forest, evergreen forest, 40–700 m alt., flowering and fruiting from April to December.

Specimens examined. Thailand, South-eastern: Kanchanaburi, Thong Pha Phum, 700 m alt., 15 Dec 1995, *FRUD & van Welzen 111* (BKF). South-western: Prachuap Khiri Khan, Pa La U. trail, 260 m alt., 20 Jun 2004, *Middleton et al. 2322* (BKF). Peninsular: Ranong, Kaper, 40 m alt., *Shimizu et al. T-26364* (AAU, BKF); *ibid.*, 500 m alt., 19 June 1932, *Kerr 21710* (BK, BM, E, K); Surat Thani, Ko Pha-ngan, 10 Nov 1927, *Put 1247* (BM, BK, E, K); *ibid.*, 50 m alt., 3 Mar 1927, *Kerr 12456* (BK, BM, E, L, K); Bandon, 2 Jan 1935, *Seidenfaden 2089* (K); Phangnga, 500 m alt., 9 Dec 1928, *S.N. 3994* (K, SING); Trang, Khao Pap Pa, 13 Mar 1974, *Larsen & Larsen 33277* (AAU); Yan Ta Khao, Sai Roong waterfall, 400 m alt., 26 Apr 1987, *Maxwell 87–434* (BKF, CMU, L); Phuket, Khao Pha Tao Non-Hunting Area, Bang Pae waterfall, 50 m alt., 14 Oct 2004, *Gardner & Sidisunthorn 44268* (QBG).

Vernacular. Fa Ta Hueng (ฟ้าตาหึ่ง)

Diagnostic characters. *Tournefortia tetrandra* can be recognised mainly by its tetramerous flower, acute apex leaf, thick leaves with minutely tuberculate on both leaf surfaces, corolla up to 8–10 mm long, fruit globose up to 8 mm in diameter. *Tournefortia tetrandra* var. *angustifolia*, there are two known duplicates of the type collection, and K [K000998147] is designated here as the lectotype because it is the best-preserved specimen.

Acknowledgements

We would like to thank the curators and staff of all the herbaria visited for providing access to the herbarium collection. We particularly thank Dr Siriporn Zungsontiporn of the National Weed Science Research Institute, Ministry of Agriculture and Cooperatives for the specimens and photographs of *Euploca ovalifolia* and Dr Pramote Triboun of the Thailand Institute of Scientific and Technological Research for his photograph of *Euploca bracteata*. This work was funded by the National Research Council of Thailand (NRCT5-RSA63018-01).

References

- Antony AA, Javad P (2020) The *Heliotropium marifolium* complex in India and five new combinations in *Euploca* (Heliotropiaceae) from *H. sect. Orthostachys* (Boraginaceae s.l.). *Nordic Journal of Botany* 38(11): 1–6. <https://doi.org/10.1111/njb.02913>
- Backer CA, Bakhuizen van den Brink Jr RC (1965) Boraginaceae In: Backer CA, Bakhuizen van den Brink Jr RC (Eds) *Flora of Java*. Vol. 2. Noordhoff, Groningen, the Netherlands, 457–464.
- Cafferty S, Jarvis EC (2004) Typification of Linnaean plant names in Boraginaceae. *Taxon* 53(3): 799–805. <https://doi.org/10.2307/4135454>

- Clarke CB (1885) Boraginaceae. In: Hooker JD (Ed.) The Flora of British India. Vol. 4. L. Reeve & Co., London, 134–179.
- Craven LA (2005) Malesian and Australian *Tournefortia* transferred to *Heliotropium* and notes on delimitation of Boraginaceae. *Blumea* 50(2): 375–381. <https://doi.org/10.3767/000651905X623049>
- Diane N, Förther H, Hilger HH (2002) A systematic analysis of *Heliotropium*, *Tournefortia*, and allied taxa of the Heliotropiaceae (Boraginaceae) based on ITS1 sequences and morphological data. *American Journal of Botany* 89(2): 287–295. <https://doi.org/10.3732/ajb.89.2.287>
- Diane N, Hilger HH, Förther H, Weigend M, Luebert, F (2016) Heliotropiaceae. In: Kadereit JW, Bittrich V (Eds) Flowering Plants. Eudicots, The Families and Genera of Vascular Plants. Vol. 14. Springer International Publishing, Switzerland, 203–211.
- Feuillet C (2016) Two new combination in *Euploca* Nutt. (Heliotropiaceae, Boraginales) and a conspectus of the species of the Guiana Shield area. *PhytoKeys* 61: 101–124. <https://doi.org/10.3897/phytokeys.61.6260>
- Fletcher HR, Kerr AFG (1951) Boraginaceae. In: Craib WG, Kerr AFG (Eds) *Florae Siamensis Enumertio, A List of the Plants Known from Siam*. Vol. 3. Siam Society, Bangkok. 77–88.
- Förther H (1998) Die infragenerische Gliederung der Gattung *Heliotropium* L. und ihre Stellung innerhalb der subfam. Heliotropioideae (Schräd.) Arn. (Boraginaceae). *Sendtnera* 5: 35–241. <https://www.biodiversitylibrary.org/part/146614>
- Frohlich MW, Thulin M, Chase MW (2020) Ninety-three new combinations in *Euploca* for species of *Heliotropium* section *Orthostachys* (Boraginaceae *sensu* APG). *Phytotaxa* 434(1): 013–021. <https://doi.org/10.11646/phytotaxa.434.1.2>
- Gagnepain F, Courchet L (1914) Boraginaceae. In: Lecomte H (Ed.) *Flore Generale de L'Indo-Chine*. Vol. 4. Masson et Cie, Paris, 197–226.
- Gagnepain F (1914) Boraginaceae nouvelles ou peu connues d'extrême-orient. *Notulae Systematicae Herbarii du Muséum de Paris* 3: 33–34.
- Hilger HH, Diane N (2003) A systematic analysis of Heliotropiaceae (Boraginales) based on trnL and ITS1 sequence data. *Botanische Jahrbücher für Systematik* 125: 19–51. <https://doi.org/10.1127/0006-8152/2003/0125-0019>
- Johnston IM (1935) Studies in the Boraginaceae. XI. 1. The species of *Tournefortia* and *Messerschmidia* in the Old World. *Journal of Arnold Arboretum* 16: 145–168.
- Johnston IM (1951) Studies in the Boraginaceae, XX representatives of three subfamilies in eastern Asia. *Journal of Arnold Arboretum* 32: 99–122.
- Luebert F, Brokamp G, Wen J, Weigend M, Hilger HH (2011) Phylogenetic relationships and morphological diversity in Neotropical *Heliotropium* (Heliotropiaceae). *Taxon* 60(3): 663–680. <https://doi.org/10.1002/tax.603004>
- Luebert F, Cecchi L, Frohlich MW, Gottschling M, Williams M, Hasenstab-Lehman KE, Hilger HH, Miller JS, Mittelbach M, Nazaire M, Nepi M, Nocentini D, Ober D, Olmstead RG, Selvi F, Simpson MG, Sutory K, Valdes B, Walden GK, Weigend M (2016). Familial classification of the Boraginales. *Taxon* 65(3): 502–522. <https://doi.org/10.12705/653.5>
- Mabberley DJ (2017) *Mabberley's Plant-Book: A portable dictionary of plants, their Classification and uses*. 4th edition. Cambridge University Press New York, 754 pp.
- Mill RR (1999) Boraginaceae. In: Grierson AJC, Long DG (Eds) *Flora of Bhutan*. Vol. 2. The Royal Botanic Garden Edinburgh, 865–881.
- Ridley NH (1923) Boraginaceae. The flora of the Malay Peninsula. Vol. 2. L. Reeve & CO.

- London, 438–442.
- Riedl H (1997) Boraginaceae. In: Kalkman C, Kirkup DW, Nooteboom HP, Stevens PF, deWilde WJJO (Eds), *Flora Malesiana, Series I*, 13: RA Leiden, The Netherlands, 43–144.
- Thiers B (2022, continuously updated). *Index Herbariorum*, a global directory of public herbaria and associated staff. New York Botanical Garden's Virtual Herbarium. The New York. <https://sweetgum.nybg.org/science/ih/>
- Zhu GL, Riedl H, Kamelin R (1995) Boraginaceae. In Wu ZY, Raven PH (Eds) *Flora of China*. Vol. 16. Science Press, Beijing, China, 329–427.




Figure 1 Morphology of *Euploca* in Thailand. **A** *E. bracteata* (R.Br.) M.W.Frohl. & M.W.Chase **B** *E. marifolia* (J.Koenig ex Retz.) Ancy & P.Javad **C–E** *E. ovalifolia* (Forssk.) Diane & Hilger: **C** inflorescence **D** dehiscing fruit separating into four nutlets. **A** Photo by Pramote Triboun **C–D** Photos by Siriporn Zungsontiporn.



Figure 2 Morphology of *Euploca* in Thailand. **A– C** *E. paniculata* (R.Br.) M.W.Frohl. & M.W.Chase: **A** habit and inflorescences **B** inflorescence **C** young fruits **D– F** *E. stigosa* (Willd.) Diane & Hilger: **D** inflorescence **E** habit **F** habit and inflorescences. Photos by Kanokorn Rueangsawang.



Figure 3 Morphology of *Heliotropium* in Thailand. **A–C** *H. foertherianum* Diane & Hilger: **A** inflorescence **B** habit **C** flowers **D–F** *H. indicum* L.: **D** inflorescences **E** habit and inflorescence **F** young fruit. Photos by Kanokorn Rueangsawang.




Figure 4 Morphology of *Tournefortia* in Thailand. **A** *T. intonsa* Kerr **B** *T. montana* Lour. Photos by Kanokorn Rueangsawang.