

Upscaling biomass production of rosmarinic acid-rich hairy root cultures of *Agastache rugosa*

Weronika Kozłowska

Wroclaw Medical University: Uniwersytet Medyczny im Piastow Slaskich we Wroclawiu

Ewelina Piątczak

Medical University of Lodz: Uniwersytet Medyczny w Lodzi

Joanna Kolniak-Ostek

Wroclaw University of Environmental and Life Sciences: Uniwersytet Przyrodniczy we Wroclawiu

Ewa Kochan

Medical University of Lodz: Uniwersytet Medyczny w Lodzi

Bartosz Pencakowski

Wroclaw Medical University: Uniwersytet Medyczny im Piastow Slaskich we Wroclawiu

Marta Stafiniak

Wroclaw Medical University: Uniwersytet Medyczny im Piastow Slaskich we Wroclawiu

Monika Bielecka

Wroclaw Medical University: Uniwersytet Medyczny im Piastow Slaskich we Wroclawiu

Bartosz J. Płachno

Jagiellonian University: Uniwersytet Jagiellonski w Krakowie

Maciej Strzemski

Medical University of Lublin: Uniwersytet Medyczny w Lublinie

Adam Matkowski

Wroclaw Medical University: Uniwersytet Medyczny im Piastow Slaskich we Wroclawiu

Sylwia Zielinska (✉ sylwia.zielinska@umw.edu.pl)

Wroclaw Medical University: Uniwersytet Medyczny im Piastow Slaskich we Wroclawiu <https://orcid.org/0000-0003-0090-5486>

Research Article

Keywords: *Agastache rugosa*, hairy roots, nutrient sprinkle bioreactor, temporary immersion system, phenolic compounds, rosmarinic acid

Posted Date: June 15th, 2023

DOI: <https://doi.org/10.21203/rs.3.rs-3035604/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Version of Record: A version of this preprint was published at Plant Cell, Tissue and Organ Culture (PCTOC) on December 19th, 2023. See the published version at <https://doi.org/10.1007/s11240-023-02626-z>.

Abstract

Agastache rugosa (Fisch. & C.A.Mey.) Kuntze known as Korean Mint is a medicinal and aromatic plant containing rosmarinic acid as a major bioactive polyphenol but its content in this herbal drug is variable. Plant *in vitro* culture is one of the approaches used to overcome the complexity of environmental factors influencing phytochemical profiles in medicinal plants.

In this study, hairy root cultures of *A. rugosa* were established through the infection of *Rhizobium rhizogenes*. Four hairy roots lines were selected on the basis of biomass production in liquid media. Transformation was confirmed by PCR using *rolC* specific primers. The hairy roots were successfully cultured in 300 mL conical flasks and scaled-up using three bioreactor types (nutrient sprinkle bioreactor – NSB and two modular temporary immersion systems – RITA[®] and Plantform[®]). The UPLC analysis of *A. rugosa* transformed roots methanol extracts showed the presence of 24 polyphenolic compounds with the predominance of rosmarinic acid (RA), which level ranged between 3.82 and 9.16 mg/g of dry weight. Nineteen compounds were identified in hairy roots growing in NSB system, 9 of them were identified in roots cultured in RITA[®] or Plantform[®] and 7 compounds were identified in roots from Erlenmeyer flasks. The *A. rhizogenes* infection (strain A4) was found to be an effective method of hairy root culture establishment of *A. rugosa*.

1. Introduction

Plant natural compounds sourcing become an issue since the need of using them has grown. Paclitaxel, shikonin, carthinin, betacyanins and many others are the most recognizable examples of specialized plant metabolites, the acquisition of which in larger quantities required many tedious biotechnological treatments.

Plant *in vitro* culture based approaches have been used since almost a century to study plant specialized metabolism and to obtain selected phytochemicals via manipulating their composition and content for pharmaceutical, cosmetic and food industry purposes. Among the commercially relevant compounds known to be produced industrially via plant biotechnology are coptisine, paclitaxel, shikonin, and several other with new examples being added on a regular basis (reviewed recently by Espinosa-Leal et al. 2018).

The difficulty in sustained and predictable recovery of these substances from naturally grown plant sources results i.a. from the complexity of their biosynthesis pathways and dynamic modifications which take place *in planta* in a response to environmental factors that often remain undefined. In this respect, the non-recombinant biotechnology offers an advantage of highly controlled experimental conditions, accelerated growth cycle, and independence on natural resources, being, however, more expensive and labor intensive than conventional cultivation. Therefore, the successful implementation of biotechnology for production of phytochemicals requires tedious optimization and upscaling experiments to fine tune the culture conditions to simultaneously achieve a possibly high biomass growth rate and superior metabolic profile. Hairy roots induced by infection of plant tissues with *Rhizobium* (*Agrobacterium*) *rhizogenes* are a well-established system for such purposes in hundreds of medicinal plant species (Malarz et al. 2023; Flores-Félix et al. 2020). The hairy root cultures are also exploited as industrial source of extract for pharmaceutical and cosmetic industries, for example ginseng, echinacea, coleus (Arpita 2021). Among the polyphenols which are considered superior antioxidants and antiinflammatory agents, rosmarinic acid (RA) is one of the common and intensively studied compounds which has been also obtained in spectacular amounts in plant cell, tissue and organ cultures, reaching over 30% of the cell dry mass in *Coleus blumei* (Qian et al. 2009; Gertlowski and Petersen 1993).

RA is characteristic for the different species of the subfamily Nepetoideae of the Lamiaceae and of numerous Boraginaceae but also occurs in other taxonomic groups (Petersen and Simmonds 2003) It is accumulated in various

proportions in different plant species, among others *Agastache rugosa* (Fisch. & C.A.Mey.) Kuntze known as Korean Mint. *A. rugosa* is a source of several phenylpropanoid derivatives, such as estragole, an essential oil constituent and rosmarinic acid (RA) among the nonvolatile compounds (Zielińska et al. 2017; Zielińska and Matkowski 2014). It is a popular medicinal and aromatic crop in East Asia, being one of the important herbs of the Traditional Korean and Chinese phytotherapy systems. The content of rosmarinic acid in *A. rugosa* has been studied in the past both in plants grown on experimental plots and *in vitro* culture. The production of this compound in plants depends on many factors, and the content may vary within large limits. Hence, *in vitro* plant organ cultures can be considered as a good source due to their stability of the production of specialized metabolites (Espinosa-Leal et al. 2018). In *A. rugosa*, an induction and culture of hairy roots was previously reported (Yeo et al. 2023; Lee et al. 2008). However, in Lee et al. (2008) the procedure for confirming the genetic transformation by molecular methods was not reported. On the other hand, in Yeo et al. (2023), the plant material was examined for the rosmarinic acid content only. However, many other phenolic compounds had been identified in this species (Zielińska et al. 2016; Zielińska and Matkowski 2014).

Therefore, the aim of the study was to establish hairy root cultures of *A. rugosa* as an efficient platform for *in vitro* production of phenolic compounds, particularly RA and evaluate, for the first time, the potential for scaling-up in various types of bioreactors. The effects of three bioreactors: two temporary immersion bioreactors (RITA® and Plantform®) and a nutrient sprinkle bioreactor (NSB) on the hairy roots biomass increases was studied followed by a detailed phytochemical characterization of the obtained plant material delivered from different bioreactor systems using UPLC-qTOF-MS/MS.

2. Materials and methods

2.1. Plant material

Two-weeks-old *A. rugosa* seedlings were used for the experiments. Seeds were collected from two-years-old plants cultivated in the Botanical Garden of Medicinal Plants at the Medical University of Wrocław, and the plant voucher specimen was deposited at reference number: L01/A.r.2/17. Nodes and leaves of aseptically grown seedlings were used as explants for the bacterial (*R. rhizogenes*) infection. Plant explants and bacteria were co-cultured on MS agar media. The roots obtained as a result of transformation were cut off and transferred to the same media supplemented with ampicillin (500 mg/l). After eliminating the bacteria, the roots were transferred into MS liquid media with all or reduced content of macro- and micronutrients and iron (for detailed description see the subsection 2.2).

2.2. Transformation of *A. rugosa* explants and hairy root cultures

The strain A4 of *R. rhizogenes* was used for the infection of *A. rugosa*. The bacterial strain is maintained in the collection of the Department of Pharmaceutical Biotechnology, Medical University of Łódź, Poland. The growth conditions for the bacterial strain was previously described by Piątczak et al. (2012).

Before the experiments, bacterial cells were transferred into the liquid culture medium. Nodes and leaf blades were wounded with a surgical steel needle submerged in bacterial suspension. Wounding with a sterile needle was treated as a negative control.

Wounded explants were incubated in the growth chamber, at 25°C in the dark on a MS agar medium (0.6% agar LAB-AGARTM (Biocorp), w/v) supplemented with sucrose (30 g/L, w/v, Pfeifer & Langen, Poznań, Poland).

The root development was monitored every seven days. After 42 days of observation, the frequency of transformation (% of hairy-root developing explants) in relation to the total number of infected explants, was determined. Then, the explants on which the roots had developed were transferred to the solid MS medium supplemented with ampicillin (500

mg/L, Sigma-Aldrich, US) to eliminate bacteria. Thereafter, the roots were cut from the explants and placed individually in 300 mL conical glass culture flasks (Schott, Germany) containing 100 mL of MS liquid medium with all or half the content of macro-, microelements and iron ($\frac{1}{2}$ MS). These flasks were placed on a laboratory shaker in the growth chamber, at the temperature 25°C in the dark. The roots were transferred every 28 days.

Overall, three root lines on MS medium, and four root lines on $\frac{1}{2}$ MS were selected on the basis of biomass production for further investigations.

2.3. Confirmation of *Ri*-plasmid integration

To confirm the plant transformation, DNA isolation from the collected material was performed, followed by the DNA polymerase chain reaction (PCR). To carry out this reaction, specific primers were used, the sequence of which was designed based on the DNA sequence of the *ro/C* gene located in the T-DNA region of the *Ri R. rhizogenes* plasmid and *virG* primers, a gene located outside the T-DNA region (Furner et al. 1986).

Three clones (K17, K19, K21) from MS medium and four (K12, K14, K16, K21) from $\frac{1}{2}$ MS medium, as well as one sample of untransformed *A. rugosa* roots derived from an *in vitro* cultures (negative control) were used in the transformation confirmation procedure. The material was triturated in liquid nitrogen in a mortar and accurately weighed approx. 100 mg into 1.5 ml test tubes. 396 μ l of extraction buffer and 4 μ l of β -mercaptoethanol were added. It was vortexed and incubated at 60°C for 30 minutes. Subsequently, 60 μ L of a 24:1 mixture of chloroform and isoamyl alcohol was added to the test tube and vortexed for approximately 10 seconds. It was centrifuged at 10,000 rpm for 3 minutes, then the supernatant was transferred to a fresh tube. The mixture of chloroform and isoamyl alcohol was added again, mixed, centrifuged and the supernatant added to the previous portion. 280 μ L of cold isopropanol was added to the supernatant, the contents were mixed gently, centrifuged at 10,000 rpm for 3 minutes. The liquid phase was removed from the tube with a pipette. 70% ethanol (500 μ L) was added to the DNA pellet of, and mixed, centrifuged at 10,000 rpm for 3 minutes, and then the ethanol was removed. The action was repeated and the DNA was allowed to dry. The dried pellets were dissolved in 50 μ l of a 10 μ g/ml RNase water solution. The solution was incubated at 37°C for one hour and then stored at -20°C. The concentration of the isolated DNA was determined using a spectrophotometer.

Then, two Master Mix mixtures were prepared consisting of: 111.6 μ l of nuclease-free water, 36 μ l of Q5 reaction buffer (5x concentrated), 3.6 μ l of free dNTPs nucleotides (10 mM), 1.8 μ l of Q5 DNA polymerase (2000 U/ml). To one of them, 9 μ l of *ro/CF* and *ro/CR* primers were added, and to the other - analogously *virGF* and *virGR*. 1 μ l of template DNA of each sample was collected into two test tubes. Then 19 μ l of Master Mix containing the *ro/C* and *virG* primers, respectively, were added to each. The contents of the tubes were mixed, centrifuged and inserted into a thermocycler T100™ Thermal Cycler (BIO-RAD Laboratories Inc) for PCR.

Nucleic acid electrophoresis was performed on an agarose gel to analyze the PCR product. A 50 ml gel containing 1.5% agarose was prepared with TBE buffer and heated in the flask to reflux temperature. Then, 5 μ l of ethidium bromide - a fluorescent dye, visible under UV light, was added to it, which allowed for the subsequent interpretation of the results. The gel was poured into a previously prepared and leveled mold and allowed to cool for 30 minutes.

Prepared samples: 10 μ l of PCR product were collected from each of 8 samples into new tubes. Then 2 μ l of loading buffer was added, mixed and centrifuged. The gel comb was removed from the set gel, then the gel and the tray were placed in the electrophoresis apparatus Sub-Cell GT (BIO-RAD Laboratories Inc). TBE buffer was poured into the apparatus so as to cover the gel. 10 μ l of the mixture of PCR product and loading buffer were taken and then applied to the wells. The electrodes were connected and the separation was started on 110V for 30 minutes.

2.4. Hairy root biomass production in culture flasks

Measurements of the biomass growth were made for the four clones selected on the basis on preliminary observations: K17, K19 from MS medium and K12, K14 from ½ MS medium without growth regulators. For better characteristics of hairy roots growth, apart from biomass, growth rate (μ) as well as time required for double the biomass (doubling time) were calculated for three intensively growing hairy root lines (K12, K17 and K19).

The root growth rate was calculated according to the formula:

$$\mu = \frac{2,303(lgx - lgx_o)}{\Delta t}$$

μ – culture growth rate [day⁻¹]

x – mean fresh weight [g/flask] in time t

x_o – mean fresh weight [g/flask] in time t_o

t_o – day of the beginning of the exponential growth phase of the culture

t – the day when the exponential growth phase of the culture ends

$\Delta t = t - t_o$

Doubling time (dt) was calculated using the formula:

$$dt = \frac{2.303 \cdot lg2}{\mu}$$

μ - culture growth rate [day⁻¹]

On the basis of the growth, morphology as well as rosmarinic acid content, studied in our preliminary experiments (unpublished), for further experiments K12 clone was chosen.

2.5. Hairy root biomass production in different bioreactor systems

To increase the scale of culture, hairy roots of *A. rugosa* were cultured in three types of bioreactors: nutrient sprinkle bioreactor (NSB), and two commercially available temporary immersion systems: Plantform® (PlantForm AB, Sweden) and RITA® (Récipient à Immersion Temporaire Automatique, VITROPIC, France). For comparison, the hairy roots were also cultured in conical flasks (300 mL of volume, 80 mL of medium) in the same medium as bioreactor cultures for the same period of time (17 days).

The NSB system (volume 5.0 L, working volume 3.0 L) has been described in details by Piątczak et al. (2014). The bioreactor contained 1000 mL of liquid medium. The medium was supplied to the spraying nozzle by a CH-8604 peristaltic pump (Chemap AG, Switzerland). The operating time of the pump was 40 s with 3.0 min breaks (60 mL of medium/1 cycle of pump working). The Plantform® bioreactor (size 180 x 150 x 150 mm) containing 500 mL of liquid medium was serially connected with an Optima pump (Hagen, Canada) with an inlet capacity of 0.6 m³/h. The 800 mL round RITA® bioreactor contained 250 mL of the liquid medium, which delivery was controlled by a DT4.4 pressure pump (Becker, Germany) with a capacity of 4.2 m³/h and pressure of 1000 mbar. The operating time of the pumps in the both temporary bioreactors was 10 min every 80 min.

The hairy root cultures in each bioreactor were cultured in liquid MS medium with reduced to half the content of macro- and micronutrients ($1/2$ MS) supplemented with 3% of sucrose without growth regulators. In the study, hairy roots from 4 subsequent subcultures were used. The inoculum for all the cultures derived from 14th day of culture. The mean fresh weight (FW) of inoculum in NSB system was 12.67 ± 3.08 g/L (1.18 ± 0.1 g/L DW), in RITA® was 12.46 ± 0.5 g/L FW (1.13 ± 0.11 g/L DW), in Plantform® was 7.53 ± 0.28 g/L FW (0.74 ± 0.04 g/L DW) and in 300 mL culture flasks was 15.85 ± 0.47 g/L FW (1.50 ± 0.05 g/L DW).

All the cultures were cultured at $26 \pm 2^\circ\text{C}$ in the darkness for 17 days. At the end of the culture period, a biomass of the culture, means of fresh weight (FW) and dry weight (DW) of culture (g/L) and a growth index (GI), calculated as final fresh or dry weights - initial fresh or dry weight/initial fresh or dry weight, were recorded.

2.6. Phytochemical analysis

2.6.1. Extracts preparation

100 mg lyophilized and grounded plant material was extracted twice using 2.5 ml solution of 0.1% formic acid and methanol (v/v) in a 4:1 ratio every time. The extraction procedure was performed according to previously published methodology (Zielińska et al. 2019).

2.6.2. Chromatographic analysis

The separation of polyphenolic compounds was carried out using a UPLC BEH C18 column ($1.7 \mu\text{m}$, 2.1×100 mm, Waters Corporation, Milford, MA). The mobile phase consisted of solvent A (0.1% formic acid in LC–MS grade water, v/v, Merck, Germany) and solvent B (0.1% formic acid in LC–MS grade acetonitrile, Merck Germany). The chromatographic procedure as well as quality and quantity analyses were performed according to previously published methodology (Zielińska et al. 2019). Compounds were optimized to their estimated molecular masses in the negative mode, before and after fragmentation. The authentic standards were used for the comparison of retentions times and spectra of analyzed compounds. The calibration curve for acacetin was used to quantify tilianin. Caffeic, chlorogenic, rosmarinic, *p*-coumaric and feruloylquinic acids as well as acacetin, apigenin, and phloridzin were quantified with their own standards. The quantities were expressed in mg of a compound per g of plant material dry weight (mg/g DW).

2.7. Statistical analysis

Calculations of arithmetic means and standard errors (SE) were made using the EXCEL 2010. Each chromatographic analysis was performed six times. Results were reported as mean values $\pm$ SE. Experiments in bioreactor systems were repeated four times. Statistical significance was calculated with STATISTICA 13.3 version (TIBCO) using Tukey HSD or Kruskal-Wallis tests at $p \leq 0.05$.

2.8. Scanning electron microscopy

For SEM, the hairy roots fragments were prepared according to previously published methodology (Piątczak et al. 2020) using a Hitachi S-4700 scanning electron microscope (Hitachi, Tokyo, Japan), housed in the Institute of Geological Sciences, Jagiellonian University in Kraków).

3. Results

3.1. Induction of hairy roots

Hairy root cultures of *A. rugosa* were established through the infection with *R. rhizogenes* strain (A4). The ability to form roots at the site of infection depended on the type of explant. Leaf blades, for which the transformation frequency

was 55%, turned out to be the best explants. In case of nodes, none of the explants responded. Roots were formed through direct rhizogenesis. They were thin, delicate, light-beige-colored (Fig. 1).

An intensive root growth and elongation, as well as numerous axillary branched roots development were observed on both liquid media. The most intensively growing four clones K17, K19 (on MS medium), and K12, K14 (on ½ MS medium) were selected for further investigation on rosmarinic acid production. For comparison, untransformed roots were also cultured on liquid MS and ½ MS media, in which no further growth was observed. The MS medium with half the content of macro-, microelements and iron turned out to be more effective for the growth of root cultures.

3.2. Confirmation of transformation

Transformation status of the roots was confirmed by PCR using *rolC* and *virG* specific primers.

PCR reaction was performed to show the presence of a bacterial *A. rhizogenes* plasmid fragment (*rolC* gene) in the genome of *A. rugosa* hairy roots. DNA was isolated from several clones: K17, K19, K21 (from MS medium) and K12, K14, K16, K21 (from ½ MS medium) and from untransformed *A. rugosa* roots derived from an *in vitro* cultures (negative control). Primers sequence was designed based on the DNA sequence of the *rolC* gene, located in the T-DNA region of the Ri *R. rhizogenes* plasmid. Primers for *virG*, a gene located in the plasmid outside the T-DNA region not involved in the genetic transformation, were also used (positive control). An electroforetogram showing the results of the PCR reaction is shown in Fig. 2.

On the basis of the electrophoretograms, the genetic transformation of K17, K19, K12 and K14 clones was confirmed. In samples containing isolated DNA from K21 clones (both MS and ½ MS medium) and K16, the *virG* gene was additionally identified, the presence of which was most likely due to contamination of the plant material with *R. rhizogenes* bacteria, therefore further research was carried out only on the aforementioned four clones. In the control sample containing DNA isolated from untransformed *A. rugosa* roots, none of the identified genes were present.

3.3. Hairy root biomass production in Erlenmeyer flasks

For the three intensively growing clones with transformation confirmation (K19 and K17 on MS medium and K12 on ½ MS medium), an average growth rate (μ) and doubling time (dt) were calculated. Regardless of the type of clone and culture medium, the growth cycle lasted 43–50 days. The most intensive growth of cultures were observed between 30th and 40th day of culture. The K19 clone cultured on MS medium achieved a slightly higher biomass than K12 clone, but without the significant difference (at $p \leq 0.05$) (Table 1) but both the average growth rate and the time required to double the biomass (dt) was shorter for K12 clone ($\mu = 0.156$; $dt = 4.463$, respectively) than for K19 ($\mu = 0.141$; $dt = 4.906$, respectively).

Table 1
Fresh (FW) weights (g/flask) of *A. rugosa* hairy root lines after 30 days of growth in 300 mL Erlenmeyer flasks in liquid MS and ½ MS media without growth regulators in darkness.

Hairy root clone	FW [g/flask] ± SE
K12	4.0 ± 0.115 ^a
K14	0.60 ± 0.163 ^b
K17	1.65 ± 0.113 ^c
K19	4.05 ± 0.646 ^a

Means designed with the same letter are not statistically different at $p \leq 0.05$ in Tukey's test.

3.4. Hairy root biomass production in three types of bioreactors

With an aim to test the possibility to scale up, the hairy roots (K12) were cultured for 17 days in three types of bioreactors. Two of them (RITA® and Plantform®) are commercially available temporary immersion systems, where the explants are immersed in the liquid medium only by the short, pre-programmed period of time (in our study 10 min of immersion with 80 min. breaks). In the NSB system, the explants are situated at the stainless net in the middle of bioreactor and they are sprinkled with the liquid medium with 3 min intervals. Liquid ½ MS medium without growth regulators was used in the all types of bioreactors. For comparison, the hairy roots were also cultured in Erlenmeyer flasks in the same medium. It was observed that hairy roots cultured in RITA® and NSB bioreactors after 17 days of culture have occupied all available space in each bioreactor. They were thin, brown in the middle and creamy-white at their distal edges. Additionally, roots growing in NSB and Plantform® bioreactors created thin, white, fluffy hair in upper part of culture (Fig. 3, 4). The highest biomass (FW and DW) was recorded for hairy roots growing in RITA® system (160 g/L of FW and 16 g/L of DW). It was almost 1.5 times higher than the biomass of roots growing in 300 mL glass flasks and NSB system, respectively. The lowest biomass and the slowest growth of the hairy roots were observed in Plantform® system (26 g/L of FW and 3 g/L of DW). The biomass increases were 6-times lower in terms of FW and 5.5-times lower in terms of DW than the biomass of hairy roots growing in the RITA and 3–4 times lower than the biomass of roots growing in the flasks and NSB system (Fig. 5A).

After a consideration of the inoculum size, the highest growth index – GI was observed in the RITA® system (GI = 11.8 for FW and 13.5 for DW) followed by the NSB system (GI = 7.5 for FW and 7.3 for DW) and 300 mL flasks (GI = 5.5 for FW and 5.2 for DW). GI in Plantform® bioreactor was 1.7–2.2-times, 2.4–2.5-times and 4.5–4.7 times lower than GI values for DW and FW for hairy roots cultured in flasks, NSB and RITA®, respectively (Fig. 5B).

3.5. Phytochemical analysis

The selected hairy root lines were used for the extraction and UPLC analysis of phenolic compounds. Out of the four examined hairy root lines, only one (K12) was highly productive. The line was growing in liquid ½ MS medium and when maintained in the 300 mL flasks, the rosmarinic acid (RA) content was 20 to even 40 times higher than in the other lines growing in MS media (data not shown). Therefore, the K12 line was chosen for further experiments with bioreactors. The selected line was continuously cultured in the 300 mL flasks as well as in NSB and two temporary immersed systems (RITA® and Plantform®). The chromatographic analysis shown the presence of polyphenolic compounds and their derivatives (Table 2, Fig. 6).

The poorest phytochemical profile shown the hairy roots cultured in the flasks, consisting of only seven detected phenolic compounds. The major constituent was RA, but its amount was rather low (4.6 mg/g DW). The second in abundance phenolic compound found in hairy roots from culture flasks was a flavonoid – apigenin derivative (peak no 21). Its amount (0.6 mg/g DW) was the highest among all tested culture systems. The other phenolic compounds produced by the roots in the flasks were: tilianin, phloridzin, RA methyl ester, two caffeic acid derivatives, which were present in lower amounts (0.001–0.07 mg/g DW). Interestingly, one of the caffeic acid derivatives (peak no 11) was detected only in roots cultured in the flasks, but its amount was also low (0.029 mg/g DW).

Phytochemical profiles of phenolic compounds in RITA® and Plantform® were similar and consisting of nine compounds in low amounts. Only two compounds: RA and the apigenin derivative were present in higher levels (3.82–4.46 mg/g DW and 0.24–0.46 mg/g DW, respectively). Two compounds: RA hexoside (peak no 15) and RA isomer I (peak no 18) were present only in roots from RITA® and Plantform®, but not in flasks or NSB system.

The most effective for phenolic compounds accumulation in hairy roots of *A. rugosa* was NSB. After 17 days of culture in this type of bioreactor, 19 phenolic compounds were identified: 13 phenolic acids derivatives and 5 flavonoid derivatives, including acacetin and apigenin derivatives and one dihydrochalcone - phloridzin. Among these compounds, 14 were detected only in the NSB. There were mainly caffeic and caffeoylquinic acids derivatives, acacetin and apigenin derivatives. Similarly to other described culture systems, roots in the NSB produced RA as the major metabolite, but the amount (9.1 mg/g DW) was 2 times higher than in other culture systems (3.8–4.6 mg/g DW). Moreover, the total amounts of phenolic acid and flavonoid derivatives (10.193 and 1.941 mg/g DW, respectively) were also higher than in other tested culture systems (Table 2, Fig. 6).

Table 2

Identification and content of metabolites (mg/g DW) in *A. rugosa* hairy roots (K12 clone) after 17 days of growth in different culture systems in liquid ½ MS medium without growth regulators.

Peak	Compound	Rt (min)						
no			MS	MS/MS	NSB	FLASKS	RITA®	PLANTFORM®
Means ± SE (mg/g DW)								
1.	Caffeoylquinic acid (neochlorogenic acid)	2.53	353	179	0.006 ± 0.0007 ^a	-	-	-
2.	Caffeic acid hexoside I	2.85	341	179	0.009 ± 0.0009 ^a	-	-	-
3.	Caffeic acid hexoside II	3.25	341	179	0.017 ± 0.0007 ^a	-	-	-
4.	<i>p</i> -coumaric acid	3.63	325	163	0.002 ± 0.0002 ^a	-	-	-
5.	Dicaffeoyl-hexose isomer I	3.77	503	341, 179, 161, 135	0.010 ± 0.001 ^a	-	-	-
6.	Caffeic acid hexoside III	3.92	341	179	0.004 ± 0.0002 ^a	-	0.0203 ± 0.00264 ^b	0.0143 ± 0.00450 ^c
7.	Caffeic acid derivative I	4.22	445	341, 179, 161, 135	0.052 ± 0.007 ^a	-	-	-
8.	Dicaffeoyl hexose isomer II	4.48	503	341, 179, 161, 135	0.025 ± 0.002 ^a	-	-	-
9.	Tilianin	4.57	445	299, 136	0.096 ± 0.007 ^a	0.00547 ± 0.00123 ^b	0.00633 ± 0.00101 ^c	0.00572 ± 0.00078 ^{b, c}
10.	Caffeoylquinic acid (cryptochlorogenic acid)	5.13	353	179, 161	0.736 ± 0.030 ^a	-	-	-
11.	Caffeic acid derivative II	5.59	486	341, 281, 179, 161	-	0.0288 ± 0.00232 ^a	-	-
12.	Caffeic acid derivative III	5.77	445	341, 179, 161, 135	-	0.0011 ± 0.00000 ^a	0.00056 ± 0.000000 ^b	0.00049 ± 0.000150 ^c
13.	Apigenin glucoside	6.30	431	269	0.591 ± 0.028 ^a	-	-	-

Peak	Compound	Rt (min)						
no			MS	MS/MS	NSB	FLASKS	RITA®	PLANTFORM®
14.	Feruloyl-quinic acid	6.45	385	193, 177	0.006 ± 0.0002 ^a	-	-	-
15	Rosmarinic acid hexoside	6.63	521	359, 197, 179	-	-	0.0167 ± 0.00190 ^a	0.0167 ± 0.00188 ^a
16.	Acacetin acylglycosyl isomer I	6.91	473	311, 269, 161	0.034 ± 0.006 ^a	-	-	-
17.	Phloridzin	7.12	435	272, 179	0.023 ± 0.0006 ^a	0.0096 ± 0.00087 ^b	0.0257 ± 0.00284 ^a	0.0228 ± 0.00118 ^a
18.	Rosmarinic acid isomer I	7.94	357	197, 179	-	-	0.0131 ± 0.00323 ^a	0.0132 ± 0.00188 ^a
19.	Rosmarinic acid isomer II	8.47	357	197, 179	0.164 ± 0.037 ^a	-	-	-
20.	Rosmarinic acid	8.70	359	197, 179, 161	9.159 ± 0.252 ^a	4.648 ± .00715 ^b	4.485 ± 0.2520 ^b	3.817 ± 0.3115 ^c
21.	Apigenin derivative	9.28	487	269	-	0.6248 ± 0.0193 ^a	0.4549 ± 0.0645 ^b	0.2520 ± 0.0420 ^c
22.	Rosmarinic acid methyl ester	9.53	373	358, 196	0.003 ± 0.0002 ^a	0.0709 ± 0.00732 ^b	0.0779 ± 0.01022 ^b	0.0594 ± 0.0109 ^c
23.	Acacetin acylglycosyl isomer II	10.08	487	269, 182	1.057 ± 0.028 ^a	-	-	-
24.	Apigenin glucoside	10.71	431	269	0.163 ± 0.013 ^a	-	-	-

NSB – nutrient sprinkle bioreactor; means with the same letter in the upper superscript within lines are not statistically different at $p \leq 0.05$ in Kruskal-Wallis non-parametric test.

4. Discussion

In several previous studies large differences in the RA content were observed in *A. rugosa* plants from different locations and different experimental treatments (Lee et al. 2008; Tuan et al. 2012; Desta et al. 2015; Zielińska et al. 2016; Zielińska et al. 2019; Lam et al. 2020 a, b, c). In the plants cultivated in Korea, the RA content did not exceed 50 µg/g DW in all plant organs examined individually (Tuan et al. 2012; Desta et al. 2015). In turn, plants of the species cultivated in Poland produced much larger amounts of RA in leaves and inflorescences (0.03–10.76, 0.06–5.27 mg/g DW, respectively) (Zielińska et al. 2016). Moreover, samples derived from *in vitro* shoot cultures were found to be a rich source of RA (nearly 25 mg/g DW dependent on the treatment) (Zielińska et al. 2019). The Korean group (Lee et al. 2008) reported 116.3 mg/g DW of RA in hairy root cultures obtained for the first time. Due to the huge differences in the content of RA in the plant material obtained by various methods, we made an attempt to culture *A. rugosa* hairy roots with three different *in vitro* systems and verify it in terms of the content of the metabolite and other phenolic

compounds. Rosmarinic acid (RA) can be biosynthesized in undifferentiated cells unlike many other specialized metabolites that are produced and stored in developed organs in highly specialized secretory structures. RA was efficiently produced by various *in vitro* cultures of many plant species, such as *Coleus blumei* Benth. (Bauer et al. 2004; Petersen 1991), *Anchusa officinalis* L. (Su et al. 1993), *A. rugosa* (Xu et al. 2008), *Lavandula vera* DC. (Georgiev et al. 2006, Georgiev et al. 2009), *Salvia miltiorrhiza* (Dong et al. 2010), *Salvia officinalis* (Santos-Gomes et al. 2003). *Anchusa officinalis* and *Coleus blumei* suspension cell cultures were used in the experiments to discover the enzymes involved in RA biosynthesis. At the early stage of the pathway, which is the conversion of phenylalanine to cinnamic acid, activity is undertaken by enzymes fundamental to the synthesis of all phenylpropanoids (e.g. flavonoids or lignins) (Petersen and Simmonds 2003). The correlation between PAL activity and RA accumulation has been known for a long time (Razzaque and Ellis 1977). In isolated *Coleus blumei* protoplasts grown in suspension, a higher content of this metabolite was found in the vacuoles than in the cytoplasm (Häusler et al. 1993).

In our study, for hairy roots induction in *A. rugosa*, the *Rhizobium rhizogenes* strain A4 was chosen based on earlier experiments using several other plant species, such as *Rehmannia elata* (Piątczak et al. 2019), *Salvia bulleyana* (Wojciechowska et al. 2020) or *Salvia viridis* (Grzegorzczak-Karolak et al. 2018), where the bacterial strain was successful in formation of hairy roots. Similarly to the results of Chaudhuri et al. (2005), Wojciechowska et al. (2019), Dehdashti et al. (2017), and Verma et al. (2007) we observed in the present study that the type of explant infected by the bacteria was crucial for hairy roots formation frequency. In all the cited above reports leaves were more susceptible to infection than stems or nodes. However, in many other plant species, the nodes/stems were more effective in hairy root induction than leaves (Piątczak et al. 2019; Grzegorzczak-Karolak et al. 2018). The differences may correspond to the physiological state of plant tissues, especially with the fact that leaves as younger tissues than shoots, may be more susceptible for the *Rhizobium* infection. Larger number of meristematic cells in leaves compared to stems, with higher proliferative potential may also be one of the explanation. Hairy root induction can be also influenced by the individual susceptibility of plant species (De Cleene and De Ley 1981).

The hairy root lines, described in the study, with confirmed transformation status, differed in the level of fresh biomass increasing. The fastest growing lines were K12 and K19, which biomass accumulation was 6.67 times higher than in the slowest growing line (K14). The differences in the growth within different hairy root lines of the same plant species have been earlier described for example in *Rehmannia glutinosa* (Piątczak et al. 2012) and *Salvia viridis* (Grzegorzczak-Karolak et al. 2018). It could be related to differences in the presence or expression level of the *rol* genes inserted from the *Rhizobium Ri* plasmid into the plant genome. It is known that *rol* genes could affect the growth of roots by activating transcription genes (Ono and Tian 2011).

Consequently, we attempted to scale-up the *A. rugosa* hairy root culture, for the first time, in two temporary immersion bioreactors (RITA® oraz Plantform®) and in the nutrient sprinkle bioreactor (NSB) and to examine the influence of the type of bioreactor on the biomass (both fresh and dry weight) accumulation as well as on phytochemical profile of hairy roots. In our preliminary studies (unpublished data) we have observed the best growth and morphology of roots on the 17th day of culture, therefore in this study, all results were recorded after 17 days of hairy roots growth.

The obtained results indicated that bioreactor systems, especially RITA® and NSB, efficiently promote growth of *A. rugosa* hairy roots. Growth index was 1.5-5 times higher than for other culture systems (300 mL flasks and Plantform®). NSB or RITA® systems have been earlier reported to be effective for growth of other plants, for example *Schisandra chinensis* microshoots (Szopa et al. 2017, 2019) and *Dracocephalum forrestii* transformed shoots (Weremczuk-Jeżyna et al. 2020). The high biomass accumulation could be caused by better absorption of nutrients and plant growth regulators from the liquid medium in such types of bioreactors (Preil 2005, Mehrotra et al. 2007). Pavlov and Bley (2006) have also observed that temporary immersion systems, like RITA® could reduce the risk of morphology abnormalities occurring as a result of shear forces (Szopa et al. 2017). Similarly to results obtained in the

present study, other authors also reported advantageous biomass growth in RITA® bioreactors in comparison with nutrient sprinkle bioreactor, for example in shoots of *Rhododendron tomentosum* Harmaja (Jasionek et al. 2017) or shoots of *Schisandra chinensis* (Szopa et al. 2017, 2019).

On the other hand, lower growth parameters were observed for hairy roots of *A. rugosa* growing in Plantform® system. After 17 days of culture, growth indexes for roots growing in Plantform® were 3 and 5 times lower than the parameters for roots cultured in NSB and RITA®, respectively. It may be a result of lower oxygen availability while being flooded in this type of bioreactor. Therefore, this parameter turned out to be a main limiting factor for large scale cultivation of *A. rugosa* hairy roots. It may be assumed that the culture conditions in Plantform® bioreactor are not as advantageous for hairy roots growth and metabolite production as in other bioreactors used. Plantform® bioreactor has been designed and used earlier only for shoot cultures, especially for crops like *Olea europaea* (Benelli i De Carlo 2018) or *Capparis spinosa* (Gianguzzi i wsp. 2019), not for hairy root cultures.

Phytochemical analysis

In the present study, nutrient sprinkle bioreactor (NSB) and two temporary immersion systems (RITA® and Plantform®) have been used for metabolite production in hairy roots of *A. rugosa*. It was found that NSB was the most effective system for polyphenols accumulation during 17 days of culture in liquid ½ MS medium without growth regulators. Hairy roots growing in the NSB accumulated more compounds than roots growing in two other bioreactors. It was observed that among 19 phenolic compounds detected in extracts from roots cultured in NSB, 13 compounds (10 phenolic acid derivatives and 3 flavonoids derivatives) were found only in this type, whereas in hairy roots growing in RITA®, only 9 compounds were detected. However, the amounts of these metabolites was rather low and did not exceed 0.4 mg/g DW. Regardless of the bioreactor system used, the predominant metabolite was RA, which level varied from 3.8 mg/g DW in roots growing in Plantform® to 9.16 mg/g DW in the NSB. The highest amount (9.16 mg/g DW) of RA was similar to the level of the metabolite present in leaves of two years old plants growing in the field (Zielińska et al. 2016). Nonetheless, this level of RA in hairy roots was achieved in significantly shorter time – 17 days. Earlier, other biotechnological treatments, such as precursor feeding, plant growth regulators supplementation or different illumination regime were found to be also effective for promoting RA accumulation (over 20 mg/g DW) (Zielińska et al. 2019), but taking into account the dry biomass (8.71 g), which is possible to achieve during one culture cycle in NSB, it is possible to obtain almost 80 mg of RA during 17 days of culture. Additionally, it is also possible to obtain other metabolites in high amounts, for example acacetin acylglycosyl isomer II (9.2 mg) and cryptochlorogenic acid (6 mg) during single cycle (17 days). These two compounds were produced only in NSB. Moreover, the phytochemical profile of the hairy roots in NSB was richer than in other cultures. For example, in shoot cultures only 7 metabolites were identified (6 phenolic acids derivatives and one flavonoid) (Zielińska et al. 2019). Therefore, taking into consideration all these results, the NSB bioreactor turned out to be effective and feasible system for polyphenolic compounds production in hairy roots of *A. rugosa*. It is possible to make an effort to further increase the metabolite production in the hairy roots by elicitation or precursor feeding. Earlier, Lee et al. (2008) reported 116.3 mg/g DW of RA in hairy root cultures of this species, but the transformation of the cultures has not been confirmed by molecular methods. NSB system has been also suitable for the growth and polyphenolic compounds production in *Dracocephalum forrestii* transformed and untransformed shoots (Weremczuk-Jeżyna et al. 2019, 2020), RA production in shoots and hairy roots of *Salvia officinalis* (Grzegorzczuk 2010), polyphenolic compounds production in shoots of *Scutellaria alpina* (Grzegorzczuk-Karolak et al. 2017), phenolic and flavonoid production in shoots of *Rehmannia glutinosa* (Piątczak et al. 2014), lignin production in *Schisandra chinensis* (Szopa et al. 2017), secoiridoid production in shoots of *Centaureum erythraea* (Piątczak et al. 2005). It is suggested that the high metabolite production in this type of bioreactor is a result of the optimal medium circulation as well as nutrient and oxygen supply and reduced shear forces which often destroy delicate structure of hairy roots (Ziv 2005).

5. Conclusions

The infection of *Rhizobium rhizogenes* strain A4 was found to be an effective method of *A. rugosa* hairy root culture establishment. The scale-up of *A. rugosa* hairy root culture allowed to obtain a significant amounts of plant material rich in bioactive metabolites. The most effective system for the growth of biomass was RITA, where biomass of roots increased almost 15 times after 17 days of culture, but for phenolic compounds production, the most effective was NSB system. Further research is necessary to elucidate exogenous factors influencing the phytochemical composition and optimize the biomass growth with possibly highest accumulation of desired constituents.

Declarations

Conflict of interest: Authors declare none conflict of interest

Authors Contribution Statement

All authors contributed to the study conception and design. Material preparation, data collection and analysis were performed by [Weronika Kozłowska], [Sylwia Zielińska], [Ewelina Piątczak], [Ewa Kochan], [Joanna Kolniak-Ostek], [Marta Stafiniak], [Bartosz Pencakowski], [Maciej Strzemski], [Monika Bielecka] and [Bartosz Jan Płachno]. The first draft of the manuscript was written by [Sylwia Zielińska], [Ewelina Piątczak], [Weronika Kozłowska], and [Adam Matkowski] and all authors commented on previous versions of the manuscript. All authors read and approved the final manuscript.

Acknowledgment

This research was financially supported by Wroclaw Medical University grant no. SUBK.D030.23.006 to Weronika Kozłowska.

The Authors would like to express sincere thanks to Mrs J. Bursy, M. Barwińska and K. Radecka for their help in methodology and photographic documentation.

References

1. Arpita R (2021) Hairy Root Culture an Alternative for Bioactive Compound Production from Medicinal Plants. Curr. Pharm. Biotechnol. 22:136-149. <https://doi.org/10.2174/1389201021666201229110625>
2. Desta KT, Kim GS, Kim YH, Lee WS, Lee SJ, Jin JS, Abd El-Aty AM, Shin HC, Shim JH, Shin SC (2015) The polyphenolic profiles and antioxidant effects of *Agastache rugosa* Kuntze (Banga) flower, leaf, stem and root. Biomed. Chromatogr. 30(2):225-231. <https://doi.org/10.1002/bmc.3539>
3. Do TM, Choi M, Kim JK, Kim YJ, Park C, Park CH, Park NI, Kim C, Sathasivam R, Park SU (2023) Impact of light and dark treatment on phenylpropanoid pathway genes, primary and secondary metabolites in *Agastache rugosa* transgenic hairy root cultures by overexpressing *Arabidopsis* transcription factor AtMYB12. Life. 13:1042. <https://doi.org/10.3390/life13041042>
4. Espinosa-Leal CA, Puente-Garza CA, García-Lara S (2018) In vitro plant tissue culture: means for production of biological active compounds. Planta. 248:1-18. <https://doi.org/10.1007/s00425-018-2910-1>
5. Flores-Félix JD, Menéndez E, Peix A, García-Fraile P, Velázquez E (2020) History and current taxonomic status of genus *Agrobacterium*. Syst Appl Microbiol. 43:126046. <https://doi.org/10.1016/j.syapm.2019.126046>
6. Furner IJ, Huffman GA, Amasino RM, Garfinkel DJ, Gordon MP, Nester EW (1986) An *Agrobacterium* transformation in the evolution of the genus *Nicotiana*. Nature, 319:422-427. <https://doi.org/10.1038/319422a0>

7. Gertlowski C, Petersen M (1993) Influence of the carbon source on growth and rosmarinic acid production in suspension cultures of *Coleus blumei*. *Plant Cell Tissue Organ Cult.* 34:183-190.
<https://doi.org/10.1007/BF00036100>
8. Grzegorzczak I, Wysokińska H (2010) Antioxidant compounds in *Salvia officinalis* L. shoots and hairy roots cultures in the nutrient sprinkle bioreactor. *Acta Soc. Bot. Pol.* 1:7–10. <https://doi.org/10.5586/asbp.2010.001>
9. Grzegorzczak-Karolak I, Rytczak P, Bielecki S, Wysokińska H (2017) The influence of liquid systems for shoot multiplication, secondary metabolite production and plant regeneration of *Scutellaria alpina*. *Plant Cell Tissue Organ Cult.* 128:479–486. <https://doi.org/10.1007/s11240-016-1126-y>
10. Kim HK, Lee HK, Shin CG, Huh H (1999) HIV integrase inhibitory activity of *Agastache rugosa*. *Archiv Pharm Res* 22:520-523. <https://doi.org/10.1007/BF02979163>
11. Kim Y, Zhang D, Yang D (2015) Biosynthesis and biotechnological production of ginsenosides. *Biotechnol. Adv.* 33:717-735. <https://doi.org/10.1016/j.biotechadv.2015.03.001>
12. Lee SY, Xu H, Kim YK, Park SU (2008) Rosmarinic acid production in hairy root cultures of *Agastache rugosa* Kuntze. *World J Microbiol Biotechnol.* 24:969–972. <https://doi.org/10.1007/s11274-007-9560-y>
13. Oh SR, Jung KY, Lee HK (1996) *In vitro* anticomplementary activity of phenylpropanoids from *Agastache rugosa*. *Kor J Pharmacogn* 27:20-25
14. Malarz J, Yudina YV, Stojakowska A (2023) Hairy Root Cultures as a Source of Phenolic Antioxidants: Simple Phenolics, Phenolic Acids, Phenylethanoids, and Hydroxycinnamates. *Int. J. Mol. Sci.* 24:6920.
<https://doi.org/10.3390/ijms24086920>
15. Mehrotra S, Goel KM, Kukreja AK, Mishra BN (2007) Efficiency of liquid culture systems over conventional micropropagation. A progress towards commercialization. *Afr. J. Biotechnol.* 6:1484–1492.
16. Petersen M, Simmonds M (2003) Rosmarinic acid. *Phytochemistry.* 62:121-125. [https://doi.org/10.1016/S0031-9422\(02\)00513-7](https://doi.org/10.1016/S0031-9422(02)00513-7)
17. Piątczak E, Chmiel A, Wysokińska H (2005) Mist trickling bioreactor for *Centaurea erythraea* Rafn growth of shoots and production of secoiridoids. *Biotechnol Lett.* 27:721–724. <https://doi.org/10.1007/s10529-005-5189-9>
18. Piątczak E, Królicka A, Wielanek M, Wysokińska H (2012) Hairy root cultures of *Rehmannia glutinosa* and production of iridoid and phenylethanoid glycosides. *Acta Physiol. Plant.* 34:2215–2224.
<https://doi.org/10.1007/s11738-012-1022-y>
19. Piątczak E, Grzegorzczak-Karolak I, Wysokińska H (2014) Micropropagation of *Rehmannia glutinosa* Libosch.: Production of phenolics and flavonoids and evaluation of antioxidant activity. *Acta Physiol. Plant.* 36:1693–1702.
<https://doi.org/10.1007/s11738-014-1544-6>
20. Piątczak E, Jeleń A, Makowczyńska J, Zielińska S, Kuźma Ł, Balcerczak E (2019) Establishment of hairy root cultures of *Rehmannia elata* N.E. Brown ex Prain and production of iridoid and phenylethanoid glycosides. *Ind. Crops Prod.* 137:308–314. <https://doi.org/10.1016/j.indcrop.2019.05.022>
21. Piątczak E, Kuźma Ł, Kozłowska W, Lisiecki P, Szemraj M, Płachno BJ, Gonciarz W, Chmiela M, Matkowski A, Zielińska S (2020). Phenylethanoid and iridoid glycosides production in *Rehmannia elata* N.E. *Ind. Crops Prod.* 158: 113050
22. Preil W (2005) General introduction: A personal reflection on the use of liquid media for *in vitro* culture, in: Hvoslef-Eide, A.K., Preil, W. (Eds.), *Liquid Culture Systems for In Vitro Plant Propagation*. Springer: Dordrecht, The Netherlands, pp.1-18.
23. Qian J, Guiping L, Xiujun L, Xincan H, Hongmei L (2009) Influence of growth regulators and sucrose concentrations on growth and rosmarinic acid production in calli and suspension cultures of *Coleus blumei*. *Nat. Prod. Res.* 23:127-137. <https://doi.org/10.1080/14786410801890338>

24. Shi M, Liao P, Nile SH, Georgiev MI, Kai G (2021) Biotechnological Exploration of Transformed Root Culture for Value-Added Products. *Trends Biotechnol.* 39:137-149. <https://doi.org/10.1016/j.tibtech.2020.06.012>
25. Szopa A, Kokotkiewicz A, Luczkiewicz M, Ekiert H (2017) *Schisandra* lignans production regulated by different bioreactor type. *J. Biotechnol.* 247:11–17. <https://doi.org/10.1016/j.jbiotec.2017.02.007>
26. Tuan PA, Park WT, Xu H, Park NI, Park SU (2012) Accumulation of tilianin and rosmarinic acid and expression of phenylpropanoid biosynthetic genes in *Agastache rugosa*. *J. Agric. Food Chem.* 60:5945–5951. <https://doi.org/10.1021/jf300833m>
27. Wawrosch C, Zotchev SB (2021) Production of bioactive plant secondary metabolites through in vitro technologies-status and outlook. *Appl Microbiol Biotechnol.* 105:6649-6668. <https://doi.org/10.1007/s00253-021-11539-w>
28. Weremczuk-Jeżyna I, Kochan E, Szymczyk P, Lisiecki P, Kuźma Ł, Grzegorzczak-Karolak I (2019) The antioxidant and antimicrobial properties of phenol-rich extracts of *Dracocephalum forrestii* W. W. Smith shoot cultures grown in the nutrient sprinkle bioreactor. *Phytochem. Lett.* 30:254–260. <https://doi.org/10.1016/j.phytol.2019.01.032>
29. Weremczuk-Jeżyna I, Lisiecki P, Gonciarz W, Kuźma Ł, Szemraj M, Chmiela M, Grzegorzczak-Karolak I (2020) Transformed shoots of *Dracocephalum forrestii* W.W. Smith from different bioreactor systems as a rich source of natural phenolic compounds. *Molecules.* 25:4533. <https://doi.org/10.3390/molecules25194533>
30. Yeo HJ, Kwon MJ, Han SY, Jeong JC, Kim CY, Park SU, Park CH (2023). Effects of carbohydrates on rosmarinic acid production and in vitro antimicrobial activities in hairy root cultures of *Agastache rugosa*. *Plants.* 12(4):797. <https://doi.org/10.3390/plants12040797>
31. Zielińska S, Matkowski A (2014) Phytochemistry and bioactivity of aromatic and medicinal plants from the genus *Agastache* (Lamiaceae). *Phytochem. Rev.* 13:391–416. <https://doi.org/10.1007/s11101-014-9349-1>
32. Zielińska S, Kolniak-Ostek J, Dziadas M, Oszmianski J, Matkowski A (2016) Characterization of polyphenols in *Agastache rugosa* leaves and inflorescences by UPLC–qTOF–MS following FCPC separation. *J. Liq. Chromatogr. Relat. Technol.* 39:209–219. <https://doi.org/10.1080/10826076.2016.1147461>
33. Zielińska S, Dąbrowska M, Kozłowska W, Kalebka D, Abel R, Dryś A, Szumny A, Matkowski A (2017) Ontogenetic and trans-generational variation of essential oil composition in *Agastache rugosa*. *Ind. Crops Prod.* 97:612–619. <https://doi.org/10.1016/j.indcrop.2017.01.009>
34. Zielińska S, Dryś A, Piątczak E, Kolniak-Ostek J, Podgórska M, Oszmiański J, Matkowski A (2019) Effect of LED illumination and amino acid supplementation on phenolic compounds profile in *Agastache rugosa* *in vitro* cultures. *Phytochem. Lett.* 31:12–19. <https://doi.org/10.1016/j.phytol.2019.02.029>
35. Zielińska S, Piątczak E, Kozłowska W, Bohater A, Jezierska-Domaradzka A, Kolniak-Ostek J, Matkowski A (2020) LED illumination and plant growth regulators' effects on growth and phenolic acids accumulation in *Moluccella laevis* L. *in vitro* cultures. *Acta Physiol Plant*, 42:72. <https://doi.org/10.1007/s11738-020-03060-w>
36. Ziv M (2005) Simple bioreactors for mass propagation of plants. *Plant Cell Tissue Organ Cult.* 81:277–285. <https://doi.org/10.1007/s11240-004-6649-y>

Figures



Figure 1

A. rugosa hairy roots (K12 clone) cultured in $\frac{1}{2}$ MS liquid medium without growth regulators after 7 days (A), 21 days (B), 28 days (C, D) in 300 mL conical flasks; scanning of electron micrographs (E); scale: 1 cm. Photos A-D were taken by J. Bursy and photo E was taken by B.J. Plachno




Figure 2

Hairy roots of *A. rugosa* cultured in NSB system in 0th day (inoculum) (A), in 17th day (B) . Hairy roots were cultured in liquid ½ MS medium without growth regulators. Photos A-B were taken by M. Barwińska



Figure 3

Hairy roots of *A. rugosa* cultured in NSB system in 0th day (inoculum) (A), in 17th day (B) . Hairy roots were cultured in liquid ½ MS medium without growth regulators. Photos A-B were taken by M. Barwińska




Figure 4

Hairy roots of *A. rugosa* cultured in temporary immersion systems: Plantform® (A, B, C) and RITA® (D, E, F) in 0th day of culture (inoculation) – A, D and in 17th day of culture – B, C, E, F in liquid ½ MS medium without growth regulators. Scales – 1 cm. Photos A-F were taken by K. Radecka.




Figure 5

Biomass (A) and growth index - GI (B) for FW and DW in g/L of *A. rugosa* hairy roots (K12 line) after 17 days of growth in $1/2$ MS liquid medium without growth regulators in different bioreactor systems. Bars indicated with the same letter within each series were not significantly different in Tukey test at $p \leq 0.05$.




Figure 6

UPLC chromatogram of phenolic compounds (acquired at 320 nm) in extracts of *A. rugosa* transformed roots cultured in NSB for 28 days.