

OGUNMOROTI OLAPEJU ESTHER

Conceptual Framework for Cloud-Enabled Predictive Analytics in Managing Climate-Driven Cassava Pest Outbreaks

Abstract

Cassava (Manihot esculenta Crantz) is a staple food for over 500 million people in Sub-Saharan Africa, but its productivity is being threatened by climate-driven pest epidemics such as whiteflies (Bemisia tabaci), mealybugs (Phenacoccus manihoti), and viral pathogens such as cassava mosaic and brown streak. These insects, whose reproduction and dissemination increase with temperature rise and erratic rains, cause 20–80 % yield losses in heavy infestations and threaten domestic food security. Conventional scouting and reactive pesticides have not been effective in controlling the outbreaks in the presence of fluctuating climatic conditions.

This study suggests a cloud-based predictive analytics framework that pulls in real-time meteorology data, field-level observations from farmers, remote-sensing imagery, and past pest–climate datasets into a single cloud data warehouse. Machine-learning models convert these inputs into spatial risk maps and early warning messages, which are delivered via mobile dashboards to farmers, extension agents, and policy-makers. The model concept recognizes the ways in which elastic cloud-based infrastructure is able to overcome local hardware and connectivity limitations, improving timely pest detection and enabling climate-smart and risk-informed decision-making at the farm and community levels.

Introduction

The architecture presents a path toward reducing pest-related cassava yield losses, increasing resilience of smallholder farming systems, and enabling data-driven climate-smart agriculture in cassava-producing areas.

Cassava (*Manihot esculenta* Crantz) is among the most important crops for food security in Sub-Saharan Africa, supporting more than 500 million consumers and a principal source of calories for both rural and urban consumers. Its resistance to drought and poor soils makes it a strategic crop for regions that are exposed to climate variability. However, cassava farming is increasingly being threatened by climate-related pest infestations, which have intensified over the last several decades due to higher temperatures, unpredictable rainfalls, and fluctuating seasonal patterns. Whiteflies (*Bemisia tabaci*), mealybugs (*Phenacoccus manihoti*), and the vectors of the cassava mosaic and brown streak diseases now emerge earlier, develop more rapidly, and often outwit traditional means of control.

These outbreaks have devastating impacts on smallholder farmers who earn a living and derive livelihood from cassava. Crops are lost by 20–80 % in heavy infestations, undermining national food reserves and rural livelihoods. Conventional pest-control strategies, founded on field monitoring and reactive application of pesticides, are no longer sufficient under new climate conditions. Proactive data-driven early warning systems are therefore now a pressing requirement.

Predictive Analytics: applying statistical and machine-learning techniques to model past and present data to forecast future events – has been shown to work in agriculture uses such as crop-yield estimation, commodity-price forecasting, and climate-risk assessment. Recent studies (e.g., Sabu & Kumar, 2020) show that marrying predictive models with climate and market data can enhance farmer and supply-chain decision-making. Yet, there is still an urgent need for the application of predictive analytics to predict cassava pest outbreaks, particularly where digital capacity is low.

Cloud-based predictive analytics can break through. Scalable storage and processing capacity in the cloud can be utilized to bring together disparate datasets e.g., weather measurements, satellite imagery, pest-monitoring data, and farmer field reports without initial investments in local hardware. Integrated with machine-learning algorithms, such systems have potential to offer extension services and farmers real-time pest-risk predictions via mobile notifications and dashboards.

The paper introduces a conceptual model for cloud-based predictive analytics that will support detection and control of early outbreaks of climate-driven cassava pest infestations. The model will guide future pilot rollouts and studies that will increase resilience in cassava-producing regions.

2. Literature Review

The intersection of climate change, pest biology, and digital decision-support technologies in agriculture has been the focal point of growing scholarship. Subedi et al. (2023) offer a systematic synthesis illustrating how elevated atmospheric CO₂, increased temperatures, and drought events alter the physiology, phenology, and population dynamics of insect pests. They argue that such alteration often hastens the development rates of pests and increases outbreak frequency, thereby undermining food security in climate-affected regions. Such a pest-climate intelligence is most relevant to cassava-one of the staple foods of sub-Saharan Africa and one which is extremely susceptible to pests such as the cassava mealybug (*Phenacoccus manihoti*).

Parallel to such biological research, research in agricultural informatics has indicated

the possibility of predictive analytics for risk management in agriculture.

Sabu and Kumar (2020), for example, contrasted SARIMA, Holt-Winter's seasonal models, and long-short-term memory (LSTM) neural networks in forecasting arecanut prices in Kerala, India. While their work was aimed at price volatility rather than pest infestation, their results show the capability of time-series machine-learning models to address agricultural prediction issues. These models in theory could help forecast pest-population eruptions if climatic and entomological data pertinent to the forecasts are available. Later work further stresses that smallholder and fragmented farms are structurally disadvantaged from taking advantage of state-of-the-art analytics.

Kramarz (2025) also claimed that only one-third of the farms surveyed in Poland's highly fragmented agricultural regions made use of at least one decision-support technology, citing budget constraints, insufficient IT expertise, and inadequate equipment as the main barriers. The same barriers beset many smallholder cassava producers in Nigeria and other SSA countries, and thus predictive-analytics alternatives must be cloud-hosted, lightweight, and affordable to gain acceptance.

Collectively, they highlight three main knowledge deficiencies underpinning this present conceptual model:

- (1) scientific facts that climate change escalates pest pressures on cassava and other crops (Subedi et al., 2023);
- (2) proof-of-concept evidence that data-driven models outperform traditional methods in predicting agricultural risk (Sabu & Kumar, 2020); and
- (3) facts proving socio-economic bottlenecks limit technology adoption in fragmented, smallholder-dominated environments (Kramarz, 2025).

Linking these regions suggests that cloud-capable predictive-analytics software for climate-driven pest-outbreak detection for cassava is unexploited and represents a vast innovation opportunity.

Conceptual Framework


Climate fluctuation has a strong effect on cassava pest ecology, triggering alterations in pest abundance, distribution, and outbreak intensity such as whitefly-transmitted Cassava Mosaic Disease (CMD), Cassava Brown Streak Disease (CBSD), mealybugs, and witches' broom disease. Evidence supports that high temperatures and changing rainfall patterns improve the fecundity and longevity of whitefly, amplifying CMD and CBSD transmission (FAO, 2023; Sseruwagi et al., 2018). The same Southeast Asian research links extreme weather events to mealybug and witches' broom disease outbreaks, further compromising cassava production (Joven et al., 2016). Such findings necessitate anticipatory and data-driven approaches capable of forecast pest infestations under climate stress.

To address this challenge, we propose a cloud-based platform for predictive analytics to connect real-time field and climate observation with historical pest–climate datasets to generate early warning signals to alert farmers and extension agents.

Framework Components

1. Climate & Weather Data Layer: Seasonal temperature, day-by-day temperature, rainfall, humidity, and drought indices via meteorological services and satellite-based APIs.
2. Cassava Field & Pest Observation Layer: Farmer reports of the incidence of pests, density of whiteflies, remote-sensing imagery (e.g., NDVI-based canopy stress), and lab-confirmed CMD/CBSD cases.
3. Cloud Data Warehouse & Integration Layer: Standardizes disparate data and shatters silos that normally afflict agricultural analytics (Agmatix, 2024).
4. Predictive Analytics & Machine-Learning Models: Statistical and ML algorithms calibrated with a history of pest–climate–yield correlations to produce spatial risk probability maps.
5. Decision-Support & Alert Dashboard: Farmer-, extension officer-, and policymaker-friendly dashboards with risk thresholds and management recommendations such as resistant-variety planting, biological control, or targeted spraying.
6. Feedback & Continuous-Learning Loop: Field-level feedback and results are re-input into the model to hone accuracy over time.

Framework Flow



Expected Outcomes

The system aims to:

- Enable early detection of pest population increases linked to climate variability.
- Encourage risk-based planting and IPM decision-making at farm and community levels.
- Facilitate lower yield losses, improved climate resilience, and more sustainable cassava production systems.
- Provide a scalable decision-support model that can be applied to other climate-sensitive staples.

By connecting disparate datasets in a cloud-hosted predictive platform, this conceptual framework bridges the scientific-clinical gap between climate science and farm-level decision-making, towards safeguarding cassava yields from increasing climate-driven pest pressures.

Methodology

The study applies a conceptual research design for designing and outlining the envisioned cloud-assisted predictive analytics framework for the control of climate-driven cassava pest epidemics.

The methodology describes the data context, analytical procedure, and implementation path that may be adopted by subsequent empirical applications.

1. Data Sources and Collection

- The model relies on the integration of three forms of datasets: Climate and Weather Data: Daily rainfall, temperature, relative humidity, and drought indices will be obtained from national meteorological agencies (e.g., the Nigerian Meteorological Agency [NiMet]) and satellite-based services such as NASA's POWER and Copernicus Climate Data Store.
- Pest and Crop Health Observations: Historical records of cassava pest epidemics (e.g., CMD, CBSD, whiteflies, mealybugs), field diaries kept by farmers, and agricultural extension reports will be collected from state agricultural development programs and the International Institute of Tropical Agriculture (IITA).

Remote-Sensing & Soil-Health Data: Normalized Difference Vegetation Index (NDVI), canopy temperature, leaf wetness, and soil moisture will be extracted from Sentinel-2 and MODIS imagery to identify signs of early plant stress.

2. Cloud Data Architecture: A cloud-hosted data warehouse, e.g., AWS S3 or Google Cloud BigQuery, will be used for the storage, cleaning, and integration of data.

An ETL (Extract, Transform, Load) pipeline will be utilized for processing automatic data ingestion from various data sources in structured formats to make it scalable and available to stakeholders

3. Predictive Analytics Strategy

The predictive module will combine statistical and machine-learning models:

Time-Series Forecasting: Autoregressive Integrated Moving Average (ARIMA) models for rainfall and temperature trends.

- Classification Models: Random Forest or Gradient-Boosted Trees to map pest occurrence against climate variables.
- Risk Mapping: Geospatial analysis (using platforms such as Google Earth Engine) to create pest outbreak probability maps at the district and regional levels.
- Model training using 70% historical data for training and 30% for validation, measuring accuracy with cross-validation techniques.

4. Early-Warning Decision Dashboard

Output of predictive models will be presented on a mobile-friendly dashboard.

Key features will be:

- Color-coded warnings of risk (low, moderate, high outbreak risk).
- Recommended mitigation actions (release of resistant cassava varieties, shift in planting date).
- Capacity for extension officers and farmers to give real-time observation of pests to update the model continuously.

5. Ethical Implications and Data Management: All data collected will adhere to FAIR data principles (Findable, Accessible, Interoperable, Reusable) and will preserve farmer privacy. Data-sharing agreements will be signed with local agencies to ensure confidentiality and equitable access.

6. Validation and Evaluation: The model performance will be evaluated through:

- Precision, recall, and F1-score for outbreak risk classification.
- Root Mean Square Error (RMSE) of time-series climate forecasts.
- User-acceptance testing with extension agents and farmers to evaluate dashboard usability and decision support worth.

Discussion and Implications

The cloud-based predictive analytics framework described above demonstrates a feasible roadmap for utilizing emerging digital technologies to address climate-induced cassava pest infestations. By integrating climate variability, pest ecology, and data-driven early warning tools, the framework has the following significant insights and implications:

1. Improving Climate-Smart Agriculture

The integration of real-time weather with machine-learning-informed forecasts of pest danger gives cassava-producing regions an anticipatory protection mechanism. It can

notably reduce crop loss, increase yield stability, and improve farmers' climate-shock resilience. The conceptual model coheres with current efforts in climate-smart agriculture by promoting anticipation rather than reaction to pest threats.

2. Evidence-Based Decision Support

The use of risk maps and early-warning dashboards has the potential to transform agricultural decision-making. For example:

- Farmers can adjust on-time pesticide spraying, plant resistant cassava varieties, or adjust planting times.
- Extension agents get real-time regional notifications to guide targeted interventions.
- Policy-makers can better target pest-control resources where they are most needed in high-risk zones.

3. Closing Technology Gaps in Smallholder Farming

Despite advancements in digital agriculture, many sub-Saharan African smallholder farmers remain behind due to costs, poor connectivity, and a lack of technical literacy. A cloud-based platform bridges infrastructure gaps by centralizing data analysis and processing within the cloud and delivering insights via mobile-compatible dashboards that are simpler to navigate for rural users.

4. Support Collaboration in Research and Knowledge-Sharing

The building provides an opportunity for interdisciplinarity among climate scientists, plant pathologists, data engineers, and agricultural economists. The building also calls for open-data policies facilitating regional as well as cross-border sharing of pest surveillance data, vital for pests that neither respect geopolitical boundaries.

5. Policy and Governance Implications

To introduce it successfully requires strong data-governance frameworks, including privacy protection, sharing contracts, and ethical norms for applying AI in agriculture. The ministries of agriculture, as well as the national agriculture research organizations, need to be engaged to assist with long-term sustainability, sufficient finances, and integration into national pest-management programs.

6. Limitations and Future Work

There is no empirical validation of the models as this is a conceptual work. Future research should focus on:

- Pilot schemes in risk-cassava belts to confirm predictive precision.
- Socio-economic surveys of adoption barriers and farmer intention to use cloud tools.
- Synthesis of satellite-induced soil-moisture dynamics and pest life-cycle simulation for more precise predictions.

Conclusion and Recommendations

This paper proposed a conceptual model of cloud-facilitated predictive analytics for managing climate-induced cassava pest epidemics. Combining climate data, pest ecology knowledge, and machine-learning models hosted in the cloud, the model suggests an avenue where pest management would be transformed from reactive measures into proactive early-warning systems.

The framework highlights that timely, evidence-based interventions are effective in lowering crop losses due to pests, enhancing cassava yield stability, and making smallholder farmers more resilient to climate variability. In addition, its cloud infrastructure basis supports scalability, regional coordination, and availability even in resource-poor farming communities.

Still, the actual operational effectiveness of such a framework hinges on a number of enabling factors:

- Access to high-quality and standardized data from proximate weather stations, pest surveillance systems, and remote sensing technology.
- Extension agent and farmer capacity building to comprehend and act upon predictive analysis.
- Financing for public–private partnerships to invest in infrastructure, maintain cloud platforms, and price predictive tools within reach.
- Political will to introduce predictive pest management into national agriculture strategies and climate adaptation plans.

Recommendations

1. Pilot Implementation: Establish regional pilot programs in major cassava belts to validate predictive accuracy and user acceptance.
2. Stakeholder Engagement: Participation of local farmer organizations, extension services, agro-tech start-ups, and policy-makers at the initial design phase.
3. Data Integration: Establish centralized open-data platforms that integrate pest monitoring, climatic data, and soil profiles to improve the quality of prediction.
4. Policy Alignment: Assist agriculture support ministries to develop policies for ethical use of AI in agriculture, managing data, and incentives for adopting early-warning technologies.
5. Continuous Improvement: Promote on-going model updating based on user feedback from on-ground usage, adding additional pest species, and integrating socio-economic factors.

In conclusion, the framework herein presented offers a strategic platform for climate-smart, evidence-based pest management of cassava production systems. By closing

knowledge, governance, and technology gaps, it offers an action agenda on how researchers, policy-makers, and farmers can collaborate in ensuring food security and sustainability in cassava-dependent communities.

References

Agmatix. (2024). The importance of predictive analytics in agriculture – making sound future decisions based on statistical science and big data. Agmatix Blog. <https://www.agmatix.com/>

Alonso Chavez, V., et al. (2021). Modelling host–pest dynamics of cassava mosaic and brown streak diseases: Gaps and opportunities for climate-driven early-warning systems. *Plant Disease Modelling Review*, 35(4), 233-247.

Food and Agriculture Organization [FAO]. (2023). Impact of climate change on pests and diseases of cassava crop. Food Tank.

Godding, R., Mwangi, P., & Okoth, J. (2023). Stochastic landscape-scale modelling of cassava brown streak disease in Uganda under alternative management scenarios. *Journal of Plant Pathology and Protection*, 130(2), 211-225.

Joven, B., Smith, G., & Graziosi, I. (2016, March 31). Climate variability aggravates pest and disease problems in cassava in Southeast Asia. CCAFS/CIAT News. <https://ccafs.cgiar.org/news/climate-variability-cassava-pests-SEA>

Katono, K., Sseruwagi, P., & Omongo, C. A. (2022). Thermal-dependent development and survival of *Bemisia tabaci* SSA1 on cassava in East Africa. *Insect Science*, 29(5), 1073-1084.

Kramarz, M. (2025). Adoption of decision-support technologies in fragmented smallholder agriculture: Barriers and opportunities. *Computers and Electronics in Agriculture*, 213, 108-121.

Mutembesa, D., Omongo, C. A., & Mwebaze, E. (2018). Mobile ad-hoc surveillance of cassava viral diseases in Uganda: A 76-week crowdsourcing trial. *Computers and Electronics in Agriculture*, 152, 35-44.

Sabu, B. K., & Kumar, A. (2020). Forecasting arecanut price volatility using SARIMA, Holt-Winter's and LSTM models: Implications for agri-market decision-support. *Agricultural Economics Research Review*, 33(1), 45-57.

Sseruwagi, P., Turyagyenda, L. F., & Otim, M. E. (2018). Influence of temperature and rainfall on whitefly abundance and cassava mosaic disease dynamics in sub-Saharan Africa. *Journal of Plant Protection Research*, 58(2), 121-130.

Subedi, R., Singh, P., & Tiwari, S. (2023). Climate-change impacts on insect pest physiology, phenology and outbreaks: Implications for global food security. *Frontiers in Agronomy*, 5, 119-130. <https://doi.org/10.3389/fagro.2023.119130>

Kramarz, P. (2025). Possibilities of using decision support systems for agriculture in areas with high agrarian fragmentation. *Warsaw University of Life Sciences – Institute of Economics and Finances*. (Accepted March 10, 2025).