

“Bauhinia Racemosa: Exploring its Pharmacological Powers for Optimal Health and Wellness”

Bhagyashri N. Nehete*, Darshana P. Patel, Yogesh B. Rokade, Sunil P. Pawar
¹²Student, ³Assistant Professor, ⁴Principal Department of Pharmaceutical Chemistry,
P.S.G.V.P. Mandal's College of Pharmacy, Shahada, Nandurbar, Maharashtra, India

***Corresponding Author**
E-Mail Id: nehetebhagyashri4@gmail.com

ABSTRACT

Since the Vedic era, medicinal herbs have been utilized. They have been used for many years to treat and prevent many diseases. The Bauhinia racemosa Lam is a member of the family Caesalpiniaceae. This tree, which is small and gnarly with dangling branches, is found throughout India and reaches a height of 1,650 meters in the western Himalayas. This review aims to comprehend the therapeutic potential of Bauhinia racemosa by combining traditional and medical knowledge with current scientific results. Within the Bauhinia Genus there are more than two hundred species of blooming plants; one of them is racemosa. It goes by the name “sonpatta tree” because of its gold-like therapeutic properties, nearly every single part of the plant.

Keywords: *Bauhinia racemosa, medicinal uses, phytochemistry, pharmacology caesalpiniaceae, anti-inflammatory, antipyretic*

INTRODUCTION

For a lengthy duration of time, plants have been a treasured source of Natural merchandise for retaining especially in the last ten years, with a greater focus on natural remedies and human wellness. In developing nations, the majority of rural residents almost exclusively use over-the-counter medication to treat a wide range of illnesses.

Based on the antioxidant, antibacterial, and antipyretic properties of their phytochemicals, plants should be used for medical purposes.

Similarly, because knowledge about the application of certain plant species for medicinal purposes has been passed down orally from one generation to the next, knowledge about medicinal plants has also started to fade and eventually become obsolete owing to a lack of awareness. of

the younger generations as a result of a shift in perspective and a societal perception of what is valuable. The racemosa baihinia racemosa are sweetish and acrid, used as a refrigerant,

Astringent, in the cure of headache, fever, pores and skin diseases, Blood diseases, dysentery, and diarrhea It is recommended to use a decoction of the Bark as an effective ulcer wash [2]. The herb is commonly utilized in Ayurveda to cure most malignancies in their early stages [3,4].

Numerous studies have demonstrated that B. racemosa possesses a wide range of pharmacological effects, including antitumor, anti-inflammatory, analgesic, antipyretic, antihyperglycemic, hepatoprotective, antimicrobial, antiulcer, antihistaminic, anxiolytic, and anti-HIV properties.[4] is recommended to use a

decoction of the Bark as an effective ulcer wash [2]. The herb is commonly utilized in ayurveda to cure most malignancies in their early stages [3,4].

Numerous studies have demonstrated that *B. racemosa* possesses a wide range of pharmacological effects, including antitumor, anti-inflammatory, analgesic, antipyretic, antihyperglycemic, hepatoprotective, antimicrobial, antiulcer, antihistaminic, anxiolytic, and anti-HIV properties. The medical Value of this plant attracts greater attention of the human beings due To the complications of health issues.

With the advent of Chemical composition and pharmacological research, the Clinical application of *Bauhinia racemosa* has been Hence, a growing wide variety of researchers have dedicated their spare time to this extraordinary plant in the final few decades, from botany to phytochemistry and pharmacology.

However, minimal numbers of articles have reviewed the comprehensive research of *B. racemosa*. The reachable report exhibits that there is only a concise evaluate of its photochemistry and pharmacology, which can also no longer be enough for researchers to definitely understand this plant.

Researchers have identified countless new chemical components from *B. racemosa* and should make extensive analyzes of its pharmacological effects. This article discusses the common uses, botany, phytochemistry and pharmacological properties of *B. racemosa* in a precise form

that helps to apprehend readers to a more desirable and deeper understanding of this medicinal plant and additionally furnish thorough information for higher research and development of *B. racemosa*.

The extracts of leaves and bark have proven analgesic, anti-pyretic, antimicrobial, antiinflammatory, anti-spasmodic, and anti-helminthic efficacy.

The plant *bauhinia racemosa* is recognized as Banraj (Bengali), Asundro (Gujrat), Jhinjeri, Kachnal, ashta (Hindi), Banne (Kannada), Katapuli (Malaylam), Apta (Marathi), Ombaroda (Oria), Svetakanchana (Sanskrit), Araivatta-atthi (Tamil), Ari (Telugu), Kosundra (Punjabi). Bark bluish black, rough, pinkish crimson inside, turning brown on exposure; Leaves broader than lengthy , rigidly Coriaceous , barely cordate , clothed more or less densely below with gray pubescence; vegetation yellow, in Short-pedunched, lax, terminal and leaf-opposed racemes; pods falcate, glabrous ,turgid; seeds dark reddish Brown.

The bark yield a robust cordage fibre, additionally used for torches and sluggish matches. It is notably astringent and is used for dysentery. It showed full-size anti-inflammatory and cholagogue activities. Octacosane, β - Amylin, and β -sitosterole have been isolated from the bark.

The leaves are given with onion for Diaehoea;they are also suggested to be anthelmintic. The leaves are used in manufacturing of bidis.




Fig. 1: Plant of Bauhinia Racemosa.

CLASSIFICATION

Botanical Name: Bauhinia racemosa lam.

Local Name: English (Burmese Silk Orchid), Gujarati (Apto), Hindi (Kachnar), Kannada (Aapta, Aralukadumandara), Sona Patti.

Kingdome- Planate

Division – Angiosperms

Class- Eudicots

Order- fabales

Family- Fabaceae

Subfamily- Caesalpinoideae, [6]

Genus- Bauhinia

Species -*B. racemosa*.

HABITATES AND DISTRIBUTION

It is veining and evergreen plant which is having vigorous Growth wanting sturdy support. It is Very common in foothills up to 1000m. There Global Distribution India and Sri Lanka Indian Distribution State – Kerala, Tamil, Maharashtra etc.[8] Throughout India, besides Jammu.

NUTRITIONALSIGNIFICANCE

Bauhinia racemosa seeds are rich in calcium, potassium, magnesium, zinc, manganese and iron. Bauhinia racemosa seed protein is dominated by glutelins, but

less by albumins and globulins. Essential amino acids like isoleucine, lysine, phenylalanine and tyrosine are excessive whereas the contents of sulphur amino acids are limiting in the seed proteins. The fatty acids, linoleic, oleic and palmitic acid are noticeably higher in the seed lipids [8,9,10]

MORPHOLOGY

It is a small, broad leafed tree, up to 12 meter Bark grey to black, rough, thinly scaly, with number of vertical cracks. Leaves simple, bilobed, alternate, stipules small caduceus. Flowers bisexual, 10-12 mm across, yellowish white, in terminal & leaf opposed few flowered Racemosa. Fruit pods 15-22*1.5-2 cm, oblong, blackish brown, turgid, apex horned indehiscent. Seeds 10-20 ovoid.

COLLECTION PERIOD

Flowering season: March-June, **Fruiting period:** November and December, **Pale:** Blooms in sunny conditions and blooms best in good sunlight. **Moisture:** Water well and regularly, keeping it evenly moist. You need extra water in hot seasons

and less in colder climates. While the plant is drought resistant also.



Fig. 2: Leaf of *Bauhinia racemosa*.

DISCRIPTION

Trees, deciduous, small, up to 15 m. Bark black, rough; branches Spreading or pendulous, zigzag, thin, smooth. Stipules decaying; petiole 0.8-1.2 cm; Leaf blade very orbicular, 1.5-4 × 2.2-6 cm, 7-9 grooved, leathery, abaxially hairy or glabrous, abaxially glabrous, base cordate, apex about 1/3 bifid, apices rounded. It is an inflorescence, lateral or terminal shield, c. 20-flowered; stem short; bracts and bracteoles linear. Flower buds obovoid, pubescent, apices protruding.. turbinate is Short. Calyx split spathaceously at a thesis. Petals yellowish subequal, oblanceolate, 8-10 Mm, subsessile. Fertile stamens 10, unequal; filaments 6-7 mm; anthers small, ca. three mm. Ovary stalked, glabrous; stigmas subsessile, peltate, small. There Legume linear-cylindric, 15-20 × 1.8-2.2 cm; valves woody, glabrous. Seeds 12-20, darkish brownish, ellipsoid, 8-10 mm In diam. Fl. Apr-May, fr. Jun-Aug. Bauhinia timber usually reach a peak of 6-12 m and Their branches spread 3-6 m The leaves are usually 10-15 cm in diameter. [12] The five-petalled plants are 7.5-12.5 cm in

diameter, usually red, pink, purple, orange or yellow in color and often fragrant... Tree starts flowering in late winter and often Continues to flower into early summer.[13]

USEFUL PART OF PLANT

The components which are historically used of these B.racemose. It is used in Traditional medication for the remedy of various ailments. The stem bark of the plant is an Astringent and is used in the remedy of headache, fever, pores and skin diseases, and tumors, ailments Of the blood, dysentery and diarrhoea.[14] Pharmacological studies of the plant printed that The ethanol extract of leaves of B. racemosa introduced analgesic, antipyretic, anti- Inflammatory, antispasmodic,[15] and antimicrobial activity. [16] The fresh flower buds of this Plant showed antiulcer activity. [17] Cytotoxicity against CA-9 KB in telephone culture, hypotensive, And hypothermic things to do had been additionally mentioned from the hydroalcoholic extract of B. Racemosa. [18]

Table 1: Part of *Bauhinia racemosa lam.* Used for medicinal purpose.

Sr. No.	Parts of Plants	Medicinal Preparation
1	Stem Bark	Astringent, headache, fever, Skin disease & tumor, Disease of the blood, Dysentery and diarrhoea, anthelmintic, antidiabetic, antineoplastic
2	Leaves	ntipyretic, analgesic, anti-inflammatory, antispasmodic and antimicrobial activity
3	Flower	Antiulcer activity, Cytotoxicity, hypotensive and hypothermic activity, Haemorrhages.
4	Seeds	Antibacterial and antimicrobial activity.

PHYTOCHEMISTRY

Phytochemical screening of the plant leaves exhibits the presence of carbohydrates [19-20], alkaloids [19-20], steroids [19,21,20], glycosides [21,22], tannins [23,22], saponins [25,26,27], phenolic compounds [23,21,20], flavonoids [19,23,21,22,,24], protein [27,30], oil and fat [27,28]. Bioassay guided fractionation of ethanolic extract of the leaves led to the isolation of galactolipid and catechin class of the compounds from the most lively n-butanol fraction [24]. Phytochemical screening of the seed reveals the presence of palmitic and stearic acids [29]. It also carries phenolics, flavonoids, saponins, glycosides, tannins and proteins [22]. De-O-methylracemosol used to be acquired From the column chromatographic separation Of a benzene extract of the roots, recrystallized From benzene as brown crystals [30]. Racemosol and de-O-methylracemosol have been synthesized by way of way of Sae-Lao et al [31]. The form of pacharin, Previously isolated from the heartwood of B. Racemosa Lamk has been established as 1,7- dihydroxy-3-methoxy-2-methyl-dibenzo(2,3-6,7)oxepin by way of a find out about of its chemical and spectroscopic properties, together with X-ray evaluation [28,32]. Resveratrol (trans-3,5,4'-trihydroxystilbene), a phytoalexin is bought from various flowers and additionally from B. racemosa [33]

Crude fibre, fats and carbohydrate contents have been discovered to be 1.91%, 15.53% and 36.12% respectively in B. racemosa seeds. Its saponification and acid values were determined to be 343.72. 5.64 [34]. Crude proteins, crude lipids, ash and nitrogen free extractives represent 19.84%, 9. racemose Lamk; while in the germplasm of Mundanthurai Wildlife Sanctuary they are 19.31%, 8.94%, 3.81% and 61.30% respectively. Caloric values were determined as 407.64 Kcal (germplasm from Ayyanarkoil Forest) and 402.90 Kcal (germplasm from Mundanthurai Wildlife Sanctuary). Essential amino acids such as isoleucine, tyrosine, phenylalanine and lysine have been found to be abundant in the seed proteins of all germplasm. More fatty acids, palmitic, oleic and linoleic acids were found in the seed lipids of each germplasm. Both germplasms appeared to be rich sources of calcium, potassium, magnesium, zinc, manganese and iron. Antinutrients such as total free phenols, tannins, %, and 60.65%, respectively in Ayyanarkoil Forest germplasm of the tribal pulse, B. L-DOPA and phytohaemagglutinating undertaking had been additionally investigated [35] Dahikar et al. (2011) confirmed that petroleum ether extract of B. racemosa (PEEBR) had solely alkaloids and steroids taking place in high concentration [19]. The chloroform extract (CEBR) and ethyl acetate extract (EAEBR of B. racemosa had similar photochemical constituents, alkaloids and steroids in greater

concentration, carbohydrates, flavonoids, glycosides and saponins. Methanol extract of *B. racemosa* (MEBR) had alkaloids, steroids, flavonoids, saponins and tannins, however no protein. Kumar et al. (2010) found the presence of protein, oil and fats, phenolic compounds, flavonoids, alkaloids, saponins, tannins and carbohydrates as principal phytochemical corporations in crude MEBR and aqueous extract of *B. racemosa* (AEBR) leaf [23]. Manohar et al. (2011) located the presence of carbohydrates, glycosides, alkaloids, phytosterols, saponins, tannins, phenolics, flavanoids, fixed oils and fats as phytochemicals in crude MEBR and AEBR leaf [21]. Anjaneyulu et al. (1986) stated a new tetracyclic phenol, racemosol

from the hexane extract (HEBR) of the heart-wood of *B. racemosa* Lamk [28]. Racemosol was once crystallized as deep pink prisms from chloroform. It is soluble in alkali giving yellow coloration which reverts to the pink colour on acidification. Racemosol diacetate and racemosol dimethylether was synthesized by him. Prakash and Khosa (1976) mentioned octacosane, β -amyrin and β -sitosterol in PEEBR stem bark from column chromatographic resolution over Brockmann alumina [41]. Elution of the column with petroleum ether (60°-80°) furnished octacosane, with benzene β -amyrin and with benzene:chloroform (3:1) β -sitosterol.

Table 2: Active constituent present in part of *B. racemosa* lam.

S. No.	Parts of plant	Phytoconstituents
1	Leaves	Flavonol (kaempferol & Quercetin) and coumarins (Scopoletin & scopolin).
2	Stem Bark	B-sitosterol and β -amyrin.
3	Hard wood	Stilbenes (resveratrol), Dibenzoepin
4	Flower	Flavonoids, terpenoids, and steroids.

TRADITIONAL USES

Bauhinia racemosa is one of the plant life oftentimes used by way of Ayurvedic medical practitioner for the prevention and therapy of cancer[36]. Traditionally, mixture of its bark and *Bridelia retusa* is given orally to female to boost sterility in tribal region of southern Rajasthan and act as contraceptive [37,38].

The people and the locales of the Shiwalik Himalaya of Uttarakhand make use of it in the structure of one-of-a-kind merchandise as ethnomedicine, fodder, food, fibre etc. It is commercially exploited with the aid of drug sellers and it come in threatened categories. Therefore, there is an urgent want of its conservation for sustainable development [39]. This is one of the plant life eaten by way of the Rhesus monkey

(*Macaca mulatta*) and Hanuman langur (*Presbytis entellus*) in Himachal Pradesh [40].

Its stem bark is saved in house as snake Repellent. Leaves are chewed through two humans and air is blown in patient's ear, after which a glass of water is given to drink to cure scorpion chunk [41]. Pounded bark powder of Kachnal Tree, in 10 g quantity, or decoction Witwatersrand in 20 ml quantity, is used twice per day for continual dysentery and diarrhoea in the tribals of Sonaghati of Sonbhadra district, Uttar Pradesh, India [42,32]. It is used in headache also [32]. Whole plant juices either wild or cultivated, administered with vegetable oil orally is used in leucorrhoea by way of 5% people of Sariska vicinity of Rajasthan [28]. For the treatment of

jaundice, Water extracts of its bark, leaves and roots are taken two times each day after meal for 2-4 weeks in the region of Jalgaon, Dhuleand, Nandurbar districts of Maharashtra and also the stem bark of *Woodfordia fruticosa* (L.) Kurz floor with its bark and *Oroxylum indicum* are taken as poultice administered in two spoonfuls banana fruit twice a day for 5-7 days in Adilabad district of Andhra Pradesh [43]. Leaves of *B. racemosa*, *Aloe vera* and *P. murex* overwhelmed together and mixed with water given to animals three times a day can relief food poisoning in cattle [44].

INDUSTRIAL USES

Medicinal flowers are the richest bio useful resource of tablets for regular structures of medicine, present day medicines, nutraceuticals, meals supplements, people medicines, Pharmaceutical intermediates and chemical entities for artificial drugs. The first step in the Value addition of medicinal bio assets is the manufacturing of herbal drug preparations, using A range of methods from simple normal applied sciences to advanced extraction techniques.

The extract is further processed to be integrated in any dosage form such as capsules and Capsules. With the increasing demand for natural medicinal products, nutraceuticals, and Natural products for pharmaceuticals, and herbal products for health care all over the world, Medicinal plant extract manufacturers and vital oil manufacturers have commenced the usage of the most appropriate extraction technologies in order to produce extracts and essential oils of defined quality with the least variations from batch to batch. [45].

PHARMACOLOGICAL ACTIVITY

Bauhinia racemosa indicates a range of pharmacological Activities like Chemoprotective antitumor activity, antidiabetic activity, Hepetoprotective, Anti-inflammatory activity, Antimutagen activity, antidiabetic activity, Antinociceptor, Antioxidant, antihyperlipidemic pastime and many others due to its presence of quite a number active constituent All over the parts of plant. By considering the ethno medicinal background and it has been concluded that this plant carries some phytochemical materials which are accountable for more than a few activity which has been pronounced such as, quite a number pharmacological activity.

Anti-Diabetic Activity

Diabetes is a metabolic sickness where in human physique does Not produce or properly uses insulin, a hormone that is required to convert sugar, starches, And different meals into energy. Diabetes consequences in odd tiers of glucose in the Bloodstream.[46] Alloxan precipitated hyperglycaemic rats showed a good sized reduce (In body weight on days 10, 15, 20 of the experiment.

Daily oral Treatment with each extracts Showed extensive amplify (body weight at the stop of the experiment as Compared to diabetic manipulate group. The present learn about shown the therapy of alloxan Induced rats with both methanol and aqueous extracts of *Bauhinia racemosa* for 20 days could fix the normal biotransformation via transferring the stability of carbohydrate Metabolism. Improved pancreatic exocrine activities can be ascribed to insulin secretion from Existing residual Beta-cells of islets or due to enhanced transport of blood glucose to Peripheral. The rats were divided in to seven companies and each crew consists of six animals. The serum was once analyzed

for blood glucose level (GOP-POD method).[47,48]

Analgesic Activity

A lookup in 2005 has proved that Methanol mixed With *B. racemosa* stem bark possessed remarkable Analgesic efficacy at the dose of two hundred mg/kg. In writhing Method, the inhibition exercise is shown to be 53.41%. In Hot plate method, the rate of reaction time significantly Increased at the dose of two hundred mg/kg, which improved From 9.36 ± 0.74 to 16.57 ± 1.34 . Aspirin and morphine are used as control pills to determine the ability of Stem bark of *B. racemosa*. This is tested on acetic acid-Induced writhing and the warm plate method pain-induced Mice as well.

Aspirin is used as a correspondent to anal-Gesic motion in acetic acid-induced writhing mice, whereas in the hot plate process, morphine is used as an Analgesic agent in pain-induced mice. This experiment Also examined analgesic efficacy the usage of a combination of MEBR and authorized drugs. There, the combinations Exposed more strong analgesic activities than the individual doses of accredited tablets and prepared extracts. Especially, the combos of aspirin and MEBR exhibited notable writhes inhibition percentages such as 71.36% in a hundred 50 mg/kg, 77.04% in a hundred 100 mg/kg, and 84.02% in a hundred 200 mg/kg. To determine the analgesic activity, the stem bark of *B. racemosa* used to be subjected to an analgesic experiment On tail lesion-induced albino Wister rats [49]. Analgin is Used as a testing drug to examine the effectiveness of The *B. racemosa* stem bark with that of a commercial Drug.

The extracts are prepared the usage of aqueous and ethanolic solvents and examined at different doses with each

Solvent at 100 mg/kg and 2 hundred mg/kg respectively. BoriKar *et al.* [49] stated that *B. racemosa* stem bark displayed widespread analgesic efficacy in Wister rat tail Lesions when delivered with alcohol. They also stated that Rats dealt with aqueous extracts at a dose of one hundred mg Had no analgesic activity in tails. Eventually, 200 mg of *B. racemosa* alcoholic extract has a better analgesic characteristic comparable to 10 mg of analgin. In their experiment, the aqueous extract did not possess analgesic action due to a lack of sufficient dose. Chandrasekar and Prasanna [50] tested the analgesic potency of *B. racemosa* stem bark when combined with petroleum ether. In order to consider the activity, the researchers used the animals in two techniques, such as acetic acid-induced writhing and thermal stimulus-induced ache technique (hot plate methods).

The animals were categorized into three organizations in each acetic acid induced writhing and hot plate method. In their trial, Aspirin was used as a manage agent to the writhing pain- induced rats by acetic acid, whereas Pentazocin used to be used as a control agent to the pain-induced rats by means of warm plate method. Eventually, Chandrashekar and Prasanna [50] reported that the petroleum ether blended stem bark of *Bracemosa* had no tremendous analgesic results in animal model. From the conclusion of preceding studies, we realized that it exposes efficient activities when the extract combined with the high polar solvents such methanol and alcoholic solvents.

Antiulcer Activity

The antiulcer activity of aqueous and alcoholic extracts of the stem bark Of *Bauhinia racemosa* Lam was once observed as per the technique described through Sahani et al, (1990).[51] To find out about the antiulcer activity, the ulcer

no. and ulcer rating have been calculated and Analyzed for the extensive discount in the direction of normal. There was full-size reduction in the Ulcer no. on administration of aqueous (200mg/kg body weight) and alcoholic extracts (100 And 200mg/kg body weight) of *Bauhinia racemosa* (Pin contrast with manage Group).[52]

Anti-HIV-1 Activity

B. racemosa extracts were minimally poisonous and confirmed anti-HIV-1 Activity. Ethyl acetate fraction from methanol (80%) extract of *B. racemosa* had a larger Cytotoxic effect. It was significantly distinctive from that of the different extracts.[53] The anti- HIV-1 endeavor assay used to be performed via syncytia formation. These consequences may additionally be explained by The presence of the following phytochemical in the methanol extract; tannins, flavonoids, Carbohydrates, terpenes and coumarins. These phytochemical outcomes are similar to these Obtained via Prakash and Khosa, (1976) and Kumar et al. (2007), remoted from one of a kind components Of the plant such, as β -sitosterol and β -amyrin had been from the stem bark two flavonols (kaempferol and quercetin) and two coumarins (scopoletin and scopolin) from the leaves of The plant . Stilbene (resveratrol) was once separated from the heart wooden of *B. racemosa*. [54].typhimurium, but moderate inhibitory effect on *Escherichia coli*, *Enterobacter aerogenes*.

Anti-histaminic Activity

Anthelmintic pastime of stem bark, leaves, seeds and root of *B. Racemosa* had been one at a time studied the usage of Indian person earthworms, *Pheretima posthuma*. Results confirmed that the leaves of *B. Racemosa* took less Time to motive paralysis and dying of the earthworms. The petroleum ether, chloroform, ethyl

acetate and Methanol extracts of leaves had been also studied for their anthelmintic activity, which concerned determination of the time of paralysis and time of loss of life of the worms. Results confirmed that the petroleum ether extract of Leaves of *B. Racemosa* at 60 mg/ml used to be most mighty as compared to different extracts and popular drug Albendazole. The order of efficiency was observed as petroleum ether > ethyl acetate > methano.37 order of potency was located As petroleum ether > ethyl acetate > methanol > chloroform extract. It can be concluded that anthelmintic Activity of the leaves of *B. Racemosa* is due to the lively ideas present in the petroleum ether and ethyl Acetate extracts .[57]l > Chloroform extract. It can be concluded that anthelmintic endeavor of the leaves of *B. Racemosol* and de-0-methylracemosol exhibited cytotoxicity against KB mobilephone line (EC90 at 15.0 μ g/ml and 5.6 μ g/ml, respectively) and BC telephone line (EC90 at 6.1 μ g/ml and 3.6 μ g/ml respectively) exhibiting moderate Antimalarial undertaking in opposition to parasite plasmodium falcipararum (EC50 at 0.9 μ g/ml and 2 μ g/ml, respectively). [58]

Antixiity Activity

The MEBR, extract administered orally to swiss albino mice at the dose 150mg/kg and 300mg/kg. Increase In the time spent and the elevated plus-maze. It additionally extended the time spent by means of mice in the illuminated side Of the light-dark test, showed widespread increase in nose poking and decese locomotion in holr board test Also brought about giant discount in freezing time in evaluation with manage group. That point out MEBR is An fantastic anxiolytic agent. [59]

Antioxident Activity

The leaves of AEBR confirmed properly anti- oxidant endeavor in distinctive

machine of assay. IC₉₀ price found have been 1739, 536,216 and 797 µg/ml for hydroxyl radial-scavenging, superoxide radial- scavenging, DPPH radial scavenging and lipid peroxidation preventive activities, respectively. Total antioxidant capability used to be discovered to be 16.5 and 58.1 Gallic acid equivalent, AAE (µg/mg plant material) respectively. Phenol and flavonoid content was once found to be 150 GAE and thirteen Catechin Equivalent (µg/mg plant material) respectively. [60]

MARKEDTED FORMULATION

Masanumas² is the method of the second month of pregnancy. Each coated tablet contains Ashmantak (*Bauhinia racemosa*) bark extract, Black sesame (*Sesum indicum*) seed extract, Manjishtha (*Rubia cordifolia*) stem extract and Shatavari (*Asparagus recemosus*) root extract 60 mg. Each pack contains one hundred and twenty tabs. Dose: 2 tablets with lunch and 2 tablets with dinner. Indications: routine abortions, recurrent abortions, overestimated pregnancies, pregnancies caused by excessive risk[61]..

TOXICOLOGICAL EFFECTS

MEBR bark administered at the doses of one hundred and 200 mg/kg bw twice a week for thirteen weeks to Swiss albino mice did no longer set off any poisonous effects. Adverse impact used to be noted at the dose of 400 mg/kg bw. Its administration at the dose of 400 mg/kg bw considerably accelerated the stages of bilirubin, serum enzymes (SGOT, SGPT and ALP) and altered the level of urea, uric acid, creatinine, cholesterol, glucose and hematological parameters (hemoglobin content, complete count of RBC and WBC, differential be counted of leucocytes such as neutrophil, lymphocyte, monocyte, hematocrit, imply cellphone volume, suggest corpuscular hemoglobin,

imply corpuscular hemoglobin attention and platelet count) [62].

CONCLUSION

An natural medicinal drug is crude tablets bought from vegetable or plant which are the beginning of Herbal capsules utilized for the remedy of disease states, frequently of a chronic nature or to reap Or maintain a condition of elevated health. Herbal medicine uses are based on historic Medicinal practices. India is richly endowed with a huge variety of plant life having medicinal Value. The time period “herbal drugs” denotes flora or plant parts that have been converted into Phytopharmaceuticals through potential of easy approaches involving harvesting, drying, and Storage.[63,64] The plant processing encompasses drying, mechanical disruption, and solvent Extraction such as aqueous or organic solvent, e.g., ethanol, and will influence the remaining Quality of the natural product. Analytical approaches can be used to decide the lively Constituents that are existing in natural substances. [65] Herbal medicine makes use of are primarily based on Historical medicinal practices. Historical practices determine the way herbal drugs are Formulated and used. In some cases e.g. China, there are well-defined methods that are Well documented in pharmacopoeias courting lower back almost 2000 years and other monographs. The *Bauhinia* genuses are more than 2 hundred of modern medicine. *Bauhinia racemosa* Has a variety of regular medicinal uses, which on occasion require that it be blended with Other plant or natural ingredients. By considering the ethnomedicinal background and a number of Research articles on *Bauhinia racemosa* lam, it has been concluded that this plantincorporates Some necessary phytochemicals parts displaying a number of pharmacological activities such As Anti-inflammatory activity,

Chemoprotective Antitumor activity,
Antiulcer activity, Antidiabetic activity,
Antinociceptor activity,
Antihyperlipidemic activity, Antioxidant
Activity etc.

REFERENCES

1. Ushie, O.A, Nyong, B. E., Tabe, N.N, Jones ,B.B, Dogara, F. (2019). Bioactivity of active extract of leaf of bauhinia racemosa. *International journal of clinical chemistry and laboratory medicine*, 5(2), 19-24.
2. Panda, P., Das, D., Dash, P., & Ghosh, G. (2015). Therapeutic potential of Bauhinia racemosa-a mini review. *Int J Pharm Sci Rev Res*, 32(2), 169-179.
3. Kirtikar, K. R., & Basu, B. D. (1999). *Indian medicinal plants. vol 3*. International Book Distributors.
4. Egerton, A. (1955). Scientific Research in India. *Journal of the Royal Society of Arts*, 103(4962), 799-825.
5. Sharanabasappa, G. K., Santosh, M. K., Shaila, D., Seetharam, Y. N., & Sanjeevarao, I. (2007). Phytochemical studies on Bauhinia racemosa lam. Bauhinia purpurea Linn. And hardwickia binata roxb. *E-journal of Chemistry*, 4(1), 21-31.
6. Linn and Hardwickia binata Roxb. *E-J Chem* 4(1), 21-31.
7. Sinou, C., Forest, F., Lewis, G. P., & Bruneau, A. (2009). The genus Bauhinia sl (Leguminosae): a phylogeny based on the plastid trnL-trnF region. *Botany* 87: 947-960.
8. Bauhinia, L. Genus: Germplasm Resources Information Network. United States. Department of Agriculture.
9. Sasidharan, N. (2006). Illustrated manual on tree flora of Kerala supplemented with computer-aided identification. *Kerala Forest Research Institute, Peechi, Kerala. 698p.*
10. Davey, M. S., Atlee, C., Ashok, S. R. S., & Bharathi, M. F. (2011). Antianxiety effect of methanolic extract of B. racemosa (lamk) stems bark in mice. *Int J Pharma and Bio Sci*, 2, 217-224.
11. Gupta, M., Mazumder, U. K., Kumar, R. S., & Kumar, T. S. (2004). Antitumor activity and antioxidant role of Bauhinia racemosa against Ehrlich ascites carcinoma in Swiss albino mice. *Acta Pharmacologica Sinica*, 25, 1070-1076.
12. Nirmal, S. A., Dhasade, V. V., Laware, R. B., Rathi, R. A., & Kuchekar, B. S. (2011). Antihistaminic effect of Bauhinia racemosa leaves. *Journal of Young Pharmacists*, 3(2), 129-131.
13. Lenoir, R. (1924). Lamarck. *The Monist*, 187-235.
14. Bauhinia, L. Genus: Germplasm Resources Information Network. United States. Department of Agriculture.
15. Kirtikar, K. (1975). R. and Basu, BD, Indian Medicinal Plants II. *International Book Distributors, Dehradun*. 842-844.
16. El-Khatiba, A. S. (1995). Evaluation of some pharmacological properties of different extract of Bauhinia racemosa leaf and Bassia muricata whole plant. *Bull. Fac. Pharm. Cairo. Univ.*, 33, 59-65.
17. Ali, M. S., Azhar, I., Amtul, Z., Ahmad, V. U., & Usmanhane, K. (1999). Antimicrobial screening of some Caesalpiniaceae. *Fitoterapia*, 70(3), 299-304.
18. Akhtar, A. H., & Ahmad, K. U. (1995). Anti-ulcerogenic evaluation of the methanolic extracts of some indigenous medicinal plants of Pakistan in aspirin-ulcerated

- rats. *Journal of Ethnopharmacology*, 46(1), 1-6.
19. Dhar, M. L., Dhar, M. M., Dhawan, B. N., Mehrotra, B. N., & Ray, C. (1968). Screening of Indian plants for biological activity: Part I.
20. Dahikar, S. B., Bhutada, S. A., Tambekar, D. H., Vibhute, S. K., & Kasture, S. B. (2011). In-vitro antibacterial efficacy of solvent extracts of leaves of *Bauhinia racemosa* Lam.(Caesalpiniaceae) against enteric bacterial pathogens. *International Journal of Pharmaceutical Sciences and Drug Research*, 3(1), 32-34.
21. Chavan, R. T., & Kadam, A. S. (2012). Preliminary phytochemistry and antimicrobial activity of bark of *Bauhinia racemosa* Lamk. *Curr Bot*, 3(4), 31-33.
22. Manohar, P., Rajesham, V. V., Ramesh, M., Kiran, K. S., & Prasanna, K. J. (2011). Pharmacognostical, Phytochemical and antimicrobial activity of *Bauhinia racemosa* leaves. *Journal of pharmaceutical biology*, 1, 10-14.
23. Sharanabasappa, G. K., Santosh, M. K., Shaila, D., Seetharam, Y. N., & Sanjeevarao, I. (2007). Phytochemical studies on *Bauhinia racemosa* lam. *Bauhinia purpurea* Linn. And *hardwickia binata* roxb. *E-journal of Chemistry*, 4(1), 21-31.
24. Kumar, G., Karthik, L., & Rao, K. B. (2010). Phytochemical composition and in vitro antimicrobial activity of *Bauhinia racemosa* Lamk (Caesalpiniaceae). *International Journal of Pharmaceutical Sciences and Research*, 1(11), 51.
25. Sashidhara, K. V., Singh, S. P., Misra, S., Gupta, J., & Misra-Bhattacharya, S. (2012). Galactolipids from *Bauhinia racemosa* as a new class of antifilarial agents against human lymphatic filarial parasite, *Brugia malayi*. *European journal of medicinal chemistry*, 50, 230-235.
26. Upadhyay, B., Singh, K. P., & Kumar, A. (2011). Ethno-veterinary uses and informants consensus factor of medicinal plants of Sariska region, Rajasthan, India. *Journal of Ethnopharmacology*, 133(1), 14-25.
27. Singh, A. K., Raghubanshi, A. S., & Singh, J. S. (2002). Medical ethnobotany of the tribals of Sonaghati of Sonbhadra district, Uttar Pradesh, India. *Journal of ethnopharmacology*, 81(1), 31-41.
28. Singh, A. K., Raghubanshi, A. S., & Singh, J. S. (2002). Medical ethnobotany of the tribals of Sonaghati of Sonbhadra district, Uttar Pradesh, India. *Journal of ethnopharmacology*, 81(1), 31-41.
29. Anjaneyulu, A. S. R., Reddy, A. R., Reddy, D. S. K., Ward, R. S., Adhikesavalu, D., & Cameron, T. S. (1984). Pacharin: a new dibenzo (2, 3-6, 7) oxepin derivative from *Bauhinia racemosa* Lamk. *Tetrahedron*, 40(21), 4245-4252.
30. Mohan, V. R., & Janardhanan, K. (1995). Chemical analysis and nutritional assessment of lesser known pulses of the genus, *Mucuna*. *Food chemistry*, 52(3), 275-280.
31. Prabhakar, P., Gandhidasan, R., Raman, P. V., Krishnasamy, N. R., & Nanduri, S. (1994). De-o-methylracemosol: a tetracyclic 2, 2-dimethylchroman from the roots of *Bauhinia racemosa*. *Phytochemistry*, 36(3), 817-818.
32. Sae-Lao, P., Kittakoo, P., & Rajviroongit, S. (2006). Total synthesis of racemosol and de-O-methylracemosol, potent cyclooxygenase (COX) inhibitors

- and antimalarial agents. *Tetrahedron letters*, 47(3), 345-348.
33. Reddy, M. V., Reddy, M. K., Gunasekar, D., Caux, C., & Bodo, B. (2003). A flavanone and a dihydrodibenzoxepin from *Bauhinia variegata*. *Phytochemistry*, 64(4), 879-882.
 34. Aggarwal, B. B., Bhardwaj, A., Aggarwal, R. S., Seeram, N. P., Shishodia, S., & Takada, Y. (2004). Role of resveratrol in prevention and therapy of cancer: preclinical and clinical studies. *Anticancer research*, 24(5A), 2783-2840.
 35. Amoo, I. A., & Moza, L. (1999). Extraction and Physico-chemical properties of oils from *Bauhinia racemosa* seeds. *Rivista Italiana delle Sostanze Grasse*, 76(9), 399-400.
 36. Mohan, V. R., & Janardhanan, K. (1994). Chemical and nutritional evaluation of two germplasms of the tribal pulse, *Bauhinia racemosa* Lamk. *Plant Foods for Human Nutrition*, 46, 367-374.
 37. Garodia, P., Ichikawa, H., Malani, N., Sethi, G., & Aggarwal, B. B. (2007). From ancient medicine to modern medicine: ayurvedic concepts of health and their role in inflammation and cancer. *J Soc Integr Oncol*, 5(1), 25-37.
 38. Jain, A., Katewa, S. S., Chaudhary, B. L., & Galav, P. (2004). Folk herbal medicines used in birth control and sexual diseases by tribals of southern Rajasthan, India. *Journal of ethnopharmacology*, 90(1), 171-177.
 39. Jain, A., Katewa, S. S., Galav, P. K., & Sharma, P. (2005). Medicinal plant diversity of Sitamata wildlife sanctuary, Rajasthan, India. *Journal of ethnopharmacology*, 102(2), 143-157.
 40. Jyotsana, S., Gaur, R. D., & Painuli, R. M. (2011). Conservation status and diversity of some important plants in the Shiwalik Himalaya of Uttarakhand, India. *International Journal of Medicinal and Aromatic Plants*, 1(2), 75-82.
 41. Pirta, R. S., Gadgil, M., & Kharshikar, A. V. (1997). Management of the rhesus monkey *Macaca mulatta* and Hanuman langur *Presbytis entellus* in Himachal Pradesh, India. *Biological Conservation*, 79(1), 97-106.
 42. Jagtap, S. D., Deokule, S. S., & Bhosle, S. V. (2006). Some unique ethnomedicinal uses of plants used by the Korku tribe of Amravati district of Maharashtra, India. *Journal of ethnopharmacology*, 107(3), 463-469.
 43. Singh, A. K., Raghubanshi, A. S., & Singh, J. S. (2002). Medical ethnobotany of the tribals of Sonaghati of Sonbhadra district, Uttar Pradesh, India. *Journal of ethnopharmacology*, 81(1), 31-41.
 44. Sharma, J., Gairola, S., Gaur, R. D., & Painuli, R. M. (2012). The treatment of jaundice with medicinal plants in indigenous communities of the Sub-Himalayan region of Uttarakhand, India. *Journal of ethnopharmacology*, 143(1), 262-291.
 45. Patel, D. K., Laloo, D., Kumar, R., & Hemalatha, S. (2011). *Pedaliu murex* Linn.: an overview of its phytopharmacological aspects. *Asian Pacific journal of tropical medicine*, 4(9), 748-755.
 46. Gennaro, L. (2008). Extraction technology for medicinal and aromatic Plant. *United Nations Industrial Development Organization and the International Centre for Science and High Technology*, 6-8.

47. Finkel, R., Clark, M. A., & Cubeddu, L. X. (Eds.). (2009). *Pharmacology*. Lippincott Williams & Wilkins.
48. AC01271235, A. (Ed.). (1986). *Webster's medical desk dictionary*. Merriam-Webster.
49. Viswanathaswamy, A. H. M., Koti, B. C., Gore, A., Thippeswamy, A. H. M., & Kulkarni, R. V. (2011). Antihyperglycemic and antihyperlipidemic activity of *Plectranthus amboinicus* on normal and alloxan-induced diabetic rats. *Indian journal of pharmaceutical sciences*, 73(2), 139.
50. Borikar, V. I., Jangde, C. R., Rekhe, D. S., & Philip, P. (2009). Study of Analgesic activity of *Bauhinia racemosa* lam in Rats. *Veterinary World*, 2(4), 135.
51. Chandrashekar, K. S. (2010). Lack of analgesic activity of petroleum ether extract of *Bauhinia racemosa* Lam in rats and mice. *Journal of Pharmaceutical Negative Results*, 1(1), 22-23.
52. Desai, V. G. (1975). Aushadhi sangraha. *Shri Gajanan Book Depot, Dadar, Mumbai*, 188.
53. Borrelli, F., & Izzo, A. A. (2000). The plant kingdom as a source of anti-ulcer remedies. *Phytotherapy research: An international journal devoted to pharmacological and toxicological evaluation of natural product derivatives*, 14(8), 581-591.
54. Akhtar, A. H., & Ahmad, K. U. (1995). Anti-ulcerogenic evaluation of the methanolic extracts of some indigenous medicinal plants of Pakistan in aspirin-ulcerated rats. *Journal of Ethnopharmacology*, 46(1), 1-6.
55. Andersson, N., & Cockcroft, A. (2012). Choice-disability and HIV infection: a cross sectional study of HIV status in Botswana, Namibia and Swaziland. *AIDS and Behavior*, 16, 189-198.
56. Khandelwal, K. (2008). *Practical pharmacognosy*. Pragati Books Pvt. Ltd..
57. NCCLS, N. (1999). Performance standards for antimicrobial susceptibility testing; ninth informational supplement. *Approved standard M100-S9*. NCCLS, Wayne, Pa.
58. Panda, P., Das, D., Dash, P., & Ghosh, G. (2015). Therapeutic potential of *Bauhinia racemosa*-a mini review. *Int J Pharm Sci Rev Res*, 32(2), 169-179.
59. Azizur, R. M., Mehnaz, K., Hussain Arshad, A. M., & Mohd, K. (2015). Phytochemistry and pharmacology of traditionally Used tropical medicinal plant *Bauhinia racemosa* (Lam.). *The pharma research: An International Journal of Pharmacy Research*, 13(1), 26-41.
60. Davey, M. S., Atlee, C., Ashok, S. R. S., & Bharathi, M. F. (2011). Antianxiety effect of methanolic extract of *B. racemosa* (lamk) stems bark in mice. *Int J Pharma and Bio Sci*, 2, 217-224.
61. Gupta, M., Mazumder, U. K., Siva, K. T., Gomathi, P., & SAMBATH, K. R. (2004). Antioxidant and hepatoprotective effects of *bauhinia racemosa* against paracetamol and carbon tetrachloride induced liver damage in rats.
62. Azizur, R. M., Mehnaz, K., Hussain Arshad, A. M., & Mohd, K. (2015). Phytochemistry and pharmacology of traditionally Used tropical medicinal plant *Bauhinia racemosa* (Lam.). *The pharma research: An International Journal of Pharmacy Research*, 13(1), 26-41.
63. Kumar, R. S., Gupta, M., Mazumdar, U. K., Rajeshwar, Y., Kumar, T. S., Gomathi, P., & Roy, R. (2005).

- Effects of methanol extracts of *Caesalpinia bonducella* and *Bauhinia racemosa* on hematology and hepatorenal function in mice. *The Journal of toxicological sciences*, 30(4), 265-274.
64. World Health Organization. (1998). *Quality control methods for medicinal plant materials*. World Health Organization.
65. River Charles, Toxicology Studies
66. Verma, S., & Singh, S. P. (2008). Current and future status of herbal medicines. *Veterinary world*, 1(11), 347.