

Taxonomic notes on the genus *Itea* (Iteaceae)

Zhu-Qiu Song^{1,2}, Bu-Yun Zhang^{1,2}¹ Key Laboratory of Plant Resources Conservation and Sustainable Utilization, South China Botanical Garden, Chinese Academy of Sciences, Guangzhou, China² South China National Botanical Garden, Guangzhou, China

Corresponding author: Zhu-Qiu Song (songzhuqiu@scbg.ac.cn)

Abstract

The genus *Itea* (Iteaceae) is recognised as a genus with about 21 extant species of shrubs and trees. Within the genus, most species have oblong to elliptical leaves. The lanceolate and oblanceolate leaves are only found in three species, viz *Itea amoena* Chun, *Itea riparia* Collett & Hemsl. and *Itea tenuinervia* S. Y. Liu. The results of our examination of literature, specimens and living plants in the wild have shown that *Itea tenuinervia* is conspecific with *Itea riparia* and is here reduced to a synonym of the latter species. The morphological description, colour pictures, voucher specimens, distribution map of *Itea riparia* and its related *Itea amoena*, together with the morphological comparisons between the two species, are provided in this study.

Key words: China, lectotypification, synonym, taxonomy

Introduction

Itea L. (Iteaceae) was first described based on the species *I. virginica* L. from eastern North America (Linnaeus 1753) and later it was reported in Asia, for example, *I. macrophylla* Wall. from tropical Asia (Wallich 1824), *I. nutans* Royle from western Himalaya (Royle 1835) and *I. japonica* Oliv. from Japan (Oliver 1865). *Itea* was previously divided into two sections based on its deciduous character, i.e. sect. *Deciduae* Engl. (= sect. *Itea*) and sect. *Sempervirentes* Engl. (Engler 1891). The former section comprises only two species with deciduous leaves, *I. virginica* and *I. japonica*, and the latter includes the other species with evergreen leaves. Recently, *Choristylis rhamnoides* Harv. (Harvey 1842) from southeastern Africa was transferred to *Itea* as the third section, i.e. sect. *Choristylis* (Harv.) Jordaan (Kubitzki 2007; Jordaan 2012). Thus, *Itea* was considered as an unusual genus with an East Asian-eastern North American-southeastern African disjunction pattern (Kubitzki 2007; Tian et al. 2021). The genus comprises about 21 extant species of shrubs and trees, including one in Africa, one in North America and 19 species in temperate, subtropical and tropical Asia (Tian et al. 2021; POWO 2023).

The leaf shapes of the genus *Itea* are usually oblong to elliptical, while the lanceolate and oblanceolate leaves only appear in three species, including *I. riparia* Collett & Hemsl. (Collett and Hemsley 1890), *I. amoena* Chun (Chun 1934) and *I. tenuinervia* S.Y. Liu (Liu 2001; Tian et al. 2021). *Itea riparia* was

Academic editor: Bing Liu
Received: 26 December 2023
Accepted: 21 February 2024
Published: 12 March 2024**Citation:** Song Z-Q, Zhang B-Y (2024)
Taxonomic notes on the genus
Itea (Iteaceae). PhytoKeys 239:
59–72. <https://doi.org/10.3897/phytokeys.239.117851>**Copyright:** © Zhu-Qiu Song & Bu-Yun Zhang.
This is an open access article distributed under
terms of the Creative Commons Attribution
License (Attribution 4.0 International –
CC BY 4.0).

first published based on specimens collected on river-banks from Shan Hills, southern Shan States, Myanmar and it was characterised by narrow leaves, straight branches and erect racemes (Collett and Hemsley 1890). Gagnepain (1916) originally described *I. thorelii* based two gatherings from Vietnam and Laos and considered that it is very similar to *I. riparia*, but is different from the latter in the leaf width, the shape of anthers and the length of inflorescence, flower, stamen and ovary. Craib (1931) observed that some specimens of *I. riparia* from Thailand match *I. thorelii* in some respects. In China, *I. riparia* was recorded in Yunnan and *I. thorelii* was reported in Yunnan and Guangxi (Wu 1977; Fang and Qin 1985; Jin 1995), but the two species were merged in the Flora of China due to absence of obvious different characters (Jin and Ohba 2001). The earlier name *I. riparia* was adopted as its accepted name (Jin and Ohba 2001; Esser 2005; Chen and Liu 2015). The species grows on river-banks or near streams.

Chun (1934) described another *Itea* species with narrow leaves, *Itea amoena*, based on specimens from southern Guangxi, China and noted that it is well characterised by narrow, long-acuminate leaves and by long lanceolate calyx lobes. The species was previously identified as "*Itea riparia* var. *auminata* Hu var. nov.", based on *Ren-Chang Ching* 8059 (Chun 1934) and recognised as "*Itea lanceolata* Merr. n. sp." based on *Kuan-Kwang Tsoong* 1868 (Jin 1995). However, the latter two names have never been validly published. This species grows near streams, just like *I. riparia* (Fig. 5). Besides the habitat, *I. amoena* and *I. riparia* are also similar in the habit of evergreen shrubs, the lanceolate leaves and terminal inflorescences. Liu (2001) described *Itea tenuinervia* on the basis of a flowering collection found near streams in Shuolong Town, Daxin County, southwestern Guangxi, i.e. *Shou-Yang Liu* 2893 (Fig. 1A) and it was morphologically compared with *I. riparia* and *I. thorelii* in the protologue. However, our examination of literature and specimens showed that there are no obvious differences between *I. tenuinervia* and *I. riparia* (including *I. thorelii*). Thus, this study treats *Itea tenuinervia* as a new synonym of *I. riparia* and discusses the morphological differences between *I. riparia* and *I. amoena*.

Materials and methods

The specimens of the genus *Itea* kept in the Herbaria HITBC, IBSC and KUN have been examined by visiting these Herbaria and the images of *Itea* specimens deposited in the Herbaria E, GXMG, GXMI, IBK, IMDY, L, MW, N, NAS, NF, P, PE, SZ, TCD and US were also studied through the databases of specimens, such as Chinese Virtual Herbarium (CVH, <https://www.cvh.ac.cn>), Global Biodiversity Information Facility (GBIF, <https://www.gbif.org>), RBGE Herbarium Catalogue (<https://data.rbge.org.uk/search/herbarium/>), Naturalis BioPortal (<https://bioportal.naturalis.nl/>), Muséum National D'Histoire Naturelle (<https://science.mnhn.fr/all/search>) and Smithsonian National Museum of Natural History (<https://collections.nmnh.si.edu/search/botany>). Acronyms for the herbaria follow the Index Herbariorum (Thiers 2023). We also observed the living plants of the relevant species in the wild, including in the type localities of *I. tenuinervia* and *I. amoena*, as well as multiple localities of *I. riparia*.

Taxonomic treatment

1. *Itea riparia* Collett & Hemsl., J. Linn. Soc., Bot. 28: 57. 1890.

Figs 1–4

= *Itea thorelii* Gagnep., Notul. Syst. (Paris) 3: 222. 1916. Type. Laos. Luang-prabang, 1866–1868, *Thorel* 3474 (lectotype: P00709632, photo!, designated by Lecompte 1965: 40; isoelectotypes: P00709633 photo!, P00709634 photo!).

= *Itea tenuinervia* S. Y. Liu, Guihaia 21(1): 35. 2001. Type. China. Guangxi Province, Daxin County, Shuolong Town, by stream, 14 Jan 2000, *Shou-Yang Liu* 2893 (holotype: GXCM; isotype: IBK00190789!), syn. nov.

Type. Myanmar. Southern Shan States, Shan hills, on river-banks, 2000–4000 feet (ca. 700–1400 m) elev., *H. Collett s.n.* (holotype, K).

Description. Shrub, evergreen, usually 0.5–2 m tall, sometimes up to 7 m. Stipules subulate, 0.2–0.7 mm long, caducous. Leaves alternate, both surfaces glabrous, chartaceous, lanceolate or oblanceolate, 3.5–13 cm long, 1–5 cm wide, length/width ratio is 2.6–5.2 (mean = 3.6), acute or slightly acuminate at apex, cuneate at base, distantly shallowly subglandular crenate-serrate along upper margin; mid-vein raised on both surfaces, lateral veins in 4–7 pairs, arcuate ascending, lateral and reticulate veins visible and impressed above surface; petiole 0.4–1.2 cm long, raised above. Pseudoracemes, terminal or sometimes also axillary, usually solitary, sometimes with short branches at base, 4–18 cm long, rachis puberulous, usually 2- or 3-flowered per node; bracts at base of pedicel, subulate, 2–3 mm long, 0.3–0.5 mm wide, deciduous. Flowers white; pedicel erect, puberulous, 3–11 mm long; calyx shallowly cupular; calyx lobes 5, triangular, green, 1.5–2 mm long, 1.2–1.5 mm wide at base, puberulous; petals 5, white, triangular-lanceolate, acute at apex, widest at base, ca. 5 mm long, 1.5–1.9 mm wide at base, glabrous, erect-spreading at anthesis, becoming green and thickened at fruiting; stamens 5, alternating with the petals, shorter than petals, 3.3–3.6 mm long; filaments glabrous, ca. 0.8 mm wide; anthers oblong, ca. 0.6 mm long, 0.4 mm wide; floral disc annular, fleshy, slightly yellow; ovary semi-inferior, consisting of two carpels diverging in the middle and united at the top; stigma capitate. Capsule glabrous, 5–8 mm long, consisting of two obviously divergent carpels, with persistent sepals and petals, turning brown and dehiscing along the ventral suture at mature. Seed numerous, white at immature and brown at mature, 1–1.7 mm across; testa slightly granular.

Distribution. The species is distributed in China, Laos, Myanmar, Thailand and Vietnam (Fig. 6). It usually occurs near streams under forests, with elevation range from 200 to 1400 m.

Specimens examined. CHINA. Guangxi: Daxin, 15 May 2008, *Guangxi Exped. Inst. Bot.* 498 (IBK00278403), 17 Sept 2004, *Shou-Yang Liu* F000749 (GXMI014126), 400 m elev., 9 Jun 2023, 22.854261°N, 106.725634°E, *Zhu-Qiu Song et al.* JZ20230549 (IBSC); Jingxi, 457 m elev., 30 June 2013, *En-De Liu* 3996 (KUN1241727), 500 m elev., 10 May 2023, 22.991658°N, 106.674116°E, *Bu-Yun Zhang et al.* JZ20230371 (IBSC), 500 m elev., 9 Jun 2023, 23.009193°N, 106.659319°E, *Zhu-Qiu Song et al.* JZ20230553 (IBSC); Napo, 450 m elev., 18 May 1943, *Pu-Chin Tsoong & Ko-Zen Kuang* 356 (PE00865551, PE00865552), 15 Apr 1977, *Yuan Lin et al.* 3-5104 (GXMI014127, GXMI014128). Yunnan:

Cangyuan, 700 m elev., 29 May 1974, *Yan-Hui Li* 11737 (HITBC015745, HITBC015750, IBSC0273493, KUN0477791, KUN0540565, KUN0540567), 790 m elev., 15 May 1979, *Yan-Hui Li* 20907 (HITBC015744, HITBC015751, KUN0540559, KUN0540568); Jinghong, 540 m elev., 7 May 1955, *Kuo-Mei Feng* 20754 (IBSC0273491, KUN0477788, KUN0477789, PE00865549, SZ00179768), 30 Aug 2019, *Yun-Hong Tan* ML56 (XSBN002276); Malipo, 1100 m elev., 21 Jun 1940, *Chi-Wu Wang* 86321 (KUN0477792, PE00865553), 1000 m elev., 3 Feb 1940, *Chi-Wu Wang* 86599 (KUN0477790); Mengla, 17 Jan 1976, *Lai-Yun Xiao* 11269 (HITBC099861, IBSC0273492), 680 m elev., 18 Dec 1986, *Shao-Rong Guo* 592 (IMDY0008066), 850 m elev., 11 Feb 1960, *Yan-Hui Li* 2813 (HITBC015743, IBK00170997, IBSC0273495, KUN0477787), 560 m elev., 14 Apr 1961, *Yan-Hui Li* 3049 (HITBC015746), 11 May 1961, *Yan-Hui Li* 3222 (HITBC015747, IBSC0273494, KUN0477785), Jun 1967, *Yunnan First Section* 138 (PE00865550), 600 m elev., 2 Sept 2023, 21.989364°N, 101.368001°E, *Zhu-Qiu Song et al.* JZ20231022 (IBSC); Zhenkang, 12 Apr 1976, *Yan-Hui Li* 20116 (HITBC015748, HITBC015749, KUN0477786). **LAOS.** Champasak: Paksong, 29 Apr 2013, *Hang Sun et al.* *sun-hang*16056 (KUN1256568, KUN1256569); Luang Namtha: Nateuy, 1130 m elev., 21 Apr 2006, *En-De Liu* 1686 (KUN1204594); Luang Prabang: 1866–1868, *Thorel* 3474 (P00709632, P00709633, P00709634); Phongsal: 30 Apr 1993, *Guo-Da Tao et al.* 930162 (HITBC059236, HITBC059244); Vientiane: Mt. Phu Kao Kwai, 700–800 m elev., 31 Dec 1993, *N. Fukuoka & H. Koyama* L-65039 (L4134635); Without precise locality, *H. Bon* s.n. (P03179260). **MYANMAR.** Maymyo: 760 m elev., 21 Apr 1916, *A. Rodger* 510 (NAS00337458), 1066 m elev., 8 May 1912, *J. H. Lace* 5802 (E00397666), 1066 m elev., 4 May 1912, *J. H. Lace* s.n. (E00397667); Taunggyi: 1219 m elev., Apr 1939, *F. G. Dickason* 8273 (E01033109). **THAILAND.** Chaiyaphum: Phu Khieo, 600–700 m elev., 1972, *K. Larsen et al.* 31288 (L1861211), Ban Nam Phrom, 600 m elev., 24 May 1974, *R. Geesink et al.* 6886 (L1861215, P00392452); Chiang Mai: Thawatchai, Kasem, 700 m elev., 7 May 1991, *Guo-Da Tao et al.* 653 (KUN0540552), Me Ka Pak Bank, 1300 m elev., 21 Apr 1928, *H. B. Garret* 505 (L1861224, L1861225, P03179261), Mae Rim District, 600 m elev., 12 Jun 2009, *J. F. Maxwell* 09-162 (L2058714, P00911865), Doi Sutep, Mae Sa Falls, 10 Apr 1989, *J. F. Maxwell* 89-446 (L1861217), Doi Intson National Park, 650 m elev., 4 May 1997, *J. F. Maxwell* 97-443 (L4160466), Doi Inthanon, 1400 m elev., 3 Jan 1975, *R. Geesink et al.* 8048 (L1861214, P03371228), Doi Inthanon, 920–960 m elev., 21 Jul 1988, *S. Tsugaru* T-61686 (L4161221), 18 Aug 1970, *Voravut* 8 (L1861218); Kanchanaburi: Sangklaburi, Toong Yai Narssuan Wildlife Reserve, 200 m elev., 12 Jan 1994, *J. F. Maxwell* 94-10 (L4134592), near Khwae Noi River, Pompee Village, 250 m elev., 25 Mar 1968, *C. F. van Beusekom & C. Phengkhilai* 108 (L1861220), 800 m elev., 30 Apr 2004, *M. van de Bult* 796 (L4160559); Nakhon Ratchasima: Khao Yai, 800 m elev., 25 Mar 1968, *B. Hansen et al.* 11340 (L1861219, L1861221); Phrae: Mae Bhaem stream, 440 m elev., 5 Jan 1972, *C. F. van Beusekom et al.* 4661 (L1861223); Tak: Pha Charoen Waterfall National Park, 20 June 2005, *Pooma et al.* 5329 (L4160075), Lansang National Park, 600 m elev., 21 Jul 1973, *G. Murata et al.* 16617 (L1861212, P03179258), Lan Sang National Park, 250–400 m elev., 30 Aug 1967, *M. Tagawa et al.* 8558 (L1861213), Larn Sarn Nat. Park, 350 m elev., 29 May 1973, *R. Geesink et al.* 5532 (L1861222, P03179259); Without precise locality, Apr 1921, *A. F. G. Kerr* 5273 (TCD0016587), Mar 1924, *A. F. G. Kerr* 8763 (TCD0016588). **VIETNAM.** Dak Lak: Lak, Chu Yang Sin National Park, 900 m elev., 23 Mar 2013, *M.S. Nuraliev* 724 (MW0738997); Hoa



Figure 1. *Itea riparia* A isotype of *Itea tenuinervia* (Shou-Yang Liu 2893, IBK00190789, <https://www.cvh.ac.cn/spms/detail.php?id=c1cf1dd4>) B topotype of *Itea tenuinervia* (Shou-Yang Liu F000749, GXMI014126), showing two divergent carpels per fruit C specimen of *Itea riparia* from Myanmar (J. H. Lace s.n., E00397667, <https://data.rbge.org.uk/herb/E00397667>), showing two divergent carpels per fruit D lectotype of *Itea thorelii* (Thorel 3474, P00709632, <https://science.mnhn.fr/institution/mnhn/collection/p/item/p00709632>), showing two divergent carpels per fruit.




Figure 2. *Itea riparia* from Jingxi County, Guangxi, China **A** habitat **B–C** terminal inflorescences **D** leaves in adaxial and abaxial views **E** inflorescences **F** flowers with the different parts separated **G** infructescence, showing two divergent carpels per fruit and persistent sepals and petals **H** fruits and seeds. Voucher specimens: *Bu-Yun Zhang et al.* JZ20230371 (fl., IBSC), *Zhu-Qiu Song et al.* JZ20230553 (fr., IBSC). Scale bars: 2 cm (**B, D**); 5 mm (**F**); 1 cm (**H**).




Figure 3. *Itea riparia* from Daxin County, Guangxi, China **A** habitat **B** leaves in adaxial and abaxial views **C** fruiting branchlets **D** infructescence **E** fruits, showing two divergent carpels per fruit and becoming green and thickened petals and persistent sepals and petals. Voucher specimens: Zhu-Qiu Song et al. JZ20230549 (IBSC). Scale bar: 2 cm (**B**).

Binh: Yen Thuy, Cuc Phuong National Park, 12 Sept 2002, M. V. Xinh MVX 241 (L1861216), Barrage de Cho-bo (Rivière Noire), sur les roches calcaires atteintes par les hautes eaux, 18 Oct 1888, Balansa 3152 (P00709635).

Taxonomic notes. In the protologue, Liu (2001) considered that *I. tenuinervia* is distinguished from *I. riparia* in the height (1–2 m vs. 6 m), leaf shape (long elliptical, rarely oblanceolate vs. oblong, elliptical to lanceolate), pedicel length (10 mm vs. 3–5 mm) and petal length (6 mm vs. 4 mm) and it is distinguished from *I. thorelii* in the leaf shape (long elliptical, rarely oblanceolate vs.

oblanceolate), inflorescence type (racemes vs. racemose panicles), bract shape (triangular-lanceolate vs. linear-lanceolate), petal shape (narrowly lanceolate var. oblong-triangular) and pedicel length (10 mm vs. 4–6 mm). Subsequently, the species was accepted by Chen and Liu (2015). However, our examination of literature and specimens showed that there are no obvious differences between *I. tenuinervia* and *I. riparia* (including *I. thorelii*) in the characters above (see Table 1, Fig. 1). In the protologue, Liu (2001) did not describe the fruits of




Figure 4. *Itea riparia* from Mengla County, Yunnan, China **A** habitat **B** habit **C** leaves **D** fruiting branchlet **E** part of infructescence **F** fruits and seeds, showing two divergent carpels per fruit. Voucher specimens: Zhu-Qiu Song et al. JZ20231022 (IBSC). Scale bars: 2 cm (**C**); 5 mm (**F**).

I. tenuinervia, but he collected a fruiting gathering at the type locality in 2004, i.e. *Shou-Yang Liu F000749*, which shows two obviously divergent carpels per fruit (Fig. 1B). The fruits with two obviously divergent carpels are very special in the genus *Itea*, because most species bear two partly or completely fused carpels before dehiscence (Jordaan 2012) and only *Itea riparia* (including *I. thorelii*) was found to have two obviously divergent carpels (see Lecompte (1965: Pl. IV 8–10); Wu (1977: Pl. 26)). Thus, in the fruit character, *I. tenuinervia* is very similar to *I. riparia* as well. Furthermore, our observations of living plants in the wild, including in the type locality of *I. tenuinervia*, also showed that *I. tenuinervia* is consistent with *I. riparia* in the morphological characters and habit (Figs 2–4).

Table 1. Comparison between *Itea tenuinervia* and *Itea riparia*.

Characters	<i>Itea tenuinervia</i>	<i>Itea riparia</i>
Habit	shrubs 1–2 m	shrubs 1–6 m tall
Branchlet	green, striate, glabrous	yellow-green, striate, glabrous
Leaf arrangement	alternate, usually crowded at upper and lower parts of branchlets	alternate, usually crowded at apex of branchlets, subclustered
Leaf texture	thinly leathery	thinly leathery
Leaf hairiness	glabrous	both surfaces glabrous
Leaf shape	long elliptic, rarely oblanceolate, base cuneate, apex acuminate, margin incurved glandular dentate	elliptic to lanceolate or obovate-elliptic, base cuneate, apex acute or acuminate, margin curved cartilaginous glandular dentate
Leaf size	5–9 × 1.7–2.2 cm	5–13 × 1.5–5 cm
Lateral veins	4–5 pairs, confluent near margin, raised abaxially, slightly sunken adaxially	4–7 pairs, arcuate, confluent near margin, slightly raised abaxially, obscure adaxially
Tertiary veins	invisible above and visible below	reticulate, invisible above
Petiole	5 mm, glabrous	4–12 mm, glabrous
Stipules	ca. 1.5 mm long, caducous	0.2–0.7 mm long
Inflorescence	racemes terminal, 8–14 cm; rachis puberulous; flowers numerous, rather crowded, often 2–3-clustered	racemes terminal, 4–18 cm; rachis puberulous; flowers numerous, rather crowded, often 3-clustered
Bracts and bracteoles	bracts triangular-lanceolate, 3–5 mm; bracteoles 1–1.5 mm long	bracts ca. 3 by 0.5 mm, caducous, bracteoles to 1 mm long, similar
Pedicels	(6–) 10 mm	3–6 mm, puberulous
Calyx	shallowly cupular; lobes triangular, ca. 2 mm	shallowly cupular; lobes erect at anthesis, triangular-lanceolate, 1.5–4 mm
Petals	5, white, sometimes slightly pink, narrowly lanceolate, ca. 6 mm long, erect at anthesis	5, white, distinctly larger than the sepals, 4–5 by 1.5–2 mm, erect
Stamens	5, situated opposite the calyx lobes; filaments 5 mm long, slightly wider below the middle, glabrous; anthers ovoid, dorsifixed.	5, situated opposite the calyx lobes; filaments 2.5–3 mm long, glabrous; anthers ca. 1 mm long, dorsifixed, ovoid.
Disc	Ring-like, yellow	Ring-like
Ovary	semi-inferior, consisting of two carpels diverging above the base at the end of flowering, with a capitate stigma	superior or semi-inferior, glabrous, consisting of two carpels diverging in the middle and united at the top in a short style, surmounted by a capitate stigma
Fruit	capsules ovoid-conical, with very divergent carpels, except at the base; dehiscence by a slit corresponding to the ventral suture (based on a topotype, <i>Shou-Yang Liu F000749</i> , Fig. 1B)	capsules ovoid-conical, ca. 5 mm, glabrous, with very divergent carpels, except at the base; dehiscence by a slit corresponding to the ventral suture
Phenology	Fl. Jan., fr. Sept.	Fl. and fr. May–Feb.
Distribution	Guangxi, China	Myanmar, Thailand, Laos, Vietnam, China (Guangxi, Yunnan)
References	Liu (2001)	Lecompte (1965), Jin and Ohba (2001), Esser (2005)

2. *Itea amoena* Chun, Sunyatsenia 1(4): 238. 1934.

Fig. 5

Type. CHINA. Guangxi Province, Shangsi County, Shup-man-tai Shan [Shi Wan Da Shan], in shaded ravine, about 1200 ft. (ca. 400 m), 29 Jul 1933, *Ching-Lieh Tso* 23439 (lectotype, designated here: IBSC0004326, islectotypes: IBSC0004324, IBSC0004325, PE00864804).

Description. Shrub, evergreen, usually 0.2–0.5 m, sometimes up to 2 m tall, glabrous, except puberulent on inflorescences. Stipules subulate, ca. 2 mm long, caducous. Leaves alternate, both surfaces glabrous, thinly leathery, narrowly lanceolate, 6.5–13.9 cm long, 1.2–2.2 cm wide, length/width ratio is 4.1–9.5 (mean = 6.5), acuminate or gradually acute at apex, cuneate or sub-obtuse at base, distantly shallowly subglandular crenate-serrate along upper margin; mid-vein raised on both surfaces, lateral veins in 6–8 pairs, arcuate ascending, lateral and reticulate veins obviously visible and prominently elevated above surface; petiole 1–1.5 cm long, narrowly grooved above. Pseudoracemes, terminal or sometimes also axillary, solitary, 6–24 cm long, rachis puberulous, usually 2- or 3-flowered per node; bracts at base of pedicel, subulate, ca. 1–2 mm long, deciduous. Flowers white; pedicel erect, puberulous, 5–9 mm long; calyx shallowly cupular; calyx lobes 5, narrowly lanceolate, green, 2–2.5 mm long, 0.8–1 mm wide at base, puberulous; petals 5, white, reflexed at anthesis, deciduous at fruiting, triangular-lanceolate, acute at apex, widest at base, ca. 5 mm long, ca. 1.5 mm wide at base, glabrous; stamens 5, alternating with the petals, shorter than petals, ca. 4 mm long; filaments glabrous, ca. 0.3–0.4 mm wide; anthers oblong, ca. 0.5 mm long, 0.3 mm wide; floral disc annular, fleshy, slightly yellow; ovary semi-inferior, narrowly grooved, consisting of two united carpels; stigma capitate. Capsule suboblong-conical, 5–6 mm, glabrous, turning brown and dehiscing septicidally at mature, with persistent sepals. Seed numerous.

Distribution. The species is only found in Guangxi, southern China (Fig. 6). It usually occurs near streams under evergreen forests, with elevation range from 90 to 800 m.

Specimens examined. CHINA. Guangxi: Dongxing City, Malu, 107°58'56"E, 21°41'8"N, 90 m elev., 12 Oct 2018, *Dongxing Exped.* 450681181012069LY (GXMG0206492, IBK00418425), 21 Jun 1919, *Kuan-Kwang Tsoong* 1868 (PE00864807), 26 Sept 1973, *Chou-Fen Liang* 33611 (IBK00170636); Fangchenggang City, Pingwang, 15 Jul 2007, *Feng-Juan Mou* 127 (IBSC0712193), Banba, 15 Oct 1973, *Chou-Fen Liang* 33875 (IBK00170634), Nanguai, 11 Nov 1958, *Chao-Chien Chang* 13197 (IBK00170635, IBSC0272608), Naliang, 24–31 Jul 1936, *Wai-Tak Tsang* 26510 (IBSC0272609, P03179693), Naliang, 105 m elev., 13 Apr 1956, *Hepu Pl. Exped.* 2365 (IBSC0272605, PE00864802), Naliang, 107°47.563'E, 21°47.218'N, 140 m elev., 13 Jan 2016, *Pu Zou & Kai Xu* zp57 (IBSC0824452); Nalungou, 350 m elev., 21 Jul 1982, *Peng-Cheng Huang* 216 (NF2011745, NF2011746); Shangsi County, Shi Wan Da Shan, 325 m elev., 9 Nov 1958, *Chao-Chien Chang* 12233 (IBSC0272601), Wangle, 107°49'36"E, 21°53'24"N, 221 m elev., 1 Apr 2009, *Wei-Bin Xu et al.* liuyan0208 (KUN0906334, KUN0902457), Shi Wan Da Shan, 8 Jul 1937, *H. Y. Liang* 69500 (IBK00170632, IBK00170633, IBSC0272602, PE00864803), Shi Wan Da Shan, 11–30 Jul 1934, *Wai-Tak Tsang* 23804 (IBSC0272603), 15–24 Aug 1936, *Wai-Tak Tsang* 26675




Figure 5. *Itea amoena* from Shangsi County, Guangxi, China **A** habitat **B** leaves in adaxial and abaxial views **C** flowering branchlet **D** branchlet **E** part of inflorescences, showing usually 2 or 3 flowers per node **F** part of inflorescences, showing floral petals reflexed at anthesis **G** flowers with the different parts separated **H** flowers at the end of flowering, showing two fused carpels per fruit and deciduous petals **I** mature fruits. Voucher specimens: Zhu-Qiu Song et al. JZ20230531, JZ20230537 (IBSC). Scale bars: 2 cm (**B**); 4 cm (**C**); 5 mm (**G**).




Figure 6. Distribution map of *Itea riparia* (green circles) and *Itea amoena* (red squares).

(IBSC0272604, P03179692), Seh-feng Dar Shan [Shi Wan Da Shan], 2500 feet (ca. 800 m) elev., 21 Oct 1928, *Ren-Chang Ching* 8059 (IBSC0272600, N117066318, PE00864805, PE00864806, US03684023), 250 m elev., 14 Aug 1986, *Beijing Youth Exped.* 718 (PE01869678), 90 m elev., 23 Dec 1943, *S. H. Chun* 3304 (IBSC0272607), 170–300 m elev., 24 May 1944, *Shao-Hing Chun* 5209 (IBSC0272606), 140 m elev., 8 Jun 2023, 21.097495°N, 108.257582°E, *Zhu-Qiu Song et al.* JZ20230531 (IBSC), 320 m elev., 8 Jun 2023, 21.882874°N, 107.8346332°E, *Zhu-Qiu Song et al.* JZ20230537 (IBSC).

Taxonomic notes. *Itea amoena* can be readily distinguished from *I. riparia* in its long narrower leaves (length: width = 4.1–9.5 and ratio mean = 6.5 vs. 2.6–5.2 and 3.6), raised lateral and reticulate veins in the adaxial leaf surface (vs. impressed veins), floral petals reflexed at anthesis (vs. floral petals erect at anthesis), floral petals deciduous at fruiting (vs. becoming green and thickened at fruiting), lanceolate calyx lobes (vs. triangular calyx lobes) and fruits with two fused carpels (vs. two obviously divergent carpels). In distribution, *I. amoena* is restricted to southern Guangxi and it does not overlap with *I. riparia* (Fig. 6).

Acknowledgements

We thank the curators and staff of Herbaria E, GXMG, GXMI, HITBC, IBK, IBSC, IMDY, KUN, L, MW, N, NAS, NF, P, PE, SZ, TCD and US for hosting our visits or providing specimen images online.

Additional information

Conflict of interest

The authors have declared that no competing interests exist.

Ethical statement

No ethical statement was reported.

Funding

This research is supported by the Guangdong Provincial Forestry Bureau and Key-Area Research and Development Program of Guangdong Province (Grant no. 2022B1111230001).

Author contributions

Conceptualization: ZQS. Investigation: ZQS, BYZ. Methodology: ZQS, BYZ. Writing – original draft: ZQS.

Author ORCIDs

Zhu-Qiu Song  <https://orcid.org/0000-0002-7763-2431>

Bu-Yun Zhang  <https://orcid.org/0000-0003-1201-7638>

Data availability

All of the data that support the findings of this study are available in the main text.

References

- Chen ZD, Liu B (2015) Iteaceae. In: Zhu XY, Chen ZD, Liu B (Eds) Species catalogue of China volume 1 plants: Spermatophytes (IV), Angiosperms (Paeoniaceae–Polygalaceae). Science Press, Beijing, 11–13.
- Chun WY (1934) Contributions to the Flora of Kwangtung and South-Eastern China (1). Sunyatsenia 1(4): 209–317.
- Collett H, Hemsley WB (1890) On a collection of plants from Upper Burma and the Shan States. Journal of the Linnean Society. Botany 28: 1–150. <https://doi.org/10.1111/j.1095-8339.1890.tb01452.x>
- Craib WG (1931) Florae Siamensis Enumeratio: a list of the plants known from Siam with records of their occurrence, Vol. 1 Polypetalae. Bangkok Times Press Ltd., Bangkok, 809 pp.
- Engler A (1891) Saxifragaceae. In: Engler A, Prantl K (Eds) Die natürlichen Pflanzenfamilien Teil III. Abteilung 2a. Wilhelm Englimann, Leipzig, 41–93.
- Esser HJ (2005) Iteaceae. In: Santisuk T, Larsen K (Eds) Flora of Thailand, Vol. 9, Part 1. Forest Herbarium, Royal Forest Department, Bangkok, 36–38.
- Fang D, Qin N (1985) Some newly recorded plants from Guangxi. Guihaia 5(2): 91–100.
- Gagnepain F (1916) Quelques Saxifragacees Nouvelles. Notulae Systematicae 3: 222–223.
- Harvey WH (1842) Descriptions of several new genera of South African plants. The London Journal of Botany 1: 18–29.
- Jin SY (1995) *Itea*. In: Lu LT, Hwang SM (Eds) Flora Reipublicae Popularis Sinicae 35(1). Science Press, Beijing, 261–278.
- Jin SY, Ohba H (2001) *Itea*. In: Wu CY, Raven P (Eds) Flora of China. Vol. 8. Science Press, Beijing & Missouri Botanical Garden Press, St. Louis, 423–428
- Jordaan M (2012) Notes on African plants. Iteaceae. Taxonomic and nomenclatural notes on the genus *Choristylis*. Bothalia 42(2): 196–200. <https://doi.org/10.4102/abc.v42i2.18>
- Kubitzki K (2007) Iteaceae. In: Kubitzki K (Ed.) The families and genera of vascular plants, Vol. IX. Springer, 202–204. https://doi.org/10.1007/978-3-540-32219-1_25

- Lecompte O (1965) Saxifragaceae. In: Tardieu-Blot L (Ed.) Flore du Cambodge, du Laos et du Vietnam Vol. 4. Museum National D'histoire Naturelle, Paris, 3–55.
- Linnaeus C (1753) Species Plantarum. Impensis Laurentii Salvii, Holmiae, 1200 pp. <https://doi.org/10.5962/bhl.title.37656>
- Liu SY (2001) A new species of the genus *Itea* L. (Iteaceae). Guihaia 21(1): 35–36.
- Oliver D (1865) Notes upon a few of the Plants collected, chiefly near Nagasaki, Japan, and in the Islands of the Korean archipelago, in the years 1862–63, by Mr. Richard Oldham, late Botanical Collector attached to the Royal Gardens, Kew. Journal of the Linnean Society of London, Botany 9(35): 163–170. <https://doi.org/10.1111/j.1095-8339.1865.tb00021.x>
- POWO (2023) Plants of the World Online. Facilitated by the Royal Botanic Gardens, Kew. <http://www.plantsoftheworldonline.org/> [Retrieved 21 June 2023]
- Royle JF (1835) Illustrations of the Botany and other branches of the natural history of the Himalayan Mountains, Vol.1. London, Wm. H. Allen, 472 pp. <https://doi.org/10.5962/bhl.title.449>
- Thiers B (2023) Index Herbariorum: A global directory of public herbaria and associated staff. New York Botanical Garden's Virtual Herbarium. <http://sweetgum.nybg.org/ih/> [Accessed 26.12.2023]
- Tian YM, Huang J, Su T, Zhang ST (2021) Early Oligocene *Itea* (Iteaceae) leaves from East Asia and their biogeographic implications. Plant Diversity 43(2): 142–151. <https://doi.org/10.1016/j.pld.2020.09.006>
- Wallich N (1824) *Itea macrophylla*. In: Roxburgh W (Ed.) Flora Indica; or Descriptions of Indian Plants Vol. 2. Serampore: Printed at the Mission Press, 419–420. <https://doi.org/10.5962/bhl.title.589>
- Wu CY (1977) Flora Yunnanica, Vol. 1. Science Press, Beijing, 870 pp.