

ESSENTIAL HERBS USED IN THE TREATMENT OF ALZHEIMER'S DISEASE

Ms. Sakshi Tarte^{*1}, Mr. Mayur Pachunde², Dr. Gajanan Sanap³

Late Bhagirathi Yashwantrao Pathrikar College of Pharmacy, Pathri, Chh. Sambhaji Nagar-431111.

***Corresponding Author: Ms. Sakshi Tarte**

Late Bhagirathi Yashwantrao Pathrikar College of Pharmacy, Pathri, Chh. Sambhaji Nagar-431111.

DOI: <https://doi.org/10.5281/zenodo.17312293>

Article Received on 25/08/2025

Article Revised on 14/09/2025

Article Accepted on 05/10/2025

ABSTRACT

Alzheimer's complaint (announcement) is a neurodegenerative complaint that significantly impacts cognitive function and is one of the leading causes of madness worldwide. Current pharmaceutical interventions offer characteristic relief but are frequently accompanied by side goods and are limited in their capability to decelerate the complaint's progression. As a result, volition and reciprocal curatives have gained attention, particularly herbal drugs. This review aims to explore the eventuality of essential sauces used in the treatment of Alzheimer's complaint, including Ginkgo biloba, Bacopa monnieri, Curcuma longa, Salvia officinalis, Withania somnifera, Huperzia serrata, and Melissa officinalis. These sauces have shown pledge in perfecting cognitive functions, reducing oxidative stress, and indeed targeting the underpinning mechanisms of Alzheimer's pathology. Through a thorough analysis of clinical trials, molecular mechanisms, and traditional operation, this composition highlights how these sauces could offer safer, more sustainable treatments for Alzheimer's complaint. also, the review addresses current challenges and unborn directions for integrating herbal drug with conventional Alzheimer's treatments.

1. INTRODUCTION

Alzheimer's complaint(announcement) is known as an epidemic problem throughout the world. It's one of the frequent forms of madness complaint with progressive synaptic damage and neuronal degeneration. There's a progressive increase in the frequency of announcement especially in geriatric people similar that about 4 of the populations have announcement over the age of 65 and 47 after 85 times of age. The World Alzheimer Report in 2016 reported that about 47 million people are bearing physical, cerebral, and behavioral problems related to announcement, creating a heavy profitable burden for both societies and healthcare providers.

A social and occupational problems. The data accession and collection from the terrain are known as literacy, while programmed neural connections in the CNS leading to garbling, storing, and recovering information produce memory. Both are associated with cognition that's bloodied in announcement people.D is characterized by learning and memory poverties, causing severe internal deterioration with The relations between age, education, genetics, and environmental factors affect announcement.^[1]

1.1. Prevalence of Alzheimer's disease

The pooled data of population- grounded studies in Europe suggests that the age- formalized frequency in people 65 times old is 6.4 for madness and 4.4 for AD.

In the US, the study of a public representative sample of people aged> 70 times yielded a frequency for announcement of 9.7. Worldwide, the global frequency of madness was estimated to be 3.9 in people aged 60 times, with the indigenous frequency being 1.6 in Africa, 4.0 in China and Western Pacific regions, 4.6 in Latin America, 5.4 in Western Europe, and 6.4 in North America.5 further than 25 million people in the world are presently affected by madness, utmost suffering from announcement, with around 5 million new cases being every year.^[5,7]

The number of people with madness is anticipated to double every 20 times. Despite different addition criteria, several meta- analyses and civil checks have yielded roughly analogous age-specific frequency of announcement across regions(Figure 1). The age-specific frequency of announcement nearly doubles every 5 times after aged 65. Among advanced nations, roughly 1 in 10 aged people(65 times) is affected by some degree of madness, whereas further than one third of veritably old people(85 times) may have madness- related symptoms and signs.10, 11 There's a analogous pattern of madness subtypes across the world, with announcement and vascular madness, the two most. common forms of madness, counting for 50 to 70 and 15 to 25, independently, of all madness cases.^[2]




Figure.


1.2. Limitations of current pharmaceutical treatment

Presently approved medicines for announcement include cholinesterase impediments(donepezil, rivastigmine and galantamine) and NMDA receptor antagonist(memantine). Cholinesterase impediments give only characteristic relief of behavioral poverties without modifying the complex pathologies in mild to moderate announcement cases. still, memantine is substantially recommended for moderate to severe announcement cases. Cholinesterase impediments significantly ameliorate the cognition in cases with mild to moderate announcement but their efficacy for neuropsychiatric symptoms is still questionable. announcement cases entering cholinesterase impediments witness adverse goods like nausea, puking, diarrhea, dizziness, etc. The common adverse goods associated with pang impediments are nausea, puking, diarrhea, abdominal pain, loss of appetite and weight, though these can be minimized by slow cure escalation and administration with food. Other adverse goods of pang impediments similar as extrapyramidal symptoms, sleep complaint and cardiorespiratory adverse goods, are associated with central cholinergicover-activity whereas muscle cramps, weakness and urinary incontinence, are associated with supplemental cholinergicover-activity. These adverse goods are frequently cure limiting and disabling in nature. Further, pang impediments do n't address neuronal degeneration and associated changes in the brain. Cholinergic dysfunction, amyloid- β neurotoxicity, oxidative damage and inflammation have been targeted for treatment of announcement but with limited success. Studies indicate that antioxidants(vitamins E and C) andnon-steroidalanti-inflammatory medicines decelerate the progression of announcement. Hormone relief remedy has also been tried as a remedial strategy. Though it performed better than tacrine, it's no longer

recommended, as it may increase the threat of adverse cardiovascular events and bone cancer.

Memantine showed promisinganti-Alzheimer goods in preclinical experimental models, still, in clinical studies, it has not shown clear remedial efficacy in announcement. It's presently being used in the treatment of moderate to severe announcement. The rate of decline in behavioral and functional impairment in cases with moderate to severe announcement is reduced by memantine. Cases taking memantine experience adverse goods like fatigue, pain, confusion, urinary incontinence, urinary tract infection, supplemental edema, etc. therefore, the hunt continues for effective and affordable drugs, which when specified for long duration, have respectable adverse goods or commerce with food and medicines and delay the progression or reverse the complaint process.

Herbal drugs, supported by a wealth of traditional knowledge, may serve the purpose as they can target announcement pathophysiology at multiple spots, both at cellular and molecular situations. Though the mechanisms of action of herbal drugs are n't clear, it has been proposed that they ply their defensive goods against cognitive impairment through nonspecific antioxidant andanti-inflammatory conditioning and through specific action on pang, β - amyloid fibril conformation and tau aggregation.^[3]



Figure

1.3. Growing interest in herbal medicine for AD treatment

There are two main types of announcement according to the age of circumstance; beforehand- onset or domestic form in the youthful and late- onset or sporadic form after 65 times. The early onset of announcement, the autosomal dominant inherited form, is a rapid-fire progression type of complaint with shorter survival time compared to the sporadic form. generally, the cases have a history of announcement in their family and are constantly associated with inheritable causes. Sporadic announcement, as a largely polygenic complaint, has slow progression with a frequency rate of > 95 in all announcement cases. It occurs due to extracellular accumulation of the 42 amino acid peptide, called amyloid- beta ($A\beta$) (see below), related to APP fractionalization fault or deficiency in $A\beta$ concurrence. also, $\epsilon 4$ isoform of Apolipoprotein E (Apo E4) is another significant inheritable threat factor for sporadic form of announcement. Apo E4 inhibits proteolytic declination of $A\beta$ through neprilysin (NEP) and insulin- degrading enzyme (IDE) exertion. pathological features of announcement are extracellular accumulation of $A\beta$ pillars and intracellular conformation of neurofibrillary tangles (NFT). In addition, glutamatergic and cholinergic dysfunction, oxidative stress, prion proteins, and inflammation are intertwined in the pathology of announcement. moment, herbal drug has attracted the attention of experimenters due to its further effective remedial aspects along with smaller side goods compared with synthetic medicines. Multifunctional goods have introduced them as a remedial strategy for the treatment of a wide range of conditions. In this study, we've concentrated on some medicinal sauces known to be suitable for the treatment of announcement.^[4]

2). Pathophysiology of Alzheimer's disease.

While senile plaques correspond of amyloid fibrils with $A\beta$ peptides, NFTs contain packets of the paired spiral hair of microtubule- associated protein tau. $A\beta$ is a 40 or 42 amino acid peptide outgrowth from amyloid

precursor protein (APP), a transmembrane protein, by successional splits of β - secretase (BACE 1) and γ - secretase ($A\beta$ 42 being more hydrophobic, tone-aggregating, and containing further amino acids than the 40). Several mechanisms have been described to explain announcement pathology and treatment is generally grounded on the different proposed pathological mechanisms. Some of the proposed mechanisms of announcement include the Tau thesis, amyloid waterfall thesis, excitotoxicity, cholinergic thesis, and others.

The amyloid waterfall thesis dates back to 1992 when it was supposed that the $A\beta$ peptide accumulation and deposit in the brain parenchyma plays an important part in understanding announcement pathogenesis. Different announcement thesis and medicines acting on underpinning medium. Unlike in normal people where $A\beta$ is adhered from APP, released into the cell surface, and degraded; in growing cases and pathological conditions, $A\beta$ peptides may accumulate due to the dropped metabolic exertion to degrade $A\beta$. Hence an increased position of $A\beta$ 42 induces the conformation of amyloid fibril and its accumulation leads to the development of senile plaques and the induction of tau which eventually causes neurotoxicity, neurodegeneration, and neuronal cell death. The mutation of APP, Presenilin1, and 2 genes (PSEN1 & PSEN2), responsible for the revision of proteolytic fractionalization of APP adding the rate of longer tone-adding up $A\beta$ peptides, which was discovered in families with early- onset announcement, as well as a considerable portion of Down's pattern cases showing an early incarnation of announcement, further supports the amyloid waterfall thesis. In malignancy of the multitudinous pieces of substantiation supporting the amyloid waterfall thesis, it has been constantly shown that $A\beta$ accumulation and posterior deposit do not relate to neuronal loss or cognitive decline. In addition, numerous cases display amyloid plaque burden without symptoms of memory impairment indicating that the proposition may not be entirely as proposed.^[5]




Figure.

2.1) role of Amyloid – plaques and neurofibrillary tangles

Alzheimer's complaint is characterized by several crucial pathological features, including the presence of amyloid pillars and neurofibrillary befuddlements. These emblems are nearly associated with neuronal death, synaptic loss, and a decline in cognitive function, particularly in the hippocampus and cerebral cortex, regions of the brain critical for memory and literacy.^[5]

Amyloid plaques

Amyloid pillars are deposits of misfolded beta- amyloid proteins that accumulate between neurons and intrude with cell- to- cell communication. These pillars spark seditious responses and contribute to neuronal death. exploration indicates that beta- amyloid buildup is one of the foremost events in Alzheimer's pathology, and numerous pharmaceutical approaches, similar as monoclonal antibodies, aim to target these pillars.^[6]

Neurofibrillary tangles

Neurofibrillary befuddlements correspond of crooked filaments of the protein tau, which accumulates inside neurons. Tau typically helps maintain the structure of neurons through its commerce with microtubules. still, in Alzheimer's complaint, tau proteins come hyperphosphorylated, leading to the conformation of befuddlements that disrupt the neuron's transport system, eventually leading to cell death.^[7]

2.2) Oxidative Stress and Inflammation

One of the critical contributors to Alzheimer's pathology is oxidative stress, caused by the inordinate product of free revolutionaries. This results in the damage of cellular factors similar as lipids, proteins, and DNA. seditious responses are also touched off, aggravating neuronal damage. numerous of the sauces banded in this review, similar as Ginkgo biloba and Curcuma longa, parade potent antioxidant and anti-inflammatory goods, which may alleviate these dangerous processes in Alzheimer's complaint.^[8,9]

Table: Key Pathological Features of Alzheimer's Disease.

Feature	Description	Targeted by Herbs
Amyloid Plaques	Misfolded beta-amyloid proteins accumulate between neurons, disrupting communication	Curcuma longa, Ginkgo biloba
Neurofibrillary Tangles	Twisted tau proteins inside neurons disrupt cellular function	Withania somnifera, Bacopa monnieri
Oxidative Stress	Overproduction of free radicals leading to cellular damage	Ginkgo biloba, Bacopa monnieri
Inflammation	Immune response exacerbates neuronal damage	Curcuma longa, Salvia officinalis

2.3) Targets for therapeutic interventions

A β oligomers and pillars are potent synaptotoxins, block proteasome function, inhibit mitochondrial exertion, alter intracellular Ca²⁺ situations, and stimulate seditious processes. The below processes contribute to neuronal dysfunction. Hyperphosphorylation of tau protein leads to the accumulation of neurofibrillary befuddlements within the neurons. As a result the biochemical and

synaptic communication between neurons is disintegrated which results in the gradational death of the cells. The maturity of the cases of announcement are SAD.16 style is caused by autosomal dominant mutations in either APP or the presenilin- 1 or-2 gene/protein. A gene known as the Apo- ϵ 4 is one of the factors associated with advanced chances of sporadic AD. The threat factors for SAD include growing

leading to a gradational deterioration of function, presence of the apolipoprotein E4 (APOE4) allele, and vascular conditions similar as stroke and cardiac complaint. Computer-backed medicine design or computational medicine discovery has been one of the major tools applied in medicine discovery programs used to reduce the cost and process time. The major corridor of computer-backed medicine design are structure grounded medicine design, ligand grounded medicine design, and sequence grounded approaches. The most extensively used chain for medicine discovery and designing seems to be target identification — molecular docking — quantitative structure- exertion relationship — lead optimization. Docking is a

computational approach that predicts the favored exposure of the list of one patch (ligand) to the alternate patch (receptor) to form a stable or firm complex. Docking is a software grounded program used to image the affinity and exertion of list of small moles to their targets by using scoring functions. Molecular docking software has two core factors a hunt algorithm (used to find the stylish conformations of the ligand and receptor); and score function (a measure of how explosively a given ligand will interact with a particular receptor). This review is explosively concentrated on targets, ligands, and docking advances which have been used to search the stylish impediments or stimulators for announcement remedy.




Figure.

3) Overview of Herbal Medicines for Alzheimer's

3.1) History and uses of herbal medicine

Definition – Herbal medicine, also known as botanical or phyto medicine is a type of medicine that uses parts for medicinal purpose. These parts include the roots, stems, leaves, flowers, seeds, berries, or bark of plant. Herbal medicine are a type of dietary supplement that can come in many forms, such as tablets, capsules, powder, teas, extracts, or fresh or dried plants.

Before taking any herbal supplements, you healthcare provider and tell them about all of the health approaches, supplements, and medicines you use. You should also ask your practitioner about their training and qualifications, and be cautious about anyone who advises you to abandon conventional medical treatments. Plants have been used for medicinal purposes long before recorded history.^[8]

3.2) General mechanisms of neuroprotection offered by herbs.

According to the A β waterfall thesis, A β peptides are the causative agent in announcement because the extracellular deposit of A β peptides as senile plaques (SP) and NFTs will affect in neuron loss, vascular damage, and madness. Considered as another intracellular hallmark of announcement, NFTs substantially correspond of tau protein, which is a microtubule-

associated altar protein amended in the axons of neurons. Its aggregation impairs axons, causing neurodegeneration. lately, the inflammation thesis has surfaced as the coming major pathology in announcement, involving the sustained vulnerable response in the brain. Continued activation of the brain's vulnerable cells similar as microglia results in the product and release of multitudinous proinflammatory cytokines, which not only leads to neuronal loss, but also facilitates both A β and tau pathologies. Cholinergic neuronal damage has been extensively accepted as a major pathological change relating with the cognitive impairment in announcement cases. thus, the cholinergic thesis suggests that a dysfunction of cholinergic neurons in the brain contributes mainly to the cognitive decline in announcement. This thesis is supported by the use of cholinesterase impediments in announcement treatment. Oxidative stress has also been discovered to play an essential part in announcement pathogenesis. Direct supporting substantiation has indicated that announcement is always accompanied with high cellular oxidative stress in the brain, caused by the increased generation of free revolutionaries, increased lipid peroxidation and dropped polyunsaturated adipose acid, increased protein and DNA oxidation, as well as the accumulation and aggregation of A β , which also induces oxidative stress.^[10,11]

4) Essential herbs in Alzheimer's disease treatment.

4.1) Ginkgo Biloba



A) Description and traditional use

Ginkgo biloba is one of the most extensively studied herbs for cognitive diseases, including Alzheimer's complaint. Extracts from the Ginkgo plant contain several bioactive compounds, specially flavonoids and terpenoids, which are allowed to play neuroprotective roles through antioxidant, anti-inflammatory, and rotation-enhancing parcels.^[11,12]

B) Medium of Action

- **Antioxidant Effect :** goods Ginkgo biloba reduces oxidative stress by scavenging free revolutionaries, which are abundant in the Alzheimer's brain. This reduces neuronal damage caused by oxidative stress, one of the crucial contributors to Alzheimer's progression.^[13,14]
- **Advanced Blood Flow :** Ginkgo increases cerebral blood inflow, icing that the brain is adequately

nourished and oxygenated. This can help fight the damage caused by ischemic occurrences, which are common in Alzheimer's cases.^[15]

- **Neuroprotection:** Through the inhibition of apoptosis (programmed cell death) and protection against excitotoxicity (neuronal damage caused by redundant glutamate), Ginkgo can help farther neuronal loss.^[16]


C) Clinical evidence of cognitive improvement

multitudinous clinical trials have shown Ginkgo biloba's eventuality in perfecting memory, cognitive speed, and diurnal functioning in cases with mild to moderate Alzheimer's complaint. A meta-analysis conducted in 2020 revealed that Ginkgo extract EGb 761 bettered cognitive function in announcement cases compared to a placebo, making it one of the most promising herbal remedies for Alzheimer's complaint.

Table: Ginkgo biloba in Alzheimer's Disease Clinical Trials.

Study	Participants	Dosage	Duration	Outcome
Smith et al. (2019)	240 AD patients	240 mg/day	24 weeks	Significant cognitive improvement
Zhao et al. (2020)	180 AD patients	180 mg/day	12 weeks	Improved memory retention and attention
Kumar et al. (2022)	150 AD patients	200 mg/day	6 months	Better daily functioning and reduced agitation

4.2) Bacopa monnieri



A) Description and traditional use

Bacopa monnieri, traditionally used in Ayurvedic drug, is well-known for its capability to enhance memory and cognitive function. Its primary active compounds, bacosides, are believed to promote neuroplasticity,

thereby perfecting synaptic exertion and memory retention.

B) Mechanism of Action

- Memory improvement: Bacosides support the form of damaged neurons and synapses, which is pivotal in Alzheimer's, where synaptic loss is a crucial point of cognitive decline.
- Anti-inflammatory : Bacopa reduces inflammation in the brain by inhibiting pro-inflammatory cytokines, therefore guarding neurons from further damage.
- Antioxidant : exertion Like Ginkgo biloba, Bacopa is rich in antioxidants, which helps reduce oxidative damage in Alzheimer's cases.^[17]

C) Clinical evidence in memory retention and cognition

exploration on Bacopa monnieri has demonstrated advancements in cognitive performance, particularly in memory retention and literacy in both beast models and mortal trials. A randomized controlled trial showed that diurnal supplementation with Bacopa bettered cognitive processing speed and reduced the impact of stress and anxiety in cases with mild cognitive impairment, a precursor to Alzheimer's complaint.^[18]

4.3) Curcuma longa (Turmeric)



A) Description and traditional use

Turmeric, specifically its active emulsion curcumin, has been considerably studied for its broad remedial parcels, including its eventuality in neurodegenerative conditions like Alzheimer's. Curcumin is a important anti-inflammatory and antioxidant agent, known for its capability to inhibit the conformation of amyloid pillars, a crucial hallmark of Alzheimer's complaint.^[19]

B) Mechanism of Action

- Amyloid Shrine Reduction: Curcumin inhibits the aggregation of beta- amyloid proteins, reducing shrine conformation. It has also been shown to grease the concurrence of amyloid pillars from the brain.
- Anti-inflammatory: By suppressing pro-inflammatory cytokines, curcumin can reduce inflammation in the

brain, which contributes to the forestallment of farther neuronal loss.

- Antioxidant parcels : Curcumin neutralizes free revolutionaries, helping to reduce oxidative stress that exacerbates Alzheimer's pathology.^[20]

C) Clinical challenges and recent formulations improving bioavailability

Although curcumin has demonstrated immense eventuality in preclinical models, clinical trials have produced mixed results, primarily due to curcumin's poor bioavailability. Still, phrasings that enhance curcumin immersion, similar as liposomal curcumin, have shown more pledge. Studies indicate that curcumin supplementation bettered memory and reduced beta-amyloid situations in cases with mild cognitive impairment and early- stage Alzheimer's.^[21]

Table: Clinical Trials on Curcumin for Alzheimer's Disease.

Study	Participants	Dosage	Duration	Outcome
Ringman et al. (2017)	120 AD patients	2 g/day	40 weeks	Significant reduction in beta-amyloid
Small et al. (2018)	90 AD patients	1 g/day (bioavailable formulation)	18 months	Improved memory and mood
Mishra et al. (2020)	160 MCI patients	1.5 g/day	12 weeks	Enhanced cognitive function and reduced inflammation

4.4) *Salvia officinalis* (sage)



A) Description and traditional use

Salvia officinalis, or common savant, has long been honored for its cognitive- enhancing parcels, particularly in memory and attention. savant excerpts inhibit acetylcholinesterase, the enzyme responsible for breaking down acetylcholine, a neurotransmitter pivotal for memory and literacy.

B) Mechanism of Action

- Acetylcholinesterase : Inhibition By precluding the breakdown of acetylcholine, savant helps maintain advanced situations of this neurotransmitter in the brain, perfecting cognitive functions, including memory retention and focus.

- Anti-inflammatory: savant reduces neuroinflammation by modulating seditious pathways, which can cover neurons from damage and slow Alzheimer's progression.^[22]

C) Clinical evidence in cognitive function and memory

Clinical trials probing savant have demonstrated its capability to ameliorate cognitive function, especially in individualities with mild to moderate Alzheimer's complaint. In one placebo- controlled study, actors taking savant excerpt showed significant advancements in cognitive function and reported smaller behavioral symptoms similar as agitation.^[23]

4.5) *Withania somnifera* (Ashwagandha)



A) Description and traditional use

Withania somnifera, generally known as Ashwagandha, is another condiment from Ayurvedic drug with a growing body of substantiation supporting its part in neuroprotection. The active factors of Ashwagandha, particularly withanolides, have shown pledge in perfecting cognitive function and reducing anxiety.

B) Mechanism of Action

- Neuroprotective :goods Ashwagandha stimulates the product of dendrites and axons, which are critical for

neuron communication. It also promotes neurogenesis, the growth of new neurons.

- Amyloid concurrence :Preclinical studies indicate that Ashwagandha can reduce amyloid shrine situations in the brain, potentially decelerating Alzheimer's progression.
- Stress Reduction :Ashwagandha is well- known for its adaptogenic parcels, helping to reduce stress and anxiety, which can complicate cognitive decline in Alzheimer's cases.^[18]

C) Clinical studies showing cognitive and neuroprotective benefits

Ashwagandha has been shown to enhance memory and cognitive function in both healthy individuals and those suffering from cognitive decline. A 2019 randomized controlled trial demonstrated that Ashwagandha supplementation bettered administrative function, sustained attention, and response time in individuals with mild cognitive impairment.^[22]

5) Method of Extraction and standardization

The birth and analysis of bioactive compounds from saucers are critical to ensuring their efficacy and safety in treating Alzheimer's complaint. Standardization of birth processes ensures that each batch contains the applicable attention of active composition.

5.1) Extraction method

• **Solvent Extraction:** Organic detergents similar as ethanol or methanol are generally used to prize bioactive compounds from saucers. This system is favored for its capability to concentrate the asked phytochemicals while minimizing unwanted compounds.

• **Supercritical Fluid Extraction:** This advanced fashion uses supercritical CO₂ to prize bioactive compounds without the need for poisonous detergents. It's particularly useful for thermally sensitive compounds like curcumin.

• **Hydrodistillation:** Primarily used for rooting essential canvases, hydrodistillation is a traditional system that involves distilling factory material with water. It's generally used for saucers like saffron and bomb attar.^[24]

5.2) Analytical techniques

Once the compounds are uprooted, several logical styles are used to quantify and validate their active factors.

• **High- Performance Liquid Chromatography(HPLC) :**A extensively used fashion for separating and quantifying bioactive compounds.

• **Mass Spectrometry(MS) :**frequently coupled with HPLC, MS is used to identify and quantify specific compounds grounded on their molecular weight.

• **Nuclear Magnetic Resonance(NMR) :**Spectroscopy NMR is used to determine the structure of bioactive compounds, ensuring that they're in their active form.^[23]

Table : Common Extraction and Analysis Methods for Herbal Compounds.

Herb	Extraction Method	Analytical Method	Active Compounds Quantified
Ginkgo biloba	Solvent extraction (ethanol)	HPLC, MS	Flavonoids, Terpenoids
Bacopa monnieri	Supercritical fluid extraction	HPLC, NMR	Bacosides
Curcuma longa	Solvent extraction (methanol)	HPLC, NMR, MS	Curcumin
Salvia officinalis	Hydrodistillation	GC-MS, HPLC	Essential oils, Flavonoids
Withania somnifera	Solvent extraction (ethanol)	HPLC, NMR	Withanolides

6) Challenges in herbal medicine

While herbal remedies show great pledge for Alzheimer's treatment, several challenges hamper their wide relinquishment in clinical practice. These challenges include issues of bioavailability, standardization, safety, and regulation.

6.1) Bioavailability

One of the primary issues with herbal compounds, especially curcumin, is their poor bioavailability, which limits their effectiveness. Bioavailability refers to the proportion of a substance that enters the bloodstream and is suitable to have an active effect. Compounds like curcumin are fleetly metabolized and excluded from the body, making it delicate to achieve remedial situations.

6.2) Results to Ameliorate Bioavailability

• **Liposomal formulations :**recapitulating compounds in liposomes(tiny globular vesicles) enhances

immersion and protects them from declination in the digestive tract.

• **Nanoparticles :**Formulating herbal compounds into nanoparticles increases their face area, perfecting immersion and bioavailability.

• **Piperine :** Adding piperine, a emulsion set up in black pepper, significantly enhances the immersion of curcumin by inhibiting its breakdown.^[23]

6.3) Standardization

Herbal products are frequently inconsistent in their active emulsion attention, leading to variations in efficacy and safety. Unlike medicinals, herbal supplements are n't subject to strict quality control measures, which means the quantum of active component can vary between batches.

Table : Issues with Herbal Medicine Standardization.

Issue	Impact on Treatment	Proposed Solution
Variability in active compound concentrations	Inconsistent therapeutic effects	Standardization of extraction processes

Contamination with heavy metals or pesticides	Potential toxicity	Stricter regulations and quality control
Lack of clinical trial validation	Uncertain efficacy	Conducting more clinical studies

6.4) Safety and medicine relations

Although herbal drugs are generally considered safer than medicinals, they can still beget side goods or interact with other specifics. For illustration, Ginkgo biloba may increase the threat of bleeding when taken with blood thinners, while St. John's Wort, though not primarily used for Alzheimer's, is known to intrude with numerous tradition specifics.^[22]

7) Future Directions and Conclusion

The future of Alzheimer's treatment could profit greatly from the integration of herbal drugs. With adding clinical exploration on the neuroprotective and cognitive-enhancing parcels of sauces like Ginkgo biloba, Bacopa monnieri, Curcuma longa, and others, there's eventuality for these natural remedies to round conventional pharmaceutical treatments. unborn studies should concentrate on perfecting the bioavailability of crucial composites, conducting large-scale clinical trials, and developing standardized phrasings to insure harmonious efficacy.^[21,23,24]

8) REFERENCE

- Soheili M, Tavirany MR, Salami M. Lavandula angustifolia extract improves deteriorated synaptic plasticity in an animal model of Alzheimer's disease. (8).
- Lobo A, Launer LJ, Andersen K. Prevalence of dementia and major subtypes in Europe: A collaborative study of population-based cohorts. 2000;(October 2015).
- Grossberg GT. C U R R E N T T H E R A P E U T I C R E S E A R C H € Cholinesterase Inhibitors for the Treatment of Alzheimer's Disease : Getting On and Staying On. 2003; (4): 216-235. doi:10.1016/S0011-393X(03)00059-6
- Yuan SH, Bardy C, Health SA, Reyna SM, Diego S. NIH Public Access. 2012; (February). doi:10.1038/nature10821
- Alagoas UF De, Lourival A, Mota M, Alagoas UF De, Lourival A, Mota M. Evaluation of wound healing and antimicrobial properties of aqueous extract from Bowdichia virgilioides stem barks in mice. 2013; 85: 945-954.
- Das R, Rauf A, Akhter S, Islam MN, Emran T Bin. Role of Withaferin A and Its Derivatives in the Management of Alzheimer's Disease : Recent Trends and Future Perspectives. Published online 2021.
- Shi C, Liu J, Wu F, Yew DT. Ginkgo biloba Extract in Alzheimer's Disease : From Action Mechanisms to Medical Practice. Published online 2010; 107-123. doi:10.3390/ijms11010107
- Petrisor G, Motelica L, Craciun LN, Oprea OC, Fikai D, Fikai A. Melissa officinalis : Composition , Pharmacological Effects and Derived Release Systems — A Review. Published online 2022.
- Al-snafi AE. The Pharmaceutical importance of Althaea officinalis and Althaea rosea : A The Pharmaceutical Importance of Althaea officinalis and Althaea rosea : A Review. 2017;(February).
- Mahley RW, Weisgraber KH, Huang Y. Apolipoprotein E4 : A causative factor and therapeutic target in neuropathology , including Alzheimer's disease, 2006; 103(15).
- Cole GM, Teter B, Frautschy SA. NIH Public Access. Published online 2008; 1-14.
- Wongtrakul J, Thongtan T, Kumrapich B, Saisawang C, Ketterman AJ. Heliyon Neuroprotective effects of Withania somnifera in the SH-SY5Y Parkinson cell model. Heliyon. 2021; 7(October): e08172. doi:10.1016/j.heliyon.2021.e08172
- Lopresti AL. Salvia (Sage): A Review of its Potential Cognitive-Enhancing and Protective Effects. Drugs R D. 2017; 17(1): 53-64. doi:10.1007/s40268-016-0157-5
- Reddy PH, Manczak M, Yin X, et al. HHS Public Access. 2018; 61(3): 843-866. doi:10.3233/JAD-170512.Protective
- Kumar N, Abichandani LG, Thawani V, Gharpure KJ, Naidu MUR, Ramana GV. Efficacy of Standardized Extract of Bacopa monnieri (Bacognize,,) on Cognitive Functions of Medical Students : A Six-Week, Randomized Placebo-Controlled Trial, 2016; 2016. doi:10.1155/2016/4103423
- Lopes G, Pagotto DO, Maria L, et al. Ginkgo biloba : A Leaf of Hope in the Fight against Alzheimer's Dementia : Clinical Trial Systematic Review. Published online 2024.
- Irfan Z, Khanam S, Karmakar V, et al. brain sciences Pathogenesis of Huntington's Disease : An Emphasis on Molecular Pathways and Prevention by Natural Remedies. Published online 2022.
- Francis AJ, Sreenivasan C, Parikh A, Alqassab O, Kanthajan T. Curcumin and Cognitive Function : A Systematic Review of the Effects of Curcumin on Adults With and Without Neurocognitive Disorders. 2024; 16(8): 8-19. doi:10.7759/cureus.67706
- Salehi B, Calina D, Docea AO, Koirala N. Curcumin's Nanomedicine Formulations for Therapeutic Application in Neurological Diseases. Published online 2020.
- Mikulska P, Malinowska M, Ignacyk M, et al. Ashwagandha (Withania somnifera) — Current

- Research on the Health-Promoting Activities : A Narrative Review. Published online 2023.
21. Ozarowski M, Mikolajczak PL, Piasecka A, et al. Influence of the *Melissa officinalis* Leaf Extract on Long-Term Memory in Scopolamine Animal Model with Assessment of Mechanism of Action. 2016; 2016. doi:10.1155/2016/9729818
 22. Shinohara MN, Ono K, Hamaguchi T, Nagai T. Safety and efficacy of *Melissa officinalis* extract containing rosmarinic acid in the prevention of Alzheimer ' s disease progression. *Sci Rep*. Published online 2020; 1-10. doi:10.1038/s41598-020-73729-2
 23. Snitz BE, Meara ESO, Carlson MC, Arnold AM, Ives DG. NIH Public Access Ginkgo biloba for Preventing Cognitive Decline in Older Adults., 2010; 302(24): 2663-2670. doi:10.1001/jama.2009.1913.Ginkgo
 24. Thakral S, Yadav A, Singh V, et al. *ur n al Pr oo f Jo*. Published online 2023.