

Research Article

Effects of Environmental and Disturbance Factors on Plant Community Distribution in Tropical Moist Afromontane Forests, South-West Ethiopia

Semegnew Tadese ¹, Teshome Soromessa ¹, and Getaneh Gebeyehu ²

¹Addis Ababa University, College of Natural and Computational Sciences, Centre for Environmental Science, Addis Ababa, Ethiopia

²Injibara University, Department of Biology, Injibara, Ethiopia

Correspondence should be addressed to Semegnew Tadese; semetade@gmail.com

Received 1 November 2022; Revised 14 February 2023; Accepted 15 March 2023; Published 6 April 2023

Academic Editor: Anna Żróbek-Sokolnik

Copyright © 2023 Semegnew Tadese et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

This study was carried out to investigate the effects of environmental and disturbance factors on plant community distribution in the Majang Forest Biosphere Reserve (MFBR) in south-west Ethiopia. A systematic sample design was conducted to collect vegetation and environmental factors in four study sites. In a nested plot design, the vegetation data were collected from 140 main plots, i.e., 400 m² (trees), 25 m² subplots (shrubs, lianas, seedlings, and saplings), and 1 m² (herbs), respectively. The plant community classification was performed using agglomerative hierarchical cluster analysis (Ward's Linkage method) in R software (version 4.0.1). The distribution of plant communities along an environmental gradient was computed using canonical correspondence analysis (CCA). In this study, a total of 15 (9.5%) endemic plant species were recorded in MFBR. Four plant community types were identified, and these were *Celtis zenkeri*-*Blighia unijugata*, *Pouteria altissima*-*Lecaniodiscus fraxinifolius*, *Antiaris toxicaria*-*Celtis toka*, and *Dracaena afromontana*-*Cyathea manniana*. Environmental and disturbance factors, such as elevations, slopes, harvesting indexes, soil pH, silt, and herbaceous cover, were the most important for determining plant community distribution in the area. Elevation and slope were found to have a significant ($P < 0.05$) negative and positive relationship with species diversity and richness, respectively. Therefore, the finding of this study provides baseline information that could be necessary for making further conservation and management in MFBR.

1. Introduction

Plant ecologists have been successful in defining the variations in species diversity of communities along environmental gradients at various spatial scales [1]. Plant community responses are well related to climatic factors at regional and global scales [2, 3], whereas at local or plot scales, topographic and edaphic factors play an important role in controlling community structure and species distribution [4–8]. As a result, environmental factors are important not only in confirming plant community structure and variability of species distribution at a spatial scale but also for providing insight into the environmental requirements of the tree species

required for successful ecological restoration and biodiversity protection [7, 8].

Environmental factors have been a cornerstone in the evolution of ecological theory [9, 10]. A variety of factors influence the spatial and temporal patterns of vegetation change, including the physical environment, land use history, prior disturbance, and initial vegetation composition [11–14]. The most important data required to understand vegetation patterns in forest landscapes are relationships between species diversity and environmental factors [15–21]. As a result, the interaction between physical environmental variables such as elevations, slopes, and anthropogenic factors may influence species diversity and composition [19, 22–29]. Elevations, slopes, and aspects are determinants

of the spatial and temporal distribution of elements such as radiation, precipitation, and temperature that determine species composition [30]. Soil heterogeneity, such as soil texture, moisture content, electric conductivity (EC), and pH, creates niches with specific conditions, which in turn affect the distribution pattern of plants [31–33].

Ethiopia has greater geographic diversity with ten eco-systems, 18 major and 49 minor agroecological zones [34]. This diverse physiographic feature is responsible for the existence of a wide range of habitats and is conducive to the survival of various plant and animal species [35]. Accordingly, Ethiopia has a diverse set of plant, animal, microbial, and genetic resources [36]. Ethiopia is recognized as a major centre of diversity and endemism for a number of plant species in the Horn of Africa region [37, 38]. The flora of Ethiopia's is diverse, with an estimated 6000–7000 species of higher plants, including 647 (10.74%) endemic taxa [36].

The study of plant community types and factors (soil, topographic, and disturbance) that influence the distribution patterns of plant communities are essential inputs for forest conservation planning and management. For instance, previous studies have reported an association between plant community distributions and variations in environmental gradients in protected forests [25], fragmented forests [39], and community-managed forests of Ethiopia [40]. However, there are limited studies on the effects of environmental and anthropogenic disturbance factors governing plant community distribution patterns in southwestern forests [41, 42] in general and the Majang Forest Biosphere Reserve (MFBR) in particular. This problem leads to some research questions: (a) What are the main community types in the Majang Forest Biosphere (MFBR)? (b) How do environmental and disturbance factors influence plant community distribution patterns in Majang Forest Biosphere Reserves? Therefore, to answer these questions, we aimed to study to (1) identify the type of plant community in MFBR and (2) determine plant community distribution patterns in response to environmental and disturbance factors (elevation, slope, disturbance, and physical and chemical composition of soil) in MFBR.

2. Materials and Methods

2.1. Description of the Study Area. The study was conducted in the Majang Forest Biosphere Reserve (MFBR), situated in the Majang zone, *Gambella People National Regional State* of Ethiopia. It has unique biogeography and shares a boundary with Sale Nono woreda of Oromia regional state and Anderacha, Yeki, Sheka, and Guraferda woreda of South-West regional state (Figure 1). It covers a total area of 233,254 ha of forest and agricultural land and rural settlements and towns. MFBR is located between the latitudes of 07°08'00"N and 07°50'00"N and the longitudes of 34°50'00"E and 35°25'00"E, with elevations ranging from 562 m to 2444 m.

The climate in the area is generally hot and humid, which is marked on most rainfall maps of Ethiopia as the wettest part of the country. The average annual rainfall is 1774 mm, and the mean monthly minimum and maximum

temperatures range between 13.9 and 31.8°C at the Tinishu Meti weather station. In Ermichi, the average annual rainfall is 2053 mm and the mean monthly minimum and maximum temperatures range between 11.8 and 29.7°C. The maximum average monthly temperature is between 29.8°C and 31.8°C in February, while the minimum is between 11.9°C and 13.9°C in July for Ermichi and Tinishu Meti, respectively. The maximum rainfall is between April and October, and the minimum rainfall is from November to March in Tinishu Meti and Ermichi (Figure 2). This temperature and rainfall variation between two stations is due to variations in the elevation gradient (NMSA 2018).

Based on the vegetation classification of Ethiopia, the main vegetation types of the Majang Biosphere is montane evergreen forests. Besides, the vegetation of this area has different categories in terms of life forms such as high natural forests, bush lands, and grasslands [43]. Some environmental factors, such as elevations, slopes, soil pH, total nitrogen, and phosphorus, vary between each study site in MFBR (Table 1).

2.2. Vegetation and Soil Data Collection. A systematic sampling design was established to arrange quadrats and transects as well as to collect vegetation data [44]. The study area was stratified into four sites using a digital elevation model (DEM) in Arc GIS software. The study sites' polygon was digitized using Google Earth by elevation classes. These were site I (<1200 m.a.s.l), site II (1200–1500 m.a.s.l), site III (1500–1800 m.a.s.l), and site IV (>1800 m.a.s.l) (Table 2). A total of 140 quadrats were established for vegetation and forest soil data collection, i.e., site I (1–45), site II (46–85), site III (86–115), and site IV (116–140).

The quadrats' X-Y coordinates were generated in Arc GIS software and loaded to a global positioning system (GPS) receiver for tracking quadrats. Later, a measuring tape was used to lay out 20 × 20 m² (400 m²) quadrats in each site in the biosphere. The sampling intervals between the transect line and the quadrats were 2 km apart [43, 52]. The sizes of the quadrats were determined based on the growth forms of plants [44], i.e., 400 m² (tree), 25 m² (shrubs and lianas), and 1 m² (herbs), respectively, in a nested plot design (Figure 3).

The tree diameters (DBH ≥ 5 cm) were determined, and their number of stems was counted and recorded in the main plots (400 m²) (Figure 3). The diameter at breast height and the height of the individual's tree species were measured using a diameter tape, clinometer, and meter tape, respectively. When the branching of multitemmed individuals occurred below DBH, the DBH of each stem was measured separately and developed into a common diameter of all stems by summing up and then taking an average diameter. To determine the diversity and estimate the abundance of shrubs and lianas, subplots (area = 25 m² each) were established (Figure 3) [44].

Seedlings with a height of less than 1.30 m and a DBH of less than 2.5 centimeters, saplings with a height of more than 1.30 m and a DBH of 2.5 to 5 cm, and trees as plants with a DBH of more than 5 cm were selected [53]. Herbaceous




FIGURE 1: Location of the study sites (sites I–IV). Source: <https://earthexplorer.usgs.gov>.




FIGURE 2: Mean monthly temperature and rainfall recorded. (a) Tinishu Meti (1987–2017) and (b) Ermichi (1987–2017) (NMSA 2018).

TABLE 1: Topographic and soil characteristics of the study sites.

Study sites	Ele (m)	Slo	Area (ha)	SP
Site I	1042 ± 42	5.3 ± 0.4	22,826.1	40
Site II	1365 ± 24	5.4 ± 0.4	25,220.5	45
Site III	1635 ± 24	7.2 ± 0.5	14,053	30
Site IV	2011 ± 42	11.1 ± 1.2	11,783.5	25

Note. Ele = elevation, Slo = slope, and SP = sample plots.

TABLE 2: List of equations used for calculation of vegetation parameters.

Vegetation parameters	Equation	Equation no.	Reference
Density	$D = n/N$	1	[45]
Species richness	$D = n/\sqrt{N}$	2	[46]
H	$H' = \sum_{i=1}^s (Pi) (\ln pi)$	3	[44, 47]
Shannon evenness	$J = H'/H_{max} = H'/\ln s$	4	[44, 47]
SSI	$S_s = (2a/(2a + b + c))$	5	[44, 48]
Beta (β) diversity	$\beta = (b + c)/(2a + b + c)$	6	[49, 50]
Harvesting index	$HI = \sum_{i=1}^s (SD * EF/LD)$	7	[51]

“a” = number of tree species common to sites A and B; “b” = number of tree species recorded only in the site A; “c” = number of tree species recorded only in site B, “N”: total number of individuals of all the species; “n”: total number of individuals of the species; Shannon diversity index (H) = where pi is the proportion of individuals found in the i^{th} species; SSI = Sorensen similarity index, H = Shannon diversity index; SD = stump density; EF = expansion factor (3.4 for tropical forest); LD = live tree density.




FIGURE 3: Sampling design used for data collection [44]. Note: subplots (1 × 1 m) are measured for herbs, grasses, and soil samples; subplots (5 × 5 m) are measured for shrubs and lianas, and the main plot (20 × 20 m) is measured for trees, saplings, and seedlings.

cover abundance was assessed in five subquadrats (1 m²) within the main plot (400 m²).

The herbaceous cover abundance from each quadrat was changed into 9 cover abundance [54] scale classes: 1 = (<0.5%), 2 = (0.5–1.5%), 3 = (1.5–3%), 4 = (3–5%), 5 = (5–12.5%), 6 = (12.5–25%), 7 = (25–50%), 8 = 50–75%, and 9 ≥ 75% [55]. Canopy openness was measured by using a densiometer located at the centre of each plot. We classified canopy openness by the following classification scheme: very dense canopy >70%, moderately dense canopy 40–70%, open canopy 10–40%, scrub <10%, and nonforest 0% [56].

Plant species were identified in the field, and for those species that were difficult to identify in the field, voucher specimens were collected with the help of local guides, coded, pressed, and dried for subsequent identification and verification at the National Herbarium (ETH), Addis Ababa

University. Nomenclature of plants in this article follows those published in the flora of Ethiopia as well as flora of Ethiopia and Eritrea [57–62].

Soil samples were taken with a soil auger from the topsoil at a depth of 0–30 cm, which is recommended as the default sampling depth for soil [63]. The soil samples were taken from five different locations in each plot, four from the quadrat’s corners and one from the quadrat’s centre. A total of 220 soil samples, 140 from forestland, 40 from farmland, and 40 from grassland, were collected, composited separately, labelled, and transported to the laboratory. To determine soil bulk density, soil was collected on the centre of the quadrats using a stainless core sampler, then placed in plastic bags, and transported to the laboratory for dry weight determination. Fresh wet soil weights were measured in the field with a kitchen balance with 0.1 g precision. A composite sample of 200 g was taken from each quadrat to analyze its chemical composition [64] in the Water Works Design and Supervision Enterprise (WWDSE) laboratory in Addis Ababa.

2.3. Environmental and Disturbance Data Collection.

Environmental factors such as elevations, slopes, and aspects were measured and recorded for each of 140 quadrats using clinometers and Garmin GPS, respectively. The elevation was coded into four elevation ranges, namely, 1 ≤ 1200 m.a.s.l, 2 = 1200–1500 m.a.s.l, 3 = 1500–1800 m.a.s.l, and 4 ≥ 1800 m.a.s.l [65]. Aspect values were collected and arranged ($N=0$, $NE=1$, $E=2$, $SE=3$, $S=4$, $SW=3.25$, $W=2.5$, and $NW=1.25$) in each quadrat [52, 66]. The slope range was classified into three major slope classes [67]. As a result, the classes were as follows: (1) flat <10, (2) intermediate 10–20, and (3) steep >20.

Human disturbance (which includes harvesting/cutting trees for fuel wood, charcoal, timber, and house construction illegally) was computed as the harvesting index. The

harvesting index was computed by counting individual stumps that were illegally logged trees within the quadrats. It is calculated from the density of individual tree stumps by following equation (7) in Table 2 [51]. Stumps are a small portion of the trunk that remains after a tree with about 5 cm chopped down [68].

2.4. Data Analysis

2.4.1. Cluster Analysis. The vegetation data, which were originally estimated in the field as a percentage of cover abundance values, were converted into Domin cover scales prior to analysis [54]. The vegetation data were examined using an agglomerative hierarchical cluster (AHC) analysis in package cluster to classify vegetation into plant community types. The similarity ratio (SR) with Ward's group linkage methods (minimum variance clustering) was used for cluster analysis. The similarity ratio is one of the similarity indices that determine how similar or dissimilar vegetation samples, quadrats, or plots are. Ward's group linkage method evaluates cluster distances using the analysis of variance methodology [52]. The goal of this strategy is to eliminate differences in total abundance between sample units (quadrats). In addition, Ward's method minimizes the total within-group mean of squares or residual sum of squares [69].

The maximum number of clusters in a data set is a central issue in partitioning clustering, such as k-means clustering, which requires the user to specify the number of clusters k to be created. In this study, the optimum number of clusters was identified using the gap statistic method and determined by plotting the gap statistic against the number of clusters (k-means). The point, where the maximum gap statistic in the plot is, determines the proper number of clusters. In this study, the maximum gap statistic value in the plot is shown at k-means=4 (the maximum number of clusters or communities) [70].

Characteristic species were determined using synoptic table analysis. Plant community types were named after two distinctive species with high synoptic cover-abundance ratings [44]. Indicator species analysis was performed using the package labs in R software. The significance of the indicator value was tested using the Monte Carlo test ($P < 0.05$) [71]. The indicator values were calculated from the product of relative abundance (specificity) and relative frequency of species (fidelity) within a community [71]. In this study, the species with the highest significant indicator value is considered an indicator species of a community.

2.4.2. Community Diversity Analysis. Species richness, evenness, Shannon diversity, and beta diversity indices were computed using R software (version 4.0.1). Species richness is usually expressed as the number of species per sample unit and calculated using equation 2 in Table 2 [49]. Shannon's index was computed using abundance and evenness of the species present. The Shannon diversity index (H') and the

Shannon evenness index (J) were calculated using equations 3 and 4 in Table 2. The Sorensen similarity index was used to assess the degree of floristic similarity between plant communities. The Sorensen similarity index was calculated using equation 5 in Table 2. Beta diversity was computed by the change in diversity of species from one community to another and calculated using equation 6 in Table 2 [49]. Pearson correlation coefficients were calculated to see the relationship between diversity indices and environmental disturbance factors in MFBR.

2.4.3. Soil Laboratory Analysis. The soil samples were analyzed in the Water Works Design and Supervision Enterprise (WWDSE) laboratory in Addis Ababa, Ethiopia. The Bouyoucos hydrometer method was used to determine soil textures. Soil pH was determined using a pH meter a 1:2.5 soil to water suspension potentiometric method [72]. The micro-Kjeldahl [73] and Walkley and Black [74] methods were used to determine total nitrogen (N) and soil organic carbon, respectively. The Bray-I method was used to determine available phosphorus, and the absorbance of the Bray-I extract was measured in a spectrophotometer at 882 nm [75]. Based on C and N concentrations, the carbon-to-nitrogen ratio (C/N) was calculated. The dry weight of each soil sample (MS) was determined using oven-drying set to 105°C for 24 h to achieve a constant weight [76]. The volume of the core sampler (VC) was determined as $VC = \pi r^2 h$, where r is the radius and h is the height of the core sampler ($VC = 3.14 \times (2.5 \text{ cm})^2 \times 5 \text{ cm} = 98.125 \text{ cm}^3$) to calculate the bulk density for each sample.

2.4.4. Ordination. The canonical correspondence analysis (CCA) ordination method was used to find out plant community distribution along the environmental gradient [77]. The selection of CCA was based on the results of detrended correspondence analysis (DCA), which showed that the longest axis of DCA for a data set was greater than 3 ($=4.88$) (Table 3). This indicates the presence of higher β -diversity or heterogeneous vegetation data due to the unimodal relationship between species and environmental variables. Permutational multivariate analysis of variance (function adonis test) was used to choose environmental factors that were relatively more relevant in explaining the species data and significance before CCA ordination [78]. Pearson's correlation coefficient was calculated to evaluate the relationship between environmental factors and plant diversity.

3. Result

3.1. Plant Diversity. In this study, a total of 15 (9.5%) endemic plant species were recorded from Ethiopia. Of the total endemic species, 9 are herbaceous, 3 are shrubs, 2 are trees, and 1 is a liana in life form (Table 4). *Rinorea friisii* and *Crotalaria rosenii* were listed as near threatened among the 15 endemic species, while eight others were listed as least concerned (Table 4).

TABLE 3: Detrended correspondence analysis result of MFBR vegetation.

DCA axes	DCA1	DCA2	DCA3	DCA4
Eigen values	0.8356	0.2267	0.1917	0.1600
Decorana values	0.8599	0.2441	0.1845	0.1449
Axis lengths	4.8857	2.2693	2.4418	2.2889

3.2. *Plant Community Types*. Four plant community types were identified using agglomerative hierarchical cluster analysis of the entire vegetation data set of MFBR (Figure 4 and Table 5). The identified communities were *Celtis zenkeri-Blighia unijugata* community (C1), *Pouteria altissima-Lecaniodiscus fraxinifolius* community (C2), *Antiaris toxicaria-Celtis toka* community (C3), and *Dracaena afromontana-Cyathea manniana* community (C4).

C1: *Celtis zenkeri-Blighia unijugata* community

This community was found in the elevation range of 830–1896 m.a.s.l and was represented by 77 species. Of these, 67 species are commonly shared with other communities, while 12 are found only in this community type. Among the total species, four species were significant indicator species, with higher indicator values than those of total species (Table 6), namely, *Blighia unijugata*, *Pouteria alnifolia*, *Margaritaria discoidea*, and *Garcinia buchananii*. The most dominant shrub species included *Argemuellera macrophylla*, *Whitfieldia elongata*, *Dracaena fragrans*, *Vernonia amygdalina*, and *Phyllanthus reticulatus*. *Leptaspis zeylanica*, *Aspilia mossambicensis*, *Asplenium bugoiense*, *Asplenium anisophyllum*, and *Pollia condensata* were the dominant herb layers in the community. The common lianas of this community were *Embelia schimperi*, *Combretum paniculatum*, *Dregea schimperi*, and *Goupia glabra*.

C2: *Pouteria altissima-Lecaniodiscus fraxinifolius* community

This community was distributed in the elevation range of 800 to 2310 m.a.s.l. and was represented by 77 species. Of these, 71 species are commonly shared with other communities, while 6 are found only in this community type. Among the total species, four species were significant indicators with higher indicator values (Table 6), namely, *Pouteria altissima*, *Lecaniodiscus fraxinifolius*, *Ficus exasperata*, and *Ritchiea albersii*. Moreover, the dominant tree species layers were *Lannea welwitschii*, *Ritchiea albersii*, *Grewia molli*, *Ficus sur*, *Ficus mucoso*, *Ficus ovata*, *Ficus exasperata*, *Ficus umbellata*, *Allophylus macrobotrys*, and *Croton sylvaticus*. *Erythrococca trichogyne*, *Whitfieldia elongata*, *Dracaena fragrans*, *Rinorea friisii*, *Vernonia amygdalina*, *Acalypha orbata*, *Pittosporum viridiflorum*, *Phyllanthus reticulatus*, and *Oxyanthus speciosus* were prominent shrubs in this community. *Jasminum dichotomum*, *Combretum paniculatum*, *Phytolacca dodecandra*, *Dregea schimperi*, *Symphytum officinale*, *Pseuderanthemum tunicatum*, *Uncaria africana*, and *Saba comorensis* were common lianas. The

TABLE 4: Endemic plants, their habits, and IUCN status in the Majang Forest Biosphere Reserve.

Plant species	Family name	IUCN category	LF
<i>Acanthopale aethiogeranica</i>	Acanthaceae	NA	S
Ensermu			
<i>Bothriocline schimperi</i> Oliv. & Hiern ex Benth	Asteraceae	LC	H
<i>Clematis longicauda</i> Steud. ex A. Rich	Ranunculaceae	LC	L
<i>Crotalaria rosenii</i> (Pax) Milne-Redh. ex Polhill	Fabaceae	NT	H
<i>Dorsetnia soerensenii</i> Friis	Moraceae	NA	H
<i>Euphorbia dumalis</i> S. Carter	Euphorbiaceae	LC	H
<i>Impatiens rothii</i> Hook.f	Balsaminaceae	NA	H
<i>Millettia ferruginea</i> (Hochst.) Bak	Fabaceae	LC	T
<i>Pycnostachys abyssinica</i> Fresen	Lamiaceae	NA	H
<i>Rinorea friisii</i> M.G.Gilbert	Violaceae	NT	S
<i>Solanecio gigas</i> (Vatke) C. Jeffrey	Asteraceae	LC	S
<i>Solanum marginatum</i> L.f	Solanaceae	NA	H
<i>Vepris dainellii</i> (Pic. Serm.) Kokwaro	Rutaceae	LC	T
<i>Vernonia filigera</i> Oliv. & Hiern	Asteraceae	LC	H
<i>Vernonia leopoldi</i> (Sch. Bip. ex Walp.) Vake	Asteraceae	LC	H

Note. LF = life form, T = tree, S = shrub, H = herb, and L = liana; IUCN threat categories (LC = least concern, NA = not available, and NT = near threatened).

herb layer was dominated by *Asplenium bugoiense* and *Aspilia mossambicensis*.

C3: *Antiaris toxicaria-Celtis toka* community

This community was found in the elevation range of 850–2300 m.a.s.l and was represented by 105 species. Of these, 92 species are commonly shared with other communities, while 13 are found only in this community type. Among the total species, four species are significant indicator species with higher indicator values (Table 6), namely, *Celtis zenkeri*, *Morus mesozygia*, *Vernonia hochstetteri*, and *Celtis toka*. Moreover, the dominant species in the tree layer were *Diospyros abyssinica*, *Morus mesozygia*, *Combretum molle*, *Apodytes dimidiata*, *Olea capensis*, *Dombeya torrida*, *Buddleja polystachya*, and *Lepisanthes senegalensis*. The dominant species in the shrub layer were *Vernonia hochstetteri*, *Acanthopale pubescens*, *Erythrococca trichogyne*, *Plectranthus* sp, and *Capparis tomentosa*. *Marantochloa leucantha*, *Solanum nigrum*, *Achyranthes aspera*, and *Asplenium aethopicum* were the dominant species in the herb layer. The dominant lianas in this community were *Peponium vogelii*,

TABLE 5: Synoptic cover value of plants in MFBR for species $\geq 1\%$ in at least one community.

Scientific name	C1	C2	C3	C4
Cluster size	36	25	44	35
<i>Celtis zenkeri</i>	3.450	2.120	2.659	0.000
<i>Blighia unijugata</i>	2.806	1.040	1.841	0.000
<i>Apodytes dimidiata</i>	2.444	2.000	1.364	0.000
<i>Pouteria alnifolia</i>	2.222	1.240	1.000	0.000
<i>Trichilia prieuriana</i>	1.694	1.200	0.864	0.000
<i>Margaritaria discoidea</i>	1.694	0.640	0.818	0.000
<i>Argomuelleria macrophylla</i>	1.611	1.320	1.000	0.000
<i>Diospyros abyssinica</i>	1.556	0.800	1.955	0.000
<i>Lannea welwitschii</i>	1.500	1.600	1.409	0.000
<i>Combretum molle</i>	1.389	0.600	0.432	0.000
<i>Pouteria altissima</i>	2.472	3.400	2.864	0.000
<i>Lecaniodiscus fraxinifolius</i>	1.472	3.220	1.955	0.000
<i>Erythrococca trichogyne</i>	1.361	2.040	0.750	0.943
<i>Apodytes dimidiata</i>	2.444	2.000	1.364	0.000
<i>Ritchiea albersii</i>	0.944	1.840	1.227	0.000
<i>Ficus exasperata</i>	1.000	1.760	0.364	0.000
<i>Antiaris toxicaria</i>	1.778	1.640	3.532	0.000
<i>Celtis toka</i>	1.417	1.200	2.909	0.000
<i>Morus mesozygia</i>	0.556	0.480	1.477	0.000
<i>Croton macrostachyus</i>	0.472	0.800	1.045	1.171
<i>Deinbollia kilimandschrica</i>	0.250	1.040	1.045	0.000
<i>Grewia mollis</i>	1.222	1.400	1.159	0.000
<i>Ficus sur</i>	0.222	1.520	1.136	0.943
<i>Dracaena afromontana</i>	0.000	0.000	0.000	3.514
<i>Cyathea manniana</i>	0.000	0.000	0.000	3.414
<i>Schefflera abyssinica</i>	0.000	0.000	0.000	2.800
<i>Trilepisium madagascariense</i>	0.000	0.000	0.000	2.686
<i>Galiniera saxifraga</i>	0.000	0.000	0.000	2.600
<i>Schefflera myriantha</i>	0.000	0.000	0.000	1.914
<i>Vernonia bipotini</i>	0.000	0.000	0.000	1.886
<i>Allophylus abyssinicus</i>	0.000	0.000	0.000	1.857
<i>Pouteria adolfi-friedericii</i>	0.000	0.000	0.000	1.829
<i>Clausena anisata</i>	0.000	0.000	0.000	1.829

Note. C1 = community one, C2 = community two, C3 = community three, and C4 = community four.




FIGURE 4: Dendrogram of the cluster analysis results of species abundance found in 140 of MFBR. C1: 1, 17, 5, 21, 8, 24, 14, 30, 31, 38, 45, 61, 86, 90, 78, 98, 7, 23, 10, 26, 15, 50, 66, 43, 59, 33, 47, 63, 40, 56, 74, 94, 83, 103, 77, and 97; C2: 2, 18, 9, 25, 13, 29, 3, 19, 11, 27, 72, 92, 85, 105, 76, 96, 4, 20, 89, 12, 28, 75, 95, 82, and 102; C3: 6, 22, 35, 16, 54, 70, 36, 53, 69, 42, 58, 49, 65, 71, 91, 79, 99, 87, 80, 100, 84, 104, 88, 32, 39, 55, 46, 62, 48, 64, 52, 68, 73, 93, 81, 101, 34, 37, 44, 60, 41, 57, 51, and 67; C4: 106, 126, 110, 130, 119, 139, 112, 132, 116, 136, 123, 109, 129, 120, 140, 117, 137, 114, 134, 125, 107, 127, 118, 138, 111, 131, 121, 115, 135, 124, 108, 128, 113, 133, and 122.

TABLE 6: Higher indicator values with significance of species in the clusters.

Species	Community	R. Ab	R. Fr	Indval	P value
<i>Blighia unijugata</i>	C1	0.493	0.778	0.384	0.001
<i>Pouteria alnifolia</i>	C1	0.498	0.667	0.332	0.001
<i>Margaritaria discoidea</i>	C1	0.537	0.583	0.314	0.001
<i>Garcinia buchananii</i>	C1	1.000	0.306	0.306	0.001
<i>Pouteria altissima</i>	C2	0.283	0.722	0.389	0.001
<i>Lecaniodiscus fraxinifolius</i>	C2	0.225	0.278	0.343	0.001
<i>Ficus exasperata</i>	C2	0.320	0.361	0.293	0.001
<i>Ritchiea albersii</i>	C2	0.235	0.306	0.275	0.001
<i>Celtis zenkeri</i>	C3	0.841	0.365	0.307	0.005
<i>Morus mesozygia</i>	C3	0.523	0.588	0.307	0.001
<i>Vernonia hochstetteri</i>	C3	0.341	0.799	0.272	0.001
<i>Celtis toka</i>	C3	0.636	0.422	0.268	0.005
<i>Dracaena afromontana</i>	C4	1.000	1.000	1.000	0.001
<i>Cyathea manniana</i>	C4	1.000	1.000	1.000	0.001
<i>Vernonia bipotini</i>	C4	1.000	0.914	0.914	0.001
<i>Schefflera abyssinica</i>	C4	1.000	0.886	0.886	0.001

Note. R.Ab = relative abundance, R.Fr = relative frequency, Indval = indicator values in the class, and indicator species in the respective community types at $P < 0.05$.

Ampelocissus bombycina, *Sericostachys scandens*, *Tacazzea apiculata*, *Paullinia pinnata*, *Lagenaria siceraria*, and *Cayratia gracilis*.

C4: *Dracaena afromontana*-*Cyathea manniana* community

The community is distributed in the elevation range of 1820–2359 m.a.s.l and was represented by 70 species in 35 plots. Of these, 35 species are commonly shared with other communities, while 35 are found only in this community type. Among the total species, four species are significant indicator species with higher indicator values (Table 6), namely, *Dracaena afromontana*, *Cyathea manniana*, *Vernonia bipotini*, *Schefflera abyssinica*, and *Pouteria adolfi-friedericii*. The dominant tree species in the community were *Allophylus abyssinicus*, *Pouteria adolfi-friedericii*, *Triumfetta tomentosa*, *Pycnostachys eminii*, *Dracaena reflexa*, *Ekebergia capensis*, *Ilex mitis*, *Elaeodendron buchanaanii*, *Albizia schimperiana*, *Dracaena afromontana*, *Cyathea manniana*, *Schefflera abyssinica*, *Trilepisium madagascariense*, *Galiniara saxifrage*, and *Schefflera myriantha*. *Justicia schimperiana*, *Acanthopale aethio-germanica*, *Clausena anisata*, *Vernonia auriculifera*, *Maesa lanceolata*, *Maytenus gracilipes*, *Clematis longicauda*, and *Vernonia bipotini* were the dominant species in the shrub layer. The dominant species in herb layer were *Alchemilla fischeri*, *Bothriocline schimperii*, *Justicia* sp, *Asplenium friesiorum*, *Setaria megaphylla*, *Asplenium sandersonii*, and *Physalis angulata*. The dominant lianas in this community were *Clematis simensis*, *Culcasia falcifolia*, *Hippocratea pallens*, and *Jasminum abyssinicum*.

3.2.1. Species Richness, Evenness, and Diversity among Plant Community Types. The overall Shannon diversity index and evenness in MFBR were 3.64 and 0.94, respectively (Table 7). The highest Shannon diversity was shown in community 1 (3.8), while the least was exhibited in community 4 (3.5). The maximum Shannon evenness was indicated in community type four (0.9), followed by community types three, two, and one (each with 0.7 evenness) (Table 8). The highest species richness was exhibited in community type one (102 species), whereas the lowest plant species richness was revealed in community type four (70) (Table 7).

Similarity coefficients of all communities range from 0.61 to 0.91. The highest Sorensen similarity was exhibited between communities 1 and 2 (0.91) followed by communities 2 and 3 (0.90). The lowest Sorensen similarity was revealed between communities 1 and 4 (0.61) and communities 2 and 4 (0.63) of the total species in MFBR (Table 8).

Sorensen similarity coefficients and beta diversity were inversely related to each other. Therefore, communities with the lowest similarity coefficients had the highest beta diversity (0.45) in communities 1 and 4, whereas communities with the highest similarity coefficients had the lowest beta diversity (0.102) (Table 8).

3.2.2. Species Richness, Evenness, and Diversity along Environmental and Disturbance Factor. Species diversity varied from 2.33 to 3.64; the higher value was recorded at the higher elevation gradient (>1800 m.a.s.l), and the lower value was recorded at the lower elevation gradient (<1200 m.a.s.l). Species richness and Shannon diversity exhibited a significant positive correlation with elevations ($r = 0.302$, $p = 0.0002$) and ($r = 0.493$, $p = 0.00005$), respectively, while evenness showed a significant negative correlation with the elevation gradient ($r = -0.385$, $p = 0.00003$) (Table 9). Similarly, the slope showed a significant positive correlation with species richness ($r = 0.262$, $p = 0.00002$) and species diversity ($r = 0.373$, $p = 0.00001$), while the slope showed a significant negative correlation with species evenness ($r = -0.181$, $p = 0.0005$) (Table 9).

The highest species diversity was shown on the intermediate slope (10–20) gradient, while the lowest diversity recorded the lower slope (<10). The herbaceous cover showed a significant positive correlation with species richness and Shannon diversity ($r = 0.137$, $p = 0.01$) and ($r = 0.171$, $p = 0.004$), respectively. Species richness and Shannon diversity showed a significant negative correlation with the harvesting index ($r = -0.243$, $p = 0.003$) and ($r = -0.376$, $p = 0.00004$), respectively. Similarly, soil pH and silt showed a significant negative relationship with species richness ($r = -0.321$, $p = 0.0001$) and ($r = -0.504$, $p = 0.0004$) and Shannon diversity ($r = -0.302$, $p = 0.0002$) and ($r = -0.458$, $p = 0.00001$), respectively. However, soil pH and silt showed a significant positive relationship with species evenness ($r = 0.365$, $p = 0.0009$) and ($r = 0.353$, $p = 0.00001$), respectively (Table 9).

3.3. Relationship between Community Types with Environmental and Disturbance Factors. The length of the first axis of DCA indicates species heterogeneity in vegetation as

TABLE 7: Species richness, evenness, and diversity indices of plant community types.

Vegetation parameters	Community type			
	C1	C2	C3	C4
Species richness	102	77	93	70
H'	3.8	3.6	3.7	3.5
Shannon evenness	0.7	0.7	0.7	0.9
Basal area (m ²)/ha ⁻¹	58.55	48.82	64.64	76.3
Density (ha ⁻¹)	1532	1018	1478	1309
Seedling density (ha ⁻¹)	3349	3087	4431	3804
Sapling density (ha ⁻¹)	1629	1457	1789	1509

Note. H' = Shannon diversity index, C1 = community 1, C2 = community 2, C3 = community 3, and C4 = community 4.

TABLE 8: Pairwise comparison of Sorensen's similarity coefficient and beta diversity among four plant communities.

Communities	C1	C2	C3	C4
C1	1			
C2	0.91 (0.102)	1		
C3	0.87 (0.13)	0.90 (0.104)	1	
C4	0.61 (0.45)	0.63 (0.43)	0.64 (0.38)	1

Note. Values outside the bracket indicate Sorensen's coefficient, while those inside the bracket indicate the beta diversity index. C1 = community 1, C2 = community 2, C3 = community 3, and C4 = community 4.

TABLE 9: Pearson correlation coefficients between diversity indices and environmental and disturbance factors in MFBR.

Environmental disturbance site factors	Diversity indices		
	SR	H'	J
Hca	0.137*	0.171*	-0.030 ^{ns}
HI	-0.243***	-0.376***	-0.040 ^{ns}
Ele	0.302***	0.493***	-0.385***
Slope	0.262***	0.373***	-0.181*
Soil pH	-0.321***	-0.504***	0.365***
Silt	-0.302***	-0.458***	0.353***

Note. Hca = herbaceous cover, HI = harvesting index, Ele = elevation, SR = species richness, H' = species diversity index, and J = species evenness. Positive signs indicate a positive correlation, and negative signs indicate inverse relations. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, and ns = no significance.

a result of environmental variables (Table 3). The first axis of DCA was > 3 (4.88), which indicates a wider length of the gradient and the presence of higher β -diversity or heterogeneous vegetation data due to the unimodal relationship between species and environmental factors (Table 3). Therefore, CCA was selected to show the effect of environmental and disturbance factors on the patterns of plant community distribution.

Furthermore, significant environmental and disturbance factors ($P < 0.05$) were selected using permutational multivariate analysis of variance (function adonis test). Out of 14 environmental and disturbance factors, six factors were significant in explaining patterns of plant community distribution. More specifically, herbaceous cover ($r = 0.0409$, $p = 0.001$), harvesting index

($r = 0.0445$, $p = 0.001$), elevation ($r = 0.0906$, $p = 0.001$), slope ($r = 0.0143$, $p = 0.014$), pH ($r = 0.0127$, $p = 0.032$), and silt ($r = 0.0114$, $p = 0.0047$) exhibited a significant difference among environmental and disturbance factors (Table 10). Thus, significant environmental and disturbance factors influence the species distribution pattern in all community types.

CCA ordination was performed using data on cover abundance values of plant species in the plots and six statistically significant environmental disturbance factors. In the canonical correspondence analysis (CCA), the eigenvalues obtained in the first and second axes were 54.4 and 4.4, respectively. The correlation results of environmental disturbance factors and the CCA axis showed both a positive and negative correlation from the first to the last axis (Table 11). The elevation and slope showed a positive correlation ($r = 0.868$, $p < 0.05$) and ($r = 0.638$, $p < 0.05$), respectively, in the first axis. Among significant environmental disturbance factors, the elevation was an extremely important constraining factor followed by the slope and harvesting index (HI) on plant species distribution patterns in MFBR. On the other hand, pH and silt showed a strong negative correlation ($r = -0.907$, $p < 0.05$) and ($r = 0.792$, $p < 0.05$), respectively, in the first axis. CCA showed that the cumulative proportion and proportion explained in the first axis accounted 76.3%, respectively, which is explained as species and environmental disturbance factor variations (Table 11).

Ordination of the study plots of MFBR formed four communities based on the species composition. These four community types were separated by following the arrows of the environmental variables. Red, green, blue, and purple represent sample plots of community types 1, 2, 3, and 4, respectively (Figure 5). Community four generally occurs at a higher elevation, while species in communities one, two, and three are distributed at the middle and lower elevations. The distribution of species in communities four was influenced by elevation, slope, harvesting index, and herbaceous cover factors, whereas the distribution of species in communities one, two, and three was strongly influenced by soil pH and silt factors (Figure 5). The length and direction of the vectors in the plots and environment and disturbance biplot represent the degree of correlation between the plots and environmental disturbance factors. Communities' types one, two, and three are more closely related to one another, whereas community type four is different from the other three community types (Figure 5).

The correlation results between environmental and disturbance factors exhibited both positive and negative relationships in MFBR (Table 12). The elevation showed a significant positive correlation with slopes ($r = 0.44$), MC ($r = 0.58$), sand ($r = 0.33$), and TN ($r = 0.43$), while it showed a significant negative relationship with pH ($r = -0.76$), silt ($r = -0.90$), and clay ($r = -0.91$). Similarly, the slope showed a significant negative relationship with MC ($r = -0.38$), pH ($r = -0.46$), silt ($r = -0.42$), TN ($r = -0.47$), and clay ($r = -0.46$), while it showed a significant positive correlation with sand ($r = 0.45$). Soil pH showed a significant positive

TABLE 10: Result of the function Adonis test of significant environmental and disturbance factors in MFBR.

Variables	Df	Sums of squares	Mean squares	F model	R ²	Pr(>F)
Hca	1	1.8900	1.8898	6.9218	0.0409	0.001***
CaOp	1	0.1770	0.1766	0.6470	0.0038	0.767
HI	1	2.0610	2.0608	7.5481	0.0445	0.001***
Ele	1	4.1910	4.1912	15.3514	0.0906	0.001***
Slope	1	0.6600	0.6600	2.4173	0.0143	0.014*
Aspect	1	0.4640	0.4640	1.6994	0.0100	0.08
MC	1	0.2000	0.1998	0.7316	0.0043	0.703
pH	1	0.5860	0.5863	2.1473	0.0127	0.032*
Sand	1	0.2280	0.2284	0.8367	0.0049	0.559
Silt	1	0.5290	0.5293	1.9388	0.0114	0.047*
Clay	1	0.3360	0.3365	1.5142	0.0631	0.441
TN	1	0.4140	0.4135	1.5146	0.0089	0.137
OM	1	0.2900	0.2898	1.0614	0.0063	0.369
P	1	0.1750	0.1748	0.6403	0.0038	0.814
Residuals	126	34.4000	0.2730		0.7436	
Total	139	46.2640			1.0000	

Note. Hca = herbaceous cover, CaOp = canopy openness, HI = harvesting index, Ele = elevation, MC = moisture content of the soil, TN = total nitrogen, OM = organic matter, and P = available phosphorus.

TABLE 11: Scores of constraining variables and correlations between environmental and disturbance factors with CCA axes in MFBR.

Environmental variables	CCA1	CCA2	CCA3	CCA4
Hca	0.429	-0.128	0.160	-0.541
HI	0.522	0.028	0.207	0.335
Ele	0.868	0.066	0.100	-0.014
Slope	0.638	-0.238	0.135	-0.280
pH	-0.907	0.047	-0.029	-0.020
Silt	-0.792	0.060	-0.187	-0.243
Eigenvalue	0.544	0.044	0.040	0.033
Proportion explained	0.763	0.062	0.057	0.046
Cumulative proportion	0.763	0.825	0.882	0.928

Note. Hca = herbaceous cover, HI = harvesting index, Ele = elevation, and MFBR = Majang Forest Biosphere Reserve.

relationship with silt ($r=0.91$), TN ($r=0.93$), and clay ($r=0.91$), while it showed a significant negative relationship with sand ($r=-0.93$) (Table 12).

4. Discussion

4.1. Endemic Plant Species and Diversity. The Majang Forest Biosphere Reserve had shown high endemism as proposed by the World Conservation and Monitoring Centre (comprises high or 9.5% endemic species) (Table 4) [79]. Thus, endemism is higher than reported in other moist Afromontane forests of southwestern Ethiopia. For instance, Bonga forest recorded 13 endemic species (5.35%) [80] and Yayu forest with only 3 endemic species (1.36%) [81]. On the other hand, the number of endemic species recorded in this study is lower than in some dry Afromontane forests of Ethiopia. For example, Wof Washa forest has 29 species (12%) [82]; Chilimo forest has 18 endemic species (8.45%) [83]. The lower number of endemics in the moist Afromontane forests of the southwestern and dry Afromontane




FIGURE 5: Canonical correspondence analysis (CCA) ordination diagram of 140 plots and 6 significant environmental variables and plot number distribution in four community types.

forests of Ethiopia may be due to ecological and geographical separations [84]. Based on the plant species endemic to Ethiopia, *Rinorea friisii* and *Crotalaria rosenii* were listed as near threatened, while other eight were registered as least concerned (Table 4) [85]. This suggests that special attention needs to be given to the forest stand containing species which are threatened and least concerned.

The overall diversity investigation of MFBR showed the higher Shannon diversity index (3.64) and evenness (0.94). The value of species diversity and evenness in MFBR is greater than that of Gesha-Sayilem forest, Kibate forest, Gerba Dima forest, Wurg forest, and Agama forest, while it is lower than that of Belete forest (Table 13). This difference may be due to links with geographical locations, climatic conditions, and elevation factors.

4.2. Plant Community Types. In this study, a total of four plant communities were identified, which are rich in species composition. The major distinguishing features of the identified plant communities were the difference in dominant and indicator plant species. However, communities 1–3 share more species (>70 species) than community 4 (35 species), which could be directly related to environmental and disturbance factors that make the plant communities have their own distinct or characteristic species [42]. Almost all communities identified more than four indicator species with a significant value. Indicator plant species are plants that are easily monitored and predict the condition of the environment where they originated. Indicator species can also be a sign of a distinctive set of environmental qualities or characteristics found in a specific place [91].

Blighia unijugata and *Pouteria alnifolia* were indicator species in community 1. *Blighia unijugata* is the dominant and indicator species and is among the indicator species of moist Afromontane forests in the middle canopy [92, 93].

TABLE 12: Pearson's correlation coefficient matrix for environmental and disturbance factors ($N = 14$) in MFBR.

	Hca	CaOp	HI	Elevation	Slope	Aspect	MC	pH	Sand	Silt	TN	Clay (ns)	OM	P
Hca														
CaOp	0.27*													
HI	0.13 ^{ns}	0.05 ^{ns}												
Elevation	0.23 ^{ns}	0.10 ^{ns}	0.33***											
Slope	0.15 ^{ns}	-0.07 ^{ns}	-0.28*	0.44***										
Aspect	-0.08 ^{ns}	0.08 ^{ns}	-0.03 ^{ns}	-0.09 ^{ns}	-0.11 ^{ns}									
MC	0.18*	-0.05 ^{ns}	-0.14*	0.58***	-0.38***	-0.09 ^{ns}								
pH	-0.28**	-0.08 ^{ns}	-0.32***	-0.76***	-0.46***	0.09 ^{ns}	-0.57***							
Sand	-0.27**	0.08 ^{ns}	0.31***	0.33***	0.45***	-0.06 ^{ns}	-0.47***	-0.93***						
Silt	0.25**	-0.09 ^{ns}	-0.29*	-0.90***	-0.42***	0.06 ^{ns}	0.45***	0.91***	-0.98***					
TN	-0.24**	-0.11 ^{ns}	-0.27**	0.43***	-0.47***	0.02 ^{ns}	-0.57***	0.93***	-0.87***	0.85***				
Clay	-0.28***	-0.06 ^{ns}	-0.32***	-0.91***	-0.46***	0.06 ^{ns}	0.67***	0.91***	-0.98***	-0.52***	0.85***			
OM	0.03 ^{ns}	0.01 ^{ns}	-0.07 ^{ns}	0.32 ^{ns}	-0.11 ^{ns}	-0.03 ^{ns}	0.21*	-0.12 ^{ns}	-0.13 ^{ns}	0.12 ^{ns}	0.41 ^{ns}	0.16		
P	0.03 ^{ns}	0.10 ^{ns}	-0.03 ^{ns}	0.09 ^{ns}	0.01 ^{ns}	0.03 ^{ns}	0.08 ^{ns}	-0.09 ^{ns}	0.11 ^{ns}	0.10 ^{ns}	0.01 ^{ns}	0.01	0.18*	

Note. The magnitude indicates the degree of correlation, positive signs indicate positive correlations, and negative signs indicate inverse relations. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, and ns = no significance. N = number of variables, MFBR = Majiang Forest Biosphere Reserve, Hca = herbaceous cover, CaOp = canopy openness, HI = harvesting index, MC = moisture content of the soil, TN = total nitrogen, OM = organic matter, and P = available phosphorus.

TABLE 13: Comparisons of the Shannon diversity index and evenness of MFBR with other moist Afromontane forests in southwestern Ethiopia.

Forest area	Elevation (m)	H'	J	Sources
Gesha-Sayilem forest	1734–2803	3.56	0.84	[86]
MFBR	800–2400	3.64	0.94	Present study
Gerba Dima forest	1677–2240	3.45	0.93	[87]
Wurg forest	900–2500	3.38	0.90	[88]
Agama forest	1800–2370	3.25	0.78	[89]
Belete forest	1800–2300	3.79	0.95	[90]

Note. H' = Shannon diversity index, J = evenness, and MFBR = Majang Forest Biosphere Reserve.

Celtis zenkeri is the dominant species and significant indicator species in this community among five species. This community comprises an important fodder tree for honey production, which contributes to the means of livelihood for local communities in the area [94].

P. altissima was the dominant and significant indicator species in community 2 (Table 6) and is among the indicator species in the upper canopy of the moist Afromontane forest [92, 93], while *L. fraxinifolius* was also the dominant species and significant indicator species in the middle canopy.

A. toxicaria is the dominant species and the distinctive species of moist Afromontane forests in the middle canopy, whereas *Celtis toka* was the other dominant and significant indicator species in community 3 and is among the indicator species of moist Afromontane forests in the middle canopy [92, 93]. The community had the highest number of plant species compared to other communities. This could be due to their location away from human disturbances since they are located in difficult terrain such as sloppy and deep gorges [95, 96].

Dracaena afromontana and *Cyathea manniana* were the dominant and indicator species with a significant indicator value in this community, and they are among the distinctive species in the middle canopy of the moist Afromontane forest [92, 93]. The *D. afromontana*-*C. manniana* community type comprises different species as compared to other plant communities, which are found at higher elevations. The species composition difference in the community may be due to an environmental and anthropogenic factor [95, 96].

The Shannon diversity index was extended from 3.5 to 3.8 in four plant community types, which indicates the existence of high diversity in MFBR. The plant communities showed a minor variation in their species richness, diversity, and evenness. Somewhat, community types such as *B. unijugata*-*P. alnifolia* and *P. altissima*-*L. fraxinifolius* were higher in species richness, while community types *A. toxicaria*-*C. toka* and *D. afromontana*-*C. manniana* had the lowest species richness. This variation in species richness among communities could be due to the variations of topography, anthropogenic impact, climate, and edaphic factors [97, 98]. Each community has its own distribution area with a specific combination of environmental variables [27, 99–101].

The communities with the highest similarity coefficients had the least beta diversity, while communities with the least similarity coefficients had the highest beta diversity. The higher Sorensen similarity between communities 1 and 2 might be due to the narrow topographical distance between the two communities where most of the plots forming these communities may have relatively similar environmental factors [52, 102–105].

4.3. Species Richness, Evenness, and Diversity Indices along Environmental Gradients. We found high species richness and diversity at higher elevations than at lower elevations (Table 9). Higher species richness and diversity at higher elevations may be due to variations in climatic and edaphic factors [106]. The maximum species diversity and richness were found on intermediate sloping terrain, followed by on lower sloping terrain (Table 9). The slope has been identified as one of the topographic factors that influence species diversity and richness [95, 96, 107]. This result agrees with the findings of Wondie et al. [108], who stated that forests grow favourably on intermediate sloping terrain. The increase in species richness and diversity at intermediate sloping terrain may be due to soil moisture availability and edaphic factors (soil nutrients and low soil erosion) [109, 110].

4.4. Relationship between Community Types with Environmental and Disturbance Factors. Studies focusing on the relationship between plant communities with environmental and disturbance factors have become increasingly important in understanding the ecology of forest communities [111–115]. Plant species distribution patterns and plant community formation are highly influenced by microclimate, edaphic, and anthropogenic factors [41, 93, 116–119]. Plants form a community when a plant species repeatedly appears in a similar environment. Similarly, it is the classification of plant communities, which temporally and spatially determines the boundaries of a plant community [120–122]. Each community has its own distribution area with a specific combination or preference of environmental variables.

Relatively, the elevation was the most significant environmental factor influencing and describing the community type distribution [42, 81, 123–129]. Accordingly, the variation of plant communities was also closely related to other environmental, disturbance, and site factors, including herbaceous cover, harvesting index, slope, pH, and silt (Table 10).

Disturbance (illegal logging) affects the distribution of plant communities by hindering seedling establishment and regeneration in tropical forests [130–133]. It can also assist in the ground layer plant species growth by facilitating the availability of light [134, 135].

The correlation among environmental, disturbance, and site factors governs the pattern of species distribution in the plant communities of MFBR (Table 12). There was a positive correlation between the elevation and organic matter. The high amount of organic matter at higher elevations can be

attributed to a low rate of decomposition due to relatively low temperatures [113]. High organic matter content in soils of higher elevations was also reported in other Afromontane forests of Ethiopia [136].

5. Conclusion

This study was aimed to identify the type of plant community in MFBR and determine plant community distribution patterns in response to environmental disturbance factors (elevation, slope, disturbance, and physical and chemical composition of soil) in MFBR. MFBR is rich in endemic plant species. Our study identified four plant community types and indicated the elevation as a major environmental factor for determining the community type distribution. *Celtis zenkeri*-*Blighia unijugata*, *Pouteria altissima*-*Lecaniodiscus fraxinifolius*, *Antiaris toxicaria*-*Celtis toka*, and *Dracaena afromontana*-*Cyathea manniana* were the dominant species in the community. The highest species diversity and richness were found in community type one among the identified community types. The elevation and slope were the most significant environmental factors influencing species richness and Shannon diversity. Moreover, the slope and harvesting index were significantly positively correlated with species richness and Shannon diversity in all community types. Therefore, the findings of this study provide baseline information that should be necessary for further conservation and management practices in MFBR.

Data Availability

The data used to support the findings of this study are available from the corresponding author upon request.

Conflicts of Interest

The authors declare that there are no conflicts of interest.

Acknowledgments

The authors thank the Centre of Environmental Sciences, Addis Ababa University, for their contribution. The authors would like to thank Mr Demese Jemaneh for his contribution during data collection.

Supplementary Materials

Appendix 1: list of plant species and community types in four sites (key: C1, C2, C3, and C4 = community 1, 2, 3, and 4; Indval cls = indicator values in the class). Appendix 2: environmental factor data in MFBR. (*Supplementary Materials*)

References

- [1] S. Kumar, T. J. Stohlgren, and G. W. Chong, "Spatial heterogeneity influences native and nonnative plant species richness," *Ecology*, vol. 87, no. 12, pp. 3186–3199, 2006.
- [2] M. Toledo, M. Peña-Claros, F. Bongers et al., "Distribution patterns of tropical woody species in response to climatic and edaphic gradients," *Journal of Ecology*, vol. 100, no. 1, pp. 253–263, 2012.
- [3] S. Harrison, M. J. Spasojevic, and D. Li, "Climate and plant community diversity in space and time," *Proceedings of the National Academy of Sciences*, vol. 117, no. 9, pp. 4464–4470, 2020.
- [4] C. A. Baldeck, K. E. Harms, J. B. Yavitt et al., "Soil resources and topography shape local tree community structure in tropical forests," *Proceedings of the Royal Society B: Biological Sciences*, vol. 280, no. 1753, Article ID 20122532, 2013.
- [5] Z. H. Zhang, G. Hu, and J. Ni, "Effects of topographical and edaphic factors on the distribution of plant communities in two subtropical karst forests, southwestern China," *Journal of Mountain Science*, vol. 10, no. 1, pp. 95–104, 2013.
- [6] M. V. Odgaard, P. K. Bøcher, T. Dalgaard, J. E. Moeslund, and J.-C. Svenning, "Human-driven topographic effects on the distribution of forest in a flat, lowland agricultural region," *Journal of Geographical Sciences*, vol. 24, no. 1, pp. 76–92, 2014.
- [7] B. Fadrique and J. Homeier, "Elevation and topography influence community structure, biomass and host tree interactions of lianas in tropical montane forests of southern Ecuador," *Journal of Vegetation Science*, vol. 27, no. 5, pp. 958–968, 2016.
- [8] C. Zhang, X. Li, L. Chen, G. Xie, C. Liu, and S. Pei, "Effects of topographical and edaphic factors on tree community structure and diversity of subtropical mountain forests in the Lower Lancang River Basin," *Forests*, vol. 7, no. 12, p. 222, 2016.
- [9] J. A. Zavala-Hurtado, P. Valverde, A. Díaz-Solís, F. Vite, and E. Portilla, "Vegetation-environment relationships based on a life-forms classification in a semiarid region of tropical Mexico," *Revista de Biología Tropical*, vol. 44, pp. 581–590, 1996.
- [10] G. Öhlund, *Ecological and Evolutionary Effects of Predation in Environmental Gradients*, Umeå Universitet, Umeå, Sweden, 2012.
- [11] S. C. Goslee, W. A. Niering, D. L. Urban, and N. L. Christensen, "Influence of environment, history and vegetative interactions on stand dynamics in a Connecticut forest1," *Journal of the Torrey Botanical Society*, vol. 132, no. 3, pp. 471–482, 2005.
- [12] M. Bataineh, L. Kenefic, A. Weiskittel, R. Wagner, and J. Brissette, "Influence of partial harvesting and site factors on the abundance and composition of natural regeneration in the Acadian Forest of Maine, USA," *Forest Ecology and Management*, vol. 306, pp. 96–106, 2013.
- [13] K. Jöglste, H. Korjus, J. A. Stanturf et al., "Hemiboreal forest: natural disturbances and the importance of ecosystem legacies to management," *Ecosphere*, vol. 8, no. 2, 2017.
- [14] M. B. Hailemariam and T. D. Temam, "Pattern of plant community distribution along the elevational gradient and anthropogenic disturbance in gale forest, Ethiopia," *International Journal of Ecology*, vol. 2020, Article ID 6536374, 9 pages, 2020.
- [15] D. Sheil, R. K. Puri, I. Basuki et al., *Exploring Biological Diversity, Environment, and Local People's Perspectives in forest Landscapes: Methods for a Multidisciplinary Landscape Assessment*, CIFOR, Bogor, Indonesia, 2002.
- [16] C. T. Wang, R. J. Long, Q. J. Wang, L. M. Ding, and M. P. Wang, "Effects of altitude on plant-species diversity and productivity in an alpine meadow, Qinghai-Tibetan plateau," *Australian Journal of Botany*, vol. 55, no. 2, pp. 110–117, 2007.

- [17] M. Heydari and H. Pourbabei, "Vegetation-environmental relationships and ecological species groups of the Ilam oak forest landscape, Iran," *Caspian Journal of Environmental Sciences*, vol. 8, pp. 115–125, 2010.
- [18] E. Kouhgardi, G. Z. Amiri, K. Sagheb-Talebi, and M. Akbarzadeh, "The effects of soil characteristics and physiographic factors on the establishment and distribution of plant species in mountain forests (case study: Asalouyeh, South of Iran)," *International Journal of Biodiversity and Conservation*, vol. 3, pp. 456–466, 2011.
- [19] T. Jamil, *Models to Relate Species to Environment: A Hierarchical Statistical Approach*, Wageningen University and Research, Wageningen, Netherlands, 2012.
- [20] M. G. Turner, D. C. Donato, and W. H. Romme, "Consequences of spatial heterogeneity for ecosystem services in changing forest landscapes: priorities for future research," *Landscape Ecology*, vol. 28, no. 6, pp. 1081–1097, 2013.
- [21] T. Dorji, S. R. Moe, J. A. Klein, and Ø. Totland, "Plant species richness, evenness, and composition along environmental gradients in an alpine meadow grazing ecosystem in central Tibet, China," *Arctic Antarctic and Alpine Research*, vol. 46, no. 2, pp. 308–326, 2014.
- [22] G. C. Stevens, "The elevational gradient in altitudinal range: an extension of Rapoport's latitudinal rule to altitude," *The American Naturalist*, vol. 140, no. 6, pp. 893–911, 1992.
- [23] C. Rahbek, "The elevational gradient of species richness: a uniform pattern?" *Ecography*, vol. 18, no. 2, pp. 200–205, 1995.
- [24] G. Hadera, "A study on the ecology and management of the Dessa forest in the northeastern escarpment of Ethiopia," Unpublished M.Sc. thesis, Addis Ababa University, Addis Ababa, Ethiopia, 2000.
- [25] T. Soromessa, D. Teketay, and S. Demissew, "Ecological study of the vegetation in Gamo Gofa zone, southern Ethiopia," *Tropical Ecology*, vol. 45, pp. 209–222, 2004.
- [26] C. B. Lee, J. H. Chun, H. K. Song, and H. J. Cho, "Altitudinal patterns of plant species richness on the Baekdudaegan Mountains, South Korea: mid-domain effect, area, climate, and Rapoport's rule," *Ecological Research*, vol. 28, no. 1, pp. 67–79, 2013.
- [27] J. T. Zhang, B. Xu, and M. Li, "Vegetation patterns and species diversity along elevational and disturbance gradients in the Baihua Mountain Reserve, Beijing, China," *Mountain Research and Development*, vol. 33, no. 2, pp. 170–178, 2013.
- [28] M. Meragiaw, "Role of agroforestry and plantation on climate change mitigation and carbon sequestration in Ethiopia," *Journal of Tree Sciences*, vol. 36, pp. 1–15, 2017.
- [29] M. Girma and M. Maryo, "The diversity and composition of woody plants in chebera churcura national park (CCNP), southern Ethiopia," *Open Journal of Forestry*, vol. 8, no. 4, pp. 439–458, 2018.
- [30] A. R. Stage and C. Salas, "Interactions of elevation, aspect, and slope in models of forest species composition and productivity," *Forest Science*, vol. 53, pp. 486–492, 2007.
- [31] J. A. G. Vazquez and T. J. Givnish, "Altitudinal gradients in tropical forest composition, structure, and diversity in the Sierra de Manantlán," *Journal of Ecology*, vol. 86, no. 6, pp. 999–1020, 1998.
- [32] T. Medinski, *Soil Chemical and Physical Properties and Their Influence on the Plant Species Richness of Arid South-West Africa*, University of Stellenbosch, Stellenbosch, South Africa, 2007.
- [33] X. N. Zhang, X. D. Yang, Y. Li, X. M. He, G. H. Lv, and J. J. Yang, "Influence of edaphic factors on plant distribution and diversity in the arid area of Xinjiang, Northwest China," *Arid Land Research and Management*, vol. 32, no. 1, pp. 38–56, 2018.
- [34] M. Asefa, M. Cao, Y. He, E. Mekonnen, X. Song, and J. Yang, "Ethiopian vegetation types, climate and topography," *Plant Diversity*, vol. 42, no. 4, pp. 302–311, 2020.
- [35] M. A. Jochim, *Strategies for Survival: Cultural Behavior in an Ecological Context*, Elsevier, Amsterdam, Netherlands, 2013.
- [36] E. Kelbessa and S. Demissew, "Diversity of vascular plant taxa of the flora of Ethiopia and Eritrea," *Ethiopian Journal of Biological Sciences*, vol. 13, pp. 37–45, 2014.
- [37] I. Hedberg, I. Friis, and E. Person, *General Part and Index to Vol 1–7. Flora of Ethiopia and Eritrea Volume 8*, Addis Ababa University, Addis Ababa, Ethiopia, 2009.
- [38] A. Assefa, S. Demissew, and Z. Woldu, "Floristic composition, structure and regeneration status of M asha forest, south-west Ethiopia," *African Journal of Ecology*, vol. 52, no. 2, pp. 151–162, 2014.
- [39] R. Aerts, K. Van Overtveld, M. Haile, M. Hermy, J. Deckers, and B. Muys, "Species composition and diversity of small Afromontane forest fragments in northern Ethiopia," *Plant Ecology*, vol. 187, no. 1, pp. 127–142, 2006.
- [40] A. Abiyu, M. Lemenih, G. Gratzner, R. Aerts, D. Teketay, and G. Glatzel, "Status of native woody species diversity and soil characteristics in an enclosure and in plantations of Eucalyptus globulus and Cupressus lusitanica in Northern Ethiopia," *Mountain Research and Development*, vol. 31, pp. 144–152, 2011.
- [41] T. W. Gole, *Vegetation of the Yayu forest in SW Ethiopia: Impacts of Human Use and Implications for in Situ Conservation of Wild Coffea Arabica L. Populations*, Cuvillier Verlag, Göttingen, Germany, 2003.
- [42] F. S. Wakjira, *Biodiversity and Ecology of Afromontane Rainforests with Wild Coffea Arabica L. Populations in Ethiopia*, Cuvillier Verlag, Göttingen, Germany, 2006.
- [43] S. Tadese, T. Soromessa, T. Bekele, and G. Gebeyehu, "Woody species composition, vegetation structure, and regeneration status of Majang forest biosphere reserves in southwestern Ethiopia," *International Journal of Financial Research*, vol. 2021, Article ID 5534930, 22 pages, 2021.
- [44] M. Kent, *Vegetation Description and Data Analysis: A Practical Approach*, John Wiley & Sons, Hoboken, NJ, USA, 2012.
- [45] R. Shukla and P. Chandel, *Plant Ecology and Soil Science*, S. Chand & Company Limited, Ramnagar, New Delhi, 9th edition, 2000.
- [46] J. M. Scott, B. Csuti, J. D. Jacobi, and J. E. Estes, "Species richness," *BioScience*, vol. 37, no. 11, pp. 782–788, 1987.
- [47] A. E. Magurran, *Ecological Diversity and its Measurement*, Princeton university press, Princeton, NJ, USA, 1988.
- [48] A. Magurran, *Measuring Biological Diversity*, John Wiley & Sons, Hoboken, NJ, USA, 2004.
- [49] R. H. Whittaker, "Communities and ecosystems," 1970, <https://www.cabdirect.org/cabdirect/abstract/19740615709>.
- [50] D. Perlman and G. Adelson, *Biodiversity: Exploring Values and Priorities in Conservation*, John Wiley & Sons, Hoboken, NJ, USA, 2009.
- [51] T. Furukawa, C. Kayo, T. Kadoya et al., "Forest harvest index: accounting for global gross forest cover loss of wood production and an application of trade analysis," *Global Ecology and Conservation*, vol. 4, pp. 150–159, 2015.
- [52] A. Dibaba, *Ecology, carbon stock and formulation of allometric equation for selected tree and shrub species in Gerba*

- Dima forest, southwestern Ethiopia*, PhD. Thesis, Addis Ababa University, Addis Ababa, Ethiopia, 2018.
- [53] G. Stride, C. D. Thomas, S. Benedick et al., "Contrasting patterns of local richness of seedlings, saplings, and trees may have implications for regeneration in rainforest remnants," *Biotropica*, vol. 50, no. 6, pp. 889–897, 2018.
- [54] E. Maabel, "Transformation of cover-abundance values in phytosociology and its effects on community similarity," *Vegetatio*, vol. 39, no. 2, pp. 97–114, 1979.
- [55] E. van der Maarel and J. Franklin, *Vegetation Ecology*, John Wiley & Sons, Hoboken, NJ, USA, 2012.
- [56] E. Arets, P. Van der Meer, C. Verwer et al., *Global wood Production: Assessment of Industrial Round Wood Supply from Forest Management Systems in Different Global Regions*, Alterra, Wageningen, Netherlands, 2011.
- [57] I. Hedberg, O. Hedberg, and S. Edwards, *Flora of Ethiopia and Eritrea. Vol. 3, Pittosporaceae to Araliaceae*, National Herbarium, Addis Ababa University, Addis Ababa, Ethiopia, 1989.
- [58] S. Edwards and M. Tadess, "Flora of Ethiopia and Eritrea," *Canellaceae to Euphorbiaceae, e National Herbarium*, Vol. 2, Addis Ababa University, Addis Abeba, Ethiopia, 1995.
- [59] I. Hedberg, S. Edwards, and S. Phillips, *Flora of Ethiopia and Eritrea* Vol. 7, AAU, Jorhat, Assam, 1995.
- [60] I. Hedberg, O. Hedberg, S. Edwards, and S. Demissew, "Flora of Ethiopia and Eritrea," *Hydrocharitaceae to Arecaceae*, Vol. 6, National Herbarium, Addis Ababa University, Addis Abeba, Ethiopia, 1997.
- [61] I. Hedberg, S. Edwards, and S. Nemomissa, "Flora of Ethiopia and Eritrea," *Apiaceae to Dipsacaceae*, Vol. 4, National Herbarium, Addis Ababa University, Addis Abeba, Ethiopia, 2003.
- [62] P. V. Bruyns, "Flora of Ethiopia and Eritrea," *Apiaceae to Dipsacaceae*, Vol. 4, Uppsala University, Uppsala, Sweden, 2005.
- [63] R. P. Dick, D. R. Thomas, and J. J. Halvorson, "Standardized methods, sampling, and sample pretreatment," *Methods for Assessing Soil Quality*, vol. 49, pp. 107–121, 1997.
- [64] T. Pearson, S. Walker, and S. Brown, "Sourcebook for land use," *Land-Use Change and Forestry Projects*, vol. 29, 2005.
- [65] S. Ghafari, A. Ghorbani, M. Moameri, R. Mostafazadeh, M. Bidarlord, and A. Kakehmami, "Floristic diversity and distribution patterns along an elevational gradient in the northern part of the Ardabil province rangelands, Iran," *Mountain Research and Development*, vol. 40, no. 1, 2020.
- [66] Z. Woldu, E. Feoli, and L. Nigatu, "Partitioning an elevation gradient of vegetation from southeastern Ethiopia by probabilistic methods," *Numerical Syntaxonomy*, pp. 189–198, Springer, Berlin, Germany, 1989.
- [67] G. Gebeyehu, T. Soromessa, T. Bekele, and D. Teketay, "Carbon stocks and factors affecting their storage in dry Afromontane forests of Awi Zone, northwestern Ethiopia," *Journal of Ecology and Environment*, vol. 43, pp. 7–18, 2019.
- [68] R. Sagar, A. Raghubanshi, and J. Singh, "Tree species composition, dispersion and diversity along a disturbance gradient in a dry tropical forest region of India," *Forest Ecology and Management*, vol. 186, no. 1–3, pp. 61–71, 2003.
- [69] B. McCune, J. B. Grace, and D. L. Urban, "Analysis of ecological communities," *Journal of Experimental Marine Biology and Ecology*, vol. 289, no. 2, 2002.
- [70] R. Tibshirani, G. Walther, and T. Hastie, "Estimating the number of clusters in a data set via the gap statistic," *Journal of the Royal Statistical Society - Series B: Statistical Methodology*, vol. 63, no. 2, pp. 411–423, 2001.
- [71] M. Dufrêne and P. Legendre, "Species assemblages and indicator species: the need for a flexible asymmetrical approach," *Ecological Monographs*, vol. 67, no. 3, pp. 345–366, 1997.
- [72] D. L. Sparks, A. L. Page, P. A. Helmke, and R. H. Loeppert, *Methods of Soil Analysis, Part 3: Chemical Methods*, John Wiley & Sons, Hoboken, NJ, USA, 2020.
- [73] J. M. Bremner, *Nitrogen-total. Methods of soil analysis: Part 3 Chemical methods*, Soil Science Society of America, Madison, Wisconsin, 1996.
- [74] A. Walkley and I. A. Black, "An examination of the Degtjareff method for determining soil organic matter, and a proposed modification of the chromic acid titration method," *Soil Science*, vol. 37, no. 1, pp. 29–38, 1934.
- [75] R. H. Bray and L. T. Kurtz, "Determination of total, organic, and available forms of phosphorus in soils," *Soil Science*, vol. 59, no. 1, pp. 39–46, 1945.
- [76] T. R. Pearson, *Measurement Guidelines for the Sequestration of forest Carbon*, US Department of Agriculture, Forest Service, Northern Research Station, Washington, DC, USA, 2007.
- [77] P. Šmilauer and J. Lepš, *Multivariate Analysis of Ecological Data Using CANOCO 5*, Cambridge University Press, Cambridge, UK, 2014.
- [78] J. Oksanen, *Multivariate Analysis of Ecological Communities in R: Vegan Tutorial*, University of Oulu, Oulu, Finland, 2007.
- [79] B. Groombridge, *Global Biodiversity Status of Earths Living Resources*, Chapman and Dall, London, UK, 1992.
- [80] T. Soromessa and E. Kelbessa, "Interfaces of regeneration, structure, diversity and uses of some plant species in Bonga Forest: a reservoir for wild coffee gene pool," *Sinet: Ethiopian Journal of Science*, vol. 31, no. 2, pp. 121–134, 2011.
- [81] T. W. Gole, T. Borsch, M. Denich, and D. Teketay, "Floristic composition and environmental factors characterizing coffee forests in southwest Ethiopia," *Forest Ecology and Management*, vol. 255, no. 7, pp. 2138–2150, 2008.
- [82] D. Teketay and T. Bekele, "Floristic composition of Wof-Washa natural forest, Central Ethiopia: implications for the conservation of biodiversity," *Feddes Repertorium*, vol. 106, no. 1–5, pp. 127–147, 1995.
- [83] T. Soromessa and E. Kelbessa, "Diversity and endemism of Chilimo forest, central Ethiopia," *Bioscience Discovery*, vol. 4, pp. 1–4, 2013.
- [84] Y. O. Kidane, M. J. Steinbauer, and C. Beierkuhnlein, "Dead end for endemic plant species? A biodiversity hotspot under pressure," *Global Ecology and Conservation*, vol. 19, 2019.
- [85] J.-C. Vié, C. Hilton-Taylor, and S. N. Stuart, *Wildlife in a Changing World: An Analysis of the 2008 IUCN Red List of Threatened Species*, IUCN, Gland, Switzerland, 2009.
- [86] A. Addi, T. Soromessa, and T. Bareke, "Plant diversity and community analysis of Gesha and sayilem forest in kaffa zone, southwestern Ethiopia," *Biodiversitas*, vol. 21, pp. 2878–2888, 2020.
- [87] A. D. Hundie, T. S. Urgessa, and B. W. Dullo, *Plant Community Analysis Along Environmental Gradients In Moist Afromontane Forest Of Gerba Dima, South-Western Ethiopia*, Springer, Berlin, Germany, 2020.
- [88] G. Boz and M. Maryo, "Woody species diversity and vegetation structure of Wurg forest, southwest Ethiopia," *International Journal of Financial Research*, vol. 2020, Article ID 8823990, 17 pages, 2020.
- [89] A. Dibaba, T. Soromessa, A. Kefalew, and A. Addi, "Woody species diversity, vegetation structure, and regeneration

- status of the moist afro-montane forest of Agama in south-western Ethiopia," *International Journal of Ecology*, vol. 2020, Article ID 1629624, 10 pages, 2020.
- [90] H. Yasin, Z. Kebebew, and K. Hundera, "Woody species diversity, regeneration and socioeconomic benefits under natural forest and adjacent coffee agroforests at Belete forest, southwest Ethiopia," *Ekológia*, vol. 37, no. 4, pp. 380–391, 2018.
- [91] C. Kremen, "Assessing the indicator properties of species assemblages for natural areas monitoring," *Ecological Applications*, vol. 2, pp. 203–217, 1992.
- [92] I. Friis, "The forest Vegetation of Ethiopia," *Symbolae Botanicae Upsalienses*, Uppsala Univ, Uppsala, Sweden, 1986.
- [93] I. Friis, *Forests and forest Trees of Northeast Tropical Africa: Their Natural Habitats and Distribution Patterns in Ethiopia, Djibouti and Somalia*, HMSO, Richmond, UK, 1992.
- [94] F. Senbeta, T. W. Gole, M. Denich, and E. Kellbessa, "Diversity of useful plants in the coffee forests of Ethiopia," *Ethnobotany Research and Applications*, vol. 11, pp. 049–069, 2013.
- [95] K. A. Dwire, J. B. Kauffman, E. N. J. Brookshire, and J. E. Baham, "Plant biomass and species composition along an environmental gradient in montane riparian meadows," *Oecologia*, vol. 139, no. 2, pp. 309–317, 2004.
- [96] K. Yasuhiro, M. Hirofumi, and K. Kihachiro, "Effects of topographic heterogeneity on tree species richness and stand dynamics in a subtropical forest in Okinawa Island, southern Japan," *Journal of Ecology*, vol. 92, no. 2, pp. 230–240, 2004.
- [97] C. M. McCain and J. A. Grytnes, "Elevational Gradients in Species Richness," *eLS*, Wiley, Hoboken, NJ, USA, 2010.
- [98] M. L. Monge-González, D. Craven, T. Krömer et al., "Response of tree diversity and community composition to forest use intensity along a tropical elevational gradient," *Applied Vegetation Science*, vol. 23, no. 1, pp. 69–79, 2020.
- [99] H. R. Pulliam, "On the relationship between niche and distribution," *Ecology Letters*, vol. 3, no. 4, pp. 349–361, 2000.
- [100] S. Gairola, R. Rawal, and N. Todaria, "Forest vegetation patterns along an altitudinal gradient in sub-alpine zone of west Himalaya, India," *African Journal of Plant Science*, vol. 2, pp. 042–048, 2008.
- [101] K. Brinkmann, A. Patzelt, U. Dickhoefer, E. Schlecht, and A. Buerkert, "Vegetation patterns and diversity along an altitudinal and a grazing gradient in the Jabal al Akhdar mountain range of northern Oman," *Journal of Arid Environments*, vol. 73, no. 11, pp. 1035–1045, 2009.
- [102] J. L. Ohmann and T. A. Spies, "Regional gradient analysis and spatial pattern of woody plant communities of Oregon forests," *Ecological Monographs*, vol. 68, no. 2, pp. 151–182, 1998.
- [103] J. Terborgh and E. Andresen, "The composition of Amazonian forests: patterns at local and regional scales," *Journal of Tropical Ecology*, vol. 14, no. 5, pp. 645–664, 1998.
- [104] C. R. Pyke, R. Condit, S. Aguilar, and S. Lao, "Floristic composition across a climatic gradient in a neotropical lowland forest," *Journal of Vegetation Science*, vol. 12, no. 4, pp. 553–566, 2001.
- [105] S. A. Bohlman, W. F. Laurance, S. G. Laurance, H. E. Nascimento, P. M. Fearnside, and A. Andrade, "Importance of soils, topography and geographic distance in structuring central Amazonian tree communities," *Journal of Vegetation Science*, vol. 19, no. 6, pp. 863–874, 2008.
- [106] H. Hishe, K. Giday, and M. Haile, "The influence of physical factors on deforestation of key species and their implication for forest management in the dry Afro-montane forest of Desalegn, Northern Ethiopia," *International Journal of Science and Research*, vol. 4, pp. 2400–2407, 2015.
- [107] W. Cui and X.-X. Zheng, "Spatial heterogeneity in tree diversity and forest structure of evergreen broadleaf forests in southern China along an altitudinal gradient," *Forests*, vol. 7, no. 12, p. 216, 2016.
- [108] M. Wondie, D. Teketay, A. M. Melesse, and W. Schneider, "Relationship between topographic variables and land cover in the simen mountains national park, a world heritage site in northern Ethiopia," *International Journal of Remote Sensing Applications*, vol. 2, pp. 36–43, 2012.
- [109] W. G. D. Silva, J. P. Metzger, L. C. Bernacci, E. L. M. Catharino, G. Durigan, and S. Simões, "Relief influence on tree species richness in secondary forest fragments of Atlantic Forest, SE, Brazil," *Acta Botanica Brasiliica*, vol. 22, no. 2, pp. 589–598, 2008.
- [110] G. Woldu, N. Solomon, H. Hishe, H. Gebrewahid, M. A. Gebremedhin, and E. Birhane, "Topographic variables to determine the diversity of woody species in the enclosure of Northern Ethiopia," *Heliyon*, vol. 6, no. 1, 2020.
- [111] M.-L. Smith, "Community and edaphic analysis of upland northern hardwood communities, central Vermont, USA," *Forest Ecology and Management*, vol. 72, no. 2-3, pp. 235–249, 1995.
- [112] J. M. Olano, J. Loidi, G. Ander, and A. Escudero, "Relating variation in the understorey of beech forests to ecological factors," *Folia Geobotanica*, vol. 33, no. 1, pp. 77–86, 1998.
- [113] Y. Zhang, Y. Chen, and B. Pan, "Distribution and floristics of desert plant communities in the lower reaches of Tarim River, southern Xinjiang, People's Republic of China," *Journal of Arid Environments*, vol. 63, no. 4, pp. 772–784, 2005.
- [114] M. G. Turner, "Disturbance and landscape dynamics in a changing world," *Ecology*, vol. 91, no. 10, pp. 2833–2849, 2010.
- [115] L. Götzemberger, F. de Bello, K. A. Bräthen et al., "Ecological assembly rules in plant communities—approaches, patterns and prospects," *Biological Reviews*, vol. 87, no. 1, pp. 111–127, 2012.
- [116] O. Hedberg, "Features of afroalpine plant ecology," *Sv. växtgeografiska sällsk.*, 1964.
- [117] W. Li, L. Cui, B. Sun et al., "Distribution patterns of plant communities and their associations with environmental soil factors on the eastern shore of Lake Taihu, China," *Ecosystem Health and Sustainability*, vol. 3, no. 9, Article ID 1385004, 2017.
- [118] T. Drissen, C. Faust, R. Stadtmann, J. T. Treitler, S. Zerbe, and J. Mantilla-Contreras, "Plant composition and diversity in a semi-natural Mediterranean island landscape: the importance of environmental factors," *Plant Biosystems-An International Journal Dealing with all Aspects of Plant Biology*, vol. 153, no. 6, pp. 756–766, 2019.
- [119] C. M. Hulshof and M. J. Spasojevic, "The edaphic control of plant diversity," *Global Ecology and Biogeography*, vol. 29, no. 10, pp. 1634–1650, 2020.
- [120] M. Kent, W. J. Gill, R. E. Weaver, and R. P. Armitage, "Landscape and plant community boundaries in biogeography," *Progress in Physical Geography: Earth and Environment*, vol. 21, no. 3, pp. 315–353, 1997.
- [121] M. Lee, H. Song, and S. Yee, "Study on classification of forest vegetation of Songinbong and Taeharyong in Ullungdo: with a special reference to TWINSpan and phytosociological method," *Korean Journal of Environment and Ecology*, vol. 14, pp. 57–66, 2000.

- [122] E. van der Maarel, "Vegetation ecology—an overview," *Vegetation ecology*, vol. 3, pp. 1–51, 2005.
- [123] K. Yeshitela and T. Bekele, "Plant community analysis and ecology of Afromontane and transitional rainforest vegetation of southwestern Ethiopia," *Sinet: Ethiopian Journal of Science*, vol. 25, no. 2, pp. 155–175, 2002.
- [124] H. Kreft and W. Jetz, "Global patterns and determinants of vascular plant diversity," *Proceedings of the National Academy of Sciences*, vol. 104, no. 14, pp. 5925–5930, 2007.
- [125] H. Yineger, E. Kelbessa, T. Bekele, and E. Lulekal, "Floristic composition and structure of the dry afromontane forest at Bale Mountains National Park, Ethiopia," *Sinet: Ethiopian Journal of Science*, vol. 31, no. 2, pp. 103–120, 2011.
- [126] M. Didita, S. Nemomissa, and T. W. Gole, "Floristic and structural analysis of the woodland vegetation around Dello Menna, Southeast Ethiopia," *Journal of Forestry Research*, vol. 21, no. 4, pp. 395–408, 2010.
- [127] E. Aynekulu, R. Aerts, P. Moonen et al., "Altitudinal variation and conservation priorities of vegetation along the Great Rift Valley escarpment, northern Ethiopia," *Biodiversity & Conservation*, vol. 21, no. 10, pp. 2691–2707, 2012.
- [128] O. A. Oke and K. A. Thompson, "Distribution models for mountain plant species: the value of elevation," *Ecological Modelling*, vol. 301, pp. 72–77, 2015.
- [129] G. Haile, M. Lemenih, F. Senbeta, and F. Itanna, "Plant diversity and determinant factors across smallholder agricultural management units in Central Ethiopia," *Agroforestry Systems*, vol. 91, no. 4, pp. 677–695, 2017.
- [130] G. Tesfaye, D. Teketay, and M. Fetene, "Regeneration of fourteen tree species in Harenna forest, southeastern Ethiopia," *Flora-Morphology, Distribution, Functional Ecology of Plants*, vol. 197, no. 6, pp. 461–474, 2002.
- [131] C. Mligo, H. Lyaruu, and H. J. Ndangalasi, "The effect of anthropogenic disturbances on population structure and regeneration of *Scorodophloeus fischeri* and *Manilkara sulcata* in coastal forests of Tanzania," *Southern Forests: A Journal of Forest Science*, vol. 73, no. 1, pp. 33–40, 2011.
- [132] E. Aynekulu, R. Aerts, M. Denich et al., "Plant diversity and regeneration in a disturbed isolated dry Afromontane forest in northern Ethiopia," *Folia Geobotanica*, vol. 51, no. 2, pp. 115–127, 2016.
- [133] C. C. Jakovac, A. B. Junqueira, R. Crouzeilles, M. Peña-Claros, R. C. G. Mesquita, and F. Bongers, "The role of land-use history in driving successional pathways and its implications for the restoration of tropical forests," *Biological Reviews*, vol. 96, no. 4, pp. 1114–1134, 2021.
- [134] M. Elemans, "Light, nutrients and the growth of herbaceous forest species," *Acta Oecologica*, vol. 26, no. 3, pp. 197–202, 2004.
- [135] I. M. Pérez-Ramos, M. A. Zavala, T. Marañón, M. D. Díaz-Villa, and F. Valladares, "Dynamics of understorey herbaceous plant diversity following shrub clearing of cork oak forests: a five-year study," *Forest Ecology and Management*, vol. 255, no. 8–9, pp. 3242–3253, 2008.
- [136] T. Bekele, "Phytosociology and ecology of a humid Afromontane forest on the central plateau of Ethiopia," *Journal of Vegetation Science*, vol. 5, no. 1, pp. 87–98, 1994.