

1 **Altered stomatal patterning accompanies a trichome dimorphism in a natural**
2 **population of *Arabidopsis***

3

4

5 Noriane M. L. Simon¹, Jiro Sugisaka², Mie N. Honjo², Sverre Aarseth Tunstad¹, George
6 Tunna¹, Hiroshi Kudoh², Antony N. Dodd^{3*}

7

8

9 1. School of Biological Sciences, University of Bristol, Bristol BS8 1TQ, U.K.

10 2. Center for Ecological Research, Kyoto University, Otsu, Shiga, Japan.

11 3. John Innes Centre, Norwich NR4 7UH, U.K.

12

13 * Corresponding author; antony.dodd@jic.ac.uk; +44 (0)1603 450015.

14

15 Keywords: stomata, environmental adaptation, development, herbivory.

16

17 **Abstract**

18 Trichomes are large epidermal cells on the surface of leaves that are thought to deter
19 herbivores, yet the presence of trichomes can also negatively impact plant growth and
20 reproduction. Stomatal guard cells and trichomes have shared developmental origins, and
21 experimental manipulation of trichome formation can lead to changes in stomatal density.
22 The influence of trichome formation upon stomatal development in natural populations of
23 plants is currently unknown. Here, we show that a natural population of *Arabidopsis halleri*
24 that includes hairy (trichome-bearing) and glabrous (no trichomes) morphs has differences in
25 stomatal density that are associated with this trichome dimorphism. We found that glabrous
26 morphs had significantly greater stomatal density and stomatal index than hairy morphs.
27 One interpretation is that this arises from a trade-off between the proportions of cells that
28 have trichome and guard cell fates during leaf development. The differences in stomatal
29 density between the two morphs might have impacts upon environmental adaptation, in
30 addition to herbivory deterrence caused by trichome development.

31

32 Introduction

33 In *Arabidopsis*, trichomes are large epidermal cells that protrude from the surface of the
 34 leaves and petioles. Trichomes play important roles in both biotic defences and abiotic
 35 stress tolerance (Levin, 1973; Mauricio and Rausher, 1997; Handley et al., 2005; Dalin et al.,
 36 2008; Sletvold et al., 2010; Sletvold and Ågren, 2012; Sato and Kudoh, 2016). However,
 37 trichome development appears to impose a fitness cost on growth and reproduction
 38 (Mauricio, 1998; Sletvold et al., 2010; Kawagoe et al., 2011; Sletvold and Ågren, 2012; Sato
 39 and Kudoh, 2016). In addition to trichomes, stomatal guard cells represent another
 40 specialized cell type that is present on the leaf surface. Trichome initiation occurs prior to
 41 stomatal meristemoid development, and the patterning of trichomes and guard cells appears
 42 to be linked (Larkin et al., 1996; Glover, 2000; Bean et al., 2002; Bird and Gray, 2003;
 43 Galdon-Armero et al., 2018). Therefore, there might be a trade-off between trichome and
 44 stomatal guard cell development during leaf formation (Glover et al., 1998).

45 We wished to determine whether trichome formation might be associated with changes in
 46 stomatal patterning in natural populations of plants. To achieve this, we investigated
 47 stomatal patterning in a naturally-occurring population of *Arabidopsis halleri* subsp.
 48 *gemmifera* that includes trichome-forming and glabrous morphs (Kawagoe et al., 2011; Sato
 49 and Kudoh, 2016). These trichome morph phenotypes are heritable (Sato and Kudoh, 2015,
 50 2017). The glabrous morphs within this population harbour a large transposon-like insertion
 51 within the *GLABRA1* (*GL1*) gene (Kawagoe et al., 2011). *GL1* is also required for trichome
 52 formation in *A. thaliana*, with homozygous *gl1* mutants being glabrous (Oppenheimer et al.,
 53 1991). Our experiments provide new insights into the relationship between stomatal and
 54 trichome patterning under natural conditions.

55 **Methods**

56 *Study site and experimental model*

57 This investigation used a well-characterized population of *Arabidopsis halleri* subsp.
58 *gemmifera* that is located beside a small stream in central Honshu island, Japan (Fig. 1A)
59 (Omoide-gawa site, 35°06' N, 134°56' E; 230 m altitude) (Aikawa et al., 2010; Kudoh et al.,
60 2018). *A. halleri* is metal tolerant and grows essentially as a monoculture at this field site
61 because the water is contaminated by a historical mine (Kudoh et al., 2018). The species
62 was identified by reference to herbarium and museum specimens (Kudoh et al., 2018), and
63 a nearby population that harbours glabrous and hairy morphs supplied material for the
64 genome sequencing and annotation of *A. halleri* (Briskine et al., 2017; Sato and Kudoh,
65 2017). The only subspecies of *A. halleri* present in Japan is *A. halleri* subsp. *gemmifera*
66 (Honjo and Kudoh, 2019). Sampling occurred during September 2016 (photoperiod
67 approximately 12 h, with dawn at 05:40 and dusk at 18:10). During this season, *A. halleri*
68 bore larger rosette leaves that are well-suited for quantification of stomatal density (Fig. 1B).

69 *Stomatal density measurement*

70 Eight plants of each trichome morph (hairy or glabrous) were selected at the study site, with
71 individuals chosen such that the replicate plants were distributed evenly across the site.
72 Glabrous and hairy morphs were identified by visual inspection of the leaf surface. It is
73 thought that irradiance and ambient temperature are unlikely to influence the frequency of
74 the morphs (Sato and Kudoh, 2017), but we cannot discount the possibility of
75 microenvironment- or field site edge-effects. Stomatal density was measured by obtaining
76 impressions from the adaxial surfaces of between 3 and 5 fully-expanded rosette leaves of
77 each plant. We focused on the adaxial surface because this surface also harbours the
78 majority of the trichomes. Between the times of 12:00 and 13:00, President Plus dental
79 impression paste (Coltene) was applied to the adaxial side of each leaf to create a leaf
80 surface impression (Fig. 1C). Solidified impression paste was removed from leaves and

81 transported to the laboratory for further processing. First, each impression was assigned a
82 randomly-generated number to ensure subsequent steps were performed blind. Each leaf
83 impression was painted with transparent nail varnish (60 seconds super shine, Rimmel) that,
84 after drying, was peeled away from the dental impression paste using transparent adhesive
85 tape (Scotch Crystal). Next, the adhesive tape was used to attach the nail varnish
86 impression to a 0.8 mm – 1 mm thick microscope slide. Leaf impressions were examined
87 using an epifluorescence microscope in white light illumination mode. Images were captured
88 from the centre of each leaf half, away from the midrib, using a Hamamatsu camera and
89 Volocity software set to 20x zoom. Two images were captured from each impression, and
90 the number of stomata and pavement cells was counted in an 800 µm x 800 µm square
91 using the Fiji software to obtain cell density measures. Cell density measures were
92 expressed as per mm² (multiplication by 1.56).

93 In total, 29 and 31 leaf impressions were obtained in the field from hairy and glabrous plants,
94 respectively. This produced 58 (hairy) and 62 (glabrous) microscopy images for analysis,
95 because two images were captured from each impression. Stomatal index was calculated
96 according to Equation 1. After all measurements, data were disaggregated according to the
97 blinding/randomization scheme. The differences between hairy and glabrous plants were
98 statistically tested by nested analysis of variance, whereby the mean stomatal density or
99 index per replicate plant was nested within the hairy and glabrous morphs. Tests were
100 conducted using the R 3.6.0 software (Team, 2019) and plots generated with the beeswarm
101 R package (v0.2.3) and Inkscape v0.91. No adjustments were applied to photographs in Fig.
102 1.

$$SI = \frac{s}{s + p} \times 100$$

103 **Equation 1.** Derivation of stomatal index, where *SI* is stomatal index, *s* is the number of
104 stomata in the field of view, and *p* is the number of epidermal pavement cells in the field of
105 view.

Results

We investigated stomatal patterning in naturally-occurring hairy and glabrous morphs of *A. halleri* (Sato and Kudoh, 2016). Approximately half of the *A. halleri* population at this study site is glabrous, whilst remaining plants have trichomes (Kawagoe et al., 2011). As trichome initiation occurs prior to stomatal meristemoid formation (Larkin et al., 1996; Glover, 2000), it is likely that trichome and stomatal patterning are linked (Bean et al., 2002), so we hypothesized that this might produce a difference in stomatal density between the two trichome morphs of *A. halleri* under natural conditions.

We found that the trichome formation dimorphism was accompanied by a difference in stomatal density (Fig. 2A; Fig. S1; Supplemental Dataset S1). Fully-expanded leaves of glabrous morphs had significantly greater stomatal density on the adaxial surface compared with hairy-leaved morphs (glabrous: 30.7 ± 2.8 stomata mm^{-2} ; hairy: 23.6 ± 2.3 stomata mm^{-2} ; mean $\pm$ s.e.m) (Fig. 2A; Table S1; Supplemental Dataset S1). Furthermore, the stomatal index of the adaxial surface was significantly greater in glabrous morphs (18.04 ± 0.92) compared with hairy morphs (16.44 ± 1.03) (Fig. 2B; Table S1). The adaxial surface pavement cell density did not differ significantly between the morphs (Table S1). Although leaf widths varied significantly among plants, they did not differ significantly between the morphs (glabrous: 12.4 ± 0.7 mm; hairy: 13.5 ± 1.0 mm; Fig. S2; Supplemental Dataset S1), suggesting that the stomatal density difference between the morphs is not due to differences in leaf expansion between the morphs (Table S1). Mean stomatal density ranged from 17 – 35 stomata mm^{-2} for hairy morphs and 24 – 49 stomata mm^{-2} for glabrous morphs (Fig. 2A). This stomatal density was lower than for *Arabidopsis thaliana*, which has reported stomatal densities of 180 – 350 stomata mm^{-2} depending on background accession and growth conditions (Gray et al., 2000; Zhang et al., 2008; Franks et al., 2015). Although lower than in *A. thaliana*, our measurements of stomatal density in *A. halleri* are consistent with a previous report of stomatal density of *A. halleri* subsp. *gemmifera*, which measured adaxial stomatal density of 46 stomata mm^{-2} at 430 m altitude in autumn rosette leaves, with stomatal density

progressively increasing with greater altitudes (Aryal et al., 2018). Our field site was lower altitude (230 m), so the lower stomatal densities at our study site (Fig. 2A) are congruous with this previous study (Aryal et al., 2018).

Discussion

Glabrous plants had significantly greater stomatal density and stomatal index compared with hairy plants (Fig. 2A; Fig. 2B). As the density of surrounding pavement cells did not vary between the morphs, these differences in stomatal density and index are due to the greater density of stomata in glabrous morphs compared with hairy morphs (Fig. 2B). Our field data are consistent with a laboratory-based study in which transgenic tobacco plants expressing an *Antirrhinum myb*-like transcription factor, which caused an excess of trichomes, also had significantly reduced stomatal density (Glover et al., 1998). In a segregating tomato population, there is a negative correlation between stomatal and trichome density specifically under drought conditions (Galdon-Armero et al., 2018). Similarly, the trichome-bearing Col-0 accession of *A. thaliana* has lower stomatal density than the glabrous C24 accession (e.g. about 115 mm⁻² for Col-0 and 180 mm⁻² for C24) (Perazza et al., 1998; Lake and Woodward, 2008), although factors other than trichome density are likely to influence stomatal density between the accessions. This suggests that in natural populations of *A. halleri*, there could be a trade-off between trichome and stomatal development. Since the glabrous *gl1* mutant of *A. thaliana* has a significantly greater density of stomatal units compared with the wild type (Berger et al., 1998) and the glabrous phenotype of *A. halleri* at this study site is associated with an insertion within *GL1* (Kawagoe et al., 2011), it is possible that the *GL1* haplotype influences the stomatal density within this population of *A. halleri*.

In some cases, there does not appear to be a tradeoff between stomatal and trichome density. For example, elevated CO₂ decreases stomatal density (Woodward and Kelly, 1995), but might also reduce trichome density (Bidart-Bouzat et al., 2005). Therefore, in future, it could be informative to examine the relationship between stomatal and trichome

density under a range of different experimental conditions that apply different types of selection pressure. Furthermore, we sampled the adaxial leaf surface and it is possible that the presence of trichomes might affect stomatal density differently on the abaxial surface because, depending on environmental conditions, abaxial stomatal density of *A. halleri* can be 10—30% greater than the adaxial surface (Aryal et al., 2018).

Interestingly, trichome production appears to impose a fitness cost. For example, glabrous *A. halleri* plants have 10% greater biomass than hairy plants when grown in the absence of herbivores (Sato and Kudoh, 2016). This cost of herbivore resistance arising from trichome formation also occurs in glabrous and hairy *A. lyrata* (Løe et al., 2007; Sletvold et al., 2010) and *A. thaliana* (Mauricio and Rausher, 1997; Mauricio, 1998) under experimental conditions excluding herbivores. Whilst this fitness advantage of glabrous over hairy leaves in the absence of herbivory might be due to trichome production (Mauricio and Rausher, 1997; Mauricio, 1998; Kawagoe and Kudoh, 2010; Sletvold et al., 2010; Kawagoe et al., 2011; Sletvold and Ågren, 2012), we suggest that glabrous morphs might also gain an advantage by having a greater density or number of stomata. It has been proposed that increasing the number of stomata could increase carbon assimilation (Lawson and Blatt, 2014). For example, *Arabidopsis* overexpressing *STOMAGEN* has greater stomatal density and a 30% increase in carbon assimilation compared with the wild type. However, these lines also have higher transpiration rates and consequently lower water use efficiency (Tanaka et al., 2013). An alternative interpretation is that differences in the developmental program of the hairy and glabrous morphs might lead to differences in cell or leaf size, which ultimately causes the biomass difference between the morphs. In our samples, the width of fully expanded leaves did not differ significantly between the morphs (Fig. S2). Using these leaf width measures as a proxy for leaf size suggests that the biomass difference between the morphs is not due to leaf size differences between the morphs. However, there might be differences in other developmental characteristics that affect biomass, such as leaf thickness, which we did not compare between the morphs.

Optimal stomatal density is important to achieve high photosynthetic rates. A low stomatal density restricts CO₂ vertical diffusion through the leaf and reduces photosynthetic rates, whilst high-density stomatal clustering diminishes CO₂ diffusion and causes low carbon assimilation (Lawson and Blatt, 2014). Both *A. halleri* morphs examined are likely to be within an optimal range of stomatal densities, having evolved and survived under natural conditions. However, the higher stomatal density in the glabrous morph might contribute to its faster growth in absence of herbivory (Sato and Kudoh, 2016). In future, it would be interesting to explore this by measuring the CO₂ assimilation rate of these trichome morphs under laboratory and/or natural conditions. It would also be informative to determine whether the stomatal density difference between the two trichome morphs confers any advantages within microenvironments characterized by differences in water or light availability. The lower stomatal density of *A. halleri* compared with *A. thaliana* (Gray et al., 2000; Zhang et al., 2008; Franks et al., 2015) might reflect differences in growth conditions. An alternative explanation might relate to genome size, because there appears to be a negative correlation between genome size and stomatal density (Beaulieu et al., 2008), and the genome of *A. halleri* (250 Mb) is approximately double the size of the *A. thaliana* genome (125 Mb) (The Arabidopsis Genome, 2000; Briskine et al., 2017).

In summary, we found that a glabrous morph of *A. halleri* growing under natural conditions had greater stomatal density and stomatal index than a hairy morph. This might contribute to the reported fitness advantage of glabrous plants over hairy plants in absence of herbivores (Sato and Kudoh, 2017). This differing stomatal density phenotype might derive from the common upstream components in the pathways leading to trichome and guard cell development.

Acknowledgements

We thank Dora Cano-Ramirez, Haruki Nishio and Tasuku Ito for experimental assistance. This research was funded by the UK Biotechnology and Biological Sciences Research

212 Council (BBSRC; SWBio DTP award BB/J014400/1 and Institute Strategic Programme GEN
213 BB/P013511/1), The Royal Society (grant IE140501), and the Japan Society for Promotion
214 of Science (JSPS; CREST no. JPMJCR15O1). This research was conducted through Joint
215 Usage of the Center for Ecological Research, Kyoto University.

216 **Conflict of Interests**

217 The authors declare no competing financial interests.

218 **Author contributions**

219 NMLS, JS, MNH, SAT, GT, HK and AND performed experimentation and/or analysed data,
220 and NMLS, MNH, HK and AND interpreted findings and wrote the paper.

221 **Data availability**

222 All data generated during this study are included in the published article and Supplementary
223 Information files.

224

225 Figure legends

226 **Figure 1.** Field sampling of *Arabidopsis halleri* for stomatal density. (A) Overview of field
227 site; (B) Rosette form of *A. halleri* plants when sampling during September; (C) Leaf surface
228 impression acquisition using impression paste. The impression paste is green-coloured and
229 occupies the surface of three rosette leaves.

230 **Figure 2.** Stomatal density differs between hairy and glabrous morphs within a natural
231 population of *Arabidopsis halleri*. (A) Stomatal density and (B) stomatal index for fully
232 expanded leaves of hairy and glabrous morphs. Each red point represents the mean
233 stomatal density or stomatal index from one individual plant. The mean stomatal density and
234 stomatal index per plant was derived from 2 microscopy images analysed from 3-4 leaves of
235 each plant. The centre line of the boxplot indicates the median. $n = 8$ plants from each
236 morph; analysed by one-way nested ANOVA. *** indicates $p < 0.001$.

237 **Table S1.** Nested ANOVA analysis of (a) stomatal density, (b) stomatal index, (c) pavement
238 cell density and (d) leaf width. *Df*, degree of freedom; *, **, ***, significant at $p < 0.05$, < 0.01 ,
239 < 0.001 respectively; NS, not significant at $p \geq 0.05$.

240 **Figure S1.** Stomatal density differs between hairy and glabrous morphs within a natural
241 population of *Arabidopsis halleri*. Figure summarizes all data collected for (A) stomatal
242 density and (B) stomatal index of fully expanded leaves of hairy and glabrous morphs. Each
243 red point represents one measurement from one microscopy sample, with 2 samples
244 analyzed from 3-4 leaves from 8 individual plants of each morph. The centre line of the
245 boxplot indicates the median. Data represent a total of 58 and 62 microscopy images from
246 hairy and glabrous plants, respectively.

247 **Figure S2.** Mean width of fully-expanded leaves does not differ between glabrous and hairy
248 morphs within a natural population of *Arabidopsis halleri*. Each red point represents the
249 mean leaf width from one plant, calculated from the width of either 3 or 4 leaves per plant. 8

replicate plants were sampled for each morph. The centre line of the boxplot indicates the median. Data were analyzed by one-way nested ANOVA, using the mean leaf width per plant as the level of replication. N. S. indicates difference is not statistically significant at $p \geq 0.05$

Dataset S1. All stomatal density and leaf width data collected during experimentation.

References

- Aikawa S, Kobayashi MJ, Satake A, Shimizu KK, Kudoh H** (2010) Robust control of the seasonal expression of the *Arabidopsis FLC* gene in a fluctuating environment. *Proceedings of the National Academy of Sciences* **107**: 11632-11637
- Aryal B, Shinohara W, Honjo MN, Kudoh H** (2018) Genetic differentiation in cauline-leaf-specific wettability of a rosette-forming perennial *Arabidopsis* from two contrasting montane habitats. *Annals of Botany* **121**: 1351-1360
- Bean GJ, Marks MD, Hulskamp M, Clayton M, Croxdale JL** (2002) Tissue patterning of *Arabidopsis* cotyledons. *New Phytologist* **153**: 461-467
- Beaulieu JM, Leitch IJ, Patel S, Pendharkar A, Knight CA** (2008) Genome size is a strong predictor of cell size and stomatal density in angiosperms. *New Phytologist* **179**: 975-986
- Berger F, Linstead P, Dolan L, Haseloff J** (1998) Stomata patterning on the hypocotyl of *Arabidopsis thaliana* is controlled by genes involved in the control of root epidermis patterning. *Developmental Biology* **194**: 226-234
- Bidart-Bouzat MG, Mithen R, Berenbaum MR** (2005) Elevated CO₂ influences herbivory-induced defense responses of *Arabidopsis thaliana*. *Oecologia* **145**: 415-424
- Bird SM, Gray JE** (2003) Signals from the cuticle affect epidermal cell differentiation. *New Phytologist* **157**: 9-23
- Briskine RV, Paape T, Shimizu-Inatsugi R, Nishiyama T, Akama S, Sese J, Shimizu KK** (2017) Genome assembly and annotation of *Arabidopsis halleri*, a model for heavy

276 metal hyperaccumulation and evolutionary ecology. *Molecular Ecology Resources*
277 **17**: 1025-1036

278 **Dalin P, Agren J, Bjorkman C, Huttunen P, Karkkainen K** (2008) Leaf trichome formation
279 and plant resistance to herbivory. *In* A Schaller, ed, *Induced plant resistance to*
280 *herbivory*. Springer, pp 89-105

281 **Franks PJ, W. Doheny-Adams T, Britton-Harper ZJ, Gray JE** (2015) Increasing water-use
282 efficiency directly through genetic manipulation of stomatal density. *New Phytologist*
283 **207**: 188-195

284 **Galdon-Armero J, Fullana-Pericas M, Mulet PA, Conesa MA, Martin C, Galmes J** (2018)
285 The ratio of trichomes to stomata is associated with water use efficiency in *Solanum*
286 *lycopersicum* (tomato). *The Plant Journal* **96**: 607-619

287 **Glover BJ** (2000) Differentiation in plant epidermal cells. *Journal of Experimental Botany* **51**:
288 497-505

289 **Glover BJ, Perez-Rodriguez M, Martin C** (1998) Development of several epidermal cell
290 types can be specified by the same MYB-related plant transcription factor.
291 *Development* **125**: 3497

292 **Gray JE, Holroyd GH, van der Lee FM, Bahrami AR, Sijmons PC, Woodward FI,**
293 **Schuch W, Hetherington AM** (2000) The HIC signalling pathway links CO₂
294 perception to stomatal development. *Nature* **408**: 713-716

295 **Handley R, Ekbom B, Ågren J** (2005) Variation in trichome density and resistance against
296 a specialist insect herbivore in natural populations of *Arabidopsis thaliana*. *Ecological*
297 *Entomology* **30**: 284-292

298 **Honjo MN, Kudoh H** (2019) *Arabidopsis halleri*: a perennial model system for studying
299 population differentiation and local adaptation. *AoB PLANTS* **11**

300 **Kawagoe T, Kudoh H** (2010) Escape from floral herbivory by early flowering in *Arabidopsis*
301 *halleri* subsp. *gemmifera*. *Oecologia* **164**: 713-720

302 **Kawagoe T, Shimizu KK, Kakutani T, Kudoh H** (2011) Coexistence of trichome variation
303 in a natural plant population: A combined study using ecological and candidate gene
304 approaches. *PLoS ONE* **6**: e22184

305 **Kudoh H, Honjo MN, Nishio H, Sugisaka J** (2018) The long-term "in natura" study sites of
306 *Arabidopsis halleri* for plant transcription and epigenetic modification analyses in
307 natural environments. *In Plant Transcription Factors*. Springer, pp 41-57

308 **Lake JA, Woodward FI** (2008) Response of stomatal numbers to CO₂ and humidity: control
309 by transpiration rate and abscisic acid. *New Phytologist* **179**: 397-404

310 **Larkin JC, Young N, Prigge M, Marks MD** (1996) The control of trichome spacing and
311 number in *Arabidopsis*. *Development* **122**: 997-1005

312 **Lawson T, Blatt MR** (2014) Stomatal size, speed, and responsiveness impact on
313 photosynthesis and water use efficiency. *Plant Physiology* **164**: 1556-1570

314 **Levin DA** (1973) The role of trichomes in plant defense. *The Quarterly Review of Biology*
315 **48**: 3-15

316 **Løe G, Toräng P, Gaudeul M, Ågren J** (2007) Trichome production and spatiotemporal
317 variation in herbivory in the perennial herb *Arabidopsis lyrata*. *Oikos* **116**: 134-142

318 **Mauricio R** (1998) Costs of resistance to natural enemies in field populations of the annual
319 plant *Arabidopsis thaliana*. *The American Naturalist* **151**: 20-28

320 **Mauricio R, Rausher MD** (1997) Experimental manipulation of putative selective agents
321 provides evidence for the role of natural enemies in the evolution of plant defense.
322 *Evolution* **51**: 1435-1444

323 **Oppenheimer DG, Herman PL, Sivakumaran S, Esch J, Marks MD** (1991) A *myb* gene
324 required for leaf trichome differentiation in *Arabidopsis* is expressed in stipules. *Cell*
325 **67**: 483-493

326 **Perazza D, Vachon G, Herzog M** (1998) Gibberellins promote trichome formation by up-
327 regulating *GLABROUS1* in *Arabidopsis*. *Plant Physiology* **117**: 375-383

328 **R Core Team** (2019) R: A language and environment for statistical computing. R Foundation
329 for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>.

- 330 **Sato Y, Kudoh H** (2015) Tests of associational defence provided by hairy plants for
 331 glabrous plants of *Arabidopsis halleri* subsp. *gemmifera* against insect herbivores.
 332 Ecological Entomology **40**: 269-279
- 333 **Sato Y, Kudoh H** (2016) Associational effects against a leaf beetle mediate a minority
 334 advantage in defense and growth between hairy and glabrous plants. Evolutionary
 335 Ecology **30**: 137-154
- 336 **Sato Y, Kudoh H** (2017) Fine-scale frequency differentiation along a herbivory gradient in
 337 the trichome dimorphism of a wild *Arabidopsis*. Ecology and Evolution **7**: 2133-2141
- 338 **Sletvold N, Ågren J** (2012) Variation in tolerance to drought among Scandinavian
 339 populations of *Arabidopsis lyrata*. Evolutionary Ecology **26**: 559-577
- 340 **Sletvold N, Huttunen P, Handley R, Kärkkäinen K, Ågren J** (2010) Cost of trichome
 341 production and resistance to a specialist insect herbivore in *Arabidopsis lyrata*.
 342 Evolutionary Ecology **24**: 1307-1319
- 343 **Tanaka Y, Sugano SS, Shimada T, Hara-Nishimura I** (2013) Enhancement of leaf
 344 photosynthetic capacity through increased stomatal density in *Arabidopsis*. New
 345 Phytologist **198**: 757-764
- 346 **The Arabidopsis Genome Initiative** (2000) Analysis of the genome sequence of the
 347 flowering plant *Arabidopsis thaliana*. Nature **408**: 796-815
- 348 **Woodward FI, Kelly CK** (1995) The influence of CO₂ concentration on stomatal density.
 349 New Phytologist **131**: 311-327
- 350 **Zhang L, Hu G, Cheng Y, Huang J** (2008) Heterotrimeric G protein α and β subunits
 351 antagonistically modulate stomatal density in *Arabidopsis thaliana*. Developmental
 352 Biology **324**: 68-75

353

Fig. 1




Fig. 2

