

# Nutritional Significance and Health-Promoting Potential of Solanaceae Family Vegetables: A Comparative Review in Asia and Europe

Bipasha Mridha Ghosh<sup>1</sup>, Priyanka Jana<sup>2</sup>, Labanya Banerjee<sup>3</sup>,  
Soumili Chakroborty<sup>4</sup>

<sup>1</sup>Faculty, Basic Science, Nshm Knowledge Campus-Dgp

<sup>2,3,4</sup>Student, Dietetics & Nutrition, Nshm Knowledge Campus-Dgp

## Abstract

The Solanaceae family encompasses several globally significant vegetables—tomato (*Solanum lycopersicum*), potato (*Solanum tuberosum*), eggplant (*Solanum melongena*), and chili pepper (*Capsicum annuum*)—which collectively provide macronutrients, micronutrients and bioactive phytochemicals fundamental to human diets. This review synthesizes contemporary literature on the compositional profiles, bioavailability, health outcomes, and regional production–consumption patterns of these vegetables, with a focused comparison between Asia and Europe. We integrate agronomic, nutritional, processing, and public-health perspectives and provide practical recommendations for research, breeding, processing and policy to maximize nutritional benefits across different food systems.

**Keywords:** Solanaceae, tomato, potato, eggplant, chili pepper, nutrition, Asia, Europe, processing, bioavailability

## Introduction

Vegetables from the Solanaceae family are deeply embedded in culinary cultures around the world and have substantial implications for population nutrition. These crops serve as major dietary sources of energy (potato starch), vitamins (vitamin C, provitamin A carotenoids), minerals (potassium, magnesium), dietary fiber and numerous phytochemicals (lycopene in tomato; nasunin in eggplant; capsaicinoids in chili). Scientific literature increasingly recognizes not only their nutrient density but also the importance of processing and preparation on nutrient bioavailability and net health outcomes (Rao & Rao, 2007; Liu, 2013).

This review targets two complementary regional systems: Asia—characterized by high production volumes, smallholder diversity and culinary heterogeneity—and Europe—characterized by mechanized production, value-added processing and distinctive dietary patterns (Mediterranean vs. Northern European). The goals are to (1) summarize nutrient composition and health evidence, (2) compare production and consumption trends, (3) analyze how processing and culinary practices modify nutritional outcomes, and (4) propose actionable recommendations for widening nutritional benefits in both regions.

## Methodology

We performed a narrative synthesis of peer-reviewed articles, FAO/FAOSTAT reports, regional agricultural bulletins, and authoritative reviews up to 2025. Searches prioritized systematic reviews, meta-analyses on health outcomes, composition studies, and official statistical databases for production and per-capita supply. Selected country case studies (China, India, Japan; Spain, Italy, Germany, Netherlands) were included to illustrate regional patterns. Where direct primary data were available (FAOSTAT/Our World in Data), tables were constructed to show comparative production, nutrient composition averages from food composition tables, and per-capita vegetable supply.

## Nutrient Composition and Bioactive Constituents

An accurate assessment of nutritive value requires distinguishing proximate composition (macronutrients, moisture), micronutrients (vitamins and minerals), dietary fiber, and phytochemicals (carotenoids, phenolics, flavonoids, alkaloids).

### Potato

Potatoes are primarily a source of digestible carbohydrates (starch ~15–20% fresh weight), with moderate protein (1.8–2.5% fresh weight) and low fat. Key micronutrients include vitamin C (15–20 mg/100 g raw), B vitamins (notably B6), and high potassium (~300–420 mg/100 g). Phenolic compounds and glycoalkaloids (solanine, chaconine) are present in variable amounts depending on cultivar and tuber exposure to light; phenolics contribute antioxidant activity (Camire et al., 2009).

### Tomato

Tomatoes offer vitamin C (10–25 mg/100 g), folate, potassium (200–250 mg/100 g), and are especially valued for carotenoids—principally lycopene and  $\beta$ -carotene. Lycopene accumulates in fruit chromoplasts and is more bioavailable after thermal processing and oil co-consumption due to increased matrix disruption and lipid-mediated micellarization (Rao & Rao, 2007).

### Eggplant

Eggplant flesh is mostly water with modest fiber (2–3 g/100 g) and small amounts of vitamins and minerals. The key bioactives are anthocyanins concentrated in the peel—most notably nasunin—which have radical-scavenging properties and may protect membrane lipids from oxidative injury (Hosseini, 2011).

### Chili Pepper

Chili peppers are nutrient-dense in vitamins A (provitamin A carotenoids) and C (up to 100–242 mg/100 g in some varieties). Capsaicinoids (capsaicin, dihydrocapsaicin) impart pungency and have been studied for analgesic, thermogenic, and metabolic effects (Zimmer, 2017).

## Production and Supply: Comparative Data

Regional production and per-capita supply shape accessibility and dietary impact. FAOSTAT-derived data for 2023 indicate the following broad patterns:

- **Potato:** Global production ~383 million tonnes in 2023; China and India together account for a large share, with China producing an estimated ~93 million tonnes in 2023, making it the top producer. Europe (EU countries collectively) remains a major producer and processor. (FAO/FAOSTAT summaries).

- **Tomato:** China (~67–68 million tonnes) and India (~20–21 million tonnes) are the largest producers globally, while Italy and Spain are major European producers and important processing hubs. (FAO/Our World in Data).
- **Eggplant:** China and India produce the bulk of global eggplant output (combined ~85% of world production), with China producing ~38–39 million tonnes and India ~12–13 million tonnes in recent reports. (FAOSTAT-derived summaries).
- **Peppers (Capsicum spp.):** China leads, with Spain ranking within the top producers in Europe; exact values vary among sources but indicate a clear Asia dominance for quantity. (FAOSTAT/HortiDaily).

**Table 1. Comparative production (selected countries, 2023)**

| Crop           | China (million t)           | India (million t) | Spain (million t) | Italy (million t) | Germany/Netherlands (million t, potatoes) |
|----------------|-----------------------------|-------------------|-------------------|-------------------|-------------------------------------------|
| Potato         | 93.5                        | 57.4*             | 4.0               | 4.5               | 10–20 (combined seed/processing)          |
| Tomato         | 67.5                        | 21.0              | 6.0               | 6.5               | —                                         |
| Eggplant       | 38.3                        | 12.8              | 0.3               | 0.31              | —                                         |
| Pepper (fresh) | 36.7 (value basis ? global) |                   | 2.0–3.0           | 0.8–1.2           | —                                         |

\*India potato numbers are regionally variable; figures indicate order of magnitude. Table values are synthesized from FAO/Our World in Data and regional sources (see references and data annex).

Note: Table values are rounded and intended for comparative purposes. Exact values are accessible from FAOSTAT and Our World in Data datasets.

## Per-Capita Supply and Consumption Patterns

Per-capita vegetable supply (kg/person/year) varies widely by country and region. Our World in Data (FAO-based) shows high per-capita vegetable supply in countries such as China (>300 kg/year in some estimates) compared to European averages which vary but are substantial in Mediterranean diets. Consumption of specific Solanaceae items (e.g., tomato-based sauces in Italy vs. fresh chilies in India) reflects cultural patterns and processing infrastructure.

**Table 2. Representative per-capita vegetable supply (selected countries, kg/person/year)**

| Country | Year      | Vegetables (kg/capita/year) |
|---------|-----------|-----------------------------|
| China   | 2022–2023 | ~380–408                    |
| India   | 2022–2023 | ~200–250                    |
| Spain   | 2022      | ~220–250                    |
| Italy   | 2022      | ~260–300                    |
| Germany | 2022      | ~220–240                    |

These country-level estimates reflect supply available at the consumer end of the supply chain and do not directly equal intake (food waste and retail losses reduce actual consumption). Nevertheless, they provide a useful proxy for accessibility.

### Processing, Culinary Practices and Bioavailability

Processing intersects with nutrition in two main ways: (1) altering nutrient concentration and composition (e.g., moisture loss increases nutrient density per unit weight), and (2) changing bioavailability (e.g., thermal processing increases lycopene bioavailability in tomatoes but may degrade vitamin C).

- **Tomato processing** (heating and homogenization) breaks down cell walls and liberates carotenoids; lycopene becomes more bioavailable in sauces and paste, especially when consumed with dietary fat. Industrial tomato processing in Italy and Spain creates value-added products that are important sources of bioavailable lycopene in European diets.
- **Potato processing:** frying and pre-cooked frozen products increase energy density and fat intake. Boiling preserves more water-soluble vitamins if cooking water is consumed (e.g., in soups).
- **Eggplant:** the skin contains much of the anthocyanin content; frying versus grilling can influence oil uptake and antioxidant retention.
- **Chili peppers:** capsaicin bioactivity is fairly heat-stable; however, vitamin C content may decline with thermal processing; pickling and drying concentrate certain nutrients while reducing others.

### Health Evidence: Epidemiology and Mechanisms

A growing body of epidemiological and mechanistic research links vegetable intake to reduced risk of non-communicable diseases. While most large cohort studies analyze fruit and vegetable intake collectively, several lines of evidence support specific roles for Solanaceae components.

**Cardiovascular health:** Potassium-rich foods such as potatoes and tomatoes contribute to blood pressure regulation. Carotenoids and phenolic compounds reduce oxidative stress and improve lipid profiles in experimental and observational studies (Liu, 2013).

**Cancer prevention** Lycopene intake is inversely associated with prostate cancer risk in several meta-analyses, though heterogeneity exists across studies. Anthocyanin and other phenolic have shown chemo preventive properties in cellular and animal models; human evidence is suggestive but not conclusively causal.

**Metabolic effects** Capsaicin has been studied for thermogenic and appetite-modulating effects that could contribute modestly to energy balance and weight management, though long-term population-level effects remain uncertain (Zimmer, 2017).

**Digestive and micronutrient impacts** Dietary fiber from eggplant and potatoes supports gut motility and microbiome diversity. Moreover, processed tomato products enhance carotenoid uptake when consumed with dietary fats, thus improving micronutrient status.

**Sustainability, Post-Harvest Losses and Food Security** Sustainability considerations are central to understanding the role of Solanaceae vegetables in global food systems. Despite their nutritional and economic significance, the journey from farm to fork remains inefficient, with post-harvest losses undermining both food security and environmental sustainability.

**Asia's challenge: high losses despite high production.** Asia leads global production of potatoes, tomatoes, eggplants, and chilies, yet these gains are offset by losses ranging from 20–40% in some supply chains. Weak cold-chain networks, inadequate storage facilities, and fragmented smallholder systems contribute to these inefficiencies. The consequence is a paradox: while Asia ensures global availability of Solanaceae, local populations—particularly in low-income rural and urban areas—often face seasonal shortages and micronutrient deficiencies. Investments in cold-chain logistics, decentralized processing hubs, and farmer cooperatives could drastically reduce waste and enhance year-round access.

**Europe's contrast: efficiency with trade-offs.** Europe demonstrates comparatively lower post-harvest losses, owing to robust cold-chain systems, advanced packaging, and value-added processing. However, this efficiency often comes at a nutritional cost. The conversion of fresh produce into processed products—such as fried potato snacks or high-salt tomato sauces—extends shelf life and reduces waste but introduces dietary risks. Thus, Europe's challenge is not availability but healthfulness: the sustainability agenda must extend beyond food waste reduction to reformulating products for improved nutritional quality.

**Environmental footprint and climate pressures.** Both Asia and Europe face sustainability concerns tied to the environmental footprint of Solanaceae cultivation. Potatoes and tomatoes, in particular, are water-intensive crops. Climate change is already affecting yields through heat stress, pest pressures, and erratic rainfall. Sustainable intensification, including drip irrigation, integrated pest management, and the breeding of climate-resilient varieties, will be crucial to maintaining production while minimizing ecological costs.

**Policy and food system implications.** Reducing post-harvest losses in Asia has dual benefits: improving household food security and lowering greenhouse gas emissions from decomposing waste. In Europe, policy must shift towards nutrition-sensitive processing, with incentives for reformulated products (e.g., low-salt sauces, healthier frying methods) and clearer front-of-pack labeling. Globally, embedding Solanaceae crops into the *EAT–Lancet* framework of sustainable diets underscores the need to align production and consumption with planetary boundaries while addressing region-specific cultural practices.

**Critical gaps.** Current evidence on post-harvest losses often relies on generalized FAO estimates, which may underrepresent informal markets and smallholder losses in Asia. Similarly, sustainability assessments in Europe tend to prioritize efficiency metrics without fully accounting for health trade-offs in processed products. Addressing these gaps will require integrated food system research that combines agronomic, nutritional, and environmental dimensions.

## Discussion


The comparative analysis of Solanaceae vegetables in Asia and Europe reveals both their profound nutritional potential and the limitations of current dietary, processing, and policy practices. While potatoes, tomatoes, eggplants, and chilies are established dietary staples, the strength of evidence linking their consumption to specific health outcomes varies considerably across regions and study designs.

**Nutritional and health evidence.** Epidemiological findings suggest beneficial associations between Solanaceae intake and reduced risk of cardiovascular disease, certain cancers, and improved metabolic regulation. However, closer scrutiny reveals inconsistencies. Lycopene, for instance, is frequently associated with lower prostate cancer risk, but meta-analyses show heterogeneity and potential confounding by dietary patterns such as the Mediterranean diet. Similarly, capsaicin is widely studied for thermogenic and appetite-modulating effects, yet most evidence derives from short-term interventions; robust long-term trials remain limited. Anthocyanins from eggplant peel exhibit antioxidant properties in vitro, but human data remain scarce. Thus, while the nutritional promise of these vegetables is evident, translating mechanistic and observational findings into concrete public health recommendations requires caution.

**Regional contrasts.** Asia dominates in production volume, with China and India serving as global hubs for potato, tomato, and eggplant. Yet post-harvest losses, weak cold-chain infrastructure, and high reliance on fresh distribution constrain the net nutritional impact. In contrast, Europe emphasizes processing and

value addition—exemplified by tomato-based sauces and frozen potato products—which improves nutrient bioavailability (e.g., lycopene in processed tomato) but can also introduce negative dietary elements (e.g., excess salt, fat). These divergent systems highlight the trade-off between quantity and quality: Asia’s supply strength does not consistently translate to nutrient security, while Europe’s processing gains are sometimes offset by unhealthy formulations.

| Crop         | Asia (typical loss %) | Europe (typical loss %) | Key Drivers in Asia                             | Key Drivers in Europe                         |
|--------------|-----------------------|-------------------------|-------------------------------------------------|-----------------------------------------------|
| Potato       | 15–25%                | 5–10%                   | Inadequate cold storage, sprouting, bruising    | Overproduction glut, quality grading losses   |
| Tomato       | 25–40%                | 5–15%                   | Poor transport, crushing, lack of refrigeration | Processing efficiency vs. cosmetic standards  |
| Eggplant     | 20–35%                | 5–12%                   | High perishability, microbial spoilage          | Retail wastage due to consumer preference     |
| Chili Pepper | 20–30%                | 5–10%                   | Drying/storage inefficiencies, fungal attacks   | Minimal; efficient drying & packaging systems |



**Fig-1: An Illustrative Trend Chart (2000–2023) Showing Estimated Average Post-Harvest Losses for Solanaceae Vegetables in Asia Vs Europe**

**Policy and sustainability implications.** A critical gap in both regions lies in aligning agricultural productivity with nutrition-sensitive policies. Asia requires urgent investment in storage, transportation, and processing infrastructure to reduce post-harvest losses and enhance year-round availability. Europe must address the health risks of over-processed Solanaceae products through reformulation strategies, stricter labelling, and consumer education. Globally, breeding initiatives targeting nutrient density (e.g., high-lycopene tomatoes, low-glycoalkaloid potatoes) and climate-resilient cultivars could mitigate both health and sustainability challenges. Furthermore, both regions face the imperative of integrating

Solanaceae into broader sustainable diet frameworks, as articulated by the EAT–Lancet Commission, while also tailoring interventions to cultural and culinary contexts.

**Limitations of evidence.** Critical gaps remain in distinguishing between food supply and actual intake. FAOSTAT and other databases provide production and availability data, but do not account for retail waste, household losses, or preparation methods that shape nutrient delivery. Moreover, most health outcome studies aggregate fruits and vegetables rather than isolating Solanaceae-specific effects, which limits causal inference. Addressing these gaps requires more disaggregated dietary intake surveys, long-term cohort studies, and trials that reflect region-specific preparation methods.

## Conclusion

Solanaceae vegetables represent a cornerstone of diets in Asia and Europe, offering energy, micronutrients, and bioactive compounds with significant health-promoting potential. Yet, the evidence base is uneven: while mechanistic plausibility is strong, epidemiological and clinical data remain inconsistent. Maximizing the health and sustainability benefits of these crops requires policies that move beyond production metrics toward nutrition-sensitive food systems. In Asia, this means investing in post-harvest infrastructure and reducing losses; in Europe, reformulating processed products to retain benefits without adding risks. Bridging scientific uncertainties with targeted research, and embedding these crops within climate-resilient and culturally relevant dietary strategies, will be essential to realizing their full promise for global food and nutrition security.

## Reference:

1. Camire, M. E., Kubow, S., & Donnelly, D. J. (2009). Potatoes and human health. *Critical Reviews in Food Science and Nutrition*, 49(10), 823–840. <https://doi.org/10.1080/10408390903041996>
2. FAO. (2023). FAOSTAT statistical database. Food and Agriculture Organization of the United Nations. Retrieved from <https://www.fao.org/faostat>
3. Hosseini, F. (2011). Antioxidant properties of anthocyanins in eggplant. *Journal of Medicinal Plants Research*, 5(16), 3601–3605.
4. Liu, R. H. (2013). Health-promoting components of fruits and vegetables in the diet. *Advances in Nutrition*, 4(3), 384S–392S. <https://doi.org/10.3945/an.112.003517>
5. Rao, A. V., & Rao, L. G. (2007). Carotenoids and human health. *Pharmacological Research*, 55(3), 207–216. <https://doi.org/10.1016/j.phrs.2007.01.012>
6. Zimmer, A. R., Leonardi, B., Miron, D., Schapoval, E., de Oliveira, J. R., & Gosmann, G. (2012). Capsaicinoids: Pungency beyond Capsicum. *Molecules*, 17(1), 556–574. <https://doi.org/10.3390/molecules17010556>
7. Willett, W., Rockström, J., Loken, B., Springmann, M., et al. (2019). Food in the Anthropocene: the EAT–Lancet Commission on healthy diets from sustainable food systems. *The Lancet*, 393(10170), 447–492. [https://doi.org/10.1016/S0140-6736\(18\)31788-4](https://doi.org/10.1016/S0140-6736(18)31788-4)