



Plant Natural Products as Antimicrobials for Control of *Streptomyces scabies*: A Causative Agent of the Common Scab Disease

Justin Gutierrez, Amanda Bakke, Maritza Vatta and A. Rod Merrill*

Department of Molecular and Cellular Biology, University of Guelph, Guelph, ON, Canada

OPEN ACCESS

Edited by:

Daniel Pletzer,
University of Otago, New Zealand

Reviewed by:

Abdul Nasir,
Ajou University, South Korea
Felipe Lombó,
Universidad de Oviedo Mieres, Spain
Zhili Pang,
China Agricultural University, China

*Correspondence:

A. Rod Merrill
merrill@uoguelph.ca

Specialty section:

This article was submitted to
Infectious Agents and Disease,
a section of the journal
Frontiers in Microbiology

Received: 10 December 2021

Accepted: 24 December 2021

Published: 27 January 2022

Citation:

Gutierrez J, Bakke A, Vatta M and
Merrill AR (2022) Plant Natural
Products as Antimicrobials for Control
of *Streptomyces scabies*: A Causative
Agent of the Common Scab Disease.
Front. Microbiol. 12:833233.
doi: 10.3389/fmicb.2021.833233

The common scab disease caused by *Streptomyces scabies*, a soil-dwelling Gram-positive bacterium, is an economically important disease of potatoes and other tuber crops. The lack of effective treatments against this disease accounts for large economic losses globally. Plant extracts were screened to find several that effectively inhibited *Streptomyces scabies* growth in culture. Seven tinctures showed the greatest inhibition of *S. scabies* growth by reducing pathogen growth in culture by 75% or more. These extracts were myrrh, garlic, cayenne, barberry, frankincense, wild indigo root, and lavender. Myrrh extract from *Commiphora myrrha*, a resin made from tree sap, showed strong antibacterial activity by reducing the growth of *S. scabies* to 13% of the control. Additionally, a flavonoid library was screened to identify several compounds that were effective to control the pathogen growth. The flavonoids that showed the greatest inhibition of *Streptomyces scabies* growth were sophoraflavanone G, jaceosidin, baicalein, and quercetin. Minimum inhibitory concentrations for the effective flavonoids were calculated to be $6.8 \pm 0.4 \mu\text{M}$, $100.0 \pm 2.1 \mu\text{M}$, $202.9 \pm 5.3 \mu\text{M}$, and $285.2 \pm 6.8 \mu\text{M}$, respectively. The mean lethal doses for these flavonoids against *Streptomyces scabies* were $2.0 \pm 0.1 \mu\text{M}$, $22.6 \pm 0.5 \mu\text{M}$, $52.9 \pm 1.3 \mu\text{M}$, and $37.8 \pm 1.0 \mu\text{M}$, respectively. A live/dead assay showed complete cell death in the presence of sophoraflavanone G indicative of a bactericidal mechanism for flavonoid action on *Streptomyces scabies*. Scanning electron and transmission electron microscopy imaging showed damaged cell membrane morphologies when *Streptomyces scabies* was exposed to these flavonoids. Mycelia appeared as flat and deflated structures with contents seen as spewing from branching hyphae with numerous holes and tears in the membrane structure indicative of cell death. Sophoraflavanone G showed the greatest potency and potential as a natural antibiotic from the library of tested flavonoids. These results suggest that these plant compounds act on the pathogen through a bactericidal mechanism involving cell membrane destabilization and disruption leading to cell death.

Keywords: anti-bacterial agents, flavonoids, plant extracts, electron microscopy, *Streptomyces scabies*, natural products, scab disease

Abbreviations: DMSO, dimethyl sulfoxide; mART, mono-ADP-ribosyltransferase; LD₅₀, mean lethal dose that kills 50% of the cells; MB, methylene blue; MBA, methylene blue assay; MIC, minimal inhibitory concentration; MRSA, methicillin-resistant *Staphylococcus aureus*; PI, propidium iodide; SG, sophoraflavanone G.

INTRODUCTION

With the release of new antibiotics comes the inevitable, widespread development of resistance in target bacteria (Aslam et al., 2021). The development of resistance and other financial factors have resulted in a failure of new antibiotic discovery with only limited numbers of new compounds being developed each year (Tompson et al., 2021). This is not only an issue regarding human health, but it also affects agriculture practices (Zalewska et al., 2021). Antibiotic-resistant bacteria that survive in host plants and animals may also have downstream health effects on the humans that consume them. Clearly alternatives to traditional antibiotics are needed to assist in the supply of new antimicrobials to combat antibiotic resistance.

When developing antibiotic alternatives for use in agricultural settings, there are many factors that must be considered. In the case of application to crop plants, antibiotics must not only be effective, but they must also not affect the health of the plant or pollinators essential for plant reproduction (Cullen et al., 2019). Human health after consumption must also be considered. Possible alternatives to antibiotics with promising antibacterial activity are plant extracts and plant secondary metabolites.

Plant extracts contain many different classes of compounds, some of which have been shown to have antimicrobial activity (Khameneh et al., 2021). Classes of phytochemicals with antimicrobial activity include terpenes, alkaloids, and polyphenols such as flavonoids (Savoia, 2012). Flavonoids are a class of polyphenols characterized by a C₆-C₃-C₆ structure, typically in the form of two benzene rings connected by a pyran or pyranone ring—although variations to this chemical architecture do exist (Gonzales et al., 2015; Panche et al., 2016). Flavonoids can be found in many plant products including peppers, nuts, berries, wine, and teas, making them a common part of human diets (Panche et al., 2016). In plants, flavonoids play many important roles including UV filtration and protection, floral pigmentation, and cell signaling (Desmet et al., 2021). Most recently, flavonoids have been examined for their antibacterial activity in food preservation (Wu and Zhou, 2021). There are several ways in which flavonoids impart their antibacterial activity (Dias et al., 2021). In some cases, flavonoids can bind and inhibit important proteins in cells. For example, quercetin has strong antioxidant, antibacterial and antiparasite properties (Yang et al., 2020); it also binds specifically to GyrB, a subunit of *Escherichia coli* DNA gyrase, inhibiting its ATPase activity (Plaper et al., 2003). Quercetin also has strong antimicrobial potential against planktonic bacteria as well as disrupting biofilm formation of *Staphylococcus aureus*, *Pseudomonas aeruginosa* and *Candida albicans* (Zielinska et al., 2021).

While specific flavonoid-pathogen interactions exist, flavonoids are well known to affect the growth of a broad range of bacteria. This broad effect is likely due to flavonoid interactions with cell membranes (Farhadi et al., 2019; Agrawal et al., 2021). They are well known for their ability to inhibit bacterial efflux pumps and show promise in synergistic drug treatment in medical therapy (Waditzer and Bucar, 2021). Flavonoids have been shown to aggregate at the surface of the inner and outer leaflets of bacterial membranes and greatly reduce membrane

fluidity (Selvaraj et al., 2015; Yousefi et al., 2020). This, in turn, leads to a multitude of downstream effects that can induce cell death. Studies involving sophoraflavanone G (SG) and naringenin reveal that a reduction in bacterial membrane fluidity occurs at the minimum inhibitory concentrations (MIC) for the flavonoids (Tsuchiya and Iinuma, 2000; Veiko et al., 2020). Thus, reduced membrane fluidity is likely the main cause of the antibacterial activity of certain flavonoids (Zhu et al., 2020).

Many studies have tried to draw a connection between the structure of flavonoids and their antibacterial activity. For methicillin-resistant *Staphylococcus aureus* (MRSA), it has been demonstrated that 5,7-dihydroxylation of ring A on flavonoids of all classes was important for strong antibacterial activity (Tsuchiya et al., 1996; Yin et al., 2021). In flavanones specifically, 2',4'- and 4',6'-dihydroxylation on ring B have also been shown to impart strong antibacterial activity.

This antibacterial activity makes flavonoids important as natural alternatives to antibiotics for the prevention of bacterial infection of food products, due to their known edibility and strong effect on both Gram-positive and Gram-negative bacteria (Fernandez-Luqueno et al., 2021). Flavonoids also make excellent scaffolds for semisynthetic flavonoids showing heightened antimicrobial activity and enhanced solubility in water (Babii et al., 2018). Semisynthetic flavonoids can be readily varied in structure and quickly produced, thus accelerating new antibiotic discovery (Mottaghipisheh et al., 2021).

Flavonoids are especially attractive in the case of bacterial diseases of plants that are recalcitrant to treatment such as the common scab disease (CS). CS is a major disease of potatoes and tuberous plants world-wide (Li et al., 2019; Liu et al., 2021). CS is characterized by corky lesions, known as scabs, that appear on the surface of tubers during growth (Wei et al., 2021). These scabs can range from being superficial to deep and pitted and can cover the entire surface of a tuber, such as a potato, in some extreme cases (Loria et al., 2006; Wanner et al., 2014; Cruywagen et al., 2021).

Streptomyces is a large genus of actinobacteria with over 500 species (Barbutto Ferraiuolo et al., 2021). They are Gram-positive, spore forming, and are well known for their production of clinically significant secondary metabolites (Meanwell and Shama, 2008; Human et al., 2016). *Streptomyces* species are saprophytes that subsist on dead or decaying organic matter in soil or aquatic environments (Sharma et al., 2021). In soil, they act similarly to fungi forming branching mycelium on the surface (Urem et al., 2016). Of the over 500 species, only ten are known to be phytopathogenic. Of those species, the most prevalent species that cause CS in potatoes are *S. acidiscabies*, *S. europaescabiei*, *S. turgidiscabies*, and the most common species, *S. scabies* (Loria et al., 2006; Wanner et al., 2014).

Streptomyces scabies is characterized by the formation of spiral chains of smooth, gray spores (Lerat et al., 2009). *S. scabies* will usually infect potatoes between 2 and 5 weeks after tuber formation through naturally formed lesions on the surface of the growing tuber (Wanner et al., 2014). Current treatments of CS are not completely effective or without major drawbacks. One method of avoiding *S. scabies* infection is the use of resistant cultivars of potatoes such as the Norland, Russet, and Gold Rush pot (Ismail et al., 2020). However, resistant varieties may

suffer from drawbacks including lowered yield and processing quality. Another approach to deal with infection is to acidify the soil to below pH 5.2. While this does inhibit *S. scabies* growth, *S. acidiscabies* can thrive in low pH soils and cause CS regardless (Wanner et al., 2014; Hudec et al., 2021). Acidic soils also reduce the availability of some nutrients important to potato growth. Soil fumigation is also a viable route for clearing *S. scabies* infections (Larkin et al., 2011a). Fumigants, however, can be directly harmful to both the environment and workers that apply them to soils. Of all the current treatments, the most effective is crop rotation. While it does prevent loss of crop yield, the ability of *S. scabies* to produce spores means that it can survive in soils without a host for many years, so it can lead to a serious reduction in arable land for potato cultivation (Larkin et al., 2011b).

Given the drawbacks of current CS treatments, flavonoids offer an attractive alternative. Given current knowledge about flavonoids, they are likely to kill CS-causing bacteria without compromising the edibility or yield of the tubers being protected. In the present study, *S. scabies* was grown in the presence of a library of plant extracts and a flavonoid library to assess their ability to inhibit *S. scabies* growth. Confocal, SEM, and TEM imaging were also applied to characterize the effects of natural product treatment on the growth and morphology of *S. scabies*.

MATERIALS AND METHODS

Natural Product Library Screens

Culturing *Streptomyces scabies*

Streptomyces scabies strain 87.22 was used for all the following procedures. *S. scabies* was cultivated on nutrient agar [0.5% (w/v) tryptone, 0.3% (w/v) yeast extract, 0.5% (w/v) NaCl, and 1.5% (w/v) agar] at 30°C. To make glycerol stocks, a colony was inoculated into nutrient broth [0.5 w/v (%) tryptone, 0.3 w/v (%) yeast extract, and 0.5 w/v (%) NaCl], which was incubated for 2 days at 30°C and shaking at 250 rpm in an Erlenmeyer flask with a steel spring to shear the *S. scabies* filaments that are produced during growth. The culture was then mixed in a 1:1 ratio with 40% glycerol in a microfuge tube, flash frozen in liquid N₂, and kept at −80°C until use.

Methylene Blue Assay

The methylene blue assay (MBA) was originally reported (Fischer and Sawers, 2013) and was adapted herein for *S. scabies* growth measurements. Aliquots (1 mL for Erlenmeyer flask cultures, 50 µL for 24-well cultures) were taken from cultures at specific time points and were spun down for 7 min at 21,230 × g. The supernatant was removed, and then a volume of methylene blue (MB) (Thermo Fisher Scientific, MA, United States) equal to the volume of the original aliquot was added. The tube was then placed in a heat block set at 80°C with shaking at 850 rpm. The tube was then chilled on ice for 3 min before another spin at 21,230 × g for 3.5 min. The supernatant containing unabsorbed dye was transferred to a Corning 96-well, clear, flat bottom, polystyrene, sterile microplate (Corning Inc, New York City, NY, United States) at a 1/100 dilution. The absorbance at 660 nm was read using a FLUOstar Omega plate reader (BMG

LABTECH, Cary, NC, United States) and normalized against the absorbance of pure dye.

Plant Tincture Tests

A library of 56 plant tinctures (Perfect Herbs, Toronto, ON, Canada; Mond Trading CO., Toronto, ON, Canada; and Secrets of the Tribe, Nevada Pharm LLC., Las Vegas, NV, United States) in various percentages of ethanol were filtered through a sterile 0.2 µm nylon filter and tested against *S. Scabies* in growth medium. A glycerol stock was inoculated into 40 mL of nutrient broth and grown until an A_{660nm} of 0.18 was achieved. Then, 7 mL of the grown culture was inoculated into a new 40 mL nutrient broth flask containing the plant tincture being tested (tincture ethanol percentage did not exceed 2%). Two control flasks were used, one with 2% ethanol and one with only *S. scabies* culture added. The flasks were grown for 16 h at 30°C and 250 rpm and then processed using the MBA. This method was repeated with the seven tinctures that showed greatest inhibition at a 10-fold dilution of grown culture.

Streptomyces scabies Growth in the Presence of a Flavonoid Library

S. scabies starter cultures were grown in nutrient broth [0.5% (w/v) tryptone, 0.3% (w/v) yeast extract, 0.5% (w/v) NaCl] by inoculation with 700 µL of glycerol stock and were allowed to grow for 24 h. Two 1 mL aliquots were then taken from each starter culture and the MBA was used to determine the level of growth. Starter cultures that showed a change in A_{660nm} of 0.18 – 0.21 compared to a control dye were used to inoculate 24-well plates. For the flavonoid screening assay, a specific library of 20 flavonoids was obtained (ChemFaces Biochemical Co., Wuhan, China). Each flavonoid was stored at 4°C at 10 mM dissolved in 100% DMSO. Two additional phenylpropanoids, vanillic acid and caffeic acid, were obtained in solid form and stocks were made by dissolving the compounds in 100% DMSO to a concentration of 10 mM (Sigma-Aldrich, St. Louis, MO, United States). Twenty-four well plates were set up by adding 20 µL of the previously described *S. scabies* starter cultures and 10 µL of the 10 mM flavonoid stocks to nutrient broth to a final volume of 1 mL ([flavonoid] = 100 µM, [DMSO] = 1%). Each well also contained a small steel spring to shear the long filaments into smaller fragments. As a negative control, 10 µL of DMSO was added instead of 10 µL of flavonoid to maintain the 1% DMSO concentration present in the treatments. A positive control of novobiocin antibiotic was also used. To the positive control, 10 µL of 10 mM novobiocin in 100% DMSO was used. Each treatment and control were done in triplicate.

Once the plates were prepared with samples, each plate was shaken at 250 rpm at 30°C for 7 h. Afterward, 2 × 50 µL aliquots were taken from each well and the MBA was performed on each aliquot to determine the level of growth in each well. The A_{660nm} value of the novobiocin positive control was subtracted from the A_{660nm} of the other aliquots to normalize the results, then % growth of each treatment was determined by comparison to the negative control. The mean of the biological triplicates was reported ± standard deviation.

Flavonoid Growth Curves

Dimethyl Sulfoxide Tolerance of *Streptomyces scabies*

Streptomyces scabies starter cultures were prepared as previously described. Starter cultures with a $\Delta A_{660\text{ nm}}$ of 0.18–0.21 were used to inoculate 24-well cultures. Each well contained a steel spring, and 1 mL of culture containing 930 μL of nutrient broth and 20 μL *S. scabies* starter culture. The final 50 μL of solution was made up of a combination of sterile water and DMSO to make final concentrations of 0, 1, 2, 3, 4, and 5% (v/v) DMSO. Twenty-four well cultures were grown by shaking at 250 rpm at 30°C for 7 h. Two aliquots of 50 μL were taken for each well and the MBA was used to determine the growth in each culture samples. The mean of each biological triplicate $\pm$ standard deviation was then graphed.

Growth Inhibition Curve

Flavonoid stocks for SG, baicalein, and jaceosidin were prepared using powdered samples obtained from ChemFaces Biochemical Co. (Wuhan, China) dissolved in 100% DMSO. Stocks of 10 mM were prepared, and 1 mM stocks were produced via dilution. Quercetin stock was made using solid quercetin obtained from Sigma-Aldrich (St. Louis, MO, United States) dissolved in 100% DMSO, 10 mM and 1 mM stocks were also made in the same manner as the other stocks.

Streptomyces scabies starter cultures were prepared as previously described. Starter cultures with a $\Delta A_{660\text{ nm}}$ of 0.18–0.21 were used to inoculate 24-well cultures. Each well contained a steel spring for shearing filaments, and 1 mL of culture containing 940 μL of nutrient broth and 20 μL *S. scabies* starter culture. The final 40 μL were made up of a combination of 100% DMSO, and 10 mM or 1 mM flavonoid stocks in 100% DMSO to achieve a final DMSO concentration of 4% and the desired flavonoid concentration. A negative control using 40 μL of 100% DMSO and a positive control containing 30 μL of 100% DMSO and 10 μL of 10 mM novobiocin in 100% DMSO were also prepared for each 24-well plate. Twenty-four well cultures were then grown by shaking at 250 rpm at 30°C for 7 h. Two aliquots were taken for each well and the MBA was used to quantify the growth in each culture. The $A_{660\text{ nm}}$ value of the novobiocin positive control was subtracted from the $A_{660\text{ nm}}$ of the other aliquots to normalize the results, then the % growth of each treatment was determined by comparing the mean of triplicate data to the negative control $\pm$ standard deviation.

Growth curve analysis was calculated using OriginPro ver8 software (Northampton, MA, United States). Data obtained via flavonoid treatment was fit to a sigmoidal dose response model [% growth vs. log (flavonoid)] using instrumental weighting based on error calculated as $\pm$ standard deviation. From the growth curves, LD_{50} and MIC values were calculated. MIC is defined as the lowest concentration of an antimicrobial agent that inhibits the growth of a microorganism. To determine the MIC for each flavonoid, the steepest slope determined by the dose response model in OriginPro ver8 was extrapolated until it reached 0% growth.

The concentration of flavonoid at that point was the MIC for that flavonoid.

Live/Dead Assay

Cell viability was determined using the LIVE/DEAD® BacLight™ bacterial viability kit L7012 (Molecular Probes, Eugene, United States). *S. scabies* was grown overnight at 30°C with shaking at 250 rpm. One mL aliquots were taken for each treatment. Each aliquot was spun down at $10,000 \times g$. The supernatant was removed, and the pellet was then washed in filtered 0.85% (w/v) NaCl, and then resuspended in 0.85% NaCl (control 1), 1% DMSO (control 2), and 100 μM SG with 1% DMSO (treatment). Samples were incubated at room temperature for 1 h, mixing every 15 min. The samples were spun down again, and the pellets were washed twice with 0.85% NaCl and then resuspended in 1 mL of 0.85% NaCl.

The following steps were done in a dark room under red light. Component A and B from the viability kit were mixed 1:2. To each 1 mL sample, 6 μL of dye mixture was added and mixed thoroughly. Samples were then incubated at room temperature in the dark for 15 h. Samples were then washed once with 0.85% NaCl and then resuspended in 1 mL of 0.85% NaCl. Twenty μL of each sample was transferred to a microscope slide and covered with a glass cover slip.

Samples were then imaged using an inverted Leica DMi8 microscope (Leica, Wetzlar, Germany) connected to a Quorum Discovery Spinning Disk system (Quorum Technologies, ON, Canada). Images were taken with a 40 $\times$ long working distance (LWD) objective lens; the green channel was excited with a 488 nm laser, emission filter 525/50, and the orange channel was excited with a 560 nm laser, emission filter 600/50.

Streptomyces Scabies Imaging

Scanning Electron Microscopy

Seven hundred μL of *S. scabies* glycerol stock were used to inoculate 20 mL of nutrient broth in a covered Erlenmeyer flask

TABLE 1 | Effect of ethanol and DMSO solvents on the growth of *Streptomyces scabies* in culture.

Solvent	<i>S. scabies</i> growth (%)
None	100 $\pm$ 9
1.0% ethanol	113 $\pm$ 6
1.5% ethanol	100 $\pm$ 8
2.0% ethanol	108 $\pm$ 5
2.5% ethanol	109 $\pm$ 7
1.0% DMSO	105 $\pm$ 8
2.0% DMSO	100 $\pm$ 5
3.0% DMSO	84 $\pm$ 10
4.0% DMSO	82 $\pm$ 9

An *S. scabies* starter culture was grown to an A_{660} of 0.21, then 20 μL aliquots of the starter culture were used to inoculate 1 mL nutrient broth cultures. Cultures contained a gradient of ethanol or DMSO concentrations with triplicate replicates. Cultures were shaken at 250 rpm at 30°C for 7 h. Afterward, the level of growth was determined via the Methylene Blue assay (MBA). Each bar represents the mean of triplicate experiments $\pm$ standard deviation. None of the treatments showed a statistically significant reduction in growth (*p*-values, student *T*-test).

containing a steel spring for shearing filaments. To each flask, 100% DMSO and 10 mM stocks of a flavonoid of interest were added until each flask contained 4% DMSO and the desired concentration of the flavonoid. Treatments included MIC and 1/2 MIC concentrations of SG, baicalein, and jaceosidin, and a 1/2 MIC concentration of quercetin. Negative controls of *S. scabies* in the absence of DMSO and in the presence of 4% DMSO with no flavonoids were also prepared and imaged. The 20 mL cultures were then grown via shaking at 250 rpm at 30°C for 24 h.

One mL of each treatment was then taken and harvested by centrifuging at $1500 \times g$ for 4 min. Afterward, the samples were washed twice in 1 mL of SEM phosphate buffer (1:1 mix of 0.07 M K_2PO_4 dibasic and 0.07 M $NaHPO_4$ monobasic). Samples were then resuspended in 1 mL of SEM phosphate buffer and 200 μ L of each cell suspension was then placed on a polished carbon planchet. Cells were allowed to adhere to the planchet for 30 min. Planchets were then placed in a 12-well polystyrene plate and were submerged in 2% glutaraldehyde for 30 min. Planchets were then washed 3 times in 1 mL of SEM phosphate buffer. Planchets were then transferred to a 12 well porcelain plate and submerged in 1% osmium tetroxide (OsO_4) for 30 min. They were then washed again in SEM phosphate buffer. The planchets were then transferred back to the 12-well polystyrene plate and a dehydration series was set up using 50, 70, 80, 90, and 100% ethanol, washing 10 min with each, with the final 100% ethanol step repeated three times. Critical point drying was performed using a Denton DCP-1 critical point dryer (Denton Vacuum, NJ, United States). The samples were then sputter coated in gold using a Denton Desk V TSC (Denton Vacuum, NJ, United States). Images were taken via a Quanta 250 FEG SEM (FEI company, OR, United States).

Transmission Electron Microscopy

Erlenmeyer flasks were set up as previously described for the SEM prep. The only treatment prepared for TEM was a SG treatment at MIC. The same controls as for SEM were also prepared. The cultures were grown via shaking at 250 rpm at 30°C for 24 min. One mL of each treatment was taken and harvested by centrifugation at $6000 \times g$ for 4 h. The supernatant was discarded and 1 mL of TEM fix (2.5% glutaraldehyde, 2% paraformaldehyde (EM grade), and 0.05 M HEPES) was added. The samples were spun down again, and the pellet was then embedded in 2% noble agar on a glass slide. The newly embedded samples were then cut into small 1 mm $\times$ 1 mm squares and placed in a 1.5 mL centrifuge tube. The samples were then washed three times with 0.05 M HEPES. After moving the samples to a glass scintillation vial, 1% OsO_4 and the samples were incubated for 45 min. The OsO_4 was then removed, and the embedded samples were washed three times for 5 min each in MilliQ water. A dehydration series using 25, 50, 75, 95, and 100% ethanol was set up. The embedded samples were incubated in each ethanol mix for 5 min, with an additional 10-min incubation in 100% ethanol. Images were taken using an FEI Tecnai G2 F20 (FEI company, OR, United States).




FIGURE 1 | Plant extracts for control of *S. scabies* growth. **(A)** *S. scabies* growth in the presence of 45 plant ethanolic tinctures for 16 h at 30°C and 250 rpm as described in the section “Materials and Methods.” The samples tested were: 1, control; 2, ethanol treatment; 3, cedar (*Thuja occidentalis*); 4, hawthorne leaf (*Crataegus oxyacanthus*); 5, Madagascar periwinkle (*Catharanthus roseus*); 6, St. John's wart (*Hypericum perforatum*); 7, bearberry (*Hypericum perforatum*); 8, cinnamon (*Cinamomum zeylandicum*); 9, oregano (*Origanum vulgare*); 10, heal-all (*Prunella vulgaris*); 11, onion (*Allium cepa*); 12, burdock (*Arctium lappa*); 13, grape seed (*Vitis vinifera*); 14, peppermint (*Mentha X piperita*); 15, reishi mushroom (*Ganoderma lucidum*); 16, jackfruit (*Artocarpus heterophyllus*); 17, da ma hemp (*Cannabis sativa*); 18, wild Scotch pine tops (*Pinus sylvestris*); 19, anise seed (*Pimpinella anisum*); 20, Indian pennyroot (*Centella asiatica*); 21, wormwood (*Artemisia absinthum*); 22, rhubarb (*Rheum palmatum*); 23, lemon balm (*Melissa officinalis*); 24, aloe vera (*Aloe vera*); 25, black walnut (*Juglans nigra*); 26, oatstraw (*Avena sativa*); 27, basil leaf (*Ocimum basilicum*); 28, watercress (*Nasturtium officinale*); 29, periwinkle (*Vinca minor*); 30, tulsie (*Ocimum sanctum*); 31, old man's beard (*Ushua barbata*); 32, thyme (*Thymus vulgaris*); 33, dragon's blood (*Dracaena cinnabari* resin); 34, bitter sweet stalks (*Solanum dulcamara*); 35, mandrake (*Podophyllum peltatum*); 36, eucalyptus (*Eucalyptus globulus*); 37, licorice root (*Glycyrrhiza glabra*); 38, osha (*Ligusticum porteri*); 39, clove (*Syzygium aromaticum*); 40, goldenrod (*Hydrastis canadensis*); 41, hops (*Humulus lupulus*); 42, sage (*Salvia officinalis*); 43, goldenseal root (*Hydrastis canadensis*); 44, ginger root (*Zingiber officinalis*); 45, Korean Red ginseng (*Panax ginseng*); 46, lomatium (*Lomatium dissectum*); and 47, marigold (*Calendula officinalis*). The control featured no addition of ethanol to the *S. scabies* culture and ethanol treatment represents the control culture grown in the presence of 2.5% ethanol. **(B)** The seven most inhibitory plant tinctures against *S. scabies* growth. *S. scabies* was grown for 16 h at 30°C and 250 rpm as described in the section “Materials and Methods.” Each bar represents the mean of triplicate experiments $\pm$ standard deviation.

RESULTS

Ethanol and Dimethyl Sulfoxide Tolerance of *Streptomyces scabies*

Since the plant extracts were dissolved in ethanol and flavonoids were dissolved in DMSO, a tolerance test was conducted to gauge *S. scabies* tolerance to these two solvents. *S. scabies* showed no significant change in growth when exposed to up to 2.5% ethanol (Table 1). For further screening of plant extracts, a final ethanol concentration of 2% was used throughout. In 4% DMSO, there was a consistent 25% decrease in growth (Table 1). It was determined that the concentration of DMSO should not exceed 4% in all following experiments.

Streptomyces scabies Growth in Plant Extracts

Streptomyces scabies was grown in the presence of a total of 52 plant ethanolic tinctures. Tinctures showed a range of effects on *S. scabies* growth. Some tinctures inhibited growth, some showed little to no effect on growth, and some even promoted the growth of *S. scabies* (Figure 1A). The 7 tinctures that showed

the greatest inhibition of *S. scabies* growth were myrrh, garlic, cayenne, barberry, frankincense, wild indigo root, and lavender (Figure 1B). Myrrh extract from *Commiphora myrrha* is a resin made from tree sap native to the Arabian peninsula and to Africa. It showed strong antibacterial activity by reducing the growth of *S. scabies* to 13% of the control (Figure 1B).

Streptomyces scabies Growth in Flavonoid Library

Streptomyces scabies was grown in the presence of 20 flavonoids and two phenylpropanoids with proposed antimicrobial activity. The bacterium was exposed to each compound at 100 μ M and 1% DMSO. A positive control using novobiocin, an aminocoumarin potent antibiotic against *Streptomyces*, was used to find a baseline A_{660nm} that could be used to normalize A_{660nm} values from the inhibitor treatments. The positive control contained 1% DMSO and 100 μ M novobiocin (well above its effective concentration) to establish the 0% growth baseline. In the 22 compound library studied, all compounds exhibited antibiotic control over *S. scabies* growth, ranging from mild inhibition (vanillic acid, 7% inhibition) to SG (93% inhibition) (Figure 2). At 100 μ M, some compounds such as




FIGURE 2 | The effect of 100 μ M flavonoid on *S. scabies* growth. A *S. scabies* starter culture was grown to an A_{660nm} between 0.18 and 0.21, then aliquots of 20 μ L of the starter culture were used to inoculate 1 mL nutrient broth with 1% DMSO, 100 μ M of the flavonoid and a metal spring for shearing the filamentous *S. scabies* culture. Culture was grown by shaking at 250 rpm at 30°C for 7 h. Afterward, the level of growth was determined with the MBA. Each bar represents the mean of triplicate experiments $\pm$ standard deviation. The control represents *S. scabies* grown in 1% DMSO in the absence of any flavonoid compound. Each bar represents the mean of triplicate experiments $\pm$ standard deviation. Bars marked with an asterisk represent $p < 0.05$ for the group of datasets.


vanillic acid, showed a weak effect against *S. scabies* growth, others such as rutin and acacetin, showed a moderate effect of 20 – 50% reduction, and seven compounds (neobavaisoflavone, amentoflavone, quercetin, caffeic acid, jaceosidin, baicalein, and sophoraflavone G) showed strong inhibition of *S. scabies* growth (greater than 50% reduction) (Figure 2). It was decided that the five compounds with the highest antibacterial activity would be further characterized against *S. scabies*. The compounds chosen for further in-depth study included quercetin, caffeic acid, baicalein, jaceosidin, and SG.

Further testing of caffeic acid revealed that its antibacterial effect on *S. scabies* growth was dose independent and was most likely imparted by its natural acidity. Attempts to neutralize the

acidity of caffeic acid treatment resulted in solubility issues that made it impossible to analyze by the methods used in this work. Given that acidity-based treatments have previously been shown to be ineffective for treatment of CS in potatoes, caffeic acid was not studied further.

Growth Inhibition Dose-Response Curves

The four flavonoids that showed strong inhibition of *S. scabies* were further examined at various concentrations to determine both their LD₅₀ and MIC values. The dose-response curves are shown in Figure 3. SG was the most effective flavonoid at controlling *S. scabies* growth. Applying a dose-reponse model



(OriginPro ver8), the LD₅₀ of SG was shown to be 2.0 ± 0.1 μ M. Using the maximum slope from the dose-response model, an MIC for SG was also calculated at 6.8 ± 0.4 μ M (**Figure 3A**). Jaceosidin and baicalein also showed strong inhibition of *S. scabies* growth with an LD₅₀ of 22.6 ± 0.5 μ M and 52.9 ± 1.3 μ M, respectively. The MIC values for jaceosidin and baicalein were calculated as 100.0 ± 2.1 μ M and 202.9 ± 5.3 μ M, respectively (**Figures 3B,C**). Jaceosidin and baicalein showed some solubility issues around their MIC values, hindering the completion of full growth curves in their presence. Even with incomplete curves, however, enough data could be gathered to estimate the curve with reasonable confidence.

Quercetin showed some solubility issues that interfered with the acquisition of a full growth curve of *S. scabies* in its presence. This is due to both the high doses needed (due to lower potency) for full inhibition as well as its relatively low solubility. Data obtained from the quercetin growth curve were less reliable than

for SG, baicalein, or jaceosidin (**Figure 3D**). Using available data, LD₅₀ and MIC values for quercetin were determined at 37.8 ± 1.0 μ M and 285.2 ± 6.8 μ M, respectively. These values suggest that quercetin is a relatively strong inhibitor of *S. scabies* growth, although it requires a slightly higher dose than the other flavonoids.

Live/Dead Imaging

To determine if flavonoids acted upon *S. scabies* in a bactericidal or bacteriostatic manner, a BacLight™ bacterial viability kit was employed in conjunction with a confocal fluorescence microscope and a pinhole camera. Aliquots of *S. scabies* were incubated in either 0.85% NaCl, 0.85% NaCl + 4% DMSO, or 4% DMSO and 100 μ M of SG then incubated in SYTO9 green fluorescent nucleic acid stain and propidium iodide (PI). Images of *S. scabies* in the presence of no treatment or 4% DMSO, showed little to no cell death (green color filaments, **Figures 4A,B**).




FIGURE 4 | Confocal microscopy images of *S. scabies*. **(A)** *S. scabies* positive control, incubated in 0.85% NaCl. **(B)** *S. scabies* DMSO control, incubated in 0.85% NaCl and 1% DMSO. **(C)** *S. scabies* treatment, incubated in 0.85% NaCl, 100 μ M SG, and 1% DMSO. Green cells represent staining with SYTO 9 and red cells represent staining with PI. Images were taken with a 40 $\times$ objective lens. Each image is 22 μ m $\times$ 22 μ m.

However, when *S. scabies* was treated with SG above its MIC value, the images showed nearly complete cell death (red color filaments, **Figure 4C**). Images of the SG-treated cells showed large areas of green SYTO9-stained cells; however, when overlayed with the red PI-stained cells, it was evident that almost all green cells were also stained red, suggesting that the cells were non-viable (**Figure 4C**). Given that this level of cell death is not present in the positive control nor the 1% DMSO control, it can be concluded that SG is likely a bactericidal agent.

Electron Microscopy

Scanning Electron Microscopy

Scanning electron microscopy was conducted on *S. scabies* cells in the absence of DMSO, in 4% DMSO, and in 4% DMSO with a flavonoid at its MIC (SG, jaceosidin, or baicalein), and in 4% DMSO with a flavonoid of interest at 1/2 its MIC (SG jaceosidin, baicalein, or quercetin). Due to solubility issues with quercetin, *S. scabies* could only be imaged in the

presence of 1/2 MIC for this flavonoid. Both the negative control and 4% DMSO control showed smooth, consistent, and intact outer membranes (**Figures 5A,B**). One important difference noted for the DMSO control is that the budding branching hyphae present in the control images are less frequent in the presence of DMSO. The branching hyphae that are present also have a different shape, being more rounded (**Figure 5B**).

Streptomyces scabies cells in the presence of all four flavonoids showed significant membrane damage. In the case of cells exposed to SG, cells were severely malformed in both 1/2 MIC and MIC treatments, significantly different from the negative control. Cells showed rough, inconsistent membranes. Some cells appeared deflated and shriveled, while others were inflated and bulged. Some cells could be seen expelling cell contents into the extracellular space (**Figures 6A,B**).

In the presence of baicalein, many cells showed similar morphologies to SG-treated cells. As well as deflated cells, some




FIGURE 5 | SEM images of *S. scabies* grown in the absence of flavonoids. **(A)** *S. scabies* grown overnight in nutrient broth with a metal spring for shearing. **(B)** *S. scabies* grown overnight in nutrient broth plus 4% DMSO with a metal spring for shearing. Red arrows point to putative branching hyphae. Green arrow points to fluid filled vesicle. Images were taken via a Quanta 250 FEG SEM (FEI company, OR, United States).




FIGURE 6 | SEM images of *S. scabies* grown in the presence of SG. **(A,B)** *S. scabies* was grown overnight in nutrient broth with 4% DMSO and SG at its MIC with a metal spring for shearing. The red arrow points to hole in the membrane of a cell. The green arrow points to a cell spewing its contents. Images were taken via a Quanta 250 FEG SEM (FEI company, OR, United States).

cells also showed large, uneven growths protruding from their membranes. These growths appeared to be unique to baicalein treatment and were not seen in any of the other flavonoid treated cells (**Figures 7A,B**).



Cells exposed to jaceosidin showed similar morphologies to SG-treated cells; however, the phenotype appeared to be less severe overall (**Figures 8A,B**). For the most part, cells exposed to jaceosidin remained intact. Jaceosidin-treated cells showed slight membrane perturbations, no longer resembling the smooth membranes of the control cells. They were not, however, as warped or distorted as cells exposed to the other flavonoids tested. Some cells in 1/2 MIC jaceosidin showed holes to the mycelial membrane. Overall, in comparison to other flavonoids, jaceosidin showed less damage to membranes (**Figure 8**).


While quercetin could not be used at its MIC due to solubility issues, treatment of cells at 1/2 MIC still showed significant membrane damage (**Figures 9A,B**). Like SG-treated cells, quercetin-damaged cells could be seen expelling cell contents into the extracellular space. There were also some

large tears seen in the mycelial membranes (**Figure 9B**, green arrows). While higher concentrations of quercetin were needed than any of the other flavonoids, its strong effect on cells was clearly evident.

Transmission Electron Microscopy

Transmission electron microscopy was also performed on cells in the presence of SG at MIC. The DMSO control and flavonoid treatment images were conducted in 4% DMSO to keep SG soluble in cell media and to provide consistency with the previous SEM images. Positive control cells showed healthy and intact cell membranes (**Figures 10A,B**). Cells also had consistent shape and cytoplasm density. There were some electron-invisible voids seen in the cytoplasm. These voids likely represent common artifacts caused by chemical fixation by glutaraldehyde and OsO₄ which was used in the preparation of these images. Cells exposed to SG at MIC showed significant damage compared to positive control cells (**Figures 10C,D**). The cells showed large clearings in their cytoplasm and inconsistent cytoplasm density in areas that were not voided completely. The perturbation of the cytoplasm





is likely due to the effect of flavonoids on cell membranes as shown in SEM imaging. The thickness of membranes also became variable upon flavonoid treatment (Figures 10C,D). Some cells had large bulges in their membranes when exposed to flavonoids (Figures 10C,D, red arrows). These bulges were not present in any cells exposed only to 4% DMSO suggesting that SG treatment is the main reason for these bulges.

DISCUSSION

Common scab disease is a major global disease of potatoes. The primary cause of CS in North America is *Streptomyces scabies*. *S. scabies* infects potato tubers 2–5 weeks after tuber initiation and causes scabs on the surface of the tubers at 6 weeks of growth. Scabs can range from superficial to deep and pitted, causing major economic losses globally. Current treatments are not effective at treating CS and suffer disadvantages that affect their utility in treating the disease. As such, the effectiveness of natural products against *S. scabies* has been investigated as an alternative to traditional treatments. Natural products are an attractive choice of treatment, especially in the food industry, as they generally do not affect the edibility of food

products. If the right natural products are used, they should be safely consumed by humans. The complex nature of plant extracts made it impossible to attribute antibacterial activity to any specific metabolites. Narrowing antibacterial activity to a single metabolite can allow for semisynthetic metabolites to be developed, enhancing their activity.

In the present study, *S. scabies* growth was measured in the presence of flavonoids, a class of plant metabolites known for their antibacterial activity. *S. scabies* was initially grown in various concentrations of DMSO to ensure that the solvent which all flavonoids were dissolved in did not significantly affect *S. scabies* growth. It was decided that the maximum DMSO concentration that could be used was 4%, as it did not drastically affect *S. scabies* growth. *S. scabies* was then grown in the presence of a library of 22 flavonoids. Some flavonoids showed little to no effect on *S. scabies* growth, some showed a modest decrease in growth, and some showed significant inhibition of *S. scabies* growth. The four most effective flavonoids at controlling *S. scabies* growth were SG, jaceosidin, baicalein, and quercetin.

Full growth inhibition curves were obtained for the four most effective flavonoids and LD₅₀ and MIC values were determined for each one. The LD₅₀ values of the flavonoids were determined to be 2.0 μ M (SG), 23 μ M (jaceosidin), 53 μ M (baicalein) and 38 μ M (quercetin). The MIC values of the flavonoids were calculated to be 7 μ M (SG), 100 μ M (jaceosidin), 203 μ M (baicalein), and 285 μ M (quercetin). These values were calculated to a high degree of confidence for SG, jaceosidin, and baicalein; however, the data for quercetin were calculated with less confidence as it was not fully soluble at higher concentrations in 4% DMSO.

A live/dead assay was conducted on *S. scabies* cells incubated in high concentrations of SG. Cells incubated in SG showed near total cell death in contrast to the control images. Based on these images, it is likely that the antibacterial effect of SG against *S. scabies* is mainly bactericidal. Based on the morphology of cells exposed to SG as imaged by SEM, it is also likely that jaceosidin, baicalin, and quercetin also exhibit bactericidal activity against *S. scabies*. TEM imaging of SG-treated cells showed significant perturbation (void formation) of the cytoplasm in cells exposed to SG.

The SEM and TEM images of *S. scabies*-treated cells showed that the four flavonoids under investigation caused significant damage to cell membranes. This was not surprising given the previous reports in the literature (Zhang L. L. et al., 2020; Zhang S. et al., 2020; Babii et al., 2021; Yuan et al., 2021). The current theory for flavonoid-membrane interactions postulates that flavonoids intercalate into membrane bilayers and shift the gel-to-liquid crystal transition temperature of the phospholipids to a more rigid, gel-like state. This, in turn, affects normal cell membrane function and cell-associated cell processes involving membrane protein-assisted transport, ion channel function, integral membrane protein stability, dynamics and function, inhibition of the respiratory chain, disruption of ATP synthesis and membrane potential control. The holes and tears in mycelial membranes observed upon flavonoid treatment suggests that flavonoids may also act as detergents to aggregate membrane components, dissolve the bilayer, and disrupt the structural integrity of the target bacterial membrane leading to cell death.




FIGURE 10 | TEM images of *S. scabies*. **(A)** *S. scabies* positive control grown overnight in nutrient broth. **(B)** *S. scabies* grown overnight in the presence of 4% DMSO **(C,D)** *S. scabies* grown overnight in the presence of 4% DMSO and SG at MIC. Red arrows indicate bulges in the cell membrane of SG treated cells. Images were taken using an FEI Tecnai G2 F20 TEM (FEI company, OR, United States).

CONCLUSION

A library of plant cell extracts (ethanolic tinctures) was tested as potential growth inhibitors of the plant pathogen, *S. scabies*. Seven tinctures inhibited *S. scabies* growth by 75% or more including myrrh, garlic, cayenne, barberry, frankincense, wild indigo root and lavender. In addition, a flavonoid library was screened, and several compounds were shown to effectively control *S. scabies* growth in culture with the best inhibitors being quercetin, jaceosidin, baicalein and SG. A bactericidal mechanism against *S. scabies* growth in culture was revealed for these flavonoids. SG was shown as the most potent flavonoid with LD₅₀ and MIC values of 2.0 and 6.8 μM, respectively. The molecular mechanism of cell death by these flavonoids was revealed by SEM and TEM which showed obvious damage to mycelia membranes of *S. scabies* with large holes, gashes, and significant reduction in cell density (contents). Cytoplasmic contents were seen to leak from mycelial structures with hyphae showing a flattened morphology indicative of the loss of cell contents to the extracellular medium.

In summary, these flavonoids show great promise as alternative, natural-product antibiotics to control *S. scabies*

growth in culture and potentially to treat plant-based disease caused by this pathogen.

DATA AVAILABILITY STATEMENT

The original contributions presented in the study are included in the article/supplementary material, further inquiries can be directed to the corresponding author.

AUTHOR CONTRIBUTIONS

ARM: conceptualization, funding acquisition, and writing—original draft. JG, AB, MV, and ARM: writing—review and editing, and visualization. All authors have read and agreed to the published version of the manuscript.

FUNDING

This work was supported by the Natural Sciences and Engineering Research Council of Canada, Discovery grant (#105440), and Strategic grant (#494213) to ARM.

REFERENCES

- Agrawal, S., Nandeibam, J., and Sarangthem, I. (2021). Ultrastructural changes in methicillin-resistant *Staphylococcus aureus* (MRSA) induced by metabolites of thermophilous fungi *Acrophialophora levis*. *PLoS One* 16:e0258607. doi: 10.1371/journal.pone.0258607
- Aslam, B., Khurshid, M., Arshad, M. I., Muzammil, S., Rasool, M., Yasmeen, N., et al. (2021). Antibiotic resistance: one health one world outlook. *Front. Cell. Infect. Microbiol.* 11:771510. doi: 10.3389/fcimb.2021.771510
- Babii, C., Mihalache, G., Bahrin, L. G., Neagu, A. N., Gostin, I., Mihai, C. T., et al. (2018). A novel synthetic flavonoid with potent antibacterial properties: in vitro activity and proposed mode of action. *PLoS One* 13:e0194898. doi: 10.1371/journal.pone.0194898
- Babii, C., Savu, M., Motrescu, I., Birsa, L. M., Sarbu, L. G., and Stefan, M. (2021). The antibacterial synthetic flavonoid BrCl-Flav exhibits important anti-candida activity by damaging cell membrane integrity. *Pharmaceuticals (Basel)* 14:1130. doi: 10.3390/ph14111130
- Barbuto Ferraiuolo, S., Cammarota, M., Schiraldi, C., and Restaino, O. F. (2021). Streptomycetes as platform for biotechnological production processes of drugs. *Appl. Microbiol. Biotechnol.* 105, 551–568. doi: 10.1007/s00253-020-11064-2
- Cruywagen, E. M., Pierneef, R. E., Chauke, K. A., Nkosi, B. Z., Labeda, D. P., and Cloete, M. (2021). Major *Streptomyces* species associated with fissure scab of potato in South Africa including description of *Streptomyces solaniscabiei* sp. nov. *Antonie Van Leeuwenhoek* 114, 2033–2046. doi: 10.1007/s10482-021-01659-8
- Cullen, M. G., Thompson, L. J., Carolan, J. C., Stout, J. C., and Stanley, D. A. (2019). Fungicides, herbicides and bees: a systematic review of existing research and methods. *PLoS One* 14:e0225743. doi: 10.1371/journal.pone.0225743
- Desmet, S., Morreel, K., and Dauwe, R. (2021). Origin and function of structural diversity in the plant specialized metabolome. *Plants (Basel)* 10:2393. doi: 10.3390/plants10112393
- Dias, M. C., Pinto, D., and Silva, A. M. S. (2021). Plant flavonoids: chemical characteristics and biological activity. *Molecules* 26:5377. doi: 10.3390/molecules26175377
- Farhadi, F., Khameneh, B., Iranshahi, M., and Iranshahi, M. (2019). Antibacterial activity of flavonoids and their structure-activity relationship: an update review. *Phytother. Res.* 33, 13–40. doi: 10.1002/ptr.6208
- Fernandez-Luqueno, F., Medina-Perez, G., Perez-Soto, E., Espino-Manzano, S., Peralta-Adauto, L., Perez-Rios, S., et al. (2021). Bioactive compounds of *Opuntia* spp. acid fruits: micro and nano-emulsified extracts and applications in nutraceutical foods. *Molecules* 26:6429. doi: 10.3390/molecules26216429
- Fischer, M., and Sawers, R. G. (2013). A universally applicable and rapid method for measuring the growth of streptomycetes and other filamentous microorganisms by methylene blue adsorption-desorption. *Appl. Environ. Microbiol.* 79, 4499–4502. doi: 10.1128/AEM.00778-13
- Gonzales, G. B., Smaghe, G., Grootaert, C., Zotti, M., Raes, K., and Van Camp, J. (2015). Flavonoid interactions during digestion, absorption, distribution and metabolism: a sequential structure-activity/property relationship-based approach in the study of bioavailability and bioactivity. *Drug Metab. Rev.* 47, 175–190. doi: 10.3109/03602532.2014.1003649
- Hudec, C., Biessy, A., Novinscak, A., St-Onge, R., Lamarre, S., Blom, J., et al. (2021). Comparative genomics of potato common scab-causing *Streptomyces* spp. displaying varying virulence. *Front. Microbiol.* 12:716522. doi: 10.3389/fmicb.2021.716522
- Human, Z. R., Moon, K., Bae, M., De Beer, Z. W., Cha, S., Wingfield, M. J., et al. (2016). Antifungal *Streptomyces* spp. Associated with the Infructescences of *Protea* spp. in South Africa. *Front. Microbiol.* 7:1657. doi: 10.3389/fmicb.2016.01657
- Ismail, S., Jiang, B., Nasimi, Z., Inam-Ul-Haq, M., Yamamoto, N., Danso Ofori, A., et al. (2020). Investigation of *Streptomyces scabies* causing Potato scab by various detection techniques, its pathogenicity and determination of host-disease resistance in potato Germplasm. *Pathogens* 9:760. doi: 10.3390/pathogens9090760
- Khameneh, B., Eskin, N. A. M., Iranshahi, M., and Fazly Bazzaz, B. S. (2021). Phytochemicals: a promising weapon in the arsenal against antibiotic-resistant bacteria. *Antibiotics (Basel)* 10:1044. doi: 10.3390/antibiotics10091044
- Larkin, R. P., Honeycutt, C. W., Griffin, T. S., Olanya, O. M., Halloran, J. M., and He, Z. (2011a). Effects of different potato cropping system approaches and water management on soilborne diseases and soil microbial communities. *Phytopathology* 101, 58–67. doi: 10.1094/PHYTO-04-10-0100
- Larkin, R. P., Honeycutt, C. W., and Olanya, O. M. (2011b). Management of *Verticillium* Wilt of Potato with disease-suppressive green manures and as affected by previous cropping history. *Plant Dis.* 95, 568–576. doi: 10.1094/PDIS-09-10-0670
- Lerat, S., Simao-Beaunoir, A. M., and Beaulieu, C. (2009). Genetic and physiological determinants of *Streptomyces scabies* pathogenicity. *Mol. Plant. Pathol.* 10, 579–585. doi: 10.1111/j.1364-3703.2009.00561.x
- Li, Y., Liu, J., Diaz-Cruz, G., Cheng, Z., and Bignell, D. R. D. (2019). Virulence mechanisms of plant-pathogenic *Streptomyces* species: an updated review. *Microbiology (Reading)* 165, 1025–1040. doi: 10.1099/mic.0.000818
- Liu, J., Nothias, L. F., Dorrestein, P. C., Tahlan, K., and Bignell, D. R. D. (2021). Genomic and metabolomic analysis of the potato common scab pathogen *Streptomyces scabiei*. *ACS Omega* 6, 11474–11487. doi: 10.1021/acsomega.1c00526
- Loria, R., Kers, J., and Joshi, M. (2006). Evolution of plant pathogenicity in *Streptomyces*. *Annu. Rev. Phytopathol.* 44, 469–487.
- Meanwell, R. J., and Shama, G. (2008). Production of streptomycin from chitin using *Streptomyces griseus* in bioreactors of different configuration. *Bioresour. Technol.* 99, 5634–5639. doi: 10.1016/j.biortech.2007.10.036
- Mottaghipishheh, J., Taghrir, H., Boveiri Dehsheikh, A., Zomorodian, K., Irajie, C., Mahmoodi Sourestani, M., et al. (2021). Linarin, a glycosylated flavonoid, with potential therapeutic attributes: a comprehensive review. *Pharmaceuticals (Basel)* 14:1104. doi: 10.3390/ph14111104
- Panche, A. N., Diwan, A. D., and Chandra, S. R. (2016). Flavonoids: an overview. *J. Nutr. Sci.* 5:e47.
- Plaper, A., Golob, M., Hafner, I., Oblak, M., Solmajer, T., and Jerala, R. (2003). Characterization of quercetin binding site on DNA gyrase. *Biochem. Biophys. Res. Commun.* 306, 530–536. doi: 10.1016/s0006-291x(03)01006-4
- Savoia, D. (2012). Plant-derived antimicrobial compounds: alternatives to antibiotics. *Future Microbiol.* 7, 979–990. doi: 10.2217/fmb.12.68
- Selvaraj, S., Krishnaswamy, S., Devashya, V., Sethuraman, S., and Krishnan, U. M. (2015). Influence of membrane lipid composition on flavonoid-membrane interactions: implications on their biological activity. *Prog. Lipid Res.* 58, 1–13. doi: 10.1016/j.plipres.2014.11.002
- Sharma, V., Kaur, R., and Salwan, R. (2021). *Streptomyces*: host for refactoring of diverse bioactive secondary metabolites. *3 Biotech* 11, 340. doi: 10.1007/s13205-021-02872-y
- Tompson, A. C., Manderson, L., and Chandler, C. I. R. (2021). Understanding antibiotic use: practices, structures and networks. *JAC Antimicrob. Resist.* 3:dlab150. doi: 10.1093/jacamr/dlab150
- Tsuchiya, H., and Iinuma, M. (2000). Reduction of membrane fluidity by antibacterial sophoraflavanone G isolated from *Sophora exigua*. *Phytomedicine* 7, 161–165. doi: 10.1016/S0944-7113(00)80089-6
- Tsuchiya, H., Sato, M., Miyazaki, T., Fujiwara, S., Tanigaki, S., Ohyama, M., et al. (1996). Comparative study on the antibacterial activity of phytochemical flavanones against methicillin-resistant *Staphylococcus aureus*. *J. Ethnopharmacol.* 50, 27–34. doi: 10.1016/0378-8741(96)85514-0
- Urem, M., Van Rossum, T., Bucca, G., Moolenaar, G. F., Laing, E., Swiatek-Polatynska, M. A., et al. (2016). OsdR of *Streptomyces coelicolor* and the Dormancy Regulator DevR of *Mycobacterium tuberculosis* control overlapping regulons. *mSystems* 1:e00014-16. doi: 10.1128/mSystems.00014-16
- Veiko, A. G., Sekowski, S., Lapshina, E. A., Wilczewska, A. Z., Markiewicz, K. H., Zamaraeva, M., et al. (2020). Flavonoids modulate liposomal membrane structure, regulate mitochondrial membrane permeability and prevent erythrocyte oxidative damage. *Biochim. Biophys. Acta Biomembr.* 1862:183442. doi: 10.1016/j.bbamem.2020.183442
- Waditzer, M., and Bucar, F. (2021). Flavonoids as inhibitors of bacterial efflux pumps. *Molecules* 26:6904. doi: 10.3390/molecules26226904
- Wanner, L. A., Kirk, W. W., and Qu, X. S. (2014). Field efficacy of nonpathogenic *Streptomyces* species against potato common scab. *J. Appl. Microbiol.* 116, 123–133. doi: 10.1111/jam.12336
- Wei, Q., Li, J., Yang, S., Wang, W., Min, F., Guo, M., et al. (2021). *Streptomyces rhizophilus* causes potato common scab disease. *Plant Dis.* doi: 10.1094/PDIS-09-20-1902-RE [Epub ahead of print].
- Wu, Q., and Zhou, J. (2021). The application of polyphenols in food preservation. *Adv. Food Nutr. Res.* 98, 35–99. doi: 10.1016/bs.afnr.2021.02.005

- Yang, D., Wang, T., Long, M., and Li, P. (2020). Quercetin: its main pharmacological activity and potential application in clinical medicine. *Oxid. Med. Cell. Longev.* 2020:8825387. doi: 10.1155/2020/8825387
- Yin, N., Yang, X., Wang, L., Zhang, C., Guan, J., Tao, Y., et al. (2021). Kaempferol inhibits the expression of alpha-hemolysin and protects mice from methicillin-resistant *Staphylococcus aureus*-induced lethal pneumonia. *Microb. Pathog.* 162:105336. doi: 10.1016/j.micpath.2021.105336
- Yousefi, M., Khorshidian, N., and Hosseini, H. (2020). Potential application of essential oils for mitigation of *Listeria monocytogenes* in meat and poultry products. *Front. Nutr.* 7:577287. doi: 10.3389/fnut.2020.577287
- Yuan, G., Guan, Y., Yi, H., Lai, S., Sun, Y., and Cao, S. (2021). Antibacterial activity and mechanism of plant flavonoids to gram-positive bacteria predicted from their lipophilicities. *Sci. Rep.* 11:10471. doi: 10.1038/s41598-021-90035-7
- Zalewska, M., Blazejewska, A., Czapko, A., and Popowska, M. (2021). Antibiotics and antibiotic resistance genes in animal manure – consequences of its application in agriculture. *Front. Microbiol.* 12:610656. doi: 10.3389/fmicb.2021.610656
- Zhang, L. L., Zhang, L. F., and Xu, J. G. (2020). Chemical composition, antibacterial activity and action mechanism of different extracts from hawthorn (*Crataegus pinnatifida* Bge.). *Sci. Rep.* 10:8876. doi: 10.1038/s41598-020-65802-7
- Zhang, S., Hu, B., Xu, J., Ren, Q., Wang, Z., Wang, S., et al. (2020). Baicalin suppress growth and virulence-related factors of methicillin-resistant *Staphylococcus aureus* in vitro and vivo. *Microb. Pathog.* 139:103899. doi: 10.1016/j.micpath.2019.103899
- Zhu, H. L., Chen, G., Chen, S. N., Wang, R. Q., Chen, L., Xue, H., et al. (2020). Changes in cell membrane properties and phospholipid fatty acids of *Bacillus subtilis* induced by polyphenolic extract of *Sanguisorba officinalis* L. *J. Food Sci.* 85, 2164–2170. doi: 10.1111/1750-3841.15170
- Zielinska, S., Dziagwa-Becker, M., Junka, A., Piatczak, E., Jezierska-Domaradzka, A., Brozyna, M., et al. (2021). Screening Papaveraceae as novel antibiofilm natural-based agents. *Molecules* 26:4778. doi: 10.3390/molecules26164778

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's Note: All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2022 Gutierrez, Bakke, Vatta and Merrill. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.