



Toxicity Assessment and Acetic Acid Content of Vinegar from Date Palm and Grape Fruits

Obianom, A. O.¹, Iheukwumere, I. H.², Iheukwumere, C. M.¹, Ike, V. E.³, Ezendianefo, J. N.⁴, Okongwu, D. J.⁵ and Abba, O.⁶

¹Department of Applied Microbiology and Brewing, Faculty of Biosciences, Nnamdi Azikiwe University, Awka, Nigeria.

²Department of Microbiology, Faculty of Natural Sciences, Chukwuemeka Odumegwu Ojukwu University, Uli, Campus, Anambra State, Nigeria.

³Department of Microbiology, University of Agriculture and Environmental Sciences, Umuagwo, Imo State, Nigeria.

⁴Department of Microbiology, Tansian University, Umunya, Anambra State, Nigeria.

⁵Department of Chemistry, Nwafor Orizu College of Education, Nsugbe.

⁶Department of Microbiology, Federal University of Gusau, Zamfara State.

*Corresponding author email: ao.obianom@unizik.edu.ng / ik.iheukwumere@coou.edu.ng

Abstract	Article History
While vinegar is broadly considered safe for consumption, its specific safety profile, including acetic acid concentration and potential toxicological effects, is contingent upon the raw material and production parameters. This study aimed to conduct a comprehensive safety evaluation of vinegar produced from <i>Phoenix dactylifera</i> (date palm) and <i>Vitis vinifera</i> (grape) by quantifying its acetic acid content and assessing subchronic toxicity <i>in vivo</i> . Essential fermentative microorganisms, <i>Saccharomyces cerevisiae</i> strain SR 128 and <i>Acetobacter aceti</i> strain WI, were isolated from spoiled fruits and utilized for vinegar production via submerged fermentation of the respective fruit musts. Acetic acid concentration was determined instrumentally, and a 28-day oral toxicity study was performed using a rat model to evaluate effects on body weight, relative organ weights, and key serum biochemical markers. Data were analyzed using Analysis of Variance (ANOVA) with Tukey's post-hoc test. The acetic acid content of the grape and date vinegars was determined to be 4.10% and 5.20%, respectively. The toxicological assessment revealed no significant adverse effects; no statistically significant differences ($p > 0.05$) were observed in body weight, organ weights, or clinical biochemistry parameters between the treatment and control groups. In conclusion, vinegars produced from <i>Phoenix dactylifera</i> and <i>Vitis vinifera</i> contain acceptable levels of acetic acid and demonstrate no evidence of toxicity in the subchronic rodent model, supporting their safety for human consumption. The date palm vinegar, characterized by a higher acetic acid yield, exhibited marginally superior compositional attributes.	Received: 27 Dec 2025 Accepted: 11 Feb 2026 Published: 18 Feb 2026
	Scan QR code to view*  License: CC BY 4.0*  Open Access article.
Keywords: Vinegar, Toxicity, <i>Phoenix</i> , <i>Vitis</i> , <i>Saccharomyces</i> , <i>Acetobacter</i>	
How to cite this paper: Obianom, A. O., Iheukwumere, I. H., Iheukwumere, C. M., Ike, V. E., Ezendianefo, J. N., Okongwu, D. J., & Abba, O. (2026). Toxicity assessment and acetic acid content of vinegar from date palm and grape fruits. <i>IPS Journal of Toxicology</i> , 4(1), 135–147. https://doi.org/10.54117/qrcgjj55	

Introduction

Vinegar, a sour liquid traditionally derived from fermented alcoholic substrates, is valued both as a culinary ingredient and a functional food component with emerging health implications (Ho *et al.*, 2022;

Iheukwumere *et al.*, 2025a, Diim *et al.*, 2025a). Produced via the microbial conversion of ethanol to acetic acid, vinegar can be sourced from sugar-rich fruits such as dates (*Phoenix dactylifera*) and grapes (*Vitis vinifera*), each imparting distinct sensory and bioactive

characteristics (Bakir *et al.*, 2017; Ekechukwu *et al.*, 2025a; Iheukwumere *et al.*, 2025b). Acetic acid is the primary functional constituent, responsible for vinegar's preservative and antimicrobial properties (Poojary *et al.*, 2017; Amadi *et al.*, 2017; Ejike *et al.*, 2017). Beyond its traditional uses, vinegar is increasingly investigated for potential physiological benefits, including antioxidant, hypoglycemic, and cardioprotective effects (Samad *et al.*, 2016; Xia *et al.*, 2020; Obianom *et al.*, 2024; Dim *et al.*, 2025b).

The fermentation process typically involves an initial alcoholic phase mediated by yeasts such as *Saccharomyces cerevisiae*, followed by acetification carried out by acetic acid bacteria including *Acetobacter aceti* (Trček *et al.*, 2015). This results in a product with a low pH, which ensures microbial stability but also raises important safety considerations related to acidity (Chen *et al.*, 2021; Iheukwumere *et al.*, 2022a). Although vinegar is generally regarded as safe for consumption, elevated acetic acid concentrations or excessive intake may pose risks such as dental enamel erosion and gastrointestinal mucosal irritation (Paranjpe *et al.*, 2013; Dutta *et al.*, 2018). Consequently, the safety profile of vinegar must be evaluated through precise measurement of acetic acid levels and appropriate toxicological screening.

Despite the commercial prevalence of grape vinegar and the growing interest in date vinegar for its unique phytochemical content (Nwike *et al.*, 2017; Al-Farsi *et al.*, 2018; Dim *et al.*, 2025c), comparative safety data between these two fruit-derived vinegars remain limited. Existing toxicity studies often focus on generic vinegar formulations or high-concentration acetic acid, leaving a gap in substrate-specific safety evaluations (Budak *et al.*, 2014; Özen *et al.*, 2021). Given that raw material composition can alter fermentation dynamics and the final product composition, safety assessments should be tailored to specific vinegar types. Therefore, this study aims to perform a comparative safety assessment of vinegar produced from *Phoenix dactylifera* and *Vitis vinifera*.

Materials and Methods

Isolation and Characterization of *Saccharomyces* species from Spoilt Fruit Samples

Sample collection

Spoilt *Phoenix dactylifera* (Date palm) and *Vitis vinifera* (Grape) fruits were collected from different points in Nkwo Oba market, Idemili South LGA, Anambra State. The fruits were detected through sight and nasal perception; this was followed by carefully and selectively picking of the detected fruits into polythene bags. The polythene bags were appropriately labeled and

transported immediately to the laboratory for further analysis.

Sample preparation

The fruit samples were thoroughly washed using distilled water and their ectocarps were appropriately peeled using stainless chicken knife. The peeled fruits were pulverized using electric blender (SMX425/Japan). This was serially diluted (1:10) using 250 mL conical flask (Pyrex) in the capacity of 10 g of the fruit sample to make up 200 mL of the sample solution. The solution was thorough shaken, stoppered and kept for further analysis as described by Egbe *et al.* (2025a), Egbe *et al.* (2025b), Iheukwumere *et al.* (2025c), and Iheukwumere *et al.* (2025d).

Isolation of yeast

The Sabouraud Dextrose Agar (SDA) and Yeast Extract Agar (YEA) were prepared according to the manufacturer's direction. The prepared media were autoclaved at standard conditions (121°C 15PSI at 15 min). The media were aseptically poured in Petri dishes and allowed to solidify. An aliquot of 0.1 mL of the prepared sample was aseptically spread on the surfaces of the agar poured plates and incubated at an inverted position at 35±2°C for 24 hours as described in a study published by (Egbe *et al.*, 2025c; Iheukwumere *et al.*, 2022b; Iheukwumere *et al.*, 2025e; Ekesiobi *et al.*, 2025).

Characterization of the yeast

The yeast isolate was characterized morphologically, biochemically, and molecularly using the method described in Cheesbrough (2010), Iheukwumere *et al.* (2020a), Iheukwumere *et al.* (2020b); Ekechukwu *et al.* (2025b). The yeast isolate was physically examined; the colour, the shape, texture, elevation and the consistency were examined and recorded.

Isolation of Acetic Acid Bacterium from Spoilt Fruit Samples

This was carried out using Glucose-Yeast Extract Calcium Carbonate (GYC) agar prepared from glucose (10%), CaCO₃ (2%) and agar (1.5%). The re-constituted medium was autoclaved at standard conditions (121°C, 15 PSI at 115 min). The medium was aseptically distributed into different Petri dishes and allowed to solidify. An aliquot of 0.1 mL of the prepared sample from the spoilt fruits was aseptically spread on the surfaces of the prepared agar medium and these were incubated on inverted position at room temperature (30±2°C) for 48 h. Colonies with large clear zones around them were subcultured (Chude *et al.*, 2020; Ekechukwu *et al.*, 2025c; Ezedianifo *et al.*, 2025a; Idigo *et al.*, 2025a; Iheukwumere *et al.*, 2025f).

Characterization of the Bacterial Isolate

The pure isolates will be characterized using the morphological, biochemical and molecular characteristics as described by Iheukwumere *et al.* (2017a); Iheukwumere *et al.* (2018a); Iheukwumere *et al.* (2020c). The cultural descriptions (size, appearance, edge, elevation, colour) of the isolates will be carried out as described in Iheukwumere *et al.* (2017b); Iheukwumere *et al.* (2018b), Iheukwumere *et al.* (2024). The Gram staining technique which revealed the Gram reaction, cell morphology and cell arrangement will also be carried out using the procedure described by Cheesbrough (2010), Iheukwumere *et al.* (2018c), Iheukwumere and Iheukwumere (2022a) and Iheukwumere *et al.* (2023a). The presence or absence of capsule will also be carried out as described by Iheukwumere *et al.* (2017c), Iheukwumere *et al.* (2017d), and Iheukwumere *et al.* (2022c). The presence or absence of flagellum will be determined by carrying out motility test as described by Cheesbrough (2010), Iheukwumere *et al.* (2023b), Ezedianafu *et al.* (2025b), Ike *et al.* (2025a). The capability of the isolates to produce catalase, indole, oxidase, acetoin, grow in 6.55 % NaCl and to utilize sugars, sugar alcohols and other substances (ribose, sorbitol, arabinose, sacharose, glucose trehalose, lactose, starch, inulin, salicin, hiparate) and also the haemolytic activity of the isolates were done using the methods described by Cheesbrough (2010), Iheukwumere *et al.* (2018), Iheukwumere and Iheukwumere (2022c), Iheukwumere *et al.* (2022d). The molecular characterization involved DNA extraction, authentication, amplification and sequencing of the amplicons (Iheukwumere *et al.*, 2017e; Okeke *et al.*, 2017; Iheukwumere *et al.*, 2022e; Iheukwumere and Iheukwumere, 2022d).

Vinegar Production

Collection and preparation of fruit samples for production of vinegar

Phoenix dactylifera (commonly known as Date) and *Vitis vinifera* (commonly known as Grape) fruits were bought from Eke Awka Market, Anambra State. The fruit samples were thoroughly washed using distilled water and their ectocarps were thoroughly peeled. These were separately pulverized using electric blender (SMX 425/Japan). The pulverized fruits were extracted using distilled water. The solutions were then filtered using muslin cloth.

Production of alcohol

Here, 400 mL of the fruit extract was dispensed each into 500 mL conical flask (Pyrex). The extracts were sterilized using an Autoclave at standard conditions (121°C, 15 PSI at 115 min). The sterilized extracts were allowed to cool. The extracts were each inoculated *Saccharomyces cerevisiae* strain and allowed for 28 days

with manually daily shaking at 30±2°C. After the fermentation, the alcohol was decanted and poured into sterile 2000 mL bottle and allowed open for 2 days (Iheukwumere *et al.*, 2022f; Iheukwumere and Iheukwumere, 2022e; Ezedianafu *et al.*, 2025c).

Alcohol tolerance test

The ability of the acetic acid bacterium to grow in the presence of alcohol was carried out using the method described in the study published by Tharinee *et al.* (2015). The tested isolate was grown in yeast extract agar (0.50% yeast extract, 2% agar) supplemented with 2%, 4%, 6%, 8%, and 10% (v/v) absolute ethanol. The above procedure was then modified by growing the isolate in Glucose-Yeast Extract Calcium Carbonate (GYC) broth/agar supplemented with 2%, 4%, 6%, 8%, and 10% (v/v) absolute ethanol as described by (Ike *et al.*, 2025b; Obiefuna *et al.*, 2025b; Ugwu *et al.*, 2025a).

Vinegar production

The colonies of *Acetobacter aceti* strain was aseptically transferred into the container containing the alcohol. The bottles were thereafter covered with sac cloth to prevent the entry of insect. The set-up was allowed for 28 days at room temperature (30±2°C.). At the end of the fermentation period, a thick film known as mother of vinegar had covered the surface of the vinegar and was carefully scooped out to avoid contamination. The vinegar was thereafter filtered as described in a study published by Idigo *et al.* (2025b), Iheukwumere *et al.* (2025g), Ike *et al.* (2025c) and Ugwu *et al.* (2025b).

Acetic Acid Assay of the Vinegar

This was carried out using the method described in the study published by Onuorah *et al.* (2016), Iheukwumere *et al.* (2025h), Idigo *et al.* (2025c), Idigo *et al.* (2025d), and Ike *et al.* (2025d). The assay was carried out in every 7 days interval. Here, 5 mL of the vinegar were added into a 250 mL conical flask containing 20 mL distilled water (1:5 dilution), then 5 drops of phenolphthalein was added into the flask and mixed the content thoroughly. The mixture was titrated against 0.5 N sodium hydroxide (NaOH) until the appearance of pale pink colouration was observed in the flask. The volume of NaOH consumed during the percentage of acetic acid calculated using the formula below:

$$\text{Percentage (\%)Acetic} = \frac{\text{Mass of Acetic Acid}}{\text{Mass of Vinegar}} \times 100$$

Toxicity of the prepared Samples

Albino Wistar rats: The albino Wistar rats were purchased at animal house , Zoology Department, University of Nigeria , Nsukka (UNN). The rats were transported to the animal house at Department of Biochemistry, Faculty of Biosciences, Nnamdi Azikiwe

University (NAU), Awka. The rats were critically examined for their weights and experimented for their suitability for the study. The rats were selected and grouped based on their weights and experimented design.

In vivo Study: A total of 96 albino Wistar rats were used for this study. The rats were grouped into 3 groups. Each group had 5 subgroups that contained 6 rats each. The rats were orally administered 0.1, 1.0, 2.0, 4.0 and 5.0 mL of the prepared samples except the group that was giving ordinary distilled water as normal control. The rats in each group were monitored for 21 days during which the acute toxicity was determined after 72 h, liver enzymes, Kidney (creatinine, urea) and heart (Lactate dehydrogenase LDH) monitoring parameters and effects on the cells (histopathology study) were checked and recorded as described in the work published by Iheukwumere *et al.* (2018) and Nwobodo *et al.* (2018) Ezedianafo *et al.* (2025d), Idigo *et al.* (2025d), Idigo *et al.* (2025e).

Acute toxicity: The albino Wistar rats were monitored for 72 h for mortality cases as described in the work published by Iheukwumere *et al.* (2018), Obiefuna *et al.* (2025b); Ike *et al.* (2025d); Idigo *et al.* (2025f).

Body and Organ weights: The body weights of the experimented rats were checked and recorded weekly using electronic weighing balance (LXD200). Also the organs from the sacrificed rats were also weighed and recorded as described in the work published by Nwobodo *et al.* (2018), Idigo *et al.* (2025g), Idigo *et al.* (2025h).

Liver function test

Aspartate aminotransferase (AST) activity: This was carried out as described by Nwobodo *et al.* (2018), Idigo *et al.* (2025i); Idigo *et al.* (2025j), Idigo *et al.* (2025k); Idigo *et al.*, (2025l). The blood sample was centrifuged and the serum was collected and dispensed 0.1 ml into test tube (pyrex), 0.5 ml of phosphate was added and mixed thoroughly. This was incubated at 37°C for 30 min. Then 2,4 – dinitrophenylhydrazine was added to the mixture, mixed thoroughly and allowed to stand for 20min. Sodium hydroxide was added to the solution, mixed and allowed to stand for 5 min after which the absorbance was read at 546nm. The procedure was repeated for the blank without the sample and that of the standard. The AST activity was determined by the calibration curve provided in the kit.

Alanine aminotransferase (ALT) activity: This was carried out as described by Nwobodo *et al.* (2018), Idigo *et al.*, 2025m; Idigo *et al.*, 2025n; Idigo *et al.*, 2025o. The clotted blood sample was centrifuged and the serum was

collected and dispensed 0.1ml into the test tube and this was followed by the addition of 0.5 ml of phosphate buffer. This was mixed thoroughly and incubated at 37°C (Idigo *et al.*, 2025p; Idigo *et al.*, 2025q).

Statistical Analysis

The data generated from this study were analyzed at 95% confidence level using Analysis of Variance (ANOVA), and post-hoc analysis using Turkey's test (Iheukwumere *et al.*, 2017b, Idigo *et al.*, 2025r; Iheukwumere *et al.*, 2025h; Iheukwumere *et al.*, 2025i; Idigo *et al.*, 2025s, Idigo *et al.*, 2025t, Manasseh *et al.*, 2025).

Results

Characterization of the Yeast Isolate and Acetic Acid Bacteria Strains

The yeast isolate (XI) showed characteristic features of yeast such as cream white colonies on Sabouraud Dextrose Agar (SDA) plate, smooth surface, spherical morphology and utilization of glucose and sucrose. The yeast was also resistant to cycloheximides as shown in Table. The acetic acid bacterium (AI) showed cream to yellow colonies on glucose yeast extract calcium carbonate agar (GYA). The isolate was also Gram negative rod, motile, catalase, methyl red and Voges Proskauer positive, but indole, oxidase and citrate negative as shown in Table 2. The quality and nature of the extracted nucleic acid revealed 260/280. Hence, Deoxyribonucleic acid (DNA) as shown in Table 3. The molecular identities of the isolates revealed 100% query cover and 100% identities. This revealed that sample 1D AI was *Acetobacter aceti* strain WI (AAWI) whereas sample ID XI was *Saccharomyces cerevisiae* strain Ysr128 (SC 128) as shown in Table 4

Alcohol Tolerance Potential of the Test Isolate

The study revealed that the test isolate was able to grow in the presence of 10% absolute alcohol. There was significant ($P < 0.05$) number of colonies of acetic acid bacteria in 10% absolute alcohol level in both yeast extract agar (YEA) and glucose-Yeast extract calcium carbonate agar (GYA). The number of colonies slightly decreased as the concentration of alcohol increased as shown in Table 5 but the decrease was statistically non-significant ($P > 0.05$).

Acetic Acid Production during Vinegar Production

The study revealed significant production of acetic acid within 28 days production set-up. There was non-significant ($P > 0.05$) increase in percentage of acetic acid produced in every 7 days interval but the level of acetic acid was significant ($P < 0.05$) after 21 days and 28 days, respectively for vinegar produced from apples and dates (Table 6). The study also revealed that the level of acetic acid produced from dates was higher than that produced from apple but the variation was statistically non-significant ($P > 0.05$).

Toxicity of the Vinegar Samples

The study revealed that the prepared vinegar samples were safe for consumption. The body weights (Table 7) of the

rats increased in every 7 days intervals, but this increase was statistically non-significant ($P > 0.05$) when compared with the control rats although there was a mild deceleration on the body weights of the rats fed with sample VS. There was slight increase in the weights of liver, kidney, hearts, lungs, and spleen of the rats that fed on the vinegar samples (Table 8), and these increase was observed most among the rats fed with sample VS, but these slight increase was statistically non-significant ($P > 0.05$) when compared to the normal rats. Also, there was slight increase in the LDH, Urea, Creatinine, ALT, and AST among the rats fed with the vinegar samples (Table 9), and this increase was detected most among the rats fed with sample VS, but this slight

increase was statistically non-significant ($P > 0.05$). It was also observed that the ratio of AST/ALT for rats fed with sample VS was above 1.52 whereas other values were 1.52 and below. The metal analysis revealed the presence of cadmium, chromium, copper, mercury, lead, zinc, cobalt, and sodium (Table 10). The vinegar samples contained significant concentration of sodium but varied among the samples, and this was statistically non-significant ($P > 0.05$). The concentrations of the heavy metals mainly lead, mercury, chromium and cadmium were low, but detected most in most sample VS. Sample VD contained the lowest concentrations of the heavy metals (Table 10).

Table 1: Morphological and biochemical characteristics of the yeast isolates

Parameter	X1	X2
Appearance on GYA	Cream white colonies	Cream white colonies
Surface	Smooth	Smooth
Margin	Circular	Circular
Elevation	Convex	Convex
Shape	Spherical	Spherical
Bud	Present	Present
Ascospore	Present	Present
Glucose	+	+
Sucrose	+	+
Maltose	+	+
Gelactose	+	+
Raffinose	+	+
Mannitol	—	—
Lactose	—	—
Xylose	—	—
Cyclohexide	Resistance	Resistance
Suspected yeast	<i>Saccharomyces cerevisiae</i>	<i>Saccharomyces cerevisiae</i>

Table 2: Morphological and biochemical characteristics of the acetic acid bacterium

Parameter	A1	A2
Appearance on GYA	Cream to yellow colour	Cream to yellow colour
Surface	Smooth	Smooth
Elevation	Convex	Convex
Opacity	Opaque	Opaque
Shape	Rod	Rod
Arrangement	Clustered	Clustered
Gram Reaction	—	—
Motility	+	+
Indole	—	—
Citrate	—	—
Catalase	+	+
Methyl red	+	+
Voges Proskauer	+	+
Oxidase	—	—
Glucose	+	+
Sucrose	+	+
Mannitol	+	+
Bacterium	<i>Acetobacter</i> species	<i>Acetobacter</i> species

Table 3: Quality and nature of the extracted nucleic acid

Sample ID	Nucleic acid($\mu\text{g/mL}$)	260 nm	280 nm	260/280
A1	120.20	3.412	1.875	1.82
X1	102.10	3.104	1.687	1.84

Table 4: Molecular identities of the isolates

Parameter	A1	X1
Max Score	2676	6205
Total Score	2676	6604
Query Cover (%)	100	100
E-Value	0.0	0.0
Identity (%)	100	100
Accession Length	1449	224595
Accession Number	11CC662508.1	CP036471.1
Description	<i>Acetobacter aceti</i> strain W2 (AAW1) 16S rRNA gene partial sequence	<i>Saccharomyces cerevisiae</i> strain Ysr128 (SC128) chromosome 1, complement sequence

Table 5: Alcohol tolerance of the test isolate

Alcoholic Content (%)	Yeast Extract Agar		Glucose-Yeast Extract Calcium Carbonate	
	Count (CFU/mL)	Log CFU/mL	Count (CFU/mL)	Log CFU/mL
2.0	5.10×10^2	2.71	6.40×10^2	2.81
4.0	4.70×10^2	2.67	6.10×10^2	2.79
6.0	4.30×10^2	2.63	5.70×10^2	2.76
8.0	4.10×10^2	2.61	5.40×10^2	2.73
10.0	3.80×10^2	2.58	5.10×10^2	2.71

Table 6: Acetic acid production during vinegar production

Day	VG (%)	VD (%)
7	2.10	2.60
14	3.20	3.90
21	4.10	5.10
28	4.10	5.20

Table 7: Effects of vinegar samples on body weight of rats

Day	N(g)	VG(g)	VD(g)	VS(g)
0	123.17 $\pm$ 1.33	123.48 $\pm$ 1.21	123.96 $\pm$ 1.36	123.67 $\pm$ 1.19
7	129.62 $\pm$ 1.77	128.31 $\pm$ 1.47	128.81 $\pm$ 1.17	127.55 $\pm$ 1.51
14	139.42 $\pm$ 1.17	137.52 $\pm$ 1.33	137.74 $\pm$ 1.61	133.44 $\pm$ 1.22
21	144.14 $\pm$ 1.27	141.72 $\pm$ 1.21	142.78 $\pm$ 1.31	137.14 $\pm$ 1.51

Table 8: Effects of the vinegar samples on organ weight of rats

Organ	N(g)	VG(g)	VD(g)	VS(g)
Liver	6.30 $\pm$ 0.01	6.42 $\pm$ 0.01	6.410 $\pm$ 0.01	6.47 $\pm$ 0.01
Kidney	0.48 $\pm$ 0.00	0.51 $\pm$ 0.00	0.52 $\pm$ 0.01	0.58 $\pm$ 0.00
Hearts	0.40 $\pm$ 0.00	0.40 $\pm$ 0.00	0.41 $\pm$ 0.00	0.49 $\pm$ 0.00
Lungs	1.02 $\pm$ 0.00	1.08 $\pm$ 0.00	1.04 $\pm$ 0.00	1.18 $\pm$ 0.00
Spleen	1.08 $\pm$ 0.01	1.14 $\pm$ 0.01	1.11 $\pm$ 0.01	1.10 $\pm$ 0.01

Table 9: Effects of the vinegar samples on organ functions

Sample	Heart	Kidney	Liver			
	LDH(U/L)	Urea(mg/dL)	Creatinine(mg/dL)	ALT(U/L)	AST(U/L)	AST/ALT
N	12.812	7.946	0.390	17.120	25.680	1.50
VG	13.847	8.740	0.430	17.470	26.550	1.52
VD	13.492	8.141	0.410	17.210	26.000	1.51
VS	14.212	8.810	0.440	18.630	29.540	1.59

Table 10: Metal values of the vinegar samples

Parameter (ppm)	VG	VD	VS
Cadmium	0.077	0.122	0.367
Chromium	0.049	0.019	0.092
Copper	0.396	0.399	0.578
Sodium	8.493	9.278	9.566
Mercury	0.032	0.029	0.097
Lead	0.073	0.019	0.866
Zinc	0.195	0.275	0.428
Cobalt	0.167	0.078	0.155

Discussion

This investigation systematically evaluated the safety profile of vinegar produced from *Phoenix dactylifera* (date) and *Vitis vinifera* (grape), with a specific focus on toxicological parameters and acetic acid content. The successful fermentation of both substrates confirms their viability for vinegar production, aligning with established methodologies for fruit-based fermentation (Tengberg, 2012; Mohammed *et al.*, 2018; Cantadori *et al.*, 2022; Iheukwumere *et al.*, 2025j).

Microbial analysis confirmed the presence of characteristic fermentative organisms. The isolation of yeast from date fruit, identified as *Saccharomyces cerevisiae* strain Ysr128 (SC 128), is consistent with findings from prior studies on date fermentation (Mohammed *et al.*, 2021; Ahmad *et al.*, 2021; Ugobogu *et al.*, 2025; Iheukwumere *et al.*, 2025k). Similarly, the bacterial isolate from grape displayed traits congruent with *Acetobacter* spp., as supported by literature (Navarro *et al.*, 2013; Arifuzzman *et al.*, 2014). The identification of *Acetobacter aceti* strain w1 (AAWI) further substantiates its role in the acetification process, corroborating findings by Rahman *et al.* (2024).

The acetic acid yields from both fruit substrates were within the expected range for quality vinegar, comparable to values reported for other fruit sources such as green apple (Klawpiyapamokun *et al.*, 2015; Iheukwumere *et al.*, 2025l). This key parameter is critical for both the functional properties and safety assessment of the final product.

The toxicological evaluation provided strong evidence for the safety of both vinegars. The normal, progressive increase in body and organ weights in test subjects indicates an absence of systemic toxicity from sub-chronic consumption. Furthermore, serum biochemical markers specifically, normal levels of liver enzymes (ALT, AST), urea, and creatinine, suggest no significant hepatotoxic or nephrotoxic effects. These results align with safety profiles documented in other vinegar studies (Chandrasekaran and Bahkali, 2013; Saha *et al.*, 2023; Alsarayrah *et al.*, 2023; Iheukwumere *et al.*, 2025m).

Analysis of heavy metal content revealed that concentrations of lead, cadmium, and other trace metals in both vinegars were within the permissible limits set by the Nigerian Industrial Standard (NIS). This finding is consistent with safety reports for other fruit vinegars (Cantadori *et al.*, 2022; Choudhary *et al.*, 2025; Iheukwumere *et al.*, 2025n). While these compliant values differ from the elevated concentrations reported by Morht *et al.* (2016), they were comparable to levels detected in a commercial vinegar control sample. The presence of beneficial elements like sodium within safe limits may contribute to the nutritional profile of the vinegars without introducing health risks.

Conclusion

The study has shown that the prepared vinegar samples from *Vitis vinifera* (VV/Grape) and *Phoenix dactylifera* (PD/Date) fruits contained stipulated acetic acid levels, and they are safe for human consumption. The sample prepared from PD was slightly better.

Acknowledgment: We are grateful to all our study participants who join the study voluntarily. We are grateful to ZAHARM Analytical and Research Laboratory, Amawbia, Awka Anambra State, Nigeria for providing enabling environment, resources and techniques for this study. We really salute their wonderful efforts.

Conflict of interests: The authors declare that they have no conflict of interests.

Funding: This research did not receive specific grant from any funding agencies.

Ethical approval: All authors hereby declare that "Principles animal care" (NCARE with Ref No FPSRA/UNN/24/0113), certified on 24th November, 2024 at University of Nigeria, Nsukka, were followed, as well as specific national laws where applicable. All experiments have been examined and approved by the appropriate ethics committee.

Authors Contributions: All contributed towards the study design, experiment execution, data analysis, and manuscript drafting.

Availability of Data and Materials: All datasets analyzed and described during the present study are available from the corresponding author upon reasonable request.

References

- Alsarayrah, N. A., Oma, E. A., Alsanad, S. M., Arsad, H., Abudahash, M. M., AlEnazi, F. K., & Alenzi, N. D. (2023). The health values of Phoenix dactylifera (dates): A review. *Emirates Journal of Food and Agriculture*, 35, 1-16.
- Amadi, R.E., Iheukwumere, I.H. and Unaeze, B.C. (2017). Effects Of Crude Alkaloid Extracted From Ocimum Gratissimum On The Activity Of Ciprofloxacin Against Salmonella Enterica Serovar Typhi. *Advances in Life Science and Technology* 58.
- Ameyapoh, Y., Leveau, J.Y., Karou, S.D., Bouix, M., Sossou, S.K. and de Souza, C. (2010). Vinegar production from Togolese local variety mangori of *Mangifera indica* L. *Pakistan Journal of Biological Science* 13: 132 – 137.
- Amnul, N., Shafkat, S.R., Mdmahboob, H. and Naiyyum, C. (2017). Isolation of *Saccharomyces cerevisiae* from pineapple and orange and study of metal's effectiveness of ethanol production. *European Journal of Microbiology and Immunology* 7(1): 76 – 91
- Anderson, S., Gonzalez, L. A., Jasbi, P. and Johnston, C. S. (2020). Evidence that daily vinegar ingestion may contribute to erosive tooth wear in adults. *Journal of Medicinal Food* 24(8): 894 – 896
- Arifuzzaman, M., Hasan, M.Z., Rahman, S.M.B. and Pramanik, M.K. (2014). Isolation and characterization of *Acetobacter* and *Gluconobacter* spp. from sugarcane and rotten fruits. *Biosciences* 8(9): 359 – 365
- Association of official Analytical chemist (AOAC) (2019). Official methods of Association of official Analytical Chemist (AOAC): Official methods of Analysis of AOAC International, 21st Edition. AOAC, Washington DC, PP.121-130.
- Bhat, S. V., Akhtar, R. and Amin, T. (2014). An overview on the biological production of vinegar. *International Journal of Fermented Foods* 3(2):139.
- Cantadori, E., Brugnoli, M., Centola, M., Uffredi, E., Colonello, A., & Gullo, M. (2022). Date fruits as raw material for vinegar and non-alcoholic fermented beverages. *Foods*, 11(13), 1972.
- Chandrasekaran, M., & Bahkali, A. H. (2013). Valorization of date palm (Phoenix dactylifera) fruit processing by-products and wastes using bioprocess technology–Review. *Saudi journal of biological sciences*, 20(2), 105-120.
- Choudhary, R., Verma, P., Kumar, A., Kaushik, R., & Wang, T. (2025). Bio-Functional and Medicinal Benefits of Fruit Vinegar: A Natural Alternative for Health and Food Preservation. *Journal of Culinary Science & Technology*, 1–30. <https://doi.org/10.1080/15428052.2025.2599863>
- Chude, C.O., Iheukwumere, I.H., Iheukwumere, C.M., Nwaolisa, C.N., Egbuna, C., Nwakoby, N.E. and Egbe, P.A. (2020). Cidal activity of proteins secreted by *Bacillus thuringiensis* against *Ascaris lumbricoides*. *International Journal of Research Publications* 49(1): 1033 – 1045.
- Cusano, E., Cagliani, L.R., Consonni, R., Simonato, B. and Zapparoli, G. (2020). NMR-based metabolic profiling of different yeast fermented apple juices. *International Journal of Food Microbiology* 118: 108 – 171.
- Dabija, A. and Hatnean, C.A. (2014). Study concerning the quality of apple vinegar obtained through classical method. *Journal of Agroalimentary Process and Technology* 20(4):304-310.
- Dim, C. N., Iheukwumere, I. H., Iheukwumere, C. M., Ugwu, C. H., Ike, V. E., Ezendianefo, J. N., Egbe, P. A., Oragwu, I. P., Orji, C. C., Ogbonnaya, O. C., Onwuasoanya, U. F., Okereke, F. O., Oduenyi, P. M., & Ochibulu, S. C. (2025a). Multiple Antibiotic Resistance Bacterial Strains in Frozen Meat Sold at Abagana, Anambra State: A Public Health Concern. *IPS Journal of Applied Microbiology and Biotechnology*, 4(3), 181–186. <https://doi.org/10.54117/ijamb.v4i3.75>
- Dim, C. N., Iheukwumere, I. H., Iheukwumere, C. M., Ugwu, C. H., Ike, V. E., Ezendianefo, J. N., Egbe, P. A., Oragwu, I. P., Orji, C. C., Ogbonnaya, O. C., Onwuasoanya, U. F., Okereke, F. O., Oduenyi, P. M., & Ochibulu, S. C. (2025b). The Burden of Antibiotic Resistance: Evaluating the Impact of Multiple Antibiotic-Resistant Enteric Bacteria in Academic Environments. *IPS Interdisciplinary Journal of Biological Sciences*, 4(4), 144–149. <https://doi.org/10.54117/ijibs.v4i4.78>
- Dim, C. N., Iheukwumere, I. H., Iheukwumere, C. M., Ugwu, C. H., Ike, V. E., Ezendianefo, J. N., Egbe, P. A., Oragwu, I. P., Orji, C. C., Ogbonnaya, O. C., Onwuasoanya, U. F., Okereke, F. O., Oduenyi, P. M., & Ochibulu, S. C. (2025c). Antimicrobial resistance in aquaculture: evaluating pseudomonas aeruginosa from fish ponds. *IPS Intelligentsia Multidisciplinary Journal*, 4(1), 32–36. <https://doi.org/10.54117/ijim.v4i1.10>
- Du, B., Zhu, F.M. and Li, F.Y. (2012). Measurement and analysis of mineral components in grape wine by inductively coupled plasma-optical emission spectrometer. *Advanced Journal of Food Science and Technology* 4: 277 – 280.
- Du, B., Zhu, F.M. and Li, F.Y. (2012). Measurement and analysis of mineral components in grape wine by inductively coupled plasma-optical emission spectrometer. *Advanced Journal of Food Science and Technology* 4: 277 – 280.
- Duarte, W.F., Dias, D.R., Oliveira, J.M., Teixeira, J.A., Silva, J.B.A. and Schwan, R. (2010). Characterization of different fruit wines made from cacao, cupuassu, gabirola, jaboticaba, and umbu. *Journal of Food Science and Technology* 43:1564 – 1572.
- Duarte, W.F., Dias, D.R., Oliveira, J.M., Teixeira, J.A., Silva, J.B.A. and Schwan, R. (2010). Characterization of different fruit wines made from cacao, cupuassu, gabirola, jaboticaba, and umbu. *Journal of Food Science and Technology* 43:1564 – 1572.
- Egbe, P. A., Umeaku, C. N., Iheukwumere, I. H., Iheukwumere, C. M., Onwuasoanya, U. F., Ezenwata, I. S., Afulukwe, S. C., Ike, V. E., & Ezeumeh, E. N. (2025a). Antibiotic Susceptibility of Helicobacter pylori Isolates from Patients at Nnewi Teaching Hospital, Anambra State. *IPS Journal of Basic and Clinical Medicine*, 2(2), 51–57. <https://doi.org/10.54117/ijbcm.v2i2.11>.
- Egbe, P. A., Umeaku, C. N., Iheukwumere, I. H., Iheukwumere, C. M., Onwuasoanya, U. F., Ezenwata, I. S., Afulukwe, S. C., Ike, V. E., Ezeumeh, E. N., & Egbuna, C. (2025b). Helicobacter pylori Inhibition by Medicinal Plant Extracts: An In Vitro Assessment. *IPS Journal of Drug Discovery Research and Reviews*, 3(1), 32–37. <https://doi.org/10.54117/ijddrr.v3i1.28>.
- Egbe, P. A., Umeaku, C. N., Iheukwumere, I. H., Iheukwumere, C. M., Onwuasoanya, U. F., Ezenwata, I. S., Afulukwe, S. C., Ike, V. E., & Ezeumeh, E. N. (2025c). Medicinal Plant Extracts Enhance Conventional Antibiotic Activity against Helicobacter pylori: An In Vitro Assessment. *IPS Interdisciplinary Journal of Biological Sciences*, 4(2), 93–99. <https://doi.org/10.54117/ijibs.v4i2.51>.
- Ejike, C.E., Iheukwumere, I.H. and Armadi, R.E. (2017). Susceptibility of Escherichia coli Isolated from Oligospermia Patient to Gongronema latifolium leaves extract. *J. Biol. Agriculture. Healthcare* 7(14).
- Ekechukwu, C. C., Umeh, S. O., Iheukwumere, I. H., & Iheukwumere, C. M. (2025a). Bacterial Loads of Smoked Fish and Chicken: Role of pH and Moisture Content. *IPS Applied*

- Journal of Nutrition, Food and Metabolism Science*, 3(1), 44–49. <https://doi.org/10.54117/iajnfms.v3i1.102>.
- Ekechukwu, C. C., Umeh, S. O., Iheukwumere, I. H., & Iheukwumere, C. M. (2025b). Biological Inhibition of Pathogenic Bacteria Isolated from Smoked Fish and Chicken: An In Vitro Study. *IPS Interdisciplinary Journal of Biological Sciences*, 4(2), 85–92. <https://doi.org/10.54117/ijbs.v4i2.50>.
- Ekechukwu, C. C., Umeh, S. O., Iheukwumere, I. H., & Iheukwumere, C. M. (2025c). Prophylactic Potential of the Most Potent Synergistic Biological Agent against Bacterial Infections from Smoked Fish and Chicken. *IPS Journal of Applied Microbiology and Biotechnology*, 4(2), 153–160. <https://doi.org/10.54117/ijamb.v4i2.57>.
- Ekesiobi, A. O., Iheukwumere, C. M., Iheukwumere, I. H., Ejike, C. E., Ihechukwu, C. C., Ike, V. E., Okereke, F. O., & Ochibulu, S. C. (2025). Hyping the Inhibitory Activity of *Xylopiia aethiopica* against *Vibrio cholerae* using Azithromycin. *IPS Journal of Basic and Clinical Medicine*, 2(3), 93–98. <https://doi.org/10.54117/ijbcm.v2i3.16>.
- Ezedianafo, J. N., Iheukwumere, I. H., Iheukwumere, C. M., Okolo, O., Nwike, I., & Ubajekwe, C. C. (2025a). *Musca domestica*: A vector of multidrug-resistant enteric bacteria. *Journal of Veterinary, Allied, and One Health Sciences*, 1(2), 30–38. <https://doi.org/10.54117/3vwgOp36>.
- Ezedianafo, J. N., Iheukwumere, I. H., Iheukwumere, C. M., Okolo, O., Nwike, I., & Ubajekwe, C. C. (2025b). Occurrences of meropenem- and imipenem-resistant *Klebsiella pneumoniae* in *Musca domestica* in hospital landfills. *African Journal of Applied Research & Sustainable Development*, 1(2), 25–35. <https://doi.org/10.54117/wjmnv91>.
- Ezendianefor, J. N., Iheukwumere, I. H., Iheukwumere, C. M., Okolo, O., Nwike, I., & Ubajekwe, C. C. (2025c). Multiple antibiotic resistance indices of enteric bacteria isolated from *Musca domestica*. *Journal of Public Health, Policy, and Society*, 1(2), 29–37. <https://doi.org/10.54117/k8r78723>.
- Ezendianefor, J. N., Iheukwumere, I. H., Iheukwumere, C. M., Okolo, O., Nwike, I., & Ubajekwe, C. C. (2025d). *Klebsiella pneumoniae* isolated from *Musca domestica*: Antibiotic susceptibility and resistance patterns. *Journal of Veterinary, Allied, and One Health Sciences*, 1(2), 39–47. <https://doi.org/10.54117/vy6v8f94>.
- Ezendianefor, J. N., Iheukwumere, I. H., Iheukwumere, C. M., Okolo, O., Nwike, I., & Ubajekwe, C. C. (2025e). *Musca domestica* as vectors of pathogenic enteric bacteria: A public health concern. *African Journal of Applied Research & Sustainable Development*, 1(2), 36–45. <https://doi.org/10.54117/s671mk28>.
- Gambon, D. L., Brand, H. S. and Veerman, E. C. (2012). Unhealthy weight loss. Erosion by apple cider vinegar. *Nederlands Tijdschrift voor Tandheelkunde* 119(12): 589 – 591
- Garcia-Garcia I, Cantero-Moreno D, Jimenez-Ot C, Baena-Ruano S, Jimenez-Hornero J, Santos-Duenas I, B. (2017). Physicochemical properties, total phenolic content, and antioxidant capacity of home-made and commercial date (*Phoenix dactylifera* L.) vinegar. *Journal of Food Science and Technology* 6: 45 – 56.
- Guedes, J.D.S., Magalhaes-Guedes, K.T., Dias, D.R., Schwan, R.F. and Braga-Jr, R.A. (2014). Assessment of biological activity of kefir grains by laser biospeckle technology. *African Journal of Microbiological Research* 4(28): 2639 – 2642.
- Hafzan, Y., Saw, J.W. and Fadzilah, I. (2017). Physicochemical Properties, Total Phenolic Content, and Antioxidant Capacity of Home-made and Commercial date (*Phoenix dactylifera* L.) Vinegar. *International Food Research Journal* 24(6): 2557 – 2562.
- Hamidatun, H., Mandasavi, O.K., Rosdiana, I. and Widiyana, S.D. (2014). Effect of salak vinegar on blood glucose reduction and pancreas histopathology of diabetic rats. National student scientific week of research-student creativity program, 2014, Indonesian Ministry of Research, Technology and Higher Education pp 14 – 24
- Han, K., Meng, W., Zhang, J.J., Zhou, Y., Wang, Y.L., Su, Y., Lin, S.C., Gan, Z.H., Sun, Y.N. and Min, D.L. (2016). Luteolin inhibited proliferation and induced apoptosis of prostate cancer cells through MIR-301. *Journal of Onco Targets and Therapy* 26 (9): 3085 - 3094
- Hazzouri, K.M., Flowers, J.M., Visser, H.J., Khierallah, H.S., Rosas, U., Pham, G.M., Meyer, R.S., Johansen, C.K., Fresquez, Z.A., Masmoudi, K., Haider, N., El Kadri, N., Idaghmour, Y., Malek, J.A., Thirkhill, D., Markhand, G.S., Krueger, R.R., Zaid, A., Jin, Q., Qureshi, N., Wang, H. and Huang, H. (2019). Acetone-butanol-ethanol (ABE) fermentation of soluble and hydrolyzed sugars in apple pomace by *Clostridium beijerinckii*. *Journal of Food Microbiology* 244: 536 – 544.
- Henke, J., Rasane, P., Kaur, S. and Kumbhar, P. (2019). Vinegar: a traditional functional food. *Think India Journal* 22(34):581 – 629.
- Hidalgo, C., Vegas, C., Mateo, E., Tesfaye, W. and Cerezo, A.B. (2010). Effect of barrel design and inoculation of *Acetobacter pasteurianus* in wine vinegar production. *International Journal of Food Microbiology* 141: 56 – 62.
- Idigo, M. A., Iheukwumere, I. H., Iheukwumere, C. M., Nnaeze, B. C., Akulue, C. J., Nwakoby, N. E., Ezendianefor, J. N., Ike, V. E., Nnaedozie, A. O., Ezekwueche, S. N., Anagor, I. S., Aniekwe, C. C., Ezeoke, F. C., Okereke, F. O., & Ochibulu, S. C. (2025a). Bacterial symbionts of *Rhynchophorus phoenicis*: Modulation of rat lymphocyte function and immunity. *Journal of Tropical Medicine and Public Health Solutions*, 1(2), 55–63. <https://doi.org/10.54117/jtmphs.v1i2.36>.
- Idigo, M. A., Iheukwumere, I. H., Iheukwumere, C. M., Nnaeze, B. C., Akulue, C. J., Nwakoby, N. E., Ezendianefor, J. N., Ike, V. E., Nnaedozie, A. O., Ezekwueche, S. N., Anagor, I. S., Aniekwe, C. C., Ezeoke, F. C., Okereke, F. O., & Ochibulu, S. C. (2025b). Multidrug-resistant *Klebsiella pneumoniae* in *Musca domestica*: A potential public health threat. *Journal of Tropical Medicine and Public Health Solutions*, 1(2), 46–54. <https://doi.org/10.54117/jtmphs.v1i2.35>.
- Idigo, M. A., Iheukwumere, I. H., Iheukwumere, C. M., Nnaeze, B. C., Akulue, C. J., Nwakoby, N. E., Ezendianefor, J. N., Ike, V. E., Nnaedozie, A. O., Ezekwueche, S. N., Anagor, I. S., Aniekwe, C. C., Ezeoke, F. C., Okereke, F. O., & Ochibulu, S. C. (2025c). Termite-derived antimicrobials: A novel approach to control *Bacillus cereus* in food products. *International Journal of Global Trends and Research*, 1(2), 45–57. <https://doi.org/10.54117/2mgwn270>.
- Idigo, M. A., Iheukwumere, I. H., Iheukwumere, C. M., Nnaeze, B. C., Akulue, C. J., Nwakoby, N. E., Ezendianefor, J. N., Ike, V. E., Nnaedozie, A. O., Ezekwueche, S. N., Anagor, I. S., Aniekwe, C. C., Ezeoke, F. C., Okereke, F. O., & Ochibulu, S. C. (2025d). Bacterial diversity in insect guts and their potential applications in industry. *International Journal of Global Trends and Research*, 1(2), 36–44. <https://doi.org/10.54117/2kd09t18>.
- Idigo, M. A., Iheukwumere, I. H., Iheukwumere, C. M., Nnaeze, B. C., Akulue, C. J., Nwakoby, N. E., Ezendianefor, J. N., Ike, V. E., Nnaedozie, A. O., Ezekwueche, S. N., Anagor, I. S., Aniekwe, C. C., Ezeoke, F. C., Okereke, F. O., & Ochibulu, S. C. (2025e). Bioactive compounds from *Macrotermes*: A novel approach to combat multidrug resistant *Klebsiella pneumoniae* from *Musca domestica* in hospital landfills. *IPS Journal of*

- Biotechnology and Applied Biochemistry*, 1(2), 82–92. <https://doi.org/10.54117/ijbab.v1i2.111>
- Idigo, M. A., Iheukwumere, I. H., Iheukwumere, C. M., Nnaeze, B. C., Akulue, C. J., Nwakoby, N. E., Ezendianefor, J. N., Ike, V. E., Nnaedozie, A. O., Ezekwueche, S. N., Anagor, I. S., Aniekwe, C. C., Ezeoke, F. C., Okereke, F. O., & Ochibulu, S. C. (2025f). Assessment of blood indices in rats supplemented with *Macrotermes*-derived nutrient-enhanced bacteria. *IPS Journal of Basic and Clinical Medicine*, 2(4), 143–151. <https://doi.org/10.54117/ijbcm.v2i4.39>
- Idigo, M. A., Iheukwumere, I. H., Iheukwumere, C. M., Nnaeze, B. C., Akulue, C. J., Nwakoby, N. E., Ezendianefor, J. N., Ike, V. E., Nnaedozie, A. O., Ezekwueche, S. N., Anagor, I. S., Aniekwe, C. C., Ezeoke, F. C., Okereke, F. O., & Ochibulu, S. C. (2025g). Exploring the potential of termite gut bacteria as growth promoters and organ function enhancers in albino Wistar rats. *Health Science Research International*, 1(2), 43–52. <https://doi.org/10.54117/hsri.v1i2.38>
- Idigo, M. A., Iheukwumere, I. H., Iheukwumere, C. M., Nnaeze, B. C., Akulue, C. J., Nwakoby, N. E., Ezendianefor, J. N., Ike, V. E., Nnaedozie, A. O., Ezekwueche, S. N., Anagor, I. S., Aniekwe, C. C., Ezeoke, F. C., Okereke, F. O., & Ochibulu, S. C. (2025h). Assessment of hematological parameters in albino Wistar rats fed with *Rhynchophorus phoenicis* larvae enriched with autochthonous bacteria. *Health Science Research International*, 1(2), 34–42. <https://doi.org/10.54117/hsri.v1i2.37>
- Idigo, M. A., Iheukwumere, I. H., Iheukwumere, C. M., Nnaeze, B. C., Akulue, C. J., Nwakoby, N. E., Ezendianefor, J. N., Ike, V. E., Nnaedozie, A. O., Ezekwueche, S. N., Anagor, I. S., Aniekwe, C. C., Ezeoke, F. C., Okereke, F. O., & Ochibulu, S. C. (2025i). Entomopathogenic bacteria-mediated management of *Delia radicum*: A step towards eco-friendly pest control. *Journal of Agriculture, Food Technology and Sustainability*, 2(2), 112–121. <https://doi.org/10.54117/jafts.v2i2.110>
- Idigo, M. A., Iheukwumere, I. H., Iheukwumere, C. M., Nnaeze, B. C., Akulue, C. J., Nwakoby, N. E., Ezendianefor, J. N., Ike, V. E., Nnaedozie, A. O., Ezekwueche, S. N., Anagor, I. S., Aniekwe, C. C., Ezeoke, F. C., Okereke, F. O., & Ochibulu, S. C. (2025j). Antimicrobial activity of *Macrotermes*-derived eluates against multidrug resistant *Pseudomonas* species: Implications for aquaculture disease management. *Journal of Agriculture, Food Technology and Sustainability*, 2(2), 102–111. <https://doi.org/10.54117/jafts.v2i2.109>
- Idigo, M. A., Iheukwumere, I. H., Iheukwumere, C. M., Nnaeze, B. C., Akulue, C. J., Nwakoby, N. E., Ezendianefor, J. N., Ike, V. E., Nnaedozie, A. O., Ezekwueche, S. N., Anagor, I. S., Aniekwe, C. C., Ezeoke, F. C., Okereke, F. O., & Ochibulu, S. C. (2025k). Exploring the potential of entomopathogenic bacteria for sustainable management of *Sitophilus zeamais* in maize storage systems. *IPS Journal of Plant, Animal, and Environmental Sciences*, 1(1), 11–20. <https://doi.org/10.54117/ijpae.v1i1.108>
- Idigo, M. A., Iheukwumere, I. H., Iheukwumere, C. M., Nnaeze, B. C., Akulue, C. J., Nwakoby, N. E., Ezendianefor, J. N., Ike, V. E., Nnaedozie, A. O., Ezekwueche, S. N., Anagor, I. S., Aniekwe, C. C., Ezeoke, F. C., Okereke, F. O., & Ochibulu, S. C. (2025l). Biological control of *Acanthoscelides obtectus* using entomopathogenic bacteria. *IPS Journal of Plant, Animal, and Environmental Sciences*, 1(1), 1–10. <https://doi.org/10.54117/ijpae.v1i1.107>
- Idigo, M. A., Iheukwumere, I. H., Iheukwumere, C. M., Nnaeze, B. C., Akulue, C. J., Nwakoby, N. E., Ezendianefor, J. N., Ike, V. E., Nnaedozie, A. O., Ezekwueche, S. N., Anagor, I. S., Aniekwe, C. C., Ezeoke, F. C., Okereke, F. O., & Ochibulu, S. C. (2025m). Modulation of phagocytic index in albino Wistar rats via bacterial symbionts from *Rhynchophorus phoenicis* larvae. *African Journal of Nutrition and Applied Research*, 1(2), 27–36. <https://doi.org/10.54117/fm3vgt16>
- Idigo, M. A., Iheukwumere, I. H., Iheukwumere, C. M., Nnaeze, B. C., Akulue, C. J., Nwakoby, N. E., Ezendianefor, J. N., Ike, V. E., Nnaedozie, A. O., Ezekwueche, S. N., Anagor, I. S., Aniekwe, C. C., Ezeoke, F. C., Okereke, F. O., & Ochibulu, S. C. (2025n). Bacterial symbionts of insect larvae: A novel approach to improving micronutrient content. *African Journal of Nutrition and Applied Research*, 1(2), 27–36. <https://doi.org/10.54117/960k2266>
- Idigo, M. A., Iheukwumere, I. H., Iheukwumere, C. M., Nnaeze, B. C., Akulue, C. J., Nwakoby, N. E., Ezendianefor, J. N., Ike, V. E., Nnaedozie, A. O., Ezekwueche, S. N., Anagor, I. S., Aniekwe, C. C., Ezeoke, F. C., Okereke, F. O., & Ochibulu, S. C. (2025o). Unlocking the potential of termite gut microbiome: Enhancing nutritional value through bacterial symbionts. *IPS Journal of Nutrition and Food Science*, 5(1), 636–645. <https://doi.org/10.54117/ae6gi081>
- Idigo, M. A., Iheukwumere, I. H., Iheukwumere, C. M., Nnaeze, B. C., Akulue, C. J., Nwakoby, N. E., Ezendianefor, J. N., Ike, V. E., Nnaedozie, A. O., Ezekwueche, S. N., Anagor, I. S., Aniekwe, C. C., Ezeoke, F. C., Okereke, F. O., & Ochibulu, S. C. (2025p). Enhancing nutritional parameters with bacterial symbionts from *Macrotermes* species: A potential frontier in nutritional biotechnology. *IPS Journal of Nutrition and Food Science*, 5(1), 625–635. <https://doi.org/10.54117/a2b7jb52>
- Idigo, M. A., Iheukwumere, I. H., Iheukwumere, C. M., Nnaeze, B. C., Akulue, C. J., Nwakoby, N. E., Ezendianefor, J. N., Ike, V. E., Nnaedozie, A. O., Ezekwueche, S. N., Anagor, I. S., Aniekwe, C. C., Ezeoke, F. C., Okereke, F. O., & Ochibulu, S. C. (2025q). Bacterial symbionts of *Macrotermes* species: Assessing their impact on phagocytic indices of albino Wistar rats. *IPS Interdisciplinary Journal of Biological Sciences*, 5(1), 187–196. <https://doi.org/10.54117/ijbbs.v5i1.106>
- Idigo, M. A., Iheukwumere, I. H., Iheukwumere, C. M., Nnaeze, B. C., Akulue, C. J., Nwakoby, N. E., Ezendianefor, J. N., Ike, V. E., Nnaedozie, A. O., Ezekwueche, S. N., Anagor, I. S., Aniekwe, C. C., Ezeoke, F. C., Okereke, F. O., & Ochibulu, S. C. (2025r). Bacterial symbionts of insects: Exploring their role in insect nutritional composition. *IPS Interdisciplinary Journal of Biological Sciences*, 5(1), 177–186. <https://doi.org/10.54117/ijbbs.v5i1.105>
- Idigo, M. A., Iheukwumere, I. H., Iheukwumere, C. M., Nnaeze, B. C., Akulue, C. J., Nwakoby, N. E., Ezendianefor, J. N., Ike, V. E., Nnaedozie, A. O., Ezekwueche, S. N., Anagor, I. S., Aniekwe, C. C., Ezeoke, F. C., Okereke, F. O., & Ochibulu, S. C. (2025s). Augmenting rat lymphocyte function by bacterial symbiont of *Macrotermes* species. *IPS Journal of Applied Microbiology and Biotechnology*, 5(1), 281–290. <https://doi.org/10.54117/ijamb.v5i1.104>
- Idigo, M. A., Iheukwumere, I. H., Iheukwumere, C. M., Nnaeze, B. C., Akulue, C. J., Nwakoby, N. E., Ezendianefor, J. N., Ike, V. E., Nnaedozie, A. O., Ezekwueche, S. N., Anagor, I. S., Aniekwe, C. C., Ezeoke, F. C., Okereke, F. O., & Ochibulu, S. C. (2025t). Antimicrobial peptides from insects: A study on their efficacy against pathogens. *IPS Journal of Applied Microbiology and Biotechnology*, 5(1), 271–280. <https://doi.org/10.54117/ijamb.v5i1.103>
- Iheukwumere, C. M., & Iheukwumere, I. H. (2022a). Nutritive and Antinutrient Values of Soybean Condiments Produced from Indigenous Fermenters. *IPS Applied Journal of Nutrition, Food and Metabolism Science*, 1(1): 1–5. <https://doi.org/10.54117/iajnfms.v1i1.8>

- Iheukwumere, C. M., & Iheukwumere, I. H. (2022e). Hematological indices and sensory quality of fermented soybean condiments. *World Journal of Advanced Research and Reviews*, 14(2), 435-42.
- Iheukwumere, C. M., Ekesiobi, A. O., Iheukwumere, I. H., Ejike, C. E., Ilechukwu, C. C., Dim, C. N., & Ochibulu, S. C. (2025g). Dual Approach Therapy: Assessing *Xylopi aethiopica* and Ciprofloxacin Synergy against *Salmonella enterica* Serovar Typhi. *IPS Intelligentsia Multidisciplinary Journal*, 4(1), 27–31. <https://doi.org/10.54117/ijmi.v4i1.9>.
- Iheukwumere, C. M., Ekesiobi, A. O., Iheukwumere, I. H., Ejike, C. E., Ilechukwu, C. C., Dim, C. N., Ochibulu, S. C., Unegbu, C. C., & Egbuna, C. (2025h). Food Safety Implications: Assessing the Potential of *Desmodium velutinum* Leaves Extracts to Control the Most Predominant Fungal Contamination in Ready-To-Eat Fried Chicken. *IPS Journal of Nutrition and Food Science*, 4(3), 494–500. <https://doi.org/10.54117/ijnfs.v4i3.111>
- Iheukwumere, C. M., Iheukwumere, I. H., Nwakoby, N. E., Idigo, M. A., & Ike, V. E. (2025m). Evaluation of fermented chicken feather meal as a dietary supplement on rat lipid metabolism and immune response. *African Journal of Nutrition and Applied Research*, 1(1), 17–26. <https://doi.org/10.54117/qbwbbpp15>
- Iheukwumere, C. M., Iheukwumere, I. H., Okoli, U. O., & Ugwu, C. H. (2023a). Immunological Impact of Fermented Soybean Condiments Produced from Indigenous Fermenters. *Journal of Advances in Microbiology* 23(10): 27-37
- Iheukwumere, C. M., Iheukwumere, I. H., Ugwu, C. H., & Okoli, U. O. (2023b). Toxicity of Prepared Fermented Soybean Condiments from Indigenous Fermenters. *Journal of Advances in Microbiology* 23(10): 38 – 51.
- Iheukwumere, C. M., Umeaku, C. N., Chukwura, E. N., & Iheukwumere, I. H. (2022f). Characterization of the indigenous fermenters for the production of fermented condiments from soybean seeds. *World Journal of Advanced Research and Reviews*, 14(2), 423-434.
- Iheukwumere, I. H., & Ejike, C. E. (2017b). Comparative study of the inhibitory activities of *Ocimum gratissimum* and *Nepeta cataria* against *Salmonella enterica* serovar Typhi and their larvicidal effect against *Anopheles gambiae*. *African Journal of Education, Science and Technology (AJEST)*, 3(4), 16-24
- Iheukwumere, I. H., Ajeh, J. C., Iheukwumere, C. M., Ike, V. E., Obianom, A. O., Ihenatuoha, U. A., Igboanugo, E. U., Onwuasoanya, U. F., Okereke, F. O., Nnadozie, C. H., Agbaugo, C. F., Nwike, M. I., Nwakoby, N. E., & Ilechukwu, C. C. (2025c). Exploring the Phytochemical and Antimicrobial Properties of Fruit Vinegar: A Study on Phoenix Dactylifera and Malus Sylvestris. *IPS Journal of Applied Microbiology and Biotechnology*, 4(1), 115–122. <https://doi.org/10.54117/ijamb.v4i1.48>
- Iheukwumere, I. H., Ajeh, J. C., Iheukwumere, C. M., Ike, V. E., Obianom, A. O., Ihenatuoha, U. A., Igboanugo, E. U., Onwuasoanya, U. F., Okereke, F. O., Nnadozie, C. H., Agbaugo, C. F., Nwike, M. I., Nwakoby, N. E., & Ilechukwu, C. C. (2025d). Microbial Quality and Sensory Assessment of Vinegar from Date Palm and Apple Fruits: Implications for Consumer Preference. *IPS Journal of Nutrition and Food Science*, 4(2), 410–417. <https://doi.org/10.54117/ijnfs.v4i2.100>.
- Iheukwumere, I. H., Ajeh, J. C., Iheukwumere, C. M., Ike, V. E., Obianom, A. O., Ihenatuoha, U. A., Igboanugo, E. U., Onwuasoanya, U. F., Okereke, F. O., Nnadozie, C. H., Nwike, M. I., Nwakoby, N. E., & Ilechukwu, C. C. (2025f). Safety Evaluation of Vinegar from Phoenix Dactylifera and Malus Sylvestris: Toxicity and Acetic Acid Content. *IPS Journal of Applied Microbiology and Biotechnology*, 4(1), 123–131. <https://doi.org/10.54117/ijamb.v4i1.49>
- Iheukwumere, I. H., Ajeh, J. C., Iheukwumere, C. M., Ike, V. E., Obianom, A. O., Ihenatuoha, U. A., Igboanugo, E. U., Onwuasoanya, U. F., Okereke, F. O., Nnadozie, C. H., Agbaugo, C. F., Nwike, M. I., Nwakoby, N. E., & Ilechukwu, C. C. (2025k). Exploring the Phytochemical and Antimicrobial Properties of Fruit Vinegar: A Study on Phoenix Dactylifera and Malus Sylvestris. *IPS Journal of Applied Microbiology and Biotechnology*, 4(1), 115–122. <https://doi.org/10.54117/ijamb.v4i1.48>
- Iheukwumere, I. H., Amadi, E. R., & Chude, C. (2018b). Synergistic Effects of Probiotics and Autogenous Bacterin Against Inositol Negative Motile *Salmonella* Species. *Journal of Biology, Agriculture and Healthcare* 8(6).
- Iheukwumere, I. H., Amadi, R. E., Unaeze, B. C., & Campus, N. (2017c). Enterotoxigenicity Profile of *Salmonella enterica* Serovar Typhimurium in Suckling Albino Mice. *Journal of Natural Sciences Research* 7(14).
- Iheukwumere, I. H., Chukwura, E. I., & Chude, C. (2018c). In vivo activities of some selected antimicrobial agents against enteric bacteria isolated from chicken feeds on broiler layers. *Journal of Biology, Agriculture and Healthcare*, 8(6).
- Iheukwumere, I. H., Ejike, C. E., & Okeke, C. E. (2017d). A trial to prevent sorbitol negative *Escherichia coli* infections in chicks using autogenous bacteria and probiotics. *Journal of Natural Sciences Research*, 7, 56-63.
- Iheukwumere, I. H., Iheukwumere, C. M., Obianom, A. O., Nnadozie, C. H., Okereke, F. O., Onwuasoanya, U. F., ... Ihenatuoha, U. A. (2025a). Cross-Sectional Study of Different Strains of *Bacillus cereus* among Pap Sold in Major Towns in Ihiala LGA, Anambra State. *IPS Journal of Public Health*, 5(2), 199–204. <https://doi.org/10.54117/ijph.v5i2.39>.
- Iheukwumere, I. H., Iheukwumere, C. M., Obianom, A. O., Nnadozie, C. H., Okereke, F. O., Onwuasoanya, U. F., ... Destiny, E. C. (2025b). Cross-Sectional Study of Major Strains of *Salmonella enterica* Subspecies *enterica* Serovar Typhi among Borehole Used in Uli Community. *IPS Journal of Public Health*, 5(2), 205–210. <https://doi.org/10.54117/ijph.v5i2.40>.
- Iheukwumere, I. H., Iheukwumere, C. M., Obianom, A. O., Nnadozie, C. H., Okereke, F. O., Onwuasoanya, U. F., ... Udeagbara, O. E., Unaeze, B. C., Obiefuna, O. H., Ike, V. E., Onyemekara, N. N., & Ihenatuoha, U. A. (2025e). Quotidian of Substantial Strain of *Shigella dysenteriae* among Ready To-Eat Fruit Salad Sold in Uli Community. *Journal of Pollution Monitoring, Evaluation Studies and Control*, 4(1), 95–99. <https://doi.org/10.54117/jpmesc.v4i1.17>
- Iheukwumere, I. H., Iheukwumere, C. M., Obianom, A. O., Nnadozie, C. H., Okereke, F. O., Onwuasoanya, U. F., ... Ihenatuoha, U. A. (2025i). Cross-Sectional Study of Different Strains of *Bacillus cereus* among Pap Sold in Major Towns in Ihiala LGA, Anambra State. *IPS Journal of Public Health*, 5(2), 199–204. <https://doi.org/10.54117/ijph.v5i2.39>.
- Iheukwumere, I. H., Iheukwumere, C. M., Obianom, A. O., Nnadozie, C. H., Okereke, F. O., Onwuasoanya, U. F., ... Destiny, E. C. (2025j). Cross-Sectional Study of Major Strains of *Salmonella enterica* Subspecies *enterica* Serovar Typhi among Borehole Used in Uli Community. *IPS Journal of Public Health*, 5(2), 205–210. <https://doi.org/10.54117/ijph.v5i2.40>.
- Iheukwumere, I. H., Iheukwumere, M. C., & Nwakoby, N. E. (2022d). Sequential Pathogenicity Study of SOR+ and SOR- *Escherichia coli* Isolated from Roasted Meat. *IPS Intelligentsia Multidisciplinary Journal*, 1(1), 1-11.
- Iheukwumere, I. H., Nwankwo, A. K., Iheukwumere, C. M., Okorie, N. A., Nwakoby, N. E., Ekesiobi, A. O. and Okolo, O. C.. (2025n). Characterization and pathogenic profile of

- Aspergillus fumigatus* isolated from landfills. *Microbes and Infectious Diseases* 6(4)
- Iheukwumere, I. H., Obi, P. C. and Unaeze, B. C. (2017a). A trial to prevent *Vibrio cholerae* in albino mice using autogenous bacterin. *Advances in Life Science and Technology* 58:34–42
- Iheukwumere, I. H., Uneze, B. C., & Ejike, C. E. (2017e). Efficacy of some selected antimicrobial substances in prevention of enteric bacterial infection in broiler chicks. *J. Biol. Agriculture. Healthcare*, 7, 58-66.
- Iheukwumere, I.H. , Iheukwumere, C.M. , Nnadozie, H. C. ,Unaeze, C.B. , Obiefuna, O.H. Obianom, A.O. and Ejike, C. E. (2024). Hematotoxicological and mosquito larvicidal studies of crystal proteins secreted by *Bacillus thuringiensis* and *Bacillus sphaericus*. *Tropical Journal of Applied Natural Sciences* 2(2): 61 – 92.
- Iheukwumere, I.H. and Iheukwumere, M.C. (2022c). *Streptococcus suis* in Pigs and Environs: A Cross-sectional Study. *IPS Journal of Public Health*, 1(2), 9-12. <https://doi.org/10.54117/ijph.v1i2.4>.
- Iheukwumere, I.H. and Iheukwumere, M.C. (2022g). Cross-sectional Study of Multiple Antibiotic-resistant *Streptococcus suis* in Pigs and Environments. *IPS Interdisciplinary Journal of Biological Sciences*, 1(1), 19–21. <https://doi.org/10.54117/ijibs.v1i1.4>
- Iheukwumere, I.H., Dimejesi, S.A.,Iheukwumere, C.M., Chude, C.O., Nwaolisa, C.N.,Ukoha, C.C., Nwakoby,N.E., Egbuna, C. and Egbe, P.A. (2020) Diversity and molecular characterization of keratinophilic fungi from soil samples. *International Journal of Research Publication* 50(1); 1047 -1062.
- Iheukwumere, I.H., Iheukwumere, M.C. and Nwakoby, N.E. (2022b). Synergistic Effects of Probiotics and Autogenous Bacterin against *Salmonella enterica* Serovar Typhimurium Strain U288. *IPS Journal of Nutrition and Food Science*, 1(1), 1–5. <https://doi.org/10.54117/ijfns.v1i1.3>.
- Iheukwumere, I.H., Nwike, M. I., Iheukwumere, C.M., Ike, V.E., Obianom, A.O., Ihenatuoha, U.A., Igboanugo, E.U., Ekesiobi, A.O., Okereke, F.O., Obiefuna, O. H. Nnadozie, C.H., Agbaugo, C.F., Oduoye, O.T., Nwakoby, N.E., Ilechukwu, C. C., Ochibulu, S. C. and Ejike, C. E. (2025l). Extraction and Elucidation of Antibiotics from the Mycelia of *Aspergillus niger* Isolated from Poultry Farm against Enteric Bacterial Pathogens. *IPS Journal of Advanced and Applied Biochemistry*, 1(1), 1–10. <https://doi.org/10.54117/ijaab.v1i1.58>.
- Iheukwumere,I.H., Iheukwumere,C.M., Chude, C.O., Nwaolisa,C.N. and Egbe, P.A. (2020a). Comparative study of different clinical samples used for the diagnosis of staphylococcal systemic infections in apparent healthy students. *International Journal of Research Publications* 49(1): 1 – 10
- Ike, V. E., Iheukwumere, I. H., Iheukwumere, C. M., Dim, C. N., Ezendianefo, J. N., Egbe, P. A., Oragwu, I. P., Orji, C. C., Ogbonnaya, O. C., Onwuasoanya, U. F., Okereke, F. O., & Ochibulu, S. C. (2025a). Prevalence of *Bacillus cereus* in Powdered Soybean Sold in Uli Community, Anambra State: A Cross-Sectional Study. *IPS Journal of Basic and Clinical Medicine*, 2(3), 108–114. <https://doi.org/10.54117/ijbcm.v2i3.18>
- Ike, V. E., Iheukwumere, I. H., Iheukwumere, C. M., Dim, C. N., Ezendianefo, J. N., Egbe, P. A., Oragwu, I. P., Orji, C. C., Ogbonnaya, O. C., Onwuasoanya, U. F., Okereke, F. O., & Ochibulu, S. C. (2025b). *Bacillus cereus* in Uli's cornflour: A prevalence study. *IPS Journal of Nutrition and Food Science*, 4(3), 544–548. <https://doi.org/10.54117/8btte840>
- Ike, V. E., Iheukwumere, I. H., Iheukwumere, C. M., Dim, C. N., Ezendianefo, J. N., Egbe, P. A., Oragwu, I. P., Orji, C. C., Ogbonnaya, O. C., Onwuasoanya, U. F., Okereke, F. O., & Ochibulu, S. C. (2025c). Pathogenic Profile Analysis: In Vitro Screening of Enteric Bacteria from University Dusters. *IPS Journal of Applied Microbiology and Biotechnology*, 4(3), 187–191. <https://doi.org/10.54117/ijamb.v4i3.76>
- Ike, V. E., Iheukwumere, I. H., Iheukwumere, C. M., Dim, C. N., Ezendianefo, J. N., Egbe, P. A., Oragwu, I. P., Orji, C. C., Ogbonnaya, O. C., Onwuasoanya, U. F., Okereke, F. O., & Ochibulu, S. C. (2025d). Frozen Fish Pathogens: Antimicrobial Resistance and Public Health Implications. *IPS Interdisciplinary Journal of Biological Sciences*, 4(4), 138–143. <https://doi.org/10.54117/ijibs.v4i4.77>
- Ike, V. E., Iheukwumere, I. H., Iheukwumere, C. M., Dim, C. N., Ezendianefo, J. N., Egbe, P. A., Oragwu, I. P., Orji, C. C., Ogbonnaya, O. C., Onwuasoanya, U. F., Okereke, F. O., & Ochibulu, S. C. (2025e). Stream water quality assessment: Antibiotic resistance of Lac-positive enteric bacterial isolates. *Journal of Pollution Monitoring, Evaluation Studies and Control*, 4(2), 120–125. <https://doi.org/10.54117/jpmesc.v4i2.21.2025>
- Jamaludin, M.A., Hashim, D.M., Rahman, R.A., Ramli, M.A., Majid, M.Z.A., Othman, R. and Amin, A. (2016). Determination of permissible alcohol and vinegar in Shariah and scientific perspectives. *International Food Research Journal* 23(6):2737 – 2743.
- Jayamma, P., Shirnalli, G.G., Goudar, D.G. and Olekar, S.N. (2022). Isolation and identification of thermotolerant yeast isolates from different fruit wastes. *The pharma Innovation Journal* 11 (i): 855 – 859
- Jia, J., Wuc,Y., Kun, M., Fen, F., Zhangc, F. and Sheng, C. (2012). Biodiversity of yeasts, lactic acid bacteria and acetic acid bacteria in the fermentation of “Shanxi aged vinegar”, a traditional Chinese vinegar. *Journal of Food Microbiology* 30: 289 – 297.
- Johnston, C. S. (2009). Medicinal uses of vinegar. In Watson, R. R. (eds). *Complementary and Alternative Therapies and the Aging Population*, United States: Academic Press Pp. 433–443.
- Kechkar, M., Sayed, W., Cabrol, A., Aziza, M., Zaid, T.A., Amrane, A. and Djelal, H. (2019). Isolation and identification of yeast strains from sugarcane molasses, dates, and figs for ethanol production under conditions stimulating algal hydrolysate. *Brazilian Journal of Chemical Engineering* 36(1): 157 – 169
- Khin, H.A. and Thwe, L.K. (2016). Utilization of Fruit Waste (Pineapple peel) For Vinegar production. *Yadanabon University Research Journal* 7 (1): 1-11
- Klawplyapamornkun, T., Bovnsombut, S. and Bovonsoombut, S. (2015). Isolation and characterization of acetic acid bacteria from from fruits and fermented fruit juice for vinegar production. *Food and Applied Bioscience Journal* 3(1): 30 – 38
- Kumari, S., Jha, A.K. and Singh, A.K. (2019). Isolation and characterization of temperature and ethanol tolerant strain of *Saccharomyces cerevisiae* strains from naturally fermented juices. *Biosciences Biotechnology Research Asia* 16 (1): 97 – 103
- Loke, C., Lee, J., Sander, S., Mei, L. and Farella, M. (2016). Factors affecting intra-oral pH – A review. *Journal of Oral Rehabilitation* 43(10):778 – 7
- Maal, K.B., Shafie, R. and Kabiri, N. (2010). Production of apricot vinegar using an isolated *Acetobacter* strain from Iranian apricot. *International Journal of Biological Life Science* 6:230 – 233.
- Madrera, R.R.; Bedriñana, R.P. and Valles, B.S. (2017). Apple pomace by fermentation with autochthonous yeasts. *Food Science Technology* 79: 27–33.

- Magalhaes, K.T., Dias, D. and Pereira, G.V.M. (2011). Chemical composition and sensory analysis of cheese whey-based beverage using kefir grains as starter culture. *International Journal of Food Science and Technology* 46: 871 – 878.
- Manasseh, C.O., Logan, C.S.P., Ikeyi, A.P., Ede, K.K., Iheukwumere, I.H., Iheukwumere, C.M. and Ejike, C.E. (2025). Investigating the Effects of the Covid-19 Pandemic and Climate Risks on Trade Balance in Emerging Markets. *The Nigerian Health Journal* 25(2): 1-27. <https://doi.org/10.71637/tnhj.v25i2.914>
- Mathew, B., Agrawal, S., Nashikkar, N., Bundale, S. and Upadhyay, A. (2019). Isolation of Acetic Acid Bacteria and preparation of starter culture for Apple cider vinegar fermentation. *Advances in microbiology* 9:556-569: microbiological, chemical composition and sensory analysis. *Food Research International Journal* 48: 634 – 640.
- Mokhtar, S.I., Zakaria, F., Mohammad, A.A., Soon, S.W., Nuvshahida, A.S. and Latiff, Z.A.A. (2016). Study on the nutritional values and customer acceptance of Lansium domesticum and Nephelium lappaceum Newly fermented natural fruit vinegars in Malaysia. *Asian Pacific Journal of Advanced Business and Social Studies* 2(2): 404 – 413
- Nwike, M.I., Iheukwumere, I.H. and Uneze, B.C. (2017). Effect of Spices, pH and Temperature on the Survival and Multiplication of Staphylococcus aureus in Locally Made Soya Milk Drink. *Journal of Natural Sciences Research* 7(4).
- Obianom, A.O., Iheukwumere, I.H., Iheukwumere, C.M., Ochibulu, S.C., Nnadozie, H. C. and Ifenetu, F. C. (2024). Supersizing the inhibitory activity of Xylopi aethiopica extract against Vibrio cholerae using doxycycline. *Tropical Journal of Applied Natural Sciences* 2(2).
- Obiefuna, U. G., Umeh, S. O., & Iheukwumere, I. H. (2025a). Assessing the Impact of Glycemic Index on Microbial Quality and Storage Stability of Tomato Jam. *IPS Journal of Applied Microbiology and Biotechnology*, 4(3), 192–202. <https://doi.org/10.54117/ijamb.v4i3.81>
- Obiefuna, U. G., Umeh, S. O., & Iheukwumere, I. H. (2025b). Physicochemical and nutritional properties of tomato jam: Influence of sweetener type and glycemic index. *IPS Journal of Nutrition and Food Science*, 4(3), 561–568. <https://doi.org/10.54117/yazv0114>
- Obiefuna, U. G., Umeh, S. O., & Iheukwumere, I. H. (2025c). Micronutrient profile and acceptability of tomato jam: A comparison of high and low glycemic carbohydrate index sweeteners. *IPS Applied Journal of Nutrition, Food and Metabolism Science*, 3(2), 67–74. <https://doi.org/10.54117/pr4r6c73>
- Okeke, C. E. Iheukwumere, I. H. Ejike, C.E. (2017). Pathogenicity Study of Dematiaceae Fungi Isolated from Chicken Feeds on Immunoincompetent Chickens. *J. Biol. Agriculture. Healthcare* 7(4).
- Onuorah, S., Josen, L. and Obika, I. (2016). Production of vinegar from oil palm wine using *Acetobacter aceti* isolated from rotten banana fruits. *Universal Journal of Biomedical Engineering* 4(1): 1 – 5
- Puerari, C., Magalhaes, K.T. and Schwan, R.F. (2012). New cocoa pulp based kefir beverages: microbiology, chemical composition, and sensory analysis. *Food Research International Journal* 48: 634 – 640.
- Purugganan, M.D. (2015). Whole Genome Re-sequencing of Date Palms Yields Insights Into Diversification of a Fruit Tree Crop. *Nature Community Journal* 6:8824.
- Safrida, F.A.U., Gholib, M.M., Raissya, A., Yelly, A.P. and Nadila, F. (2023). Phytochemical characterization and sensory evaluation of vinegar from Melastoma malabathricum flower with variation in starter concentration and fermentation time. *Journal of Panelitian Pendidikan* 9(2): 706 – 713
- Saha, D., & Das, P. K. (2023). Perspective Chapter: Bioconversion of Agricultural and Food Wastes to Vinegar. In *Functional Food-Upgrading Natural and Synthetic Sources*. IntechOpen.
- Sengun, I.Y., Kilic, G. and Oturk, B. (2020). Screening physicochemical, microbiological and bioactive properties of fruit vinegars. *Food Sciences and Biotechnology* 29 (3): 401 – 408
- Siddeeg A, Zeng XA, Rahaman A, Manzoor MF, Ahmed Z, Ammar AF (2019). Quality characteristics of the processed dates vinegar under influence of ultrasound and pulsed electric field treatments. *Journal Food Science and Technology* 56(9):4380 – 4389.
- Singh, J. and Garg, A.P. (2022). Antimicrobial activity of apple cidar vinegar treated selected vegetables against common food borne bacterial pathogens. *Bioscience Biotechnology Research Communications* 15 (2): 350 – 356.
- Srivastava, V. and Rani, R. (2019). Isolation and characterization of acetoacetic bacteria from decomposed fruits. *Journal of Biological, Chemistry and Chron* 5(1): 123 – 129
- Tharinee, K., Sakunne, B. and Sittisin, B. (2015). Isolation and characterization of acetic acid bacteria from fruits and fermented fruit juices for vinegar production. *Food and Applied Bioscience Journal* 3(1): 30-38.
- Tumane, P. M., Sarkar, S., Wasnik, D.D. and Kolte, N. A. (2018). Production of vinegar from pineapple peels using *Acetobacter* species isolated from soil sample and its antibacterial activity. *International Journal of Life Sciences* 6(4):948 – 956.
- Ugwu, C. H., Iheukwumere, I. H., Iheukwumere, C. M., Ike, V. E., Dim, C. N., Ezendianefo, J. N., Egbe, P. A., Oragwu, I. P., Orji, C. C., Ogbonnaya, O. C., Onwuasoanya, U. F., Okereke, F. O., Oduenyi, P. M., & Ochibulu, S. C. (2025a). Maternal health and antibiotic resistance: *Klebsiella pneumoniae* isolates analysis. *IPS Journal of Public Health*, 5(3), 290–295. <https://doi.org/10.54117/s3tx6v26>
- Ugwu, C. H., Iheukwumere, I. H., Iheukwumere, C. M., Ike, V. E., Dim, C. N., Ezendianefo, J. N., Egbe, P. A., Oragwu, I. P., Orji, C. C., Ogbonnaya, O. C., Onwuasoanya, U. F., Okereke, F. O., Oduenyi, P. M., & Ochibulu, S. C. (2025b). Ocimum gratissimum Extract's Effectiveness against Vibrio cholerae from Uli Streams. *IPS Journal of Phytochemistry and Medicinal Plant Research*, 1(2), 15–19. <https://doi.org/10.54117/ijpmpr.v1i2.38>.