



Canadian Journal of Forest Research

Mixedwood management positively affects forest health during insect infestations in Eastern North America

Journal:	<i>Canadian Journal of Forest Research</i>
Manuscript ID	cjfr-2020-0462.R2
Manuscript Type:	Review
Date Submitted by the Author:	13-Apr-2021
Complete List of Authors:	MacLean, David; University of New Brunswick Clark, Kenneth; USDA Forest Service, Northern Research Station
Keyword:	Insect outbreaks, associational resistance, silviculture, defoliation, tree mortality
Is the invited manuscript for consideration in a Special Issue? :	Temperate mixedwood forests: ecology and management of hardwood-softwood mixtures

SCHOLARONE™
Manuscripts

Revision Submitted to Canadian Journal of Forest Research

Mixedwood management positively affects forest health during insect infestations in Eastern North America

David A. MacLean¹ and Kenneth L. Clark²

¹ Faculty of Forestry and Environmental Management, University of New Brunswick, POB 4400, Fredericton, NB E3B 5A3, Canada; macleand@unb.ca

² USDA Forest Service, Northern Research Station, Silas Little Experimental Forest, 501 Four Mile Road, New Lisbon, NJ 08064, USA; kenneth.clark@usda.gov

Abstract

Mixedwood stands are composed of hardwoods and softwoods, with neither comprising greater than 75-80% of basal area or aboveground biomass. By conferring associational resistance and greater resilience to forests when stressed or disturbed, the more diverse composition of mixedwood stands contributes to forest health. We analyzed three examples where mixedwood stands are more resistant to insect infestations in eastern North America. In balsam fir and spruce forests, susceptibility and vulnerability to spruce budworm infestations is reduced with increasing hardwood density. In mid-Atlantic oak-dominated forests, oak mortality following gypsy moth infestations was much greater in oak-dominated than in oak-pine mixedwoods, while pine mortality during southern pine beetle infestations was greater in pine-dominated than in oak-pine mixedwoods. Overall, mixedwood stands have greater resistance to infestations of defoliators and bark beetles and recover more rapidly from disturbances, reducing economic losses associated with tree mortality and mitigating short-term impacts to ecosystem functioning resulting from insect damage, especially carbon sequestration. Finally, we discuss challenges and opportunities for mixedwood management to minimize insect damage. Management strategies that incorporate mixedwood stands may provide better continuity in supply of forest products and ecosystem services in the face of projected increases in insect infestations associated with changing climate.

Keywords: Insect outbreaks, insect epidemics, associational resistance, silviculture, defoliation, tree mortality.

Introduction

Forests composed of a diversity of tree species have a greater range of shade tolerances, growth rates, longevity, phenology, and crown and root structures than pure species stands (Kabrick et al. 2017). Such mixtures diversify ecological niches and accommodate tree species with different ecological requirements through complementary resource use (Coates and Burton 1997; Kneeshaw and Prévost 2007). Further, high diversity forests have been observed to exhibit reduced susceptibility to insect infestations and pathogens compared to those dominated by a single species (Jactel and Brockerhoff 2007; Jactel et al. 2017; Zhang et al. 2018; Fig. 1). Based on a worldwide meta-analysis of 119 studies that compared herbivory in single-species and mixed forests, Jactel and Brockerhoff (2007) concluded that herbivory was significantly reduced in more diverse forests, but this varied with the host specificity of insects. Herbivory by oligophagous species (those that consume plants of closely related genera or a single taxonomic family) was nearly always reduced in diverse forests, but the response of polyphagous species that consume a wide variety of plants was weaker (Jactel and Brockerhoff 2007). The species composition of tree mixtures was more important than species richness per se, and effects of diversity on herbivory were greater when mixed forests comprised taxonomically more distant tree species or when the proportion of non-host exceeded host trees (Jactel and Brockerhoff 2007; Fig. 1).

Diverse species composition of forests may decrease the likelihood of detection by, and/or vulnerability to, herbivores, termed associational resistance (Barbosa et al. 2009). Specific mechanisms involved in greater associational resistance to insect pests include: (i) host dilution due to intermingling with non-host species in diverse stands, affecting how insects perceive the overall quality of a stand and lowering their probability of finding and feeding on host trees; (ii) lower visual or chemical apparency of individual host trees within a stand when their frequency

is lower relative to non-host trees; and (iii) greater diversity of parasitoids or predators associated with increased diversity of predator habitats. In contrast, increased detection by and/or vulnerability to herbivores in diverse stands is an example of associational susceptibility. Associational resistance is more likely with insects while associational susceptibility is more likely with mammals (Barbosa et al. 2009).

One commonly used category of diverse forests is mixed hardwood-softwood (broadleaved-conifer) forests or “mixedwoods”, which are composed of stands where neither group comprises greater than 75 to 80% of basal area or aboveground biomass (e.g., Kabrick et al. 2017). There is considerable interest in mixedwood silviculture and management to diversify timber products and ecosystem services, and maintain diverse forest landscapes (Kenefic et al. 2021; Vickers et al. 2021). Mixedwoods comprise a large portion of North American forests, with 111 million hectares of mixedwood forest (defined as mixed coniferous and broadleaf trees, each between 25% and 75%) representing 17% of the total (North American Forest Database 2020, Vickers et al. 2021). Some mixedwood species combinations are challenging to manage (Smith 1986, Kern et al. 2021) and many have historically been managed for softwood species and products (e.g., Westveld 1930; Lieffers and Beck 1994; Etheridge et al. 2006; Kenefic et al. 2021). Mixedwood stands may be multistoried, especially in early stand development stages, or have only a single canopy layer, and structure often changes over time from an early dominant hardwood overstorey and a softwood understory, which gradually grows into the main canopy and eventually overtops the hardwood trees (Comeau 1996). This pattern contrasts with pine-dominated forests in the mid-Atlantic and mid-western regions, which are maintained by disturbance (primarily fire), and oaks and other hardwoods tend to dominate through time (Little 1998; Nowacki and Abrams 2008; Kenefic et al. 2021).

Paralleling observations in forests with high diversity, there is strong evidence that

mixedwood management has benefits in reducing insect herbivory and associated impacts compared to pure species stands, with either the softwood or hardwood component likely to experience less damage or mortality depending on the species of insect (Fig. 1). These benefits often exceed the effect expected by just replacing host with non-host species (Jactel and Brockerhoff 2007; Campbell et al. 2008; Jactel et al. 2017; Zhang et al. 2018). In addition, mixedwoods can potentially provide greater timber volume or biomass, more diverse habitats, and be more resistant or resilient to pathogens and other disturbances than pure stands (Cavard et al. 2011; Kabrick et al. 2017). Mixedwoods are also thought to be more compatible with projected future climates; all five mixedwood stand types examined by Kabrick et al. (2017) appeared to be better adapted to a changing climate than pure softwood stands.

Insect damage that results in tree growth reduction or mortality has multiple effects on community dynamics and ecosystem functioning, including succession, competition, productivity, nutrient cycling, energy flow, and carbon sequestration (Fig. 1). Most insects are specialists that target either individual (or multiple) conifer or deciduous species. Therefore, the most basic difference between mixedwood (or any diverse species stand) and pure stands is that a portion is not susceptible to damage. The non-susceptible species may undergo increased growth and reduced competition in response to the insect outbreak of host species (e.g., increased growth of hardwood species in mixed hardwood-spruce-fir stands during spruce budworm outbreaks; Hennigar et al. 2007). A variety of short-term impacts of insect infestations on ecosystem functioning are relatively well-characterized in some regions (e.g., in mid-Atlantic; Clark et al. 2010, 2014, 2018; Amiro et al. 2010; Deel et al. 2012; Renninger et al. 2014). In addition, a number of simulation models have captured the overall dynamics of carbon, hydrologic, and nutrient cycling associated with these disturbances (Medvigy et al. 2012; Kretchun et al. 2014; Xu et al. 2017). In summary, infestations of defoliators and bark beetles

initially reduce leaf area of infested stands as a function of defoliation intensity or tree and sapling mortality. This causes an immediate decline in photosynthetic capacity, which decreases net ecosystem productivity (NEP) and C sequestration, but enhances litterfall of nutrient-rich litter, and in the case of defoliators, frass. Similarly, reduced leaf area results in lower rates of evapotranspiration (Et; Clark et al. 2012). Compensatory photosynthesis by the remaining foliage, which is typically exposed to higher light levels, and the rapid cycling of nutrients from nutrient-rich frass and/or litter facilitates resprouting of new foliage (Curtis and Gough 2018; Hornslein et al. 2019). As a result, gross ecosystem productivity (GEP), Et, and ecosystem water use efficiency (WUE_e), defined as the amount of CO_2 assimilated per unit of water transpired, often recover rapidly following insect damage (Clark et al. 2014; Guerrieri et al. 2019). In contrast, tree and sapling mortality resulting from repeated defoliation or bark beetle infestations increases standing dead and coarse woody debris, resulting in increased heterotrophic respiration and a long-term depression of NEP (Renninger et al. 2014; Xu et al. 2017; Clark et al. 2018; Fig. 1). Defoliation or reduced leaf area increases light penetration into the understory, which promotes growth of advance regeneration and non-host species (e.g. Nie et al. 2019), and may make the stand more susceptible to fire (e.g. Stocks 1987) or windthrow (Taylor and MacLean 2009).

In this paper, we review the role of mixedwoods in three important insect-forest systems in eastern North America. We focus on effects on defoliation or damage, growth reduction, tree mortality, and ecosystem functioning, primarily patterns of carbon sequestration during and following insect infestations. We describe the evidence for, and mechanisms involved in, mixedwood resistance and resilience to insect damage in spruce budworm (*Choristoneura fumiferana* Clem.) in balsam fir (*Abies balsamea* L. Mill.) and spruce (*Picea* spp.) forests, gypsy moth (*Lymantria dispar* L.) in oak (*Quercus* spp.) and pine (*Pinus* spp.) forests, and southern

pine beetle (*Dendroctonus frontalis* Zimmermann) in pine, oak and other hardwood forests.

Effects of Mixedwoods on Insect Damage in Three North American Ecosystems

1. Spruce budworm and balsam fir defoliation, growth reduction and mortality

Spruce budworm causes the most damaging outbreaks in eastern North America, periodically occurring about every 30-40 years over millions of hectares of balsam fir and spruce forests. When budworm larvae defoliate a tree, they remove a portion of the photosynthetic factory as well as sites where chemicals such as growth hormones are produced, and this affects many vital functions (MacLean 1985). Repeated annual defoliations, which typically can last for 5-10 years during an outbreak, cause reduction of wood production, tree mortality, and decreased stand yield (e.g., Kulman 1971). Tree mortality usually begins after about 4 years of defoliation for balsam fir and 6-7 years for spruce (MacLean 1980). This is not new information; data on losses caused by spruce budworm were published nearly 100 years ago by Swaine and Craighead (1924) and Craighead (1925). There has been continued interest and research to the present about spruce budworm damage because of the large extent of outbreaks (covering 52 million ha of Canada in 1975; MacLean 2016), and the severe tree mortality (e.g., MacLean 1980) and timber supply and economic losses (e.g. Chang et al. 2012; Hennigar et al. 2013).

Changing stand composition by maintaining or increasing hardwoods in balsam fir stands has long been proposed as a silvicultural technique to reduce susceptibility (probability of being attacked) and vulnerability (probability of growth reduction or mortality once attacked) to spruce budworm (Westveld 1946; Blum and MacLean 1984, 1985; Carlson and Wulf 1989; Hudak 1991). Defoliation level (susceptibility) declines in the order balsam fir > white spruce (*Picea glauca* [Moench] Voss) > red spruce (*Picea rubens* Sarg.) > black spruce (*Picea mariana* [Mill.] B.S.P.) (Hennigar et al. 2008; Table 1). Thus, silvicultural recommendations to reduce spruce

budworm damage have largely been to reduce balsam fir at the stand and landscape level through spruce plantation establishment or pre-commercial thinning that favors spruce species over balsam fir (Blum and MacLean 1984, 1985). In addition, Croome (1970) and van Raalte (1972) suggested that 30-60% of stand composition in nonhost species was needed for a major reduction in susceptibility. Balsam fir mortality caused by spruce budworm defoliation declined from >80% to <50% as the percentage of hardwoods increased from 0% to 30% (Turner 1952). Mixedwood stands with at least 30% hardwoods sustained budworm-caused mortality that was 30% less than in pure fir stands (MacLean 1980).

Hypotheses proposed to explain the mechanisms resulting in lower defoliation and impacts with increased hardwood content include (i) dispersal losses of first- and second-instar spruce budworm larvae during dispersal to other hosts (Kemp and Simmons 1978; Zhang et al. 2020), both within stands and between stands if the stand edge to area ratio is high (Roland 1993; Kouki et al. 1997); (ii) a more abundant and diverse complex of budworm natural enemies, including higher tachinid parasitism of larvae and ichneumonid parasitism of pupae in fir stands isolated by hardwood forests (Cappuccino et al. 1998) and higher hymenopteran parasitism of budworm eggs in mixed fir-hardwood than in balsam fir stands (Quayle et al. 2003); and (iii) higher larval predation by a potentially more diverse community of birds at forest edges and in mixedwoods (Crawford and Jennings 1989, Drapeau et al. 2000). Quayle et al. (2003) specifically concluded that parasitism was particularly important in mixedwood stands.

Mixedwood stand composition has been shown to reduce defoliation levels during spruce budworm outbreaks. Defoliation in 25 mixed balsam fir-hardwood stands, over 5 years near the end of a spruce budworm outbreak in New Brunswick, Canada, was significantly negatively related to hardwood content (Su et al. 1996). Ten balsam fir stands with <40% hardwoods sustained 58-71% defoliation, whereas 15 stands with >80% hardwoods had only 12-15%

defoliation (Su et al. 1996). Zhang et al. (2018) also tested the influence of stand composition on defoliation, but during the first 5 years of a spruce budworm outbreak in Québec, Canada. Defoliation of fir in 27 balsam fir-hardwood plots representing softwood, mixedwood, and hardwood compositions was significantly lower as hardwood content increased. Percent hardwood content and average defoliation in softwood plots (an indicator of overall outbreak severity) were the most important variables in a Random Forests model of annual defoliation per plot (Zhang et al. 2018). Mean defoliation differed significantly among stand types, with softwood > mixedwood > hardwood in 2 years, and with softwood > hardwood in all 5 years tested. Zhang et al. (2020) determined that increasing hardwood content increased second-instar budworm larvae (L2) dispersal losses, but in their case, stand type did not significantly affect parasitism rates. L2 dispersal loss was less in softwood than in mixedwood or hardwood stand types based on measurements on ground traps and less in softwood and mixedwood than in hardwood stands based on sampling budworm larvae on regeneration (Zhang et al. 2020).

Hardwood content of surrounding forest landscapes also influences spruce budworm damage. Campbell et al. (2008) evaluated factors influencing radial growth reductions during a budworm outbreak, and found that hardwood content of surrounding (within 1 km) landscapes was a significant factor explaining variance in budworm-caused growth reductions. For example, in a peak outbreak year, mean growth reduction was ~40% in stands with <50% surrounding hardwood content versus ~20% with >50% surrounding hardwoods (Campbell et al. 2008).

Associational susceptibility has also been observed in some species mixtures during spruce budworm infestations, resulting in defoliation and subsequent growth reduction and mortality of non-preferred species. For example, the presence of balsam fir in a stand can increase defoliation of the lower susceptibility black spruce (Bognounou et al. 2017) and eastern hemlock (*Tsuga canadensis* (L.) Carrière), although usually considered non-host species for

209 spruce budworm.

210 If the management objective is primarily to produce softwood products, replacing balsam
211 fir with hardwoods also reduces the fir softwood volume, so what is the best hardwood-softwood
212 level? It depends upon outbreak severity. Needham et al. (1999) analyzed trade-offs between
213 reduced defoliation and mortality versus the amount of balsam fir volume reduction from
214 allocating growing space to hardwoods. Optimum hardwood levels varied with the severity of
215 spruce budworm attack: below a 45% defoliation level (5-year average), the reduction of fir
216 volume replaced by hardwoods exceeded the volume protected, but with defoliation >75%,
217 maximum softwood production was achieved with hardwood content comprising 50% of stand
218 volume (Needham et al. 1999). This is depicted for generalized balsam fir, mixedwood, and
219 hardwood forest classes in Fig. 2. On average, the spruce budworm infestation reduced balsam
220 fir merchantable volume by 88% in mature balsam fir stands but only by 28% in mixedwood
221 stands and balsam fir volume increased slightly during the outbreak in hardwood stands (Fig. 2).

222 Spruce budworm outbreaks influence stand species composition. In mature balsam fir
223 stands, species composition 30 years following a budworm outbreak was closely related to
224 predisturbance species composition, but in immature fir stands, hardwood composition increased
225 by up to 27% (Spence and MacLean 2012). Seedling density of advanced balsam fir regeneration
226 decreased over a 10-year outbreak period, but average height of surviving seedlings increased by
227 up to five times as the canopy cover decreased with tree mortality (Spence and MacLean 2012).

228 Spruce budworm outbreaks and mixedwood content also have major effects on forest
229 carbon. Hennigar and MacLean (2010) calculated potential spruce budworm outbreak and
230 management strategies (salvage, biological insecticide application) effects on forest and wood
231 product carbon (C) over 50 years for a 210,000-ha intensively managed forest in New
232 Brunswick, Canada. Harvest volume, forest C storage, and wood product C storage were

projected to be reduced, compared to a no outbreak scenario, by 1.34 million m³, 1.48 Mt C, and 0.26 Mt C, respectively. Modelling analyses indicated that use of salvage and harvest replanning plus a biological insecticide applied aurally to 40% of susceptible forest area were projected to reduce future spruce budworm impacts on harvest losses by 73%, forest C losses by 41%, and wood product C losses by 56% (Hennigar and MacLean 2010). Hardwood volume increases 15 years after the outbreak resulting from reduced stand competition from mortality of spruce-fir averaged one-third to one-half in immature mixedwood stands, which offset some of the C impacts. This effect was augmented by lower specific gravity of spruce (~0.37) and fir (0.33) compared with hardwood species (e.g., poplar, 0.35; white birch, 0.48; red maple, 0.49; sugar maple, ~0.55) (Green et al. 1999). As a result, C in living biomass 15-years after the outbreak was reduced by 8%, substantially less than the 24% total volume reduction. Dymond et al. (2010) also showed that a spruce budworm outbreak would significantly affect the C sink or source status of a large landscape (106,000 km² of forest in eastern Québec). Projections with a model of ecosystem C showed that spruce budworm-caused mortality decreased stand-level merchantable C stocks by up to 90% and changed the landscape from a sink to a source, due to reduced net primary production (Dymond et al. 2010). The ecosystem C stock was reduced on average by 2 Tg C yr⁻¹ for the simulated area (Dymond et al. 2010). However, Dymond et al. (2010) did not explicitly model competitive release of hardwoods following severe defoliation or reduced defoliation levels of host species associated with hardwoods.

2. Gypsy moth and oak defoliation, growth reduction and mortality

European gypsy moth is one of the more destructive forest insects in Eastern North America, defoliating an average of 400,000 ha of forest per year, with a maximum of 5 million ha in 1981 (McManus et al. 1992). Since their introduction in Massachusetts in 1869, European

gypsy moth have spread throughout the Northeast and mid-Atlantic regions of the US, to portions of the Midwest US, and to Southeastern Canada (USDA 2020). Gypsy moth defoliate primarily oaks and other hardwood species, and although severe infestations can result in complete defoliation of both hardwood and softwood species, repeated defoliation over consecutive growing seasons typically results in significant mortality of vulnerable hardwood trees and saplings (Mosher 1915; Brown et al. 1988; Davidson et al. 1999; Morin and Liebhold 2015; Table 1). Gypsy moth damage has reduced oak productivity and regeneration across much of the range where they overlap, contributing to regional oak decline (Morin and Leibhold 2015; Fei et al. 2019). Tree mortality has decreased the supply of high-quality hardwood sawlogs and resulted in increased density of less commercially important species such as American beech (*Fagus grandifolia* Ehrh.) and red maple (*Acer rubrum* L.) (www.fia.fed.us). There is continued interest in control and mitigation of gypsy moth damage, because gypsy moth have impacted less than 25% of their potential range in North America, and all vulnerable hardwood species are potentially at risk.

Stand composition alters susceptibility and mortality from gypsy moth (Mosher 1915; Clement and Munro 1917; Brown et al. 1988). Oak-dominated stands in oak-hickory and northern hardwood forests are most vulnerable, with repeated defoliation over consecutive growing seasons resulting in progressively greater levels of reduced growth and mortality (Davidson et al. 1999; Fei et al. 2019). Stands consisting of mixtures of softwoods and hardwoods demonstrate both associational resistance and associational susceptibility to gypsy moth infestations, depending on defoliation intensity and stand species composition (e.g., Jactel et al. 2017). For example, in a recent severe gypsy moth infestation on the mid-Atlantic coastal plain, approximately 40% of oak trees died following repeated defoliation over two to three years in oak-dominated stands, while oak mortality was much lower in mixedwood oak-pitch

pine stands, and minimal in pitch pine-dominated stands where post oak (*Q. stellata* Wangenh.) and white oak (*Q. alba* L.) represented < 10% of basal area (Fig. 3a; Clark et al. 2010, 2018). Further south, loblolly pine-hardwood stands experienced lower mortality from gypsy moth than oak-dominated stands, with greater mortality occurring in loblolly pine-oak stands than in loblolly pine-sweetgum (*Liquidamber styraciflua* L.) stands, but mature loblolly pine mortality was low (Eisenbies et al. 2007).

Associational susceptibility of softwood trees and saplings to gypsy moth infestations in mixedwood stands depends on the softwood species and canopy position as well as the density of vulnerable hardwoods and intensity of defoliation. For example, > 80% defoliation of eastern white pine (*P. strobus* L.) in stands containing northern red oak (*Q. rubra* L.) significantly increased mortality of intermediate and suppressed pines (Brown et al. 1988, Gottschalk and Twery 1989), while mortality of pitch (*P. rigida* Mill.) and shortleaf pine (*P. echinata* Mill.) trees and saplings was minimal in stands containing black oak (*Q. velutina* Lam.) (Campbell and Garlo 1982, Clark et al. 2018). Eastern hemlock in red oak–softwood stands were highly vulnerable to defoliation by gypsy moth, with highest mortality occurring in intermediate and suppressed trees and understory saplings (Stephens 1988). Hennigar et al. (2007) observed that gypsy moth infestations resulted in unexpectedly large growth reductions and mortality of balsam fir, white birch (*Betula papyrifera* Marsh.), and trembling aspen (*Populus tremuloides* Michx.) in stands containing northern red oak in central New Brunswick, Canada. Collectively, these results indicate that mixedwood stands often have lower tree mortality following gypsy moth infestations, but interactions between defoliation intensity and frequency, and stand species composition are important in determining patterns of defoliation, growth reduction and mortality.

Silvicultural treatments to enhance resistance to gypsy moth have been proposed since the early 1900s, but few long-term experiments testing the increased resistance of mixedwood

vs. hardwood-dominated stands have been conducted (Clement and Munro 1917; Muzika 2017). Initial prescriptions sought to shift the emphasis from eradication to mitigation by reducing the density of vulnerable species (Behre 1939). Pre-infestation crown condition is a strong indicator of mortality of vulnerable species following gypsy moth infestations, but not necessarily crown class or tree size, thus mortality is difficult to predict *a priori* in the field for all but the most stressed trees. Thinning treatments alone have been inconclusive, with some studies demonstrating reduced mortality and greater stem increment growth following gypsy moth infestations (Muzika et al. 1998) but not others (Fajvan et al. 2008). Prescriptions for sanitation thinning and pre-salvage thinning have been used to remove vulnerable species and increase stand vigor, with thinning of vulnerable species to < 50% of stand basal area described as beneficial and thinning to < 20% of stand basal area as best for reducing mortality of commercially important oaks, although these levels may be difficult and expensive to achieve (Gottschalk and Twery 1989; Gottschalk 1993).

In some cases, uneven-aged forests have greater resistance to insect infestations and other disturbances than even-aged forests (Jactel et al. 2017). However, many of the forests in the mid-Atlantic and northeastern US impacted by gypsy moth infestations are intermediate aged, following recovery from extensive logging, agricultural use and abandonment, and in some areas, severe wildfires, and thus do not encompass the full range of age classes (Pan et al. 2011; Duveneck et al. 2017; Stambaugh et al. 2018). As these forests age, simulating natural successional processes (or delaying them) through silvicultural management to create more resistant and resilient mixedwood forests is possible (Webster et al. 2018; Willis et al. 2019; Kenefic et al. 2021). Treatments combining multi-aged mixedwood management in the prescription could be particularly successful (Castagneyrol et al. 2020). In some mid-Atlantic forests, insect infestations are doing just that; for example, by promoting the regeneration and

growth of pine seedlings and saplings in previously oak-dominated stands, gypsy moth infestations and subsequent mortality may be driving the formation and persistence of uneven-aged mixedwood stands. Overstory mortality in oak-dominated stands releases pine seedlings and saplings in the understory, and a decade following gypsy moth infestation, pine trees and saplings increased from 25% to 39% of basal area, with a 10-fold increase in pine saplings (Fig. 3a). Relative basal area has converged with that characterizing oak-pine mixedwood stands (Fig. 4, left arrow). As these uneven-aged mixedwood stands recover, they are likely to be more resistant to damage from future gypsy moth infestations, much like the mixedwood stand in Fig. 3a. Overall, designing and implementing disturbance-based management strategies that mimic similar responses in other forest systems could help mitigate the impacts of future insect infestations (Webster et al. 2018; Puettmann and Messier 2020).

Gypsy moth infestations have reduced forest productivity and carbon sequestration by impacted forests throughout the northeastern and mid-Atlantic regions of the US, with impacts generally proportional to defoliation severity and the density of vulnerable species (Clark et al. 2010; Kautz et al. 2017; Xu et al. 2017; Kosiba et al. 2018). US Forest Service Forest Inventory and Analysis (FIA) data indicate that net primary productivity (NPP) in the dominant oak-hickory, mixedwood oak-pine, and pine-dominated forest types in the absence of gypsy moth infestations are similar, and net ecosystem productivity (NEP) ranges between 1.7-2.1, 1.2-1.7, and 1.0-1.6 T C ha⁻¹ yr⁻¹, respectively (<https://www.fia.fs.fed.us>). These are similar to pre-infestation annual NEP values of 1.8, 1.4 and 1.7 T C ha⁻¹ yr⁻¹ for oak-dominated, mixedwood, and pine-dominated stands in the Pinelands National Reserve of New Jersey (Table 2; Clark et al. 2010, 2018), about 20% of which were heavily to severely defoliated (New Jersey Department of Agriculture 2004-2016; USDA Forest Service 2005-2016), reducing NEP to -2.5, -0.2 and 0.4 T C ha⁻¹ yr⁻¹, respectively (Table 2). Clark et al. (2010) estimated that annual

landscape-scale C uptake was reduced by approximately 40%. Mortality in oak-dominated stands reduced oak tree and sapling basal area compared to peak pre-defoliation values and in the following years, there was no net increase in basal area or aboveground biomass (Fig. 3a). NEP has been depressed for a decade, averaging only 0.4 T C yr^{-1} , 22% of pre-defoliation values (Clark et al. 2018), and enhanced ecosystem respiration can depress NEP for up to two decades as standing dead and coarse woody debris decompose (Renninger et al. 2014).

In contrast, in mixedwood oak-pine stands, oak tree and sapling mortality was minor following infestations (Fig. 3a). Basal area of trees and saplings had increased by 22% by 2018 compared to 2004, with increases in both pine and oak tree basal resulting from ingrowth and sapling recruitment (Fig. 3a). Mixedwood oak-pine stands that incurred lower amounts of mortality and coarse woody debris recruitment exhibited lower ecosystem respiration rates and a more rapid recovery in NEP following infestations (Kretchun et al. 2014). In pine-dominated stands, oak mortality was minimal following gypsy moth infestations. While defoliation of oaks and shrubs reduced annual NEP to only 23% of values in undisturbed years, ingrowth and sapling recruitment increased basal area by 56% between 2004 and 2018 (Table 2; Fig. 3a), and NEP was similar to pre-infestation levels within two years (Clark et al. 2018). Overall, growth and NEP were greater in mixedwood stands during gypsy moth defoliation and recovered to pre-infestation levels rapidly compared to severely infested oak-dominated stands.

3. Southern pine beetle and pine mortality

Southern pine beetle is native to the southeastern USA, south through Mexico, and west to Texas, but recently its range has extended northward on the Atlantic coastal plain into New Jersey and Long Island, New York, and it has been detected further north in forests dominated by pitch pine (*P. rigida* Mill.) (Weed et al. 2013; Clark et al. 2017; Dodds et al. 2018; Heuss et al.

2019). It has also expanded westward into southern Missouri, Illinois and Ohio. Southern pine beetle infests primarily hard pines (Nowak et al. 2015; Dodds et al. 2018). On the Atlantic coastal plain, loblolly (*P. taeda* L.), shortleaf (*P. echinata* Mill.), pitch, and Virginia (*P. virginiana* Mill.) pines are susceptible (Table 1), while white pine susceptibility in northeastern forests is not as well known (Dodds et al. 2018). Further northward expansion into forests composed of red pine (*P. resinosa* Aiton) and jack pine (*P. banksiana* Lamb.) is projected in the future, following isoclines of increasing winter temperatures with changing climate (Lesk et al. 2017; Dodds et al. 2018). Pitch pine is not a particularly important commercial species in the region, but white pine is, thus further expansion of southern pine beetle into forests of the northeastern US and Canada could impact softwood sawlog and pulpwood products.

Overall stand density and the relative importance of pines are critical factors in the probability of southern pine beetle leading to pine tree and sapling mortality, with basal areas > 28 m² ha⁻¹ considered highly susceptible (Guldin 2011). Stand density (as reflected in basal area) is proportional to turbulence regimes within forest canopies, altering dispersion of aggregation pheromone released by southern pine beetle (Thistle et al. 2004; Edburg et al. 2010). The recent infestations in New Jersey and New York occurred in relatively dense pine-dominated stands that exceeded the target basal area of 18 m² ha⁻¹ that has been effective in reducing southern pine beetle damage in southern US forests (see below). Recent infestations resulted in extensive mortality of pitch, shortleaf and Virginia pine trees, averaging 94% of pine tree basal area, while pine saplings were infested less frequently, and pine sapling basal area was similar in uninfested and infested areas (Fig. 3b; Clark et al. 2017; Aoki et al. 2018). Southern pine beetle infestations (and suppression treatments) had little effect on oak trees and saplings or other hardwoods such as red maple and black gum (*Nyssa sylvatica* Marshall) (Fig. 3b). Southern pine beetle rarely impacted pines in oak-dominated stands, and tree mortality was lower in mixedwood stands, a

pattern also documented by Heuss et al. (2019) in New York, where mortality averaged 60% of total basal area in pitch pine-dominated stands, versus 50% and 35% in unmanaged and managed pine-oak stands, respectively.

Silvicultural treatments to reduce southern pine beetle damage include thinning of dense stands before outbreaks, with a target basal area for southern pines of 18 m² ha⁻¹ (Guldin 2011). In unmanaged pitch pine stands in New Jersey and New York, thinning to reduce basal area to approximately 18 m² ha⁻¹ and prescribed fire to reduce understory competition has been recommended (Dodds et al. 2018). Sanitation cutting of infected green trees and salvage cutting of dead trees impacted by southern pine beetle are often used as suppression treatments to isolate an infestation and reduce its spread (Guldin 2011). Much like the long-term effects of gypsy moth in oak-dominated stands, changes in stand composition and structure in previously pine-dominated stands infested by southern pine beetle in southern New Jersey have converged on structural attributes characterizing mixedwood stands. Relative basal area of trees and saplings in untreated pine stands are now similar to that characterizing mixed oak-pine stands (Fig. 4; right arrow). Heuss et al. (2019) reported a similar pattern following southern pine beetle infestations and pine mortality in New York.

Recent infestations of southern pine beetle in the mid-Atlantic region have significant impacts on ecosystem functioning and carbon dynamics of pine-dominated stands (Fig. 3b; Heuss et al. 2019; Clark et al. 2020). Projected leaf area and canopy nitrogen content of remaining trees and saplings in infested stands in New Jersey were less than half of pre-infestation values three to five years following extensive pine mortality (Clark et al. 2017; 2020).

Clark et al. (2014, 2018) and others have shown significant positive relationships between leaf area, canopy N content and GEP, indicating that GEP and NEP will be reduced in the decades following southern pine beetle infestations. Simulations of C dynamics indicated that

annual NEP averaged $1.8 \text{ T C ha}^{-1} \text{ yr}^{-1}$ in uninfested pine-dominated stands, compared to $-5.2 \text{ T C ha}^{-1} \text{ yr}^{-1}$ in infested stands and $-4.8 \text{ T C ha}^{-2} \text{ yr}^{-1}$ in infested stands where cut-and-leave suppression treatments had been conducted (Clark et al. 2020). In infested stands, standing dead trees and coarse woody debris increased simulated annual ecosystem respiration up to 1.2 times the values for uninfested areas, and increased heterotrophic respiration will further depress NEP (Renninger et al. 2014; Clark et al. 2018). Because reduction in GEP and increased ecosystem respiration are largely proportional to pine mortality, net productivity and NEP will recover relatively rapidly in mixedwood stands that experience lower levels of damage and mortality (Heuss et al. 2019; Clark et al. 2020).

Although seasonal differences occur, carbon and hydrologic fluxes indicate that oak-pine mixedwood stands function similarly to oak- and pine-dominated stands when values are integrated over annual time scales (Clark et al. 2010, 2014). Because oaks and other hardwoods experience lower mortality in mixedwood stands following gypsy moth infestations, and pitch and shortleaf pine experience lower mortality in mixedwood stands during southern pine beetle infestations (Clark et al. 2018; Heuss et al. 2019), mixedwood stands maintain greater continuity in ecosystem functioning through time. Both decreased loss of leaf area and foliar nutrients in the short-term and reduced production of snags and coarse wood following mortality in the long-term contribute to the maintenance of ecosystem functioning.

Principles, commonalities, and management implications

Disturbance regimes in eastern North American forests over the last two decades have been dominated by insect damage, with impacts resulting from infestations of both native and non-native insects (Lovett et al. 2016; Kautz et al. 2017; Kosiba et al. 2018; Pasquarella et al. 2018; Fei et al. 2019). In previous decades (from 1975-2000), the total (summed) area of

moderate-severe (30–100%) defoliation or beetle-killed trees was 709 million hectares, with 64% caused by spruce budworm, 27% by forest tent caterpillar (*Malacosoma disstria* Hübner), 2% by jack pine budworm (*Choristoneura pinus* Freeman), and the remaining 7% caused by 39 other recorded insect species (MacLean 2016). In many cases, mixedwood stands are less vulnerable to insect infestations than forests dominated by a single species or genus, especially for monophagous insects (Jactel and Brockerhoff 2007; Jactel et al. 2017). Where mixedwood forests consist of mixtures of vulnerable and less vulnerable species, associational resistance reduces the severity of infestations and subsequent tree and sapling mortality, although resistance can change as stands age (Castagneyrol et al. 2020). Employing disturbance-based management to modify stand species composition and structural attributes, including simulating or delaying natural successional processes, as well as multi-aged management of mixedwood stands can create more resistant and resilient forests (e.g., Webster et al. 2018; Puettmann and Messier 2020). Such carefully designed silvicultural treatments would provide lower mortality of commercially important species in some forest types (Gottshalk 1993; Guldin 2011; Heuss et al. 2018). Following infestations, lower rates of mortality in mixedwood stands result in a more rapid recovery of leaf area and productivity, as well as reduce ecosystem respiration associated with stem decomposition, and a faster recovery to pre-infestation rates of growth and NEP.

In this paper and in most other related literature, the approach to assessing effects of species composition on insect damage has been at the stand rather than the landscape level. We have demonstrated for these three insect system examples that mixedwood stands sustain lower defoliation, beetle damage, and tree mortality than conifer- or broadleaf-dominated stands. What about a mixed landscape composed of a mixture of conifer and broadleaf stands? Theoretically, the boundaries between such stands in a mixed landscape should provide similar benefits to those from a mixedwood stand, but the interiors of pure conifer and broadleaf stands should sustain

levels of damage approaching that of pure stands. Yet there may be some landscape-level synergy that exceeds the effect of boundary conditions, depending on the mechanism of the reduced damage response, whether it involves other trophic levels (e.g. parasitoids), and if so, those species' mobility. In one of the few related studies, Campbell et al. (2008) evaluated how radial growth reductions during a spruce budworm outbreak varied in response to the broadleaf content of surrounding forest landscapes (within a 1-km radius of the balsam fir stand centroid). Broadleaf content of surrounding landscapes entered into all significant regression models, explaining 12–41% of the variance in budworm-caused growth reductions. Landscape broadleaf content had stronger relationships to growth reductions among stands than did elevation or insecticide spray history (Campbell et al. 2008). Further experimental study of insect damage as a function of insect population processes and location within mixed landscapes is worthy of future study.

A number of challenges and opportunities can be identified for managing mixedwood forests to mitigate insect damage and promote forest health in forests. Land ownership presents both challenges and opportunities because management options for privately owned forests are often driven by factors that differ from those on public lands. Approximately 85% and 77% of the forested lands in the northeastern US and mid-Atlantic regions, respectively, are in private ownership. Diverse ownership results in forests being managed (or not in some cases) for a variety of goals, and not solely the production of sawlogs and pulpwood products (Shifley et al. 2014; Butler et al. 2016). Current condition of private forests is also important for future potential management strategies. Nearly half of the forested land in northeastern US forests is considered “degraded”, based on low stocking densities and poor form of commercially desirable species, and increased basal area of American beech and red maple as a result of exploitative harvesting, insect damage and changing climate (Kosiba et al. 2018; Gunn et al. 2019).

Additional interactions between forest management, altered disturbance regimes, increasing ungulate abundance and introductions of invasive species have further contributed to forest degradation. However, adopting appropriate multi-age mixedwood silvicultural systems that promote a greater diversity of tree species could reverse the erosion of ecological “memory” (*sensu* Webster et al. 2018) associated with forest degradation. Such silvicultural systems could enhance the combined adaptations of forest species that guide recovery following insect infestations and increase resistance and resilience to future infestations (Jactel et al. 2017; Kenefic et al. 2021).

Timber products markets also present some challenges and opportunities for managing mixedwood forests. Demand for softwood sawlogs and pulpwood products has decreased in many regions relative to pre-2008, although it strongly rebounded in 2020 and 2021. Global prices for pulpwood are relatively low, with abundant supply from international sources (e.g., Australia, Brazil, Chile). This has driven a recent reevaluation of short-rotation pine production in the southeastern and mid-Atlantic US for some landowners, with longer-rotation mixedwood forest management as an option (Willis et al. 2019).

Opportunities also exist for using mixedwood systems in intensive forest management. Currently, a great majority of plantation forests consist of one or a few species. If the proportion of plantation forests expands in the future in some regions or on some ownerships, carefully designed mixedwood silvicultural systems could be integrated with strategies such as assisted migration to produce more resistant and resilient plantations. Such an approach may provide additional biodiversity and economic benefits associated with increased tree species and structural diversity (resulting from vertical stratification of hardwoods and softwoods), and improved stem quality if softwoods shade the boles of hardwoods and increase branch shedding.

521 **Conclusions**

522 Insect damage is now the dominant disturbance in forests in the northeastern and mid-
523 Atlantic regions of North America. However, standing and downed deadwood are increasingly
524 being recognized as very important ecologically, and improve habitat for numerous biotic
525 species, so there is a tradeoff. Hence, there are some positives to these mortality events and
526 values/tradeoffs involved, but a problem is the large-scale nature of mortality occurring across
527 hundreds of thousands or millions of hectares. Associational resistance of host species often
528 results in mixedwood stands being more resistant to insect infestations and incurring lower rates
529 of tree and sapling mortality. Mechanisms vary among insect species, but often involve dilution
530 of preferred host trees and reduced visual or chemical apparency, higher density or diversity of
531 natural enemies, especially parasitoids, or higher dispersal losses of insects in mixedwoods than
532 in pure hardwood or softwood stands.

533 Forest inventory data, process-based productivity models, and carbon flux measurements
534 for some forest types indicated that NPP and NEP are not significantly different among
535 mixedwood stands and hardwood or softwood-dominated stands (e.g., oak-dominated,
536 mixedwood, and pine-dominated stands in the mid-Atlantic region). Long-term inventory and
537 flux data from insect-damaged stands indicate that ecosystem functioning parallels changes in
538 structure and species composition, and that carbon and hydrologic fluxes converge on those
539 characterizing oak-pine mixedwood forests in that region. Although mixedwoods are not a
540 panacea and characteristics of tree species and desired products must be considered, we believe
541 that mixedwood management can play a substantive role in reducing insect damage of host
542 species. The impetus toward ecological forestry and uneven-aged forest management that is
543 occurring in many regions provides an opportunity for commercial mixedwood management at
544 stand and landscape scales.

Acknowledgements

Funding for this work was provided in part by the U.S. Forest Service, Northern Research Station. We thank Drs. John Kabrick, Laura Kenefic, Patricia Raymond, Lance A. Vickers, S. Ellen Macdonald, and three anonymous reviewers for providing helpful comments on earlier versions of the manuscript.

Literature Cited

- Amiro, B.D., Barr, A.G., Barr, J.G., Black, T.A., Bracho, R., Brown, M., Chen, J., Clark, K.L., Davis, K.J., Desai, A.R., Dore, S., Engel, V., Fuentes, J.D., Goldstein, A.H., Goulden, M.L., Kolb, T.E., Lavigne, M.B., Law, B.E., Margolis, H.A., Martin, T., McCaughey, J.H., Misson, L., Montes-Helu, M., Noormets, A., Randerson, J.T., Starr, G., Xiao, J. 2010. Ecosystem carbon dioxide fluxes after disturbance in forests of North America. *J. Geophys. Res.: Biogeosciences*, **115**: G00K02. doi:10.1029/2010JG001390.
- Aoki, C.F., Cook, M., Dunn, J., Finley, D., Fleming, L., Yoo, R., and Ayres, M.P. 2018. Old pests in new places: Effects of stand structure and forest type on susceptibility to a bark beetle on the edge of its native range. *For. Ecol Manage.* **419**: 206-219. doi:0.1016/j.foreco.2018.03.009.
- Barbosa, P., Hines, J., Kaplan, I., Martinson, H., Szczepaniec, A., and Szendrei, Z. 2009. Associational resistance and associational susceptibility: having right or wrong neighbors. *Ann. Rev. Ecol. Evol. Syst.* **40**: 1-20. doi: 10.1146/annurev.ecolsys.110308.120242.
- Basham, J.T., Mook, P.V., and Davidson, A.G. 1953. New information concerning balsam fir decays in eastern North America. *Can. J. Bot.* **31**: 334-360. doi:10.1139/b53-027.
- Behre, C. E. 1939. The opportunity for forestry practice in the control of gypsy moth in Massachusetts woodlands. *J. For.* **37**: 546-551. doi:10.1093/jof/37.7.546.
- Blum, B.M., and MacLean, D.A. 1984. Silviculture, forest management, and the spruce budworm. P. 83–102 In *Managing the spruce budworm in eastern North America*, Schmitt, D.M., Grimbale, D.G., and Searcy, J.L. (eds.). U.S. For. Serv., Washington, DC, Agricultural Handbook 620.
- Blum, B.M., and MacLean, D.A. 1985. Potential silviculture, harvesting, and salvage practices in eastern North America. pp. 264-280. In *Recent Advances in Spruce Budworms Research*. Proc. CANUSA Spruce Budworms Research Symp. C.J. Sanders, R.W. Stark, E.J. Mullins, and J. Murphy (Eds.). Can. For. Serv., Ottawa, Ont.

- Bognounou, F., De Grandpré, L., Pureswaran, D.S., and Kneeshaw, D. 2017. Temporal variation in plant neighborhood effects on the defoliation of primary and secondary hosts by an insect pest. *Ecosphere* **8**(3): e01759. doi:10.1002/ecs2.1759.
- Brown, J.H., Cruickshank, V.B., Gould, W.P. and Husband, T.P. 1988. Impact of gypsy moth defoliation in stands containing white pine. *North. J. Appl. For* **5**: 108-111. doi:10.1093/njaf/5.2.108.
- Butler, B.J., Hewes, J.H., Dickinson, B.J., Andrejczyk, K., Butler, S.M., and Markowski-Lindsay, M. 2016. Family forest ownerships of the United States, 2013: Findings from the USDA Forest Service's national woodland owner survey. *J. For.* **114**(6): 638-647. doi:10.5849/jof.15-099.
- Campbell, R.W. and Garlo, A.S. 1982. Gypsy moth in New Jersey pine-oak. *J. For.* **80**: 89-90. doi:10.1093/jof/80.2.89.
- Campbell, E.M., MacLean, D.A., and Bergeron, Y. 2008. The severity of budworm-caused growth reductions in balsam fir/spruce stands varies with the hardwood content of surrounding forest landscapes. *For. Sci.* **54**: 195-205. doi:10.1093/forestscience/54.2.195.
- Cappuccino, N., Lavertu, D., Bergeron, Y., and Régnière, J. 1998. Spruce budworm impact, abundance and parasitism rate in a patchy landscape. *Oecologia* **114**: 236–242. doi:10.1007/s004420050441.
- Carlson, C.E., and Wulf, N.W. 1989. Silvicultural strategies to reduce stand and forest susceptibility to the western spruce budworm. *USDA For. Serv., Coop. State Res. Serv., Agric. Handb.* 676.
- Castagneyrol, B., Kozlov, M.V., Poeydebat, C., Toïgo, M. and Jactel, H., 2020. Associational resistance to a pest insect fades with time. *J. Pest Sci.* **93**: 427-437. doi:10.1007/s10340-019-01148-y.
- Cavard, X., Macdonald, S.E., Bergeron, Y. and Chen, H.Y., 2011. Importance of mixedwoods for biodiversity conservation: Evidence for understory plants, songbirds, soil fauna, and ectomycorrhizae in northern forests. *Environ. Rev.* **19**: 142-161. doi:10.1139/a11-004.
- Chang, W-Y., Lantz, V.A., Hennigar, C.R., and MacLean, D.A. 2012. Economic impacts of forest pests: a case study of spruce budworm outbreaks and control in New Brunswick, Canada. *Can. J. For. Res.* **42**: 490-505. doi:10.1139/x11-190.
- Clark, K.L., Ayres, M., Aoki, C., Wengrowski, E., and Peterken, J. 2017. Impact of Southern pine beetle on forest structure and fuel loading in a wildfire-prone landscape. *USDA For. Serv., Gen. Tech. Rep. SRS-222*. pp. 147-154.
- Clark, K.L., Gallagher, M., Aoki, C., and Ayres, M. 2020. Southern Pine Beetle: Damage and Consequences in Forests of the Mid-Atlantic Region, USA. *Tree Planter's Notes* **62**: 91-103.

- Clark, K.L., Renninger, H., Skowronski, N., Gallagher, M., and Schäfer, K.V.R. 2018. Decadal scale reduction in forest net ecosystem production following insect defoliation contrasts with short-term impacts of prescribed fires. *Forests* **9**(3): 145. doi:10.3390/f9030145.
- Clark, K.L., Skowronski, N., Gallagher, M., Renninger, H., and Schäfer, K.V.R. 2012. Effects of invasive insects and fire on forest energy exchange and evapotranspiration in the New Jersey Pinelands. *Agric. For. Meteor.* **166-167**: 50-61. doi:10.1016/j.agrformet.2012.07.007.
- Clark, K.L., Skowronski, N., Gallagher, M., Renninger, H., and Schäfer, K.V.R. 2014. Contrasting effects of invasive insects and fire on ecosystem water use efficiency. *Biogeosciences* **11**: 6509–6523. doi:10.5194/bg-11-6509-2014.
- Clark, K.L., Skowronski, N., and Hom, J. 2010. Invasive insects impact forest carbon dynamics. *Glob. Change Biol.* **16**: 88-101. doi:10.1111/j.1365-2486.2009.01983.x.
- Clement, G. E., and Munro, W. 1917. Control of the gipsy moth by forest management. US Department of Agriculture, *Agric. Bull.* No. 484. doi:10.5962/bhl.title.108601.
- Coates, K.D., and Burton, P.J. 1997. A gap-based approach for development of silvicultural systems to address ecosystem management objectives. *For. Ecol. and Manage.* **99**(3): 337-354. doi: 0.1016/S0378-1127(97)00113-8
- Comeau, P. 1996. Why conifers? In: *Silviculture of Temperate and Boreal Broadleaf-conifer Mixtures*. P.G. Comeau and K.D. Thomas, Editors. British Columbia Ministry of Forests, Research Branch, Land Management Handbook 36. Pp. 1-7.
- Craighead, F.C. 1925. Relation between mortality of trees attacked by the spruce budworm (*Cacoecia fumiferana* Clem.) and previous growth. *J. Agric. Res.* **30**: 541-555.
- Crawford, H.S., and Jennings, D.T. 1989. Predation by birds on spruce budworm: functional, numerical, and total responses. *Ecology* **70**: 152–163. doi:10.2307/1938422.
- Croome, G.C.R. 1970. A trial of selection management in a mature fir-spruce-birch forest. *For. Chron.* **46**: 317-321. doi:10.5558/tfc46317-4.
- Curtis, P.S. and Gough, C.M. 2018. Forest aging, disturbance and the carbon cycle. *New Phytologist* **219**: 1188-1193. doi:10.1111/nph.15227.
- Davidson, C.B., Gottschalk, K.W., and Johnson, J.E. 1999. Tree mortality following defoliation by the European gypsy moth (*Lymantria dispar* L.) in the United States: a review. *For. Sci.* **45**: 74-84. doi:10.1093/forestscience/45.1.74.
- Deel, L.N., McNeil, B.E., Curtis, P.G., Serbin, S.P., Singh, A., Eshleman, K.N. and Townsend, P.A. 2012. Relationship of a Landsat cumulative disturbance index to canopy nitrogen and forest structure. *Remote Sens. Envir.* **118**: 40-49. doi:10.1016/j.rse.2011.10.026.
- Dodds, K.J., Aoki, C.F., Arango-Velez, A., Cancelliere, J., D’Amato, A.W., DiGirolomo, M.F.

- and Rabaglia, R.J. 2018. Expansion of southern pine beetle into northeastern forests: Management and impact of a primary bark beetle in a new region. *J. For.* **116**: 178-191. doi: 10.1093/jofore/fvx009.
- Drapeau, P., Leduc, A., Giroux, J.-F., Savard, J.-P. L., Bergeron, Y., and Vickery, W.L. 2000. Landscape-scale disturbances and changes in bird communities of boreal mixed-wood forests. *Ecol. Monogr.* **70**: 423-444. doi:10.1890/0012-9615(2000)070[0423:LSDACI]2.0.CO;2.
- Duveneck, M.J., Thompson, J.R., Gustafson, E.J., Liang, Y., and de Bruijn, A.M. 2017. Recovery dynamics and climate change effects to future New England forests. *Landsc. Ecol.* **32**: 1385-1397. doi:10.1007/s10980-016-0415-5.
- Dymond, C., Neilson, E., Stinson, G., Porter, K., MacLean, D.A., Gray, D., Campagna, M., and Kurz, W. 2010. Future spruce budworm outbreak may create a carbon source in eastern Canadian forests. *Ecosystems* **13**: 917–931. doi:10.1007/s10021-010-9364-z.
- Edburg, S.L., Allwine, G., Lamb, B., Stock, D., Thistle, H., Peterson, H., and Strom, B. 2010. A simple model to predict scalar dispersion within a successively thinned loblolly pine canopy. *J. Appl. Meteor. Clim.* **49**: 1913-1926. doi:10.1175/2010JAMC2339.1
- Eisenbies, M.H., Davidson, C., Johnson, J., Amateis, R. and Gottschalk, K. 2007. Tree mortality in mixed pine–hardwood stands defoliated by the European gypsy moth (*Lymantria dispar* L.). *For. Sci.* **53**: 683-691. doi:10.1093/forestscience/53.6.683.
- Etheridge, D.A., MacLean, D.A., Wagner, R.G., and Wilson, J.S. 2006. Effects of intensive forest management on stand and landscape characteristics in northern New Brunswick, Canada (1945-2027). *Landsc. Ecol.* **21**: 509-524. doi:10.1007/s10980-005-2378-9.
- Fajvan, M.A., Rentch, J. and Gottschalk, K. 2008. The effects of thinning and gypsy moth defoliation on wood volume growth in oaks. *Trees* **22**: 257-268. doi:10.1007/s00468-007-0183-6.
- Fei, S., Morin, R.S., Oswalt, C.M. and Liebhold, A.M., 2019. Biomass losses resulting from insect and disease invasions in US forests. *Proc. Nat. Acad. Sci.* **116**(35): 17371-17376. doi: 10.1073/pnas.1820601116.
- Gottschalk, K.W. 1993. Silvicultural guidelines for forest stands threatened by the gypsy moth USDA For. Serv., Northeastern Forest Experiment Station. Gen. Tech. Rep. NE-171.
- Gottschalk, K.W., and Twery, M.J. 1989. Gypsy moth impacts in pine-hardwood mixtures. In: *Proceedings of Pine-Hardwood Mixtures: A Symposium on Management and Ecology of the Type*. Waldrop, T.A., ed. USDA For. Serv., Gen. Tech. Rep. SE-58. pp. 50-58.
- Green, D.W., Winandy, J.E., and Kretschmann, D.E. 1999. Mechanical properties of wood. *Wood handbook: wood as an engineering material*. USDA Forest Service, Forest Products Laboratory, Madison, Wisc. Gen. Tech. Rep. 113. pp. 4.1–4.45.

- Guerrieri, R., Belmecheri, S., Ollinger, S.V., Asbjornsen, H., Jennings, K., Xiao, J., Stocker, B.D., Martin, M., Hollinger, D.Y., Bracho-Garrillo, R., Clark, K., Dore, S., Kolb, T., J. Munger, J.W., Novick, K., and Richardson, A.D. 2019. Disentangling the role of photosynthesis and stomatal conductance on rising forest water-use efficiency. *Proc. Nat. Acad. Sci.* **116**, 16909-16914. doi:10.1073/pnas.1905912116.
- Guldin, J.M. 2011. Silvicultural considerations in managing southern pine stands in the context of southern pine beetle. In: *Southern Pine Beetle II*. USDA For. Serv., Southern Research Station. Gen. Tech. Rep. SRS-140. pp. 317-352.
- Gunn, J.S., Ducey, M.J. and Belair, E., 2019. Evaluating degradation in a North American temperate forest. *For. Ecol. Manage.* **432**: 415-426. doi:10.1016/j.foreco.2018.09.046.
- Hennigar, C.R., and MacLean, D.A. 2010. Spruce budworm and management effects on forest and wood product carbon for an intensively managed forest. *Can. J. For. Res.* **40**: 1736-1750. doi:10.1139/X10-104.
- Hennigar, C.R., MacLean, D.A., and Norfolk, C.J. 2007. Effects of gypsy moth defoliation on softwood and hardwood growth and mortality in New Brunswick, Canada. *North. J. Appl. For.* **24**: 138-145. doi:10.1093/njaf/24.2.138.
- Hennigar, C.R., MacLean, D.A., Porter, K.B., and Quiring, D.T. 2007. Optimized harvest planning under alternative foliage-protection scenarios to reduce volume losses to spruce budworm. *Can. J. For. Res.* **37**: 1755-1769. doi:10.1139/X07-001.
- Hennigar, C.R., MacLean, D.A., Quiring, D.T., and Kershaw, J.A. Jr. 2008. Differences in spruce budworm defoliation among balsam fir and white, red, and black spruce. *For. Sci.* **54**: 158-166. doi:10.1093/forestscience/54.2.158.
- Hennigar, C.R., Erdle, T.A., Gullison, J.J., and MacLean, D.A. 2013. Reexamining wood supply in light of future spruce budworm outbreaks: a case study in New Brunswick. *For. Chron.* **89**: 42-53. doi:10.5558/tfc2013-010.
- Heuss, M., D'Amato, A.W., and Dodds, K.J. 2019. Northward expansion of southern pine beetle generates significant alterations to forest structure and composition of globally rare *Pinus rigida* forests. *For. Ecol. Manage.* **434**: 119-130. doi:10.1016/j.foreco.2018.12.015.
- Hornslein, N.J, Siegert, C., and Renninger, H.J. 2019. Physiological response of mid-canopy sweetgum trees to overstory loblolly pine mortality. *Trees* **33**: 139-151. doi:10.1007/s00468-018-1764-2.
- Hudak, J. 1991. Integrated pest management and the eastern spruce budworm. *For. Ecol. Manage.* **39**: 313-337. doi:10.1016/0378-1127(91)90188-2.
- Jactel, H., and Brockerhoff, E.G. 2007. Tree diversity reduces herbivory by forest insects. *Ecol. Lett.* **10**: 835-848. doi:10.1111/j.1461-0248.2007.01073.x.

- Jactel, H., Bauhus, J., Boberg, J., Bonal, D., Castagneyrol, B., Gardiner, B., Gonzalez-Olabarria, J.R., Koricheva, J., Meurisse, N. and Brockerhoff, E.G., 2017. Tree diversity drives forest stand resistance to natural disturbances. *Curr. For. Reports* **3**(3): 223-243. doi:10.1007/s40725-017-0064-1.
- Kabrick, J.M., Clark, K.L., D'Amato, A.W., Dey, D.C., Kenefic, L.S., Kern, C.C., Knapp, B.O., MacLean, D.A., Raymond, P., and Waskiewicz, J.D. 2017. Managing hardwood-softwood mixtures for future forests in eastern North America: assessing the suitability to projected climate change. *J. For.* **115**: 190-201. doi:10.5849/jof.2016-024.
- Katovich, S.A., and Morse, F.S. 1992. White pine weevil response to oak girdling: results from a 16-year-old study. *N. J. Appl. For.* **9**: 51-54. doi:10.1093/njaf/9.2.51.
- Kautz, M., Meddens, A.J., Hall, R.J., and Arneth, A. 2017. Biotic disturbances in Northern Hemisphere forests—a synthesis of recent data, uncertainties and implications for forest monitoring and modelling. *Global Ecol. Biogeog.* **26**: 533-552. doi:10.1111/geb.12558.
- Kemp, W.P., and Simmons, G.A. 1978. The influence of stand factors on parasitism of spruce budworm eggs by *Trichogramma minutum*. *Environ. Entomol.* **7**: 685-688. doi:10.1093/ee/7.5.685.
- Kenefic, L.S., Kabrick, J.M., Knapp, B.O., Raymond, P., Clark, K.L., D'Amato, A.W., Kern, C.C., Vickers, L.A., Dey, D.C., and Rogers, N.S. 2021. Mixedwood silviculture in North America: the science and art of managing for complex, multi-species temperate forests. *Can. J. For. Res.*, In press.
- Kern, C.C, Waskiewicz, J.D., Frelich, L., Muñoz Delgado, B.L., Kenefic, L.S. Clark, K.L, and Kabrick, J.M. 2021. Understanding compositional stability in mixedwood forests of eastern North America. *Can. J. For. Res.*, In press.
- Kneeshaw, D.D., and Prévost, M. 2007. Natural canopy gap disturbances and their role in maintaining mixed-species forests of central Quebec, Canada. *Can. J. For. Res.* **37**(9): 1534-1544. doi:10.1139/X07-112.
- Kosiba, A.M., Meigs, G.W., Duncan, J.A., Pontius, J.A., Keeton, W.S., and Tait, E.R. 2018. Spatiotemporal patterns of forest damage and disturbance in the northeastern United States: 2000–2016. *For. Ecol. Manage.* **430**: 94-104. doi:10.1016/j.foreco.2018.07.047.
- Kouki, J., McCullough, D.G., and Marshall, L.D. 1997. Effect of forest stand and edge characteristics on the vulnerability of jack pine stands to jack pine budworm (*Choristoneura pinus pinus*) damage. *Can. J. For. Res.* **27**: 1765–1772. doi:10.1139/x97-149.
- Kretchun, A.M., Scheller, R.M., Lucash, M.S., Clark, K.L., Hom, J., van Tuyl, S., and Fine, M.L. 2014. Predicted effects of gypsy moth defoliation and climate change on forest carbon dynamics in the New Jersey Pine Barrens. *PLoS One* **9**: e102531. doi:10.1371/journal.pone.0102531.

- Kulman, H.M. 1971. Effects of insect defoliation on growth and mortality of trees. *Annu. Rev. Entomol.* **16**: 289-325. doi:10.1146/annurev.en.16.010171.001445.
- Lesk, C., Coffel, E., D'Amato, A.W., Dodds, K., and Horton, R. 2017. Threats to North American forests from southern pine beetle with warming winters. *Nature Climate Change* **7**: 713-717. doi:10.1038/nclimate3375.
- Lieffers, V.J., and Beck, J.A. 1994. A semi-natural approach to mixedwood management in the prairie provinces. *For. Chron.* **70**: 260-264. doi:10.5558/tfc70260-3.
- Little, S. 1998. Fire and plant succession in the New Jersey Pine Barrens. In R. T. T. Foreman (Ed.), *Pine Barrens: Ecosystem and landscape* (pp. 297–314). New York, NY: Academic Press.
- Lovett, G.M., Weiss, M., Liebhold, A.M., Holmes, T.P., Leung, B., Lambert, K.F., Orwig, D.A., Campbell, F.T., Rosenthal, J., McCullough, D.G. et al. 2016. Nonnative forest insects and pathogens in the United States: impacts and policy options. *Ecol. Applic.* **26**: 1437-1455. doi: 10.1890/15-1176.
- MacLean, D.A. 1980. Vulnerability of fir spruce stands during uncontrolled spruce budworm outbreaks: a review and discussion. *For. Chron.* **56**: 213-221. doi:10.5558/tfc56213-5.
- MacLean, D.A. 1985. Effects of spruce budworm outbreaks on forest growth and yield. pp. 148-175. In: C.J. Sanders, R.W. Stark, E.J. Mullins, and J. Murphy (Eds.). *Recent Advances in Spruce Budworms Research. Proc. CANUSA Spruce Budworms Research Symp., Can. For. Serv., Ottawa, ON, Canada.*
- MacLean, D.A. 2016. Impacts of insect outbreaks on tree mortality, productivity, and stand development. *Can. Ent.* **148**: S138-S159. doi:10.4039/tce.2015.24.
- McManus, M., Schneeberger, N., Reardon, R., and Mason, G. 1992. Gypsy moth. Forest Insect and Disease Leaflet 162. US Department of Agriculture Forest Service. https://www.fs.usda.gov/Internet/FSE_DOCUMENTS/fsbdev2_042854.pdf (accessed 15 November 2019).
- Medvigy, D., Clark, K.L., Skowronski, N.S., and Schäfer, K.V.R. 2012. Simulated impacts of insect defoliation on forest carbon dynamics. *Envir. Res. Lett.* **7**: 045703. doi:10.1088/1748-9326/7/4/045703.
- Morin, R.S. and Liebhold, A.M. 2015. Invasive forest defoliator contributes to the impending downward trend of oak dominance in eastern North America. *Forestry* **89**: 284-289. doi: 10.1093/forestry/cpv053.
- Mosher, F.H. 1915. Food plants of the gipsy moth in America US Department of Agriculture. Bull. No. 250. doi:10.5962/bhl.title.108491.
- Muzika, R.M. 2017. Opportunities for silviculture in management and restoration of forests affected by invasive species. *Biol. Invas.* **19**: 3419-3435. doi:10.1007/s10530-017-1549-3.

- Muzika, R.M., Liebhold, A.M. and Gottschalk, K.W. 1998. Effects of silvicultural management on gypsy moth dynamics and impact: an eight year study. In: Proceedings: Population Dynamics, Impacts, and Integrated Management of Forest Defoliating Insects. USDA For. Serv., Northeastern Forest Experimental Station. Gen. Tech. Rep. NE-247. pp. 261-268.
- Needham, T., Kershaw, J., MacLean, D.A., and Su, Q. 1999. Effects of mixed stand management to reduce impacts of spruce budworm defoliation on balsam fir stand-level growth and yield. *North. J. Appl. For.* **16**: 19-24. doi:10.1093/njaf/16.1.19.
- Nie, Z., MacLean, D.A., and Taylor, A.R. 2019. Disentangling variables that influence growth response of balsam fir regeneration during a spruce budworm outbreak. *For. Ecol. Manage.* 433: 13-23. doi:10.1016/j.foreco.2018.10.050.
- New Jersey Department of Agriculture. 2017. New Jersey gypsy moth aerial defoliation survey, NJ Dept. of Agriculture, Trenton, NJ. Available online: <http://www.state.nj.us/agriculture/divisions/pi/pdf/07defoliationtable.pdf> (accessed 10 January 2019).
- North American Forest Database. 2020. Interactive map and reports on forest conditions in North America. North American Forest Commission. <https://datosforestal.nfis.org/en/> (accessed 16 January 2020).
- Nowak, J.T., Meeker, J.R., Coyle, D.R., Steiner, C.A., and Brownie, C. 2015. Southern pine beetle infestations in relation to forest stand conditions, previous thinning, and prescribed burning: evaluation of the southern pine beetle prevention program. *J. For.* **113**: 454-462. doi: 10.5849/jof.15-002.
- Nowacki, G. J., and Abrams, M. D. 2008. The demise of fire and “mesophication” of forests in the eastern United States. *BioScience*. **58**: 123-138. doi:10.1641/B580207
- Pan, Y., Chen, J.M., Birdsey, R., McCullough, K., He, L., and Deng, F. 2011. Age structure and disturbance legacy of North American forests. *Biogeosciences* **8**: 715–732. doi:10.5194/bg-8-715-2011.
- Pasquarella, V.J., Elkinton, J.S., and Bradley, B.A. 2018. Extensive gypsy moth defoliation in Southern New England characterized using Landsat satellite observations. *Biol. Invas.* **20**: 3047-3053. doi:10.1007/s10530-018-1778-0.
- Puettmann, K.J., and Messier, C. 2020. Simple guidelines to prepare forests for global change: the dog and the frisbee. *Northwest Science* **93**(3-4): 209-225. doi:10.3955/046.093.0305.
- Quayle, D., Régnière, J., Cappuccino, N., and Dupont, A. 2003. Forest composition, host-population density, and parasitism of spruce budworm *Choristoneura fumiferana* eggs by *Trichogramma minutum*. *Entomol. Exp. Appl.* **107**: 215–227. doi:10.1046/j.1570-7458.2003.00056.x.
- Renninger, H.J., Carlo, N., Clark, K.L., and Schäfer, K.V.R. 2014. Modeling respiration from

- snags and coarse woody debris before and after an invasive gypsy moth disturbance. *J. Geophys. Res.: Biogeosciences* **119**: 630-644. doi:10.1002/2013JG002542.
- Roland, J. 1993. Large-scale forest fragmentation increases the duration of tent caterpillar outbreak. *Oecologia* **93**: 25–30. doi:10.1007/BF00321186.
- Shifley, S.R., Moser, W.K., Nowak, D.J., Miles, P.D., Butler, B.J., Aguilar, F.X., DeSantis, R.D. and Greenfield, E.J. 2014. Five anthropogenic factors that will radically alter forest conditions and management needs in the northern United States. *For. Sci.* **60**: 914-925. doi:10.5849/forsci.13-153.
- Smith, D.M. 1986. *The Practice of Silviculture*. Wiley, New York.
- Spence, C.E., and MacLean, D.A. 2012. Regeneration and stand development following a spruce budworm outbreak, spruce budworm-inspired harvest, and salvage harvest. *Can. J. For. Res.* **42**: 1759-1770. doi:10.1139/x2012-121.
- Stambaugh, M.C., Marschall, J.M., Abadir, E.R., Jones, B.C., Brose, P.H., Dey, D.C., and Guyette, R.P. 2018. Wave of fire: an anthropogenic signal in historical fire regimes across central Pennsylvania, USA. *Ecosphere* **9**(5): e02222. doi:10.1002/ecs2.2222.
- Stephens, G.R., 1988. Mortality, dieback, and growth of defoliated hemlock and white pine. *N. J. Appl. For.* **5**: 93-96. doi:10.1093/njaf/5.2.93.
- Stiell, W.M., and Berry, A.B. 1985. Limiting white pine weevil attacks by side shade. *For. Chron.* **61**: 5-9. doi:10.5558/tfc61005-1.
- Stocks, B.J. 1987. Fire potential in the spruce budworm-damaged forests of Ontario. *For. Chron.* **63**: 8-14. doi:10.5558/tfc63008-1.
- Su, Q., MacLean, D.A., and Needham, T.D. 1996. The influence of hardwood content on balsam fir defoliation by spruce budworm. *Can. J. For. Res.* **26**: 1620-1628. doi:10.1139/x26-182.
- Swaine, J.M., and Craighead, F.C. 1924. Studies on the spruce budworm (*Cacoecia fumiferana* Clem.). Part I. A general account of the outbreaks, injury and associated insects. *Can. Dep. Agric., Ottawa, ON, Canada. Tech Bull.* 37 (n.s.) pp. 3-27. doi:10.5962/bhl.title.63095.
- Taylor, S.L. and MacLean, D.A. 2009. Legacy of insect defoliators: increased wind-related mortality two decades after a spruce budworm outbreak. *For. Sci.* **55**: 256-267. doi:10.1093/forestscience/55.3.256.
- Thistle, H.W., Peterson, H., Allwine, G., Lamb, B., Strand, T., Holsten, E.H., and Shea, P.J. 2004. Surrogate pheromone plumes in three forest trunk spaces: composite statistics and case studies. *For. Sci.* **50**: 610-625. doi:10.1093/forestscience/50.5.610.
- Turner, K.B. 1952. The relation of mortality of balsam fir, *Abies balsamea* (L.) Mill., caused by the spruce budworm, *Choristoneura fumiferana* (Clem.) to forest composition in the Algoma

- Forest of Ontario. Can. Dep. Agric., Ottawa. Publ. 875. 107 p.
- USDA Forest Service 2005-2016. Forest Health Highlights for New Jersey. Available online: https://www.fs.fed.us/foresthealth/docs/fhh/NJ_FHH_2016.pdf. (accessed 16 January 2020).
- USDA Forest Service. 2020. Forest Inventory and Analysis Program. Available online: <http://www.fia.fs.fed.us/>. (accessed 16 January 2020).
- Van Raalte, G.D. 1972. Do I have a budworm-susceptible forest? *For. Chron.* **48**: 190-192. doi: 10.5558/tfc48190-4
- Vickers, L.A., Knapp, B.O., Kabrick, J.M., Kenefic, L.S., D'Amato, A.W., Kern, C.C., MacLean, D.A., Raymond, P., Clark, K.L., Dey, D.C., Rogers, N.S., and Waskiewicz, J.D. 2021. Northeastern U.S. mixedwoods: contemporary status, distribution, and trends. *Can. J. For. Res.* In press.
- Webster, C.R., Dickinson, Y.L., Burton, J.I., Frelich, L.E., Jenkins, M.A., Kern, C.C., Raymond, P., Saunders, M.R., Willis, J.L. 2018. Promoting and maintaining diversity in contemporary hardwood forests: Confronting contemporary drivers of change and the loss of ecological memory. *For. Ecol. Manage.* **421**: 98-108. doi:10.1016/j.foreco.2018.01.010.
- Weed, A.S., Ayres, M.P., and Hicke, J.A. 2013. Consequences of climate change for biotic disturbances in North American forests. *Ecol. Monogr.* **83**: 441-470. doi:10.1890/13-0160.1.
- Westveld, M. 1930. Suggestions for the management of spruce stands in the northeast. USDA For. Serv., Circular No. 134. 23p.
- Westveld, M. 1946. Forest management as a means of controlling spruce budworms. *J. For.* **44**: 949-953.
- Willis, J.L., Gordon, J.S., Tanger, S., Blazier, M.A., Self, A.B. and Brodbeck, A. 2019. Managing mixed stands: reassessing a forgotten stand type in the southeastern United States. *Forests* **10**: 751. doi:10.3390/f10090751.
- Xu, B., Pan, Y., Plante, A.F., McCullough, K., and Birdsey, R. 2017. Modeling forest carbon cycle using long-term carbon stock field measurement in the Delaware River Basin. *Ecosphere* **8**: e01802. doi:10.1002/ecs2.1802.
- Zhang, B., MacLean, D.A., Johns, R.C., and Eveleigh, E.S. 2018. Effects of hardwood content on balsam fir defoliation during the building phase of a spruce budworm outbreak. *Forests* **9**: 530. doi:10.3390/f9090530.
- Zhang, B., MacLean, D.A., Johns, R.C., Eveleigh, E.S., and Edwards, S. 2020. Hardwood-softwood composition influences early-instar larval dispersal mortality during a spruce budworm outbreak. *For. Ecol. Manage.* **463**: 118035. doi:10.1016/j.foreco.2020.118035.

1005 Table 1. Gradient of susceptibility of dominant tree species in the northeast to spruce budworm,
 1006 gypsy moth, and southern pine beetle.

Susceptibility	Spruce budworm	Gypsy moth	Southern pine beetle
Highest	Balsam fir White spruce Red spruce Black spruce Eastern larch Eastern hemlock	Red, white, black, and scarlet oaks Chestnut oak Sweetgum Cottonwood and Aspen Birch and Alder Sugar maple White pine Eastern hemlock Red maple Ash and Tulip poplar	Loblolly pine Pitch pine Shortleaf pine Virginia pine White pine (Jack pine)*
 Lowest Nonsusceptible	Hardwoods	Most softwoods	Hardwoods

1007 *Susceptibility to southern pine beetle not well known

Table 2. Net ecosystem production (NEP), gross ecosystem production (GEP) and evapotranspiration (Et) before and during severe defoliation by gypsy moth in the New Jersey Pinelands National Reserve. The oak stand was dominated by chestnut (*Quercus prinus*), white (*Q. alba*), black (*Q. velutina*) and scarlet oaks (*Q. coccinea*), and shortleaf (*Pinus echinata*) and pitch (*P. rigida*) pines, the mixedwood stand was co-dominated by chestnut and white oaks and pitch pine, and the pine stand was dominated by pitch pine and white and blackjack (*Quercus stellata*) oaks. Adapted from Clark et al. (2010, 2014, 2018).

Forest type	NEP (g C m ⁻² yr ⁻¹)		GEP (g C m ⁻² yr ⁻¹)		Et (mm yr ⁻¹)	
	before	during	before	During	before	during
Oak	183	-246	1503	726	637	442
Mixedwood	137	-20	1205	992	607	419
Pine	173	40	1633	1402	757	593





Fig. 1. Composition and attributes of conifer-dominated, mixedwood, and broadleaf-dominated stands. Mixedwood stands have greater diversity at higher taxonomic levels (e.g. family and above) that is more evenly distributed across co-dominant species, although not necessarily a greater number of species. Mixedwood stands typically have greater structural diversity than conifer- or broadleaf-dominated stands. Greater diversity at higher taxonomic levels confers reduced susceptibility to insect infestations, primarily through greater associational resistance. Depending on the insect, damage and tree mortality are reduced compared to conifer- or broadleaf-dominated stands. Successional changes in species composition favors either conifer or broadleaf dominance in the absence of major disturbances, depending upon forest type (indicated by the light arrows). In contrast, insect infestations and differential tree mortality can result in the formation of mixed composition stands (indicated by the darker arrows).

1030



1031

1032 **Fig. 2.** Merchantable volume of balsam fir and hardwood trees pre-infestation ('pre') and a
1033 decade following infestations by spruce budworm ('post') in balsam fir-dominated (BF), balsam
1034 fir-hardwood mixedwood (MW), and hardwood-dominated (HW) stands. Data were modified
1035 from Needham et al. (1999), where BF included 0-20% hardwood classes, MW included 30-70%
1036 hardwoods, and HW included 80-100% hardwoods.





Fig. 3. a) Basal area of oak and pine trees and saplings pre-infestation (2004; 'pre') and a decade following infestations by gypsy moth (2018; 'post') in oak-dominated, oak-pine mixedwood, and pine-dominated stands. Dominant canopy trees were similar in age. Pre-infestation census data are from Clark et al. 2010, 2018, post-infestation from same sets of plots, unpublished data. **b)** Basal area of pine, oak and other hardwood trees and saplings pre- and post-infestation of southern pine beetle in pine-dominated stands (From Clark et al. 2017, 2020).

1061



1062

1063

1064

1065

1066

1067

1068

Fig. 4. Relative basal area of an oak-dominated stand before (pre-) and following (post) severe gypsy moth defoliation, an oak-pine mixedwood stand, and pitch pine stands before (pre-) and following (post) southern pine beetle infestation in the Pinelands National Reserve of New Jersey. Arrows indicate the directional changes in relative basal area caused by insect damage; the arrow on the left indicates gypsy moth damage and the arrow on the right indicates southern pine beetle damage. Adapted from census data in Clark et al. 2017, 2018, and 2020.