

Scientific Data

<https://doi.org/10.1038/s41597-026-07080-y>

Article in Press

Chromosome-level genome assemblies of *Nicotiana attenuata* (coyote tobacco) and *Nicotiana obtusifolia* (desert tobacco)

Received: 10 November 2025

Accepted: 13 March 2026

Cite this article as: Chakraborty, A., Xu, S. Chromosome-level genome assemblies of *Nicotiana attenuata* (coyote tobacco) and *Nicotiana obtusifolia* (desert tobacco). *Sci Data* (2026). <https://doi.org/10.1038/s41597-026-07080-y>

Abhisek Chakraborty & Shuqing Xu

We are providing an unedited version of this manuscript to give early access to its findings. Before final publication, the manuscript will undergo further editing. Please note there may be errors present which affect the content, and all legal disclaimers apply.

If this paper is publishing under a Transparent Peer Review model then Peer Review reports will publish with the final article.

Chromosome-level genome assemblies of *Nicotiana attenuata* (coyote tobacco) and *Nicotiana obtusifolia* (desert tobacco)

Abhisek Chakraborty¹, Shuqing Xu^{1*}

¹ Institute of Organismic and Molecular Evolution (IomE), Johannes Gutenberg University, Mainz
Biozentrum I, Hanns-Dieter-Hüsch-Weg 15, 55128 Mainz, Germany

* correspondence: shuqing.xu@uni-mainz.de

ARTICLE IN PRESS

ABSTRACT

Nicotiana attenuata and *Nicotiana obtusifolia* are two wild tobacco species from the Solanaceae plant family, which are rich in producing diverse specialized metabolites and have been established as ecological model systems for studying plant-insect interactions. The previously published *N. attenuata* and *N. obtusifolia* genomes were draft assemblies with low assembly contiguity and a high proportion of gaps, limiting comparative genomic analysis. Here, we performed chromosome-level genome assemblies and annotations of these two species using long-read sequencing and Hi-C data. Both *N. attenuata* and *N. obtusifolia* genomes were anchored to 12 chromosomes, with assembled genome sizes of 2.2 Gb and 1.3 Gb, respectively. We achieved high BUSCO completeness at the protein level, with 99.3% for *N. attenuata* and 97.9% for *N. obtusifolia*. The reference genomes of *N. attenuata* and *N. obtusifolia* presented in this study will advance the understanding of metabolic innovations in Solanaceae plants.

BACKGROUND & SUMMARY

Nicotiana attenuata and *Nicotiana obtusifolia* are diploid ($2n = 2x = 24$) Solanaceae species, which have been developed as ecological model plants to study plant-environment interactions in nature¹⁻³. In particular, *N. attenuata* has been extensively used to understand the function, evolution and diversification of plant specialized metabolites (PSM), such as nicotine, as a defensive response to herbivory and other ecological factors, using genomics, transcriptomics, and metabolomics³⁻⁵. Further, *N. attenuata* has been used to understand the balance between plant defence and protection of autotoxicity conferred by the toxic PSMs⁶. Similarly, *N. obtusifolia* is also used to study ecological interactions of plants with pathogens, herbivores, and pollinators in natural habitats, along with *N. attenuata* and others³. Previous omics studies have used *N. obtusifolia* to understand the genetic basis of metabolic innovation and diversification of plant specialized metabolites in the *Nicotiana* genus, which are produced as a defence response to environmental stress⁷⁻⁹.

With the significance of these species as ecological models, and the necessity of future genomics-guided multi-omics strategies to further understand the metabolic innovations at the level of both plant organ and developmental stage, high-quality genomes are essential for comparative analysis¹⁰. A large number of genomic, transcriptomic, and metabolomic datasets already exist for *N. attenuata*¹¹. However, high-quality genome assemblies were still not available for both of these *Nicotiana* species. The first published draft genomes of *N. attenuata* and *N. obtusifolia* had low assembly contiguities (N50 values of 524.5 Kb and 134.1 Kb, respectively)⁹. Currently, two genome assemblies of *N. attenuata* are available, superscaffolded into 12 linkage groups using optical mapping and genetic mapping data (NCBI GenBank accessions: GCF_001879085.1 and GCA_030864195.1); however, several issues remain, including fragmented chromosomes and a high percentage of gaps (**Supplementary Table 1**).

Here, we used the existing *N. attenuata* genome assembly (GCA_030864195.1, assembled from PacBio reads)¹² to reconstruct the chromosomes based on chromatin interaction data obtained from Hi-C sequencing, and report the improved chromosome-level genome assembly of *N. attenuata* (12 chromosomes, 2.07 Gb) (**Figure 1a, Table 1**)¹³. Compared to the previously available assemblies, our Hi-C data-based assembly showed an increased genomic length covered by the chromosomes, a reduction in the number of ambiguous bases (N), a reduction in the total number and length of gaps, clean Hi-C contact maps with no evidence of mis-joins, and a better accuracy of the genome assembly (**Supplementary Table 1**). Further, we performed a *de novo* genome assembly of *N. obtusifolia* using PacBio sequencing data, and anchored it to chromosomes using Hi-C data, reporting the first chromosome-scale *N. obtusifolia* genome (12 chromosomes, 1.29 Gb) (**Figure 1b, Table 1**)¹³. The improvements in the genome assembly statistics compared to the previous contig-level draft assembly of *N. obtusifolia* are mentioned in **Supplementary Table 2**.

Repetitive regions contributed to 83.29% and 79.21% of the *N. attenuata* and *N. obtusifolia* genome assemblies, respectively. It is noteworthy to mention that the usage of long PacBio reads and the reads obtained from high-throughput chromatin conformation capture (Hi-C) technology enabled the improvement of the assembly quality of such repetitive genomes, which was a limitation in the previous studies^{9,12}. After soft-masking the genome assemblies, we identified 35,166 and 27,352 protein-coding genes in *N. attenuata* and *N. obtusifolia*, respectively, using the BRAKER3 pipeline¹⁴ (**Table 1**).

Parameters	<i>N. attenuata</i>	<i>N. obtusifolia</i>
Assembled genome size (bp)	2,185,486,753	1,304,817,465
Number of assembled sequences	1,498	71
GC (%)	40.42	39.06
Chromosome-level genome size (bp)	2,068,473,773	1,291,577,541
Total number of chromosomes	12	12
Longest chromosome length (bp)	216,162,101	118,269,720
Number of protein-coding genes	35,166	27,352
Average CDS length (bp)	1077.16	1180.68

Table 1. Genome assembly and annotation statistics of the *Nicotiana* genomes

The chromosome-level genome assemblies of *N. attenuata* and *N. obtusifolia* presented in this study will serve as a valuable dataset, facilitating a better understanding of the genomic basis of metabolic innovations not only in *Nicotiana* but also in the Solanaceae family.

METHODS

Sample preparation and sequencing

N. attenuata (NCBI TaxID: 49451) plants were grown from Max Planck Institute for Chemical Ecology (MPI-CE) seed stock accession UT31, an inbred ‘UT’ line (31st inbred generation) derived from seeds

originally collected at DI Ranch (Santa Clara, Utah, USA) in 1988. UT31 was the same *N. attenuata* accession that was used in the previous study¹². *N. obtusifolia* (NCBI TaxID: 200316) plants were grown from MPI-CE seed stock Nob02, an inbred (2 generations of inbreeding) derived from seeds originally collected at Lytle ranch preserve (Santa Clara, Utah, USA) in 2004. The *N. attenuata* and *N. obtusifolia* plants used for sequencing were made available as herbarium vouchers at the Johannes Gutenberg University Botanical Garden Herbarium (Herbarium MJG) with accessions MJG 048398 and MJG 048399, respectively (**Supplementary Figures 1-2**).

For sequencing, high-quality genomic DNA was extracted from frozen leaf samples using a modified CTAB method¹⁵. RNase A was used to remove RNA contaminants. For *N. attenuata*, a Hi-C fragment library (insert size 300-700 bp) was constructed using the Mate-pair kit after chromatin cross-linking with formaldehyde and enzymatic digestion with DpnII. The library was then sequenced using the Illumina NovaSeq platform. For *N. obtusifolia*, whole-genome sequencing was performed using the PacBio Revio platform, and for Hi-C data-based contig anchoring, a Hi-C fragment library with 300-700 bp insert size was constructed using the same methods as *N. attenuata*, and sequenced using the Illumina NovaSeq platform. PacBio Iso-seq RNA-seq library was also constructed using total RNA from the leaf sample, which was sequenced on a PacBio Sequel II platform. The sequencing was carried out at Biomarker Technologies (Beijing, China).

Genome assembly

For *N. attenuata*, 301.92 Gb clean data were generated from the Hi-C library and were mapped to the previously assembled *N. attenuata* genome (GCA_030864195.1, assembled from PacBio reads)¹² using BWA v0.7.17 with the default parameters¹⁶. Invalid read pairs, including Dangling-End and Self-cycle, Religation and Dumped products, were filtered out. LACHESIS was used to cluster, order, and orient the contigs utilising the Hi-C interaction signals¹⁷. After Hi-C data-based anchoring, 2.07 Gb (94.6%) of the total assembly (2.19 Gb) was in confirmed order and orientation, constituting the 12 chromosomes. The parameters used in LACHESIS were: CLUSTER_MIN_RE_SITES = 161; CLUSTER_MAX_LINK_DENSITY = 2; ORDER_MIN_N_RES_IN_TRUNK = 199; ORDER_MIN_N_RES_IN_SHREDS = 239.

Prior to *N. obtusifolia* *de novo* genome assembly, Illumina paired-end reads from the previous study were quality-filtered using Trimmomatic v0.39 with the parameters “SLIDINGWINDOW:4:15 MINLEN:36”, which were then used to construct the k-mer frequency-based histogram (k-mer = 31) using Jellyfish v2.2.10, and to estimate the genomic characteristics using GenomeScope v2 (**Supplementary Figure 3**)^{18–20}. For the genome assembly, high-accuracy CCS data (96.91 Gb, N50 = 20.67 Kb) were

assembled using Hifiasm v0.24.0 with the parameters: $l=0$ and $n=4$, resulting in 329 contigs (**Supplementary Table 3**)²¹. For anchoring of the contigs, 197.74 Gb clean Hi-C data was mapped to the assembly using BWA v0.7.17 with default parameters¹⁶. Similar to the *N. attenuata* genome, the invalid read pairs were filtered out, and the valid interaction read pairs were used for the clustering, ordering, and orienting of scaffolds onto chromosomes with LACHESIS¹⁷. After Hi-C data-based anchoring, 1.29 Gb (98.99%) of the total assembly (1.3 Gb) was in confirmed order and orientation, constituting the 12 chromosomes. The parameters used in LACHESIS were: CLUSTER_MIN_RE_SITES = 723; CLUSTER_MAX_LINK_DENSITY = 2; ORDER_MIN_N_RES_IN_TRUNK = 15; ORDER_MIN_N_RES_IN_SHREDS = 15.

The unplaced contigs from both *N. attenuata* and *N. obtusifolia* genomes were further analysed using BlobTools to check for contamination²². For this analysis, first, the Illumina paired-end data from the previous study⁹ were quality-filtered using Trimmomatic v0.39 with the parameters “SLIDINGWINDOW:4:15 MINLEN:36”¹⁸. The filtered paired-end reads were mapped onto the respective *Nicotiana* contigs using BWA-MEM¹⁶, and the contigs were mapped against the NCBI-nt database using BLASTN, which were then used to construct the blobplots. Only the contigs that were assigned as “Streptophyta”, “undef”, and “no-hit” were retained (**Supplementary Figures 4-5**).

Genome annotation

Both *N. attenuata* and *N. obtusifolia* genome assemblies were annotated using a similar method. Prior to coding gene prediction, the genome assemblies were used to construct the corresponding *de novo* repeat libraries using RepeatModeler v2.0.2, with “-LTRStruct” and other default parameters²³. The resultant repeat libraries were used to soft-mask the respective genome assemblies using RepeatMasker v4.1.2 (<https://www.repeatmasker.org>).

The soft-masked *Nicotiana* genome assemblies were used for the prediction of high-confidence protein-coding genes using BRAKER3¹⁴, which implements an integrated transcriptome and proteome evidence-based gene prediction in GeneMark-ETP pipeline²⁴. For transcriptome-based evidence of *N. attenuata*, the RNA-Seq data (Illumina short reads) from the previous study⁹ were quality-filtered using Trimmomatic v0.39 with the parameters “SLIDINGWINDOW:4:15 MINLEN:36”, and the filtered data were used in BRAKER3^{14,18}. Inside the BRAKER3 pipeline, AUGUSTUS²⁵ was also used for the prediction of coding genes, and TSEBRA²⁶ was used for obtaining a combined output of GeneMark-ETP and AUGUSTUS-based gene predictions.

For transcriptome-based evidence of *N. obtusifolia*, both Illumina short read RNA-Seq data from a previous study³ and the newly generated PacBio Iso-seq ccs data (292,933 reads) were used in two separate BRAKER3 analyses. First, the Illumina short reads were quality-filtered using Trimmomatic v0.39 with the parameters “SLIDINGWINDOW:4:15 MINLEN:36”, and the filtered data were used in BRAKER3^{14,18}. Next, the Iso-seq data were mapped onto the genome using Minimap2²⁷, which was then used in a separate BRAKER3 analysis. The results from the two BRAKER3 analyses were merged using TSEBRA²⁶.

In all the BRAKER3 analyses for both the *Nicotiana* species, the protein sequences from the previous draft assembly of *N. attenuata*⁹, along with species belonging to *Nicotiana* and other Solanaceae genus (only chromosome-level assemblies were considered) were used as a set of extrinsic proteome evidence in GeneMark-ETP. These Solanaceae species were - *N. sylvestris*²⁸, *N. tomentosiformis*²⁸, *N. tabacum*²⁸, *N. benthamiana*²⁹, *Capsicum annuum*³⁰, *Solanum lycopersicum*³¹, *Physalis pruinosa*³², *Datura innoxia*³³, *Lycianthes biflora*³³, *Lycium chinense*³⁴, and *Lochroma cyaneum*³⁵.

The *Nicotiana* coding gene sets were filtered to extract the longest isoforms for each gene model and to remove the coding genes with a length of <100 bp using AGAT (<https://github.com/NBISweden/AGAT>). Further, the coding genes with a repeat content of >50% were filtered out, using a method similar to a previous study³⁶. The protein-coding genes were annotated using the eggNOG-mapper genome annotation server, which uses orthology relationships to assign the KO (KEGG Orthology) terms, GO (Gene Ontology) terms, COG (Cluster of Orthologous Gene) categories, CAZy families, and Pfam domains³⁷. Additionally, the coding genes were also mapped against the UniRef90 database using Diamond v2.0.13 with the parameters: “-k 1 -e 0.00001 -f 6 --sensitive”^{38,39}.

Evaluation of genome assembly and annotation

Mercury v1.3 was used to assess assembly completeness and accuracy by comparing k-mers from the *Nicotiana* genome assemblies and the quality-filtered Illumina paired-end reads, with a k-mer size of 21, as predicted by the “best_k.sh” script^{9,40}. Further, the LTR Assembly Index (LAI) score⁴¹ was estimated for the chromosome-level genomes of *N. attenuata* and *N. obtusifolia* using GenomeTools⁴² v1.6.1 and LTR_retriever⁴³ v3.0.1.

To evaluate the completeness of the whole-genome assembly and the predicted coding gene sets, BUSCO v5.4.3 was used in “genome” and “proteins” modes, respectively, with the solanales_odb10 database⁴⁴. The quality of *Nicotiana* gene model annotations was assessed using GAQET2 to provide metrics produced by AGAT, OMark, and PSAURON^{45–47}.

DATA RECORDS

The newly generated sequencing data for *N. attenuata* have been deposited in the NCBI database with BioProject accession number PRJNA1245670. The Hi-C genomic reads of *N. attenuata* are available under SRA accession SRR32964043⁴⁸. Genome and transcriptome sequencing raw data of *N. obtusifolia* generated in this study have been deposited in the NCBI database with BioProject accession number PRJNA1332718. The SRA accessions for the *N. obtusifolia* sequencing reads are - SRR35556615 (PacBio Revio genomic reads)⁴⁹, SRR35556614 (Hi-C genomic reads)⁵⁰, and SRR35556613 (PacBio Iso-seq transcriptome reads)⁵¹. The Whole Genome Shotgun projects have been deposited at DDBJ/ENA/GenBank under the accession JBUBWS000000000⁵² and JBUCPO000000000⁵³ for *N. attenuata* and *N. obtusifolia*, respectively. The versions described in this paper are versions JBUBWS010000000 and JBUCPO010000000 for *N. attenuata* and *N. obtusifolia*, respectively. The genome assembly and annotation files are also available at Figshare^{54,55}. The mapping results of the *Nicotiana* gene sets constructed in this study against those of the previous study⁹ are also available at Figshare^{54,55}.

TECHNICAL VALIDATION

To evaluate the genome assembly quality and completeness, we implemented multiple strategies. First, the Hi-C contact map showed strong intra-chromosomal interaction signals along the diagonal, which confirms the genome structure integrity (**Figure 2**). Second, Merqury analysis results showed the k-mer completeness and QV (consensus quality) values of 96.53% and 42.13, respectively, for *N. attenuata*, and 98.35% and 48.22, respectively, for *N. obtusifolia* (**Supplementary Figures 6-7**). Third, the chromosome-level genome assemblies of *N. attenuata* and *N. obtusifolia* had LAI scores of 14.6 and 15.6, respectively, which refer to “Reference”-standard genome assemblies, according to Ou et al.⁴¹ (**Figure 3**). Fourth, BUSCO analysis showed the presence of 98.5% complete BUSCO genes in both *N. attenuata* and *N. obtusifolia* genome assemblies (**Table 2**). At the proteome level, the BUSCO completeness was 99.3% and 97.9% for *N. attenuata* and *N. obtusifolia*, respectively (**Table 2**). Fifth, the coding gene sets showed PSAURON quality scores of 96.1 and 96.7 for *N. attenuata* and *N. obtusifolia*, respectively. Detailed quality metrics for the genome annotation are provided in **Supplementary Table 4**.

Parameters	Genome	Proteins
<i>N. attenuata</i>		
Total BUSCO groups searched	5,950	5,950
Complete BUSCOs (C)	5,861 (98.5%)	5,909 (99.3%)
Complete and single-copy BUSCOs (S)	5,648 (94.9%)	5,725 (96.2%)
Complete and duplicated BUSCOs (D)	213 (3.6%)	184 (3.1%)
Fragmented BUSCOs (F)	9 (0.2%)	6 (0.1%)
Missing BUSCOs (M)	80 (1.3%)	35 (0.6%)
<i>N. obtusifolia</i>		
Total BUSCO groups searched	5,950	5,950
Complete BUSCOs (C)	5,861 (98.5%)	5,829 (97.9%)

Complete and single-copy BUSCOs (S)	5,646 (94.9%)	5,630 (94.6%)
Complete and duplicated BUSCOs (D)	215 (3.6%)	199 (3.3%)
Fragmented BUSCOs (F)	7 (0.1%)	11 (0.2%)
Missing BUSCOs (M)	82 (1.4%)	110 (1.9%)

Table 2. BUSCO completeness of the *Nicotiana* genomes

CODE AVAILABILITY

All bioinformatic analyses were performed according to the manuals provided by the software developers. The software versions and the parameters used for the analyses are mentioned in the “Methods” section. The codes used in this study are deposited in GitHub with the link - https://github.com/Xu-lab-Evolution/Nicotiana_genome_project.

ACKNOWLEDGEMENTS

We thank Dr. Danny Kessler from the Max Planck Institute for Chemical Ecology for providing the plant materials. We thank Dr. Patrycja Baraniecka and Thoomke Roth for preparing the plant materials for sequencing. We are grateful to the help from Dr. Christian Siadjeu, Dr. Ralf Omlor, and Markus Hageneuer in preparing the voucher specimens of *N. attenuata* and *N. obtusifolia*. Parts of this research were conducted using the supercomputer Mogon 2 and/or advisory services offered by Johannes Gutenberg University Mainz (hpc.uni-mainz.de), which is a member of the AHRP (Alliance for High-Performance Computing in Rhineland Palatinate, www.ahrp.info) and the Gauss Alliance eV. The authors gratefully acknowledge the computing time granted on the supercomputer MOGON 2 at Johannes Gutenberg University Mainz (hpc.uni-mainz.de).

AUTHORS CONTRIBUTIONS

A.C. conducted the research and performed the data analysis. S.X. conceived and supervised the study. Both authors have read and approved the final manuscript.

COMPETING INTERESTS

The authors declare no competing interests.

FUNDING

This work was supported by the Agence Nationale de la Recherche (ANR)–Deutsche Forschungsgemeinschaft (DFG) joint research funding for French–German Collaboration (ANR-23-CE20-0037 to E.G. and DFG project number 529944545 to S.X.).

FIGURE LEGENDS

Figure 1. Circos plots showing the genomic features (window size 100 Kb) - **a.** *N. attenuata*, **b.** *N. obtusifolia*. From the outer to inner circle – I. Chromosome length, II. Percentage of repeat regions, III. Percentage of gene regions, IV. GC content.

Figure 2. Hi-C contact maps of the genome assemblies - **a.** *N. attenuata*, **b.** *N. obtusifolia*.

Figure 3. Distribution of the LAI scores in the 12 chromosomes of - **a.** *N. attenuata*, **b.** *N. obtusifolia*.

SUPPLEMENTARY INFORMATION

Supplementary Tables 1-4, Supplementary Figures 1-7.

REFERENCES



- Schuman, M. C., Barthel, K. & Baldwin, I. T. Herbivory-induced volatiles function as defenses increasing fitness of the native plant *Nicotiana attenuata* in nature. *Elife* **2012**, (2012).
- Choung, S. *et al.* MYC2 and MYC3 orchestrate pith lignification to defend *Nicotiana attenuata* stems against a stem-boring weevil. *New Phytol.* (2025) doi:10.1111/NPH.70325.
- Zhou, W. *et al.* Evolution of herbivore-induced early defense signaling was shaped by genome-wide duplications in *Nicotiana*. *Elife* **5**, (2016).
- Xu, S. *et al.* Allelic differences of clustered terpene synthases contribute to correlated intraspecific variation of floral and herbivory-induced volatiles in a wild tobacco. *New Phytol.* **228**, 1083–1096 (2020).
- Li, D., Heiling, S., Baldwin, I. T. & Gaquerel, E. Illuminating a plant's tissue-specific metabolic diversity using computational metabolomics and information theory. *Proc. Natl. Acad. Sci. U. S. A.* **113**, E7610–E7618 (2016).
- Li, J. *et al.* Controlled hydroxylations of diterpenoids allow for plant chemical defense without autotoxicity. *Science* (80-.). **371**, 255–260 (2021).
- Elser, D. *et al.* Evolutionary metabolomics of specialized metabolism diversification in the genus *Nicotiana* highlights Nacylnornicotine innovations. *Sci. Adv.* **9**, (2023).
- Kaminski, K. P. *et al.* Alkaloid chemophenetics and transcriptomics of the *Nicotiana* genus. *Phytochemistry* **177**, 112424 (2020).
- Xu, S. *et al.* Wild tobacco genomes reveal the evolution of nicotine biosynthesis. *Proc. Natl. Acad. Sci. U. S. A.* **114**, 6133–6138 (2017).
- Xu, S. & Gaquerel, E. Evolution of plant specialized metabolites: beyond ecological drivers. *Trends Plant Sci.* **0**, (2025).
- Brockmüller, T. *et al.* *Nicotiana attenuata* Data Hub (NaDH): An integrative platform for exploring genomic, transcriptomic and metabolomic data in wild tobacco. *BMC Genomics* **18**, 1–11 (2017).
- Ray, R. *et al.* A persistent major mutation in canonical jasmonate signaling is embedded in an herbivory-elicited gene network. *Proc. Natl. Acad. Sci. U. S. A.* **120**, e2308500120 (2023).
- Krzywinski, M. *et al.* Circos: An information aesthetic for comparative genomics. *Genome Res.* **19**, 1639–1645 (2009).



14. Gabriel, L. *et al.* BRAKER3: Fully automated genome annotation using RNA-seq and protein evidence with GeneMark-ETP, AUGUSTUS, and TSEBRA. *Genome Res.* **34**, 769–777 (2024).
15. Abu Almakarem, A. S., Heilman, K. L., Conger, H. L., Shtarkman, Y. M. & Rogers, S. O. Extraction of DNA from plant and fungus tissues in situ. *BMC Res. Notes* **5**, 1–11 (2012).
16. Li, H. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM. **00**, 1–3 (2013).
17. Burton, J. N. *et al.* Chromosome-scale scaffolding of de novo genome assemblies based on chromatin interactions. *Nat. Biotechnol.* **31**, 1119–1125 (2013).
18. Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: A flexible trimmer for Illumina sequence data. *Bioinformatics* (2014) doi:10.1093/bioinformatics/btu170.
19. Marçais, G. & Kingsford, C. A fast, lock-free approach for efficient parallel counting of occurrences of k-mers. *Bioinformatics* (2011) doi:10.1093/bioinformatics/btr011.
20. Ranallo-Benavidez, T. R., Jaron, K. S. & Schatz, M. C. GenomeScope 2.0 and Smudgeplot for reference-free profiling of polyploid genomes. *Nat. Commun.* (2020) doi:10.1038/s41467-020-14998-3.
21. Cheng, H., Concepcion, G. T., Feng, X., Zhang, H. & Li, H. Haplotype-resolved de novo assembly using phased assembly graphs with hifiasm. *Nat. Methods* **18**, 170–175 (2021).
22. Challis, R., Richards, E., Rajan, J., Cochrane, G. & Blaxter, M. BlobToolKit – Interactive Quality Assessment of Genome Assemblies. *G3 Genes|Genomes|Genetics* **10**, 1361–1374 (2020).
23. Flynn, J. M. *et al.* RepeatModeler2 for automated genomic discovery of transposable element families. *Proc. Natl. Acad. Sci. U. S. A.* (2020) doi:10.1073/pnas.1921046117.
24. Brûna, T., Lomsadze, A. & Borodovsky, M. GeneMark-ETP significantly improves the accuracy of automatic annotation of large eukaryotic genomes. *Genome Res.* **34**, 757–768 (2024).
25. Stanke, M. *et al.* AUGUSTUS: ab initio prediction of alternative transcripts. *Nucleic Acids Res.* **34**, W435–W439 (2006).
26. Gabriel, L., Hoff, K. J., Brûna, T., Borodovsky, M. & Stanke, M. TSEBRA: transcript selector for BRAKER. *BMC Bioinformatics* **22**, 1–12 (2021).
27. Li, H. Minimap2: Pairwise alignment for nucleotide sequences. *Bioinformatics* (2018) doi:10.1093/bioinformatics/bty191.
28. Sierro, N., Auberson, M., Dulize, R. & Ivanov, N. V. Chromosome-level genome assemblies of *Nicotiana tabacum*, *Nicotiana glauca*, and *Nicotiana glauca*. *Sci. Data* **11**, 1–10 (2024).
29. Ko, S. R. *et al.* High-quality chromosome-level genome assembly of *Nicotiana benthamiana*. *Sci. Data* **11**, 1–8 (2024).
30. Liu, F. *et al.* Genomes of cultivated and wild *Capsicum* species provide insights into pepper domestication and population differentiation. *Nat. Commun.* **14**, 1–14 (2023).
31. Su, X. *et al.* A high-continuity and annotated tomato reference genome. *BMC Genomics* **22**, 1–12 (2021).
32. He, J. *et al.* Establishing *Physalis* as a Solanaceae model system enables genetic reevaluation of the inflated calyx syndrome. *Plant Cell* **35**, 351–368 (2023).


33. Wu, Y. *et al.* Phylogenomic discovery of deleterious mutations facilitates hybrid potato breeding. *Cell* **186**, 2313–2328.e15 (2023).
34. Yang, J. *et al.* Multiple independent losses of the biosynthetic pathway for two tropane alkaloids in the Solanaceae family. *Nat. Commun.* **14**, 1–18 (2023).
35. Powell, A. F. *et al.* Genome sequence for the blue-flowered Andean shrub *Iochroma cyaneum* reveals extensive discordance across the berry clade of Solanaceae. *Plant Genome* **15**, e20223 (2022).
36. Wang, Y. & Xu, S. A high-quality genome assembly of the waterlily aphid *Rhopalosiphum nymphaeae*. *Sci. Data* **11**, 1–6 (2024).
37. Huerta-Cepas, J. *et al.* Fast genome-wide functional annotation through orthology assignment by eggNOG-mapper. *Mol. Biol. Evol.* (2017) doi:10.1093/molbev/msx148.
38. Suzek, B. E., Wang, Y., Huang, H., McGarvey, P. B. & Wu, C. H. UniRef clusters: A comprehensive and scalable alternative for improving sequence similarity searches. *Bioinformatics* (2015) doi:10.1093/bioinformatics/btu739.
39. Buchfink, B., Xie, C. & Huson, D. H. Fast and sensitive protein alignment using DIAMOND. *Nature Methods* at <https://doi.org/10.1038/nmeth.3176> (2014).
40. Rhie, A., Walenz, B. P., Koren, S. & Phillippy, A. M. Merqury: reference-free quality, completeness, and phasing assessment for genome assemblies. *Genome Biol.* **2020 211** **21**, 245– (2020).
41. Ou, S., Chen, J. & Jiang, N. Assessing genome assembly quality using the LTR Assembly Index (LAI). *Nucleic Acids Res.* (2018) doi:10.1093/nar/gky730.
42. Gremme, G., Steinbiss, S. & Kurtz, S. Genome tools: A comprehensive software library for efficient processing of structured genome annotations. *IEEE/ACM Trans. Comput. Biol. Bioinforma.* (2013) doi:10.1109/TCBB.2013.68.
43. Ou, S. & Jiang, N. LTR_retriever: A highly accurate and sensitive program for identification of long terminal repeat retrotransposons. *Plant Physiol.* (2018) doi:10.1104/pp.17.01310.
44. Simão, F. A., Waterhouse, R. M., Ioannidis, P., Kriventseva, E. V. & Zdobnov, E. M. BUSCO: Assessing genome assembly and annotation completeness with single-copy orthologs. *Bioinformatics* (2015) doi:10.1093/bioinformatics/btv351.
45. Garcia-Carpintero, V., de Martín, I., García-Juan, S. & Bombarely, A. Structural Genome Annotation QC with GAQET2 v2. (2025) doi:10.17504/PROTOCOLS.IO.8EPV5K5RNV1B/V1.
46. Sommer, M. J., Zimin, A. V. & Salzberg, S. L. PSAURON: a tool for assessing protein annotation across a broad range of species. *NAR Genomics Bioinforma.* **7**, (2025).
47. Nevers, Y. *et al.* Quality assessment of gene repertoire annotations with OMArk. *Nat. Biotechnol.* **2024 431** **43**, 124–133 (2024).
48. NCBI sequence read archive <https://identifiers.org/ncbi/insdc.sra:SRR32964043> (2025).
49. NCBI sequence read archive <https://identifiers.org/ncbi/insdc.sra:SRR35556615> (2025).
50. NCBI sequence read archive <https://identifiers.org/ncbi/insdc.sra:SRR35556614> (2025).
51. NCBI sequence read archive <https://identifiers.org/ncbi/insdc.sra:SRR35556613> (2025).

52. NCBI GenBank <https://identifiers.org/ncbi/insdc:JBUBWS0000000000> (2026).
53. NCBI GenBank <https://identifiers.org/ncbi/insdc:JBUCPO0000000000> (2026).
54. Chakraborty, A. & Xu, S. Chromosome-level genome assembly and annotation of *Nicotiana attenuata*. figshare. Dataset. <https://doi.org/10.6084/m9.figshare.30505763.v3> (2025).
55. Chakraborty, A. & Xu, S. Chromosome-level genome assembly and annotation of *Nicotiana obtusifolia*. figshare. Dataset. <https://doi.org/10.6084/m9.figshare.30505793.v2> (2025).

ARTICLE IN PRESS

a.**b.**

a.**b.**

a.**b.**