

Tarenna pendula (Rubiaceae), a new species from Guangxi, China

Yong-Hua Qin¹, Rainer W. Bussmann^{2,3}, Zhi-Rong Liu¹, Man Li^{1,4}, Chang-Ying Xia⁵, Wang-Hui Wu^{1,4}, Sheng-Xiang Yu^{6,7,8}

1 Guangxi Forestry Inventory and Planning Institute, CN-530011, Nanning, Guangxi, China

2 Institute of Botany and Bakuriani Alpine Botanical Garden, Ilia State University, Botanical Str. 1, 0105 Tbilisi, Georgia

3 Department of Botany, State Museum for Natural History Karlsruhe, Erbprinzenstraße 13, 76133 Karlsruhe, Germany

4 Guangxi Huasen Design Consulting Co., Ltd., CN-530011, Nanning, Guangxi, China

5 College of Teacher Education, Southwest University, CN-400715, Chongqing, China

6 State Key Laboratory of Plant Diversity and Specialty Crops, Institute of Botany, Chinese Academy of Sciences, Beijing 100093, China

7 China National Botanical Garden, Beijing 100093, China

8 College of Life Sciences, University of Chinese Academy of Sciences, Beijing 100049, China

Corresponding authors: Wang-Hui Wu (wu-wanghui@163.com); Sheng-Xiang Yu (yushengxiang@ibcas.ac.cn)

Abstract

The species *Tarenna pendula* (Rubiaceae), a small shrub, is newly described and illustrated from southwestern Guangxi Autonomous Region, China. This species is similar to *Tarenna tsangii* but is readily distinguished by its asymmetrical leaves, larger vegetative leaves than inflorescence leaves, adaxially pilosulous blades, pendulous inflorescences, longer pedicels (15–35 mm vs. 4–7 mm in *T. tsangii*), shorter corolla tubes (1–1.4 cm vs. 1.8–1.9 cm), multiple ovules per locule (vs. two), and sparsely pubescent fruits containing 13–19 seeds (vs. glabrous fruit containing only four seeds).

Key words: Distribution, Ixoroideae, morphology, Pavetteae, taxonomy



Academic editor: Petra De Block

Received: 5 February 2025

Accepted: 16 April 2025

Published: 3 June 2025

Citation: Qin Y-H, Bussmann RW, Liu Z-R, Li M, Xia C-Y, Wu W-H, Yu S-X (2025) *Tarenna pendula* (Rubiaceae), a new species from Guangxi, China. PhytoKeys 257: 1–8. <https://doi.org/10.3897/phytokeys.257.148816>

Copyright: © Yong-Hua Qin et al.

This is an open access article distributed under terms of the Creative Commons Attribution License (Attribution 4.0 International – CC BY 4.0).

Introduction

Tarenna Gaertn. is among the largest genera of the tribe Pavetteae, consisting of ca. 200 species within continental Africa, Madagascar, western Indian Ocean islands, Asia, and the Pacific region (De Block et al. 2015). The genus is distributed in primary evergreen forests and scrubs, in lowlands as well as at higher elevations. Recently, the circumscription of *Tarenna* was amended by the transfer of all African, Madagascan, and western Indian Ocean island species possessing fruits with a single, usually ruminant, seed to *Coptosperma* (Degreaf et al. 2001; De Block et al. 2001). Even in this narrower sense, *Tarenna* remains variable in certain morphological characters, such as flower, fruit, seed, and pollen types (De Block and Robbrecht 1998; De Block et al. 2001), which has raised questions as to its delimitation (Bridson 1979; Smith and Darwin 1988; De Block et al. 2001, 2015). Therefore, further studies on the plant diversity and phylogeny of *Tarenna* are needed.

During field expeditions in southern China, which is a major biodiversity hotspot (Myers et al. 2000; Hou et al. 2010) distinguished by extraordinarily high vascular plant species diversity and endemism (Xu 1995; Peng et al.

2014), a distinct *Tarenna* specimen was discovered in the Defu Nature Reserve in Napo, Guangxi Autonomous Region. A critical comparison of specimens (PE, IBK, and KUN) and literature (Chen and Taylor 2011) demonstrated that this newly discovered specimen represented a new species of *Tarenna*, which we describe and illustrate as *Tarenna pendula* sp. nov.

Taxonomy

***Tarenna pendula* Y.H.Qin, S.X.Yu & W.H.Wu, sp. nov.**

urn:lsid:ipni.org:names:77361544-1

Figs 1–3

Diagnosis. Similar to *Tarenna tsangii*, but distinguished by its small form, asymmetrical leaves, vegetative leaves larger than inflorescence leaves, blades adaxially pilosulous, pendulous inflorescences, longer pedicels (15–35 mm vs. 4–7 mm in *T. tsangii*), shorter corolla tubes (1–1.4 cm vs. 1.8–1.9 cm), multiple ovules per locule (vs. 2), and sparsely pubescent fruits with 13–19 seeds (vs. glabrous fruit containing only 4 seeds).

Type. CHINA • Guangxi Autonomous Region: Baise City, Napo County, Defu Nature Reserve, bamboo understory on Baxiong Mountain, 1380 m a.s.l., 23.292437°N, 105.784845°E, 8 July 2021, *Yonghua Qin YH2021075* (holotype: PE, isotype: IBK).

Paratypes. CHINA • Guangxi Autonomous Region: Baise City, Napo County, Defu Nature Reserve, bamboo understory in evergreen broadleaf forest on southern slope of Baxiong Mountain, 1380 m a.s.l., 23.292451°N, 105.784899°E, 30 May 2023, *Yonghua Qin YH2023069* (PE, IBK); • Baise City, Napo County, Defu Nature Reserve, bamboo understory in evergreen broadleaf forest on the southern slope of Baxiong Mountain, 1380 m a.s.l., 23.292485°N, 105.7847944°E, 7 September 2023, *Yonghua Qin YH2023116* (PE, IBK).

Description. Erect shrub up to 1–2 m tall, with a single main stem; branches flattened, shortly pilosulous, first year branchlets green, becoming brown or grayish-brown with age. Leaves opposite; petiole ca. 1 cm long, pilosulous; leaf blades asymmetrical, papery and blackish-brown when dried, oblong–obovate or oblanceolate, 4–17 cm × 1.5–4 cm, vegetative leaves large, 7–17 cm × 3–4 cm, inflorescence-supporting leaf pairs small, 4–7 cm × 1.5–3.5 cm, adaxially with adnate short hairs, abaxially scabrous and sparsely puberulent to subglabrous, with pubescence denser along principal veins, base cuneate, apex acuminate or shortly acuminate; midrib flat on upper surface, prominent on lower surface; secondary veins in 5–7 pairs, prominent on lower surface; stipules triangular, 3–5 mm, acuminate or apiculate when longer, persistent. Inflorescence corymbose, axillary or sometimes terminal, pendulous, 5–7 cm × ca. 3 cm, 3-5- or sometimes 9-flowered, shortly pedunculate, peduncles 3–8 mm, inflorescence axes and pedicels gray strigillose. Rachis unbranched. Flowers opposite on rachis, pedicels 1.5–3.5 cm long, proximal pedicels long, 2.5–3 cm long, distal pedicels short, 1.5–2 cm long. Calyx green, pilosulous; hypanthium portion cylindrical–urceolate, ca. 1 mm long; limb ca. 0.5 mm long, lobed for up to 1/2 of length. Corolla white, 1.5–1.9 cm long; tube cylindrical, ca. 5 mm long, inside at the upper part near the throat with dense hairs and at the lower part glabrous or sparsely pubescent, outside glabrous; lobes 5, contorted to the left in bud, oblong,




Figure 1. *Tarenna pendula* sp. nov. **A** flowering branches **B** portion of inflorescence showing stipule **C** portion of stem showing stipule **D** portions of adaxial and abaxial leaf surfaces, (**D1**) pressure side of blade, (**D2**) low pressure side of blade **E** flower **F** stamen **G** longitudinally opened corolla showing the position of the stamens **H** pedicel, bracteoles, ovary and calyx **I** ovary, calyx, style and stigma **J** fruit **K** cross section of ovary **L** longitudinal section of fruit. Drawings by Lijie Zhu, based on type specimens.




Figure 2. *Tarenna pendula* **A** habitat **B** flowering branch **C** pendulous cyme **D** flower (**D1**) and ovary and calyx (**D2**) **E** longitudinally opened corolla **F** ovary, calyx, style and stigma **G** fruiting branch **H** blade in adaxial view (**H1**) and in abaxial view (**H2**) **I** young fruits in different developmental stages **J** unripe seeds clustered together (after removal of fruit wall) **K** unripe seeds.




Figure 3. Distribution map of *Tarenna pendula* (triangle) and *Tarenna tsangii* (dots).

10–14 × 2–3 mm, with rounded apex, adaxial surface glabrous or sparsely pubescent at base, abaxial surface glabrous. Stamens inserted in corolla throat, alternating with corolla lobes, attached to the throat by a short filament; anthers yellowish-gray, linear, 8–10 mm long, basifixed, twisted after spreading. Style-stigma complex 1–1.2 cm long, exerted for 5–7 mm from corolla tube at anthesis; style sparsely pubescent at the middle; stigma fusiform, acentric apices. Ovary small, usually cup-shaped, 0.5–1 mm long, with 2 locules, each axile placenta with multiple ovules. Fruits berry-like, subglobose, 5–6 mm in diameter, with short persistent calyx teeth or calyx scar at the apex; seeds 13–19 per fruit, elliptic, 3–3.2 × 1.8–2 mm, brownish or black, testa rugose.

Phenology. *Tarenna pendula* has been observed to flower from May to July, and to fruit from July to January.

Distribution, habitat, and ecology. *Tarenna pendula* grows in southern subtropical evergreen broadleaf forests, particularly in mixed bamboo understory (*Indosasa sinica* C.D.Chu & C.S.Chao), at 1380 m a.s.l. The species is only known from this single locality in Napo County, Guangxi Autonomous Region, China (Fig. 3).

Conservation status. *Tarenna pendula* is only known from Baxiong Mountain, Defu Nature Reserve, Napo County, Guangxi, China with about 250 individuals recorded to date. Currently the species is only known from a single location. This location is inside a well-protected nature reserve, and as such, the species does not qualify for any threat condition. However, during fieldwork we noticed that the understory of the evergreen broadleaf forests in which *T. pendula* grows

Table 1. Morphological comparison of *Tarenna pendula* and *T. tsangii*.

	<i>T. pendula</i>	<i>T. tsangii</i>
Habit	Small shrubs, 1–2 m tall	Shrubs to trees, 1–6 m tall
Leaves	Asymmetrical, oblong–obovate or oblanceolate, 4–17 × 1.5–4 cm	Symmetrical, oblong–obovate or lanceolate, 5–26 × 1.5–7 cm
Stipules	Triangular, 3–5 mm, acuminate or apiculate when longer	Triangular, 4–5 mm, acuminate or apiculate
Blades	Adaxially with adnate short hairs	Adaxially glabrous
Pedicels	15–35 mm long	4–7 mm long
Cymes	Pendulous, with 3–5 or sometimes 9 flowers	Erect, with more than 10 flowers
Corolla	Corolla 1.5–1.9 cm long, tube ca. 5 mm long	Corolla 1.8–1.9 cm long, tube ca. 8–9 mm long
Styles	1–1.2 cm long	ca. 2 cm long
Ovary	Multiple ovules per locule	Two ovules per locule
Fruits	Sparsely pubescent	Glabrous
Seeds per fruit	13–19 seeds	4 seeds

is being colonized by luxuriant bamboo, which may constitute a danger to the new species. A second possible threatening factor for this apparently rare and narrow-ranged species is a limited capacity of natural regeneration. The conservation status for *Tarenna pendula* thus needs further study. We therefore assess the species as Data Deficient.

Etymology. The specific epithet is derived from the pendulous inflorescence, which differs from several other *Tarenna* spp. in the area.

Vernacular name. *Chuixiu wukoushu* in Mandarin Chinese, which translates to ‘pendulous *Tarenna*’.

Similar species. *Tarenna pendula* is similar to *T. tsangii* but is easily distinguished by its small form. Further diagnostic morphological characters of the new species and related species are presented in Table 1. Character information about relevant species is retrieved from Flora of China (Chen and Taylor 2011).

It is challenging to distinguish different species based on morphological characters alone. However, after carefully scrutinizing the morphological characters of the new species and related species, we found that the new species is easily recognized by its pendulous cymose inflorescences and large number of seeds per fruit. Although *T. tsangii* has a much larger distribution range than *T. pendula*, their areas do not overlap. Because the distribution range of *T. pendula* is immediately adjacent to that of *T. tsangii*, we hypothesize that these two species may be allopatric. Given the complicated evolutionary relationship among *Tarenna* species, future studies should investigate the plant diversity and phylogeny of this genus in greater detail.

Acknowledgments

The authors are grateful to Ms. Xun Liang for assistance in preparing the color plate and map, Ms. Lijie Zhu for the beautiful specimen drawings, Dr. Petra De Block for great efforts on the manuscript and two anonymous reviewers.

Additional information

Conflict of interest

The authors have declared that no competing interests exist.

Ethical statement

No ethical statement was reported.

Funding

This study was supported by the Guangxi Forestry Science and Technology Project, Ecological assessment and construction of ecological security system in Xijiang Economic Belt of Guangxi.

Author contributions

Yong-Hua Qin participated in the field investigation, wrote and revised the manuscript. Rainer W. Bussmann wrote and revised the manuscript. Zhi-Rong Liu and Man Li participated in the field investigation and completed part of the data processing. Chang-Ying Xia took photographs for identification and revised the manuscript. Wang-Hui Wu participated in the field investigation and revised the manuscript. Sheng-Xiang Yu wrote and revised the manuscript.

Author ORCIDs

Yong-Hua Qin  <https://orcid.org/0000-0002-9833-4193>

Rainer W. Bussmann  <https://orcid.org/0000-0002-3524-5273>

Zhi-Rong Liu  <https://orcid.org/0009-0001-5171-5693>

Man Li  <https://orcid.org/0009-0006-6985-4882>

Chang-Ying Xia  <https://orcid.org/0000-0002-5071-0145>

Wang-Hui Wu  <https://orcid.org/0009-0000-1565-2376>

Sheng-Xiang Yu  <https://orcid.org/0000-0002-5823-2870>

Data availability

All of the data that support the findings of this study are available in the main text.

References

- Bridson DM (1979) Studies in *Tarenna* sensu lato (Rubiaceae subfamily Cinchonoideae) for part 2 of 'Flora of tropical East Africa: Rubiaceae'. Kew Bulletin 34(2): 377–401. <https://doi.org/10.2307/4110002>
- Chen T, Taylor CM (2011) *Tarenna*. In: In Wu ZY, Raven PH, Hong DY (Eds) Flora of China. Vol. 19 (Rubiaceae). Science Press, Beijing, and Missouri Botanical Garden Press, St. Louis.
- De Block P, Robbrecht E (1998) Pollen morphology of the Pavetteae (Rubiaceae, Ixoroideae) and its taxonomic significance. Grana 37(5): 260–275. <https://doi.org/10.1080/00173139809362678>
- De Block P, Degreef J, Robbrecht E (2001) Reinstatement of the Afro-Madagascan genus *Coptosperma* (Rubiaceae). Systematics and Geography of Plants 71: 455–492. <https://doi.org/10.2307/3668694>
- De Block P, Razafimandimbison SG, Janssens S, Ochoterena H, Robbrecht E, Bremer E (2015) Molecular phylogenetics and generic assessment in the tribe Pavetteae (Rubiaceae). Taxon 64(1): 79–95. <https://doi.org/10.12705/641.19>
- Degreef J, De Block P, Robbrecht E (2001) A survey of continental African *Coptosperma* (Rubiaceae). Systematics and Geography of Plants 71(2): 367–382. <https://doi.org/10.2307/3668685>
- Hou MF, Lopez-Pujol J, Qin HN, Wang LS, Liu Y (2010) Distribution pattern and conservation priorities for vascular plants in southern China: Guangxi Province as a case study. Botanical Studies 51: 377–386.

- Myers N, Mittermeier RA, Mittermeier CG, da Fonseca GAB, Kent J (2000) Biodiversity hotspots for conservation priorities. *Nature* 403(6772): 853–858. <https://doi.org/10.1038/35002501>
- Peng CI, Ku SM, Yang HA, Leong WC, Liu Y, Nguyen TH, Kono Y, Chung KF (2014) Two new species of *Begonia* sect. *Coelocentrum*, *B. guixiensis* and *B. longa*, from Sino-Vietnamese limestone karsts. *Botanical Studies* 55(1): 52. <https://doi.org/10.1186/s40529-014-0052-8>
- Smith AC, Darwin SP (1988) Family 168. Rubiaceae. In: Smith AC (Ed.) *Flora vitiensis nova: A new flora of Fiji (Spermatophytes only)*, Vol. 4. Pacific Tropical Botanical Garden, Lawaii, 143–362.
- Xu ZR (1995) A study of the vegetation and floristic affinity of the limestone forests in southern and southwestern China. *Annals of the Missouri Botanical Garden* 82(4): 570–580. <https://doi.org/10.2307/2399837>