

Research Article

Biochemical Characterization of Partially Stabilized *Parthenium hysterophorus* L. and Cow Dung Consortium and Its Effect on the Growth Performance of Maize (*Zea mays* L.)

Shetie Gatew  and Andargachew Mengistu

Department of Biology, College of Natural and Computational Sciences, Arba Minch University, Arba Minch, Ethiopia

Correspondence should be addressed to Shetie Gatew; shetieg@yahoo.com

Received 17 August 2022; Revised 26 December 2022; Accepted 27 December 2022; Published 31 December 2022

Academic Editor: Rafael Clemente

Copyright © 2022 Shetie Gatew and Andargachew Mengistu. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Parthenium (*Parthenium hysterophorus* L.) is a major anxious weed found in pastures, wastelands, and agricultural lands. It usually competes very well and reduces crop yield. Up-rooting and herbicides have long been used as management options for weeds. However, these methods are neither economically nor environmentally sound. Another approach is to make compost of it, transforming a problem into an opportunity. Four treatments viz. *Parthenium* (100%) and *Parthenium* to cow dung ratio (75 : 25, 50 : 50, 25 : 75 w/w) were partially stabilized for 60 days under aerated conditions. The physical and chemical characteristics of partially stabilized organic materials were analyzed and the effects of the partially stabilized organic materials on the germination and growth performance of maize (*Zea mays* L.) were assessed. The results revealed that pH, EC, OC, OM, TN, P availability, and C/N were not significantly different; this showed an insignificant difference ($P > 0.05$) among the treatments. However, particle densities, P availability, EC, OC, and TN were found to be higher in 75 : 25 *Parthenium* cow dung ratios than in 25 : 75 ratios. Root length, shoot length, and germination index were significantly different among the treatments ($P \leq 0.05$). The seed germination index of maize was the highest (113%) in 75 : 25 ratios of *Parthenium* and cow dung followed by 50 : 50 ratios (95%) and 25 : 75 ratios (86%); relatively, low germination indices (84.86%) were recorded from partially stabilized organic materials prepared from *Parthenium* alone. The findings revealed that partially stabilized *Parthenium* mixed with cow dung could be promising for organic farming and an option for weed management.

1. Introduction

Parthenium hysterophorus L. (hereafter *Parthenium*) is an aggressive herbaceous plant native to the American continent that has already been documented in over 45 nations around neotropical countries [1, 2]. Currently, it is widely distributed in Australia, Pakistan, India, China, and Africa [2, 3]. It is one of the major invasive species in Ethiopia [4]. The plant grows readily in any habitat because of its several adaptation mechanisms such as simple seed dispersal mechanisms, drought tolerance, rapid seed production rates, avoidance of its natural enemies, and allelopathic properties [5–7]. Allelochemicals such as sesquiterpene lactones and phenolic acids (gallic acid, chlorogenic acid, and ferulic acid) [8] found in *Parthenium* are toxic to grazing animals and

have negative effects on crop production as well as the diversity and distribution of local flora and fauna [9]. Moreover, chemicals released from *Parthenium* to the substratum via its roots have inhibitory effects on both the germination and growth of a wide variety of crops [10–12].

The plant was introduced in Ethiopia in the 1970s [13]. However, there is no complete agreement so far on when and how it was introduced into the country. There are two speculations the weed was introduced during the Ethio-Somalia War in 1976 and the weed was introduced to Ethiopia together with the wheat seed donation from abroad [13]. The plant was first reported in Ethiopia in the 1980s from the eastern (Dire Dawa and Harar) and northeastern (near Dessie) parts of the country. Both of these places were important food aid distribution centers, and there is

substantial evidence that during the 1980s famine, *Parthenium* seeds were imported from subtropical North America as a contaminant of food assistance and were distributed with the grain [14]. Since its introduction, it has expanded persistently over Ethiopia's agricultural land, woodland, poorly managed arable crop land, and rangeland [15]. It also poses a health risk to humans and animals [16]. The weed negatively impacts food security and biodiversity. It is already threatening the crop production in Ethiopia [17, 18], and its rapid spread would further aggravate the risk of expansion and sustainable production of many crops. Therefore, controlling this aggressive invasive plant is critical for increasing the agricultural output and preventing environmental destruction.

Parthenium weed is widely spread in rangelands and cultivable fields and has negatively affected the composition and diversity of the rangeland vegetation in southern Ethiopia (Figure 1). Hence, the purpose of this research was to use invasive *Parthenium* weed as an input for preparing a partially stabilized organic material and reducing the invasion of the plant within the study area.

Many countries have been trying to control weeds, especially *Parthenium*, using a variety of tactics [4, 19, 20]. However, they were not economically feasible for most small-scale farmers [21]. This is because it colonizes new lands faster than it can be controlled [2]. Moreover, most single applications were not effective or had only short-term effects [7], and this does not easily fit into the broad category of weed control in the specific cropping patterns or systems of smallholder farmers [21].

The Ethiopian government has faced many challenges in eradicating this plant. Several physical and chemical treatments employed in the past are ineffective, expensive, and environmentally unfriendly. The chemical control has supremacy over other control methods for quick relief; however, the effect is always temporary and repeated operations will not be cost-effective for smallholder farming systems [22]. The repeated application of herbicides can affect other beneficial plants in addition to changing soil characteristics and polluting the environment due to cumulative residual effects [10]. Similar to many other weeds, *Parthenium* weeds are beneficial. The search for more effective utilization of *Parthenium* will render it suitable for effective management on the one hand and it can also be used for many different purposes besides eradication on the other hand [13]. Similarly, Saini et al. [23] suggested that large-scale utilization of *Parthenium* would also be an effective method for weed management. The most promising method for controlling *Parthenium* weeds is management through utilization. Long-term control requires the combined application of many technologies [24]. Ghadge et al. [25] proved that *Parthenium* can be used for the preparation of compost and vermicompost that contain high nutrients. Ameta et al. [26] also indicated that seeds of *Parthenium* did not germinate after composting, which could also encourage end-users of the compost.

Parthenium has the potential to be used as a green manure, compost for crop plants, biopesticides, and soil amendments [27–29]. This study aimed to assess the physicochemical characteristics of partially stabilized *Parthenium* and cow dung consortium and evaluate its effect on the germination and growth performance of maize.

2. Materials and Methods

The study was conducted in Mirab Abaya District, southern Ethiopia. The *Parthenium* weed was collected from the fallow lands before the flowering stage for its soft stem and leaves and cow dung was collected from the farmers' houses. Experimental plots were established with equal sized partially stabilizing organic material pits of $1 \times 2 \times 1 \text{ m}^3$ (width $\times$ depth $\times$ length). The pits were established in dry and elevated places to prevent water runoff from entering the pits. The sidewall and surface of the pit were compacted to prevent leaching of essential nutrients from the organic material to the soil surface. The following four treatments were prepared for partially stabilized organic material production: *Parthenium* biomass + cow dung (CD) (75:25) w/w, *Parthenium* biomass + CD (50:50) w/w, *Parthenium* biomass + CD (25:75) w/w, and *Parthenium* biomass alone (100) w/w as an experimental control (Table 1). The organic materials in all piles were kept the same, i.e., 50 kg. Each treatment was performed in triplicate.

The collected *Parthenium* weed and cow dung were spread and piled layer by layer. In the stabilizing pits, sticks were placed vertically into the organic material to allow air to circulate in various layers. Sticks were placed at each corner and at the center of the pit before piling up the materials. After piling up the layers, the sticks were joggled to expand the space between the sticks and materials to create ventilation holes to keep the layers properly aerated. After three weeks of partial decomposition, the pits were turned periodically to maintain aeration and the decomposed material was mixed from the bottom to the top and vice versa. The temperature and moisture content of partially stabilized organic materials were measured periodically using a soil temperature moisture meter (RS485 4–20 mA).

The total biomass used for partial stabilizing process was 375 kg *Parthenium* and 225 kg cow dung.

2.1. Analysis of Physical Characteristics. The bulk density and moisture content of the 60th day partially stabilized *Parthenium* with different proportions of cow dung organic materials were measured. The bulk density was measured as the ratio of the dry weight of the partially stabilized organic material to the volume of the iron ring. The iron ring was filled with the partially stabilized organic material and slightly compacted to ensure the absence of large void spaces. The moisture content of the partially stabilized organic material was measured after drying at 105°C for approximately 24 h or after the partially stabilized organic material was measured at a constant weight.

FIGURE 1: Infestations of *Parthenium* weed in Mirab Abaya Woreda, south Ethiopia.

TABLE 1: The composition of raw materials in each treatment pits.

Treatments	<i>P. hysterophorus</i>		Cow dung		Total material	
	Kg	%	Kg	%	Kg	%
T1	50	100	—	—	50	100
T2	37.5	75	12.5	25	50	100
T3	25	50	25	50	50	100
T4	12.5	25	37.5	75	50	100

$$\text{Moisture content (\%)} = \frac{\text{Weight of moist organic material} - \text{weight of oven dry organic material}}{\text{Weight of oven dry organic material}} \times 100. \quad (1)$$

2.2. Chemical Analysis. Partially stabilized organic material samples were collected in plastic bags from well-mixed composites of 200 g from each experimental pit for chemical analysis. The electrical conductivity (EC), pH, organic carbon (OC), organic matter (OM) total nitrogen (TN), C/N ratio, available phosphorus (P_{avail}), and available potassium (K_{avail}) were analyzed. The pH was measured with a digital pH meter (Beckman model 72 pH meter) in a supernatant suspension of 1:2.5 soil:H₂O [30]. The electrical conductivity (EC) was measured using an EC meter based on a 1:2.5 extract from a soil:H₂O suspension [31]. The total organic carbon was determined using the Walkey-Black rapid titration method [32]. The organic matter was obtained by multiplying the percentage of carbon by 1.724 [33]. Available phosphorus was determined using the Olsen procedure [34]. Total nitrogen (TN) was measured using the Kjeldahl method [33]. Available potassium was quantified using flame photometry [33, 35].

2.3. Seed Germination Test. Maize seed germination and shoot and root growth performance tests were conducted in triplicate in five plastic pots (10.8 cm height × 20 cm diameter) with 75% of garden soil and 25% of the partially stabilized organic material from each pile and an equal amount of pure water was added every morning and evening time. Twelve uninfected healthy maize seeds were sown in cylindrical plastic pots and then placed in a shaded place. The same setup was used for garden soil as an experimental control. After 20 days, the percentage of seed germination, root and shoot lengths, and seed germination index [36] were estimated. The germination index (GI) is a measure of relative seed germination (RSG) and root growth (RRG). The root length was measured from the root collar to the root tip, and the shoot length was recorded from the root collar to the shoot apex [36].

RSG is calculated as follows:

$$\frac{\text{Mean number of seed germinated in partially stabilized organic material}'}{\text{Mean number of seed germinated in the garden soil}} \times 100. \quad (2)$$

RRG is calculated as follows:

$$\frac{\text{Mean root length of seedling in the partially stabilized organic material}}{\text{Mean root length of seedling in the garden soil}} \times 100. \quad (3)$$

Therefore, GI is the value obtained from

$$\frac{\text{RSG}}{\text{RRG}} \times 100. \quad (4)$$

3. Results and Discussion

3.1. Physical Properties of Partially Stabilized Organic Materials. The results indicated that the bulk density of the partially stabilized organic material did not show statistically significant variation among treatments ($P > 0.05$) (Table 2). The range of the bulk densities of this study ($1.23\text{--}1.27 \text{ g cm}^{-3}$) was comparable to that of the research performed by Hunt and Gilkes [37]. The bulk density (BD) of the partially stabilized organic material was suitable for root growth and soil permeability. According to Nyéki et al. [38], the optimal bulk density for the plant growth is <1.6 , <1.4 , and $<1.10 \text{ g cm}^{-3}$ in sand, silt, and clay soils, respectively.

The mean value of the moisture content was higher in *Parthenium* (100%) treatment pits and lowest in *Parthenium* + cow dung (25 : 75) (Table 2). The moisture content of this study was less than that of the research carried out by Christian et al. (1977) (45–65%).

The porosity values ranged from 0.360 ± 0.51 to $0.440 \pm 0.51\%$ (Table 2). It depends on the bulk density and moisture content of the partially stabilized organic material [39]. The porosity was under that of the result obtained by Ahn et al. [40] (0.60–0.72%). However, according to Cresswell and Hamilton [39], the typical range of porosity in compost prepared from commonly used composting materials was 0.5–0.7%.

3.2. Chemical Properties of the Partially Stabilized Organic Material. The mean pH values of the treatment pits were found in the range between 7.90 ± 0.99 and 8.30 ± 0.99 (Table 3). The partially stabilized organic material was alkaline and which is in agreement with research conducted by Travis et al. (2003). The highest pH (8.3 ± 0.99) was found in T3, *Parthenium* biomass + CD (50 : 50) w/w while the lowest value of pH (7.9 ± 0.99) was obtained for T2, from *Parthenium* biomass + CD (75 : 25). However, there was no statistically significant difference ($P > 0.05$) in the pH values among the treatments (Table 3).

The electrical conductivities (EC) of treatment pits were in the ranges between 0.86 ± 0.137 and $1.10 \pm 0.137 \text{ mScm}^{-1}$. However, the highest value of EC ($1.10 \pm 0.137 \text{ mScm}^{-1}$) was found for T2 (75% *Parthenium* with 25% cow dung), and the lowest value of EC ($0.86 \pm 0.137 \text{ mScm}^{-1}$) was found for T1 (*Parthenium* weed alone). The range of EC ($<4 \text{ mScm}^{-1}$) revealed that the partially stabilized organic material prepared from *Parthenium* and cow dung support the agro-nomic application. The result was in line with the finding of Wang et al. [41].

The total organic carbon result was found between ranges from 2.57 ± 0.443 to 3.54 ± 0.443 and the lowest value was 2.54 ± 0.443 in T4 (*Parthenium* + CD; 25 : 75), and the highest value was 3.54 ± 0.443 in T2 (*Parthenium* + CD; 75 : 25). This result was lower than that was reported by Sivakumar et al. [42]. Conversely, the highest total organic matter content ($6.11 \pm 0.766\%$) was found in T2 (*Parthenium* + CD; 75 : 25), and the lowest total organic matter ($4.41 \pm 0.766\%$) was found in T4 (*Parthenium* + CD; 25 : 75) (Table 3). Previous studies reported that the total organic matter of *Parthenium* compost was between 17.33% and 16.31% [43–46], which is higher than the result obtained in this study.

The total nitrogen content of the partially stabilized organic material ranged from 0.221 ± 0.38 in T4 to 0.306 ± 0.38 in T2 (Table 3). Travis et al. [47] reported that the total nitrogen content ranged from 0.5 to 2.5%. Low total nitrogen results from highly alkaline compost pits [48]. Composts with a relatively high pH are found to have a lower quantity of total nitrogen due to the loss of nitrogen through volatilization in the form of ammonia [47].

The highest available phosphorus content was recorded in T1 ($135.85 \pm 13.072 \text{ ppm}$), which was the organic material prepared from *Parthenium* weed alone and the lowest was in T4 ($77.10 \pm 13.072 \text{ ppm}$) (0.25% *Parthenium* with 75% cow dung). Similarly, the quantity of available potassium was higher in T1 ($403.33 \pm 34.50 \text{ ppm}$) and the lowest in T4 ($71 \pm 34.50 \text{ ppm}$) (Table 3). Available phosphorus and potassium levels were higher than those reported by Ballard et al. (2020). The exchangeable potassium contents of partially stabilized organic materials in the four treatment pits were statistically significant ($P < 0.05$) (Table 3). A study conducted in India by Channappagoudar et al. [49] indicated that composting *Parthenium* alone before the flowering stage has higher available potassium than mixing it with other organic matter, which is in agreement with the results of this study.

The C:N ratio ranged from 11.587 ± 0.005 to 11.60 ± 0.005 (Table 3). The C:N ratio is mostly used as a measure of compost maturity, stability, and N availability [50]. According to Antil et al. [51], a C:N ratio of less than 20 is thought to indicate compost maturity, whereas a C:N ratio of 15 or less is ideal [44, 50]. The results indicated that there was no statistically significant variation ($P > 0.05$) in the C:N ratio among treatments (Table 3). This showed that all treatment pits contained easily biodegradable matrices and the transformation process might have been similarly carried out in all treatment pits. The results of the C:N ratio were inconsistent with those obtained by Travis et al. (2003) who found that the C:N ratio was between 10.82 ± 0.79 and 11.42 ± 0.21 . Except for available potassium, all the chemical properties of partially stabilized organic materials showed insignificant statistical differences ($P > 0.05$) among the treatments (Table 3).

TABLE 2: The mean values of physical properties of the partially stabilized organic material.

Parameter	Treatments				P value
	T1 P (100%)	T2 P + CD (75 : 25)	T3 P + CD (50 : 50)	T4 P + CD (25 : 75)	
BD (g/cm ³)	1.25 ± 0.087	1.27 ± 0.087	1.23 ± 0.087	1.277 ± 0.087	0.982
MC (%)	35.67 ± 2.12	31.22 ± 2.12	28.80 ± 2.12	24.24 ± 2.12	0.030
p (%)	0.43 ± 0.51	0.44 ± 0.51	0.43 ± 0.51	0.36 ± 0.51	0.697

P, *Parthenium*; CD, cow dung; BD, bulk density; MC, moisture content; p (%), porosity.

TABLE 3: The mean values of chemical properties of the partially stabilized organic material.

Parameter	Treatments				P value
	T1 P (100%)	T2 P + CD (75 : 25)	T3 P + CD (50 : 50)	T4 P + CD (25 : 75)	
pH (H ₂ O)	8.06 ± 0.099	7.9 ± 0.099	8.3 ± 0.099	8.23 ± 0.099	0.078
EC (mS/cm)	0.86 ± 0.137	1.1 ± 0.137	0.9 ± 0.137	0.96 ± 0.137	0.654
OC (%)	2.80 ± 0.443	3.55 ± 0.443	3.13 ± 0.443	2.57 ± 0.443	0.463
OM (%)	4.84 ± 0.766	6.12 ± 0.766	5.40 ± 0.766	4.41 ± 0.766	0.467
TN (%)	0.242 ± 0.38	0.306 ± 0.38	0.270 ± 0.38	0.221 ± 0.38	0.475
P _{avail.} (ppm)	135.85 ± 13.07	120.80 ± 13.07	116.60 ± 13.07	77.10 ± 13.07	0.063
K _{avail.} (ppm)	403.33 ± 34.5	365 ± 34.5	230.41 ± 34.5	71 ± 34.5	0.002
C/N	11.58 ± 0.005	11.59 ± 0.005	11.6 ± 0.005	11.59 ± 0.005	0.370

P, *Parthenium*; CD, cow dung; EC, electrical conductivity; OC, organic carbon; OM, organic matter; TN, total nitrogen; P_{avail.} available phosphorus; K_{avail.} available potassium; C/N, carbon to nitrogen ratio.

TABLE 4: The mean value of shoot and root lengths and the germination index of maize seeds.

Treatments	Mean no. of seedling (n = 12)	Root length (cm)	Shoot length (cm)	GI (%)	P value
Garden soil (control)	8 ± 0.632	14.14 ± 0.458	13.03 ± 0.567	100	0.000
P compost	7 ± 0.632	14.58 ± 0.458	13.60 ± 0.567	84.86	0.000
P + CD (75 : 25)	10 ± 0.632	15.70 ± 0.458	14.65 ± 0.567	112.61	0.000
P + CD (50 : 50)	9 ± 0.632	18 ± 0.458	15 ± 0.567	95.01	0.000
P + CD (25 : 75)	8 ± 0.632	16.3 ± 0.458	15.76 ± 0.567	86.37	0.000

3.3. Seed Germination and Seedling Growth. Maize seed germination index (GI) was found to be highest (112.61%) in *Parthenium* + CD (75 : 25) and lowest (84.86%) in *Parthenium* alone (Table 4).

The seed germination test is an effective method for determining the potential toxicity of plants [36]. GI and phytotoxins have an inverse relationship; a decrease in GI reveals the presence of phytotoxins in the partially stabilized organic mixture of *Parthenium* and cow dung, whereas an increase in GI indicates the absence of phytotoxic chemicals [36, 52].

A seed germination test on maize revealed considerable variance in root length, shoot length, and germination index in 60 days partially stabilized organic materials ($P < 0.05$). The results of GI in any of the treatments were within the acceptable limit (>80%) with no toxic effect on the growth of plants as suggested by Tiquia [52] for completely matured compost [52, 53].

4. Conclusions

Parthenium is mostly known for its traits in natural ecosystems and for its adverse effects on human and animal

health. Various techniques have been used to control this toxic weed. The most promising method is to manage the weed by proper utilization. The partially stabilized *Parthenium* weed with cow dung has higher nutritional values and germination index in contrast to their controls. The results of the experiments and laboratory analyses revealed that among the treatments of partially stabilized organic materials prepared from *Parthenium* alone showed the highest value of the moisture content, available phosphorus, and available potassium although there were no significant differences among the treatments. The highest proportion of cow dung combined with *Parthenium* (75 : 25) had the lowest moisture content, porosity, organic carbon, total nitrogen, available phosphorus, and available potassium. Partially stabilized organic materials prepared from the high concentration of *Parthenium* mixed cow dung (75 : 25) showed the highest values of porosity, electric conductivity, organic carbon, total nitrogen, and organic matter, while the same showed the lowest pH. Organic carbon, total nitrogen, and organic matter values were lower in this study. This might be the short stabilization process. The seed germination and growth test on maize indicated that the partially

stabilizing unflowered *Parthenium* weed with cow dung can be used as a rich source of nutrients and its toxicity is relatively minimized. More field research studies and seed germination tests with other crops are needed to translate this research into a wider eco-friendly approach.

Based on the findings of this study, enhancing farmers' knowledge by training on quality matured compost making and working further to improve the nutrient content of *Parthenium* are essential. These practices require the integration of the community and responsible governmental and nongovernmental organizations. More field research studies and seed germination tests with other crops are recommended with *Parthenium* and cow dung matured compost.

Data Availability

All data generated and analyzed during this study are included within the article.

Consent

Not applicable.

Conflicts of Interest

The authors declare that they have no conflicts of interest.

Authors' Contributions

Shetie Gatew focused on invasive plant species and soil science, interpreted the results, analyzed the data, and reviewed and edited the manuscript. Andargachew Mengistu conceptualized, designed, and conducted the field-work, analyzed data, and drafted the manuscript. The final version of this manuscript has been communicated to and approved by the authors.

Acknowledgments

The authors are grateful for the financial assistance provided by the Arba Minch University. The Wolaita Sodo Soil Laboratory Center and the Mirab Abaya Agricultural and Natural Resource Development Office are acknowledged for their generous assistance.

References

- [1] I. J. Dale, "Parthenium weed in the americas report on the ecology of *Parthenium hysterophorus* in south central and North America," *Australian Weeds*, vol. 1, pp. 8–14, 1981.
- [2] A. A. Bajwa, B. S. Chauhan, M. Farooq, A. Shabbir, and S. W. Adkins, "What do we really know about alien plant invasion? A review of the invasion mechanism of one of the world's worst weeds," *Planta*, vol. 244, no. 1, pp. 39–57, 2016.
- [3] A. Javaid and I. Haider Khan, "Potential use of *Coronopus didymus* (L.) SM.in parthenium management," *Pakistan Journal of Weed Science Research*, vol. 27, no. 2, pp. 37–45, 2020.
- [4] M. Wakjira, G. Berecha, and B. Bulti, "Allelopathic effects of *Parthenium hysterophorus* extracts on seed germination and seedling growth of lettuce," *Tropical Science*, vol. 45, no. 4, pp. 159–162, 2005.
- [5] R. K. Kohli, D. R. Batish, H. P. Singh, and K. S. Dogra, "Status, invasiveness and environmental threats of three tropical American invasive weeds (*Parthenium hysterophorus* L ., *Ageratum conyzoides* L ., *Lantana camara* L .).India," *Biological Invasions*, vol. 8, no. 7, pp. 1501–1510, 2006.
- [6] A. Javaid, S. Shafique, and S. Shafique, "Causes of rapid spread of *Parthenium hysterophorus* L. in Pakistan and possible control measures – a review," *Pakistan Journal of Botany*, vol. 39, no. 7, pp. 2611–2618, 2007.
- [7] M. Kaur, N. K. Aggarwal, V. Kumar, and R. Dhiman, "Effects and management of *Parthenium hysterophorus* a weed of global significance," *International Scholarly Research Notices*, vol. 2014, Article ID 368647, 12 pages, 2014.
- [8] T. T. Bezuneh, "Phytochemistry and antimicrobial activity of *Parthenium hysterophorus* L," *A Review. Sci J Anal Chem*, vol. 3, pp. 30–38, 2015.
- [9] S. Talemso, A. Abreham, M. Fisseha, and B. Alemayehu, "Distribution status and the impact of parthenium weed (*Parthenium hysterophorus* L.) at Gedeo Zone (Southern Ethiopia)," *African Journal of Agricultural Research*, vol. 8, no. 4, pp. 386–397, 2013.
- [10] H. P. Singh, D. R. Batish, R. K. Kohli, D. B. Saxena, and V. Arora, "Effect of parthenin — a sesquiterpene lactone from *parthenium hysterophorus* - on early growth and physiology of *ageratum conyzoides*," *Journal of Chemical Ecology*, vol. 28, no. 11, pp. 2169–2179, 2002.
- [11] H. P. Singh, D. R. Batish, J. K. Pandher, and R. K. Kohli, "Assessment of allelopathic properties of *Parthenium hysterophorus* residues," *Agriculture, Ecosystems & Environment*, vol. 95, no. 2-3, pp. 537–541, 2003.
- [12] H. P. A. L. Singh, D. R. Batish, J. K. Pandher, and R. K. Kohli, "Phytotoxic effects of *Parthenium hysterophorus* residues on three Brassica species," *Weed Biology and Management*, vol. 5, no. 3, pp. 105–109, 2005.
- [13] T. Tefera, "Allelopathic effects of *Parthenium hysterophorus* extracts on seed germination and seedling growth of *Eragrostis tef*," *Journal of Agronomy and Crop Science*, vol. 188, pp. 306–310, 2002.
- [14] T. Tamado and P. Milberg, "Weed flora in arable fields of eastern Ethiopia with emphasis on the occurrence of *Parthenium hysterophorus*," *Weed Research*, vol. 40, no. 6, pp. 507–521, 2000.
- [15] A. Etana, E. Kelbessa, and T. Soromessa, "Impact of *Parthenium hysterophorus* L . (asteraceae) on herbaceous plant biodiversity of awash national park (ANP), Ethiopia," *Management of Biological Invasions*, vol. 2, no. 1, pp. 69–80, 2011.
- [16] S. M. Masum, M. Hasanuzzaman, and M. H. Ali, "Threats of *Parthenium hysterophorus* on agroecosystems and its management a review," *The International Journal of Agriculture and Crop Sciences*, vol. 6, no. 11, pp. 684–697, 2013.
- [17] S. Ayele, L. Nigatu, T. Tana, and S. W. Adkins, "Impact of parthenium weed (*Parthenium hysterophorus* L .) on the above-ground and soil seed bank communities of rangelands in Southeast Ethiopia," *International Research Journal of Agricultural Science and Soil Science*, vol. 3, no. 7, pp. 262–274, 2013.
- [18] D. Million, L. Nigatu, Z. Bekeko, and H. Legesse, "Distribution and impact of parthenium (*parthenium hysterophorus* L.) on weed species diversity in maize fields in western gojjam zone, amhara national regional state, Ethiopia," *International*

- Journal of Agronomy*, vol. 2021, no. 1, Article ID 368647, 10 pages, 2021.
- [19] N. Javaid, I. H. Khan, M. H. Shah, A. Javaid, and S. M. Waleed, "Herbicidal activity of *Ageratum conyzoides* against parthenium," *Pakistan Journal of Weed Science Research*, vol. 27, no. 2, pp. 137–146, 2020.
 - [20] A. Javaid, T. Mubeen, I. Haider Khan, K. Jabeen, and M. Akbar, "Herbicidal potential of *Alternaria citri ellis* and pierce metabolites against *Parthenium hysterophorus* L.," *Allelopathy Journal*, vol. 55, no. 1, pp. 25–34, 2022.
 - [21] T. Tana, "Biology and management of parthenium weed (*Parthenium hysterophorus* L.) in Eastern Ethiopia," *Swedish University of Agricultural Sciences, Department of Ecology and Crop Production Science, Doctoral thesis*, Uppsala, 2001.
 - [22] T. Tamado and P. Milberg, "Control of *Parthenium* (*Parthenium hysterophorus*) in grain Sorghum (*Sorghum bicolor*) in the smallholder farming system in Eastern Ethiopia," *Weed Technology*, vol. 18, no. 1, pp. 100–105, 2004.
 - [23] A. Saini, N. K. Aggarwal, A. Sharma, M. Kaur, and A. Yadav, "Utility potential of *Parthenium hysterophorus* for its strategic management," *Advances in Agriculture*, vol. 2014, pp. 1–16, Article ID 381859, 2014.
 - [24] S. Adkins and A. Shabbir, "Biology, ecology and management of the invasive parthenium weed (*Parthenium hysterophorus* L.)," *Pest Management Science*, vol. 70, no. 7, pp. 1023–1029, 2014.
 - [25] S. Ghadge, P. Naikwade, and B. Jadhav, "Utilization of problematic weed for improved yield of Fenugreek," *Indian Streams Research Journal*, vol. 3, no. 4, pp. 1–8, 2013.
 - [26] S. K. Ameta, M. S. Solanki, N. Sharma, and R. Ameta, "Effect of compost of *Parthenium hysterophorus* weed on germination and growth of maize," *International Journal of Applied and pure Science and Agriculture*, vol. 2, no. 8, pp. 12–19, 2016.
 - [27] A. Javaid and M. B. M. Shah, "Use of parthenium weed as green manure for maize and mungbean production," *The Philippine Agricultural Scientist*, vol. 91, no. 4, pp. 478–482, 2008.
 - [28] A. Javaid and M. B. M. Shah, "Growth and yield response of wheat to EM (Effective microorganisms) and parthenium-green manure," *African Journal of Biotechnology*, vol. 9, no. 23, pp. 3378–3381, 2010.
 - [29] P. Kishor, B. R. Maurya, and A. K. Ghosh, "Use of uprooted parthenium before flowering as composta way to reduce its Hazards worldwide," *International Journal of Soil Science*, vol. 5, no. 2, pp. 73–81, 2010.
 - [30] L. Reeuwijk van, *Procedures for Soil Analysis*, International Soil Reference and Information Centre, Wageningen, The Netherlands, 6th edition, 2002.
 - [31] E. George, R. Sommer, and J. Ryan, "Methods of soil, plant, and water analysisa manual for the west asia and North Africa regionbeirut, Lebanon," *International Center for Agricultural Research in the Dry Areas (ICARDA)*, vol. 40, pp. 83–89, 2013.
 - [32] A. Walkley and I. A. Black, "An examination of the Degtjareff method for determining soil organic matter, and a proposed modification of the chromic acid titration method," *Soil Science*, vol. 37, no. 1, pp. 29–38, 1934.
 - [33] P. Haluschak, *Laboratory Methods of Soil Analysis*, Canada-Manitoba Soil Survey, Beijing China, 2006.
 - [34] S. R. Olsen, "Estimation of available phosphorus in soils by extraction with sodium bicarbonate," *Journal of Chemical Information and Modeling*, vol. 53, no. 9, pp. 1689–1699, 1954.
 - [35] D. L. Rowell, "Soil sciencemethods & applications. Book reviews," *Journal of the Science of Food and Agriculture*, vol. 66, pp. 513–576, 1994.
 - [36] F. Zucconi, A. Pera, M. Forte, and M. Bertoldi, "Evaluating toxicity of immature compost," *Biocycle*, vol. 22, pp. 54–57, 1981.
 - [37] N. Hunt and B. Gilkes, *Farm Monitoring Handbook A Practical Down-To-Earth Manual for Farmers and Other Land Users*, University of Western Australia, Nedlands, W.A, 1992.
 - [38] A. Nyéki, G. Milics, A. J. Kovács, and M. Neményi, "Effects of soil compaction on cereal yields review," *Cereal Research Communications*, vol. 45, pp. 1–22, 2017.
 - [39] H. P. Cresswell and G. J. Hamilton, "Bulk density and pore space relations. Soil physical measurement and interpretation for land evaluation," *Australian Soil and Land Survey Handbook*, vol. 5, pp. 35–58, 2002.
 - [40] H. K. Ahn, T. L. Richard, and T. D. Glanville, "Laboratory determination of compost physical parameters for modeling of airflow characteristics," *Waste Management (Tucson, Arizona)*, vol. 28, no. 3, pp. 660–670, 2008.
 - [41] Y. Wang, J. Gong, J. Li et al., "Insights into bacterial diversity in compost core microbiome and prevalence of potential pathogenic bacteria," *Science of the Total Environment*, vol. 718, Article ID 137304, 2020.
 - [42] S. Sivakumar, P. Senthilkumar, Y. Song, and K. Maritime, "Efficiency of composting parthenium plant and neem leaves in the presence and absence of an oligochaete, *Eisenia fetida*," *Iranian Journal of Environmental Health Science & Engineering*, vol. 6, no. 3, pp. 201–208, 2009.
 - [43] R. Gupta, P. K. Mutiyar, N. K. Rawat, M. Saini, and V. Garg, "Development of a water hyacinth based vermireactor using an epigeic earthworm *Eisenia foetida*," *Bioresource Technology*, vol. 98, no. 13, pp. 2605–2610, 2007.
 - [44] D. Raj and R. S. Antil, "Evaluation of maturity and stability parameters of composts prepared from agro-industrial wastes," *Bioresource Technology*, vol. 102, no. 3, pp. 2868–2873, 2011.
 - [45] A. Yadav and V. K. Garg, "Recycling of organic wastes by employing *Eisenia fetida*," *Bioresource Technology*, vol. 102, no. 3, pp. 2874–2880, 2011.
 - [46] S. Suthar, B. Pandey, R. Gusain, and R. Z. Gaur, "Nutrient changes and biodynamics of *Eisenia fetida* during vermicomposting of water lettuce (*Pistia* sp.) biomass a noxious weed of aquatic system," *Environmental Science and Pollution Research*, vol. 10, no. 1, 2016.
 - [47] J. W. Travis, N. Halbrecht, B. Hed, and I. Chen, *A Practical Guide to the Application of Compost in Vineyards*, Penn State University, University Park, Pennsylvania, PA, 2003.
 - [48] T. Mengistu, H. Gebrekidan, K. Kibret, K. Woldetsadik, B. Shimelis, and H. Yadav, "Comparative effectiveness of different composting methods on the stabilization, maturation and sanitization of municipal organic solid wastes and dried faecal sludge mixtures," *Environmental Systems Research*, vol. 6, no. 1, pp. 5–16, 2017.
 - [49] B. B. Channappagoudar, N. R. Biradar, J. B. Patil, and C. A. A. Gasimani, "Utilization of weed biomass as an organic source in sorghum," *Karnataka J. Agric. Sci.*, vol. 20, pp. 245–248, 2007.
 - [50] M. P. Bernal, J. A. Alburquerque, and R. Moral, "Composting of animal manures and chemical criteria for compost maturity assessment. A review," *Bioresource Technology*, vol. 100, no. 22, pp. 5444–5453, 2009.
 - [51] R. S. Antil, D. Raj, R. P. Narwal, and J. P. Singh, "Evaluation of maturity and stability parameters of composts prepared from organic wastes and their response to wheat," *Waste and Biomass Valorization*, vol. 4, no. 1, pp. 95–104, 2013.

- [52] S. M. Tiquia, "Microbiological parameters as indicators of compost maturity," *Journal of Applied Microbiology*, vol. 99, no. 4, pp. 816–828, 2005.
- [53] F. Sellami, S. Hachicha, M. Chtourou, K. Medhioub, and E. Ammar, "Maturity assessment of composted olive mill wastes using UV spectra and humification parameters," *Bioresource Technology*, vol. 99, no. 15, pp. 6900–6907, 2008.