Canadian Science Publishing | Canadian Journal of Forest Research

The impact of climate on the dominant height and climate thresholds for *P. elliottii*, *P. taeda*, *P. patula* and *P. patula x P. tecunumanii* plantation forests

Journal:	*Canadian Journal of Forest Research*
Manuscript ID	cjfr-2024-0005.R1
Manuscript Type:	Research Article
Date Submitted by the Author:	15-Mar-2024
Complete List of Authors:	van der Merwe, Jaco-Pierre; University of British Columbia, Department of Wood Science; York Timbers, Research Division Lindner, Gerard; York Timbers, Research Division; York Timbers, Planning Division; Stellenbosch University, Department of Forestry; York Timbers, Tree Breeding Division Ernst, Yolandi; University of the Witwatersrand, School of Animal, Plant and Environmental Sciences, Faculty of Science Germishuizen, Ilaria; KZN, Institute for Commercial Forestry Research van Heerden, Elane; York Timbers, Planning Division Zandberg, Hugo; York Timbers, Planning Division Sibiya, Gabigabi; York Timbers, Planning Division Raboho, Ndamulelo; York Timbers, Planning Division De Waal, Lizette; York Timbers, Research Division; York Timbers, Tree Breeding Division Ndlovu, Thabani; York Timbers, Research Division; York Timbers, Tree Breeding Division Mabaso, Leonard; York Timbers, Research Division; York Timbers, Tree Breeding Division Zondach, Niel; York Timbers, Research Division; York Timbers, Tree Breeding Division Christie, Nanette; University of Pretoria, Department Biochemistry Genetics and Microbiology Clarke, Charlie; CharlieC Consulting Mansfield, Shawn; University of British Columbia, Department of Wood Science; University of British Columbia, Department of Botany
Is the manuscript for consideration in a Special Issue or Collection?:	Not applicable (regular submission)
Keyword:	Climate, Tree height, Pinus elliottii, Pinus taeda, Pinus patula

© The Author(s) or their Institution(s)

SCHOLARONE™
Manuscripts

© The Author(s) or their Institution(s)

The impact of climate on the dominant height and climate thresholds for *P. elliottii*, *P. taeda*, *P. patula* and *P. patula x P. tecunumanii* plantation forests

Jaco-Pierre van der Merwe[a,b], Gerard Lindner[b,c,d,g], Yolandi Ernst[e], Ilaria Germishuizen[f], Elane van Heerden[c], Hugo Zandberg[c], Gabigabi Sibiya[c], Ndamulelo Raboho[c], Lizette De Waal[b,g], Thabani Ndlovu[b,g], Leonard Mabaso[b,g], Niel Zondach[b,g], Nanette Christie[h], Charlie Clarke[i], Shawn D. Mansfield[a,j] *

[a] Department of Wood Science, University of British Columbia, Vancouver, BC, Canada, V6T 1Z4, Canada.
[b] Research Division, York Timbers, Sabie, Mpumalanga, South Africa.
[c] Planning Division, York Timbers, Sabie, Mpumalanga
[d] Stellenbosch University, Department of Forestry, Stellenbosch, Western Cape, South Africa.
[e] School of Animal, Plant and Environmental Sciences, Faculty of Science, University of the Witwatersrand, Johannesburg, Gauteng, South Africa.
[f] Institute for Commercial Forestry Research, Pietermaritzburg, KZN, South Africa.
[g] Tree Breeding Division, York Timbers, Sabie, Mpumalanga, South Africa.
[h] Department Biochemistry, Genetics and Microbiology, University of Pretoria, Pretoria, Gauteng, South Africa.
[i] CharlieC Consulting, Cheltenham, Gloucestershire, GL54 4NU, United Kingdom.
[j] Department of Botany, University of British Columbia, Vancouver, BC, Canada, V6T 1Z4, Canada.

* Corresponding author, email: shawn.mansfield@ubc.ca

Draft

Abstract

Plantation forests were introduced into South Africa to satisfy the regional demand for wood-based products, which are largely found in the Mpumalanga province. To better understand the impacts of climate on the dominant height growth of commercially important plantation species, enumeration data collected from 2012 to 2023 was scrutinised, representing 34,740 plots in 2,082 plantation compartments. Statistically significant multiple linear regression models were developed to predict dominant heights for *Pinus elliottii*, *P. taeda*, *P. patula* and *P. patula x P. tecunumanii* hybrids using climate variables as independent factors. Mean annual maximum temperature was statistically significant when modelling dominant height for *P. elliottii,* while winter maximum temperature was significant for *P. taeda*, *P. patula,* and *P. patula x P. tecunumanii*. Rainfall was found to be significant for *P. elliottii* and *P. taeda*, while spring rainfall was found to be more important when modelling dominant height

© The Author(s) or their Institution(s)

for *P. patula*. Interestingly autumn rainfall was found to have a negative impact on dominant height growth of *P. taeda* and *P. patula*, while rainfall was not significant in *P. patula x P. tecunumanii*. The findings imply that the southern pines have water thresholds, while Mexican pines are more temperature limited than water limited.

Key words: Climate, Tree height, *Pinus elliottii*, *Pinus taeda*, *Pinus patula*, *Pinus tecunumanii*

Introduction

South Africa is a water scarce country with limited native forests for wood production (Van Wilgen and Richardson, 2012). Cumulatively, *P. elliottii*, *P. taeda,* and *P. patula*, represent 82% of the planted pine plantation forests in South Africa (Forest Economic Services CC. 2020). More recently, large areas have been established with *P. patula x P. tecunumanii* hybrids, due to their fast growth and natural resistance to the pathogen *Fusarium circinatum*, which causes large scale mortality in new plantings of *P. patula* (Hodge and Dvorak 2007; Kanzler et al. 2014; Mitchell et al. 2012).

Pine species represent 49% of the plantation forests in South Africa, of which the majority are located in the Mpumalanga province (Forest Economic Services CC. 2020; Van der Merwe et al. 2023a). The Mpumalanga forestry region is climatically diverse and can be broadly divided into the Lowveld and Highveld forestry regions. The Lowveld has irregular topography, is fractioned by the Drakensberg Mountain range, and receives more rainfall annually due to its closer proximity to the eastern seaboard (Muller et al. 2017; Van der Merwe et al. 2023c). The Highveld region has more uniform terrain conditions, little or no mountains, and comparatively receives less rainfall (Van der Merwe et al. 2023c).

Draft

© The Author(s) or their Institution(s)

Tree height growth occurs when resources required to maintain tree roots, stem, and leaves are satisfied, and is therefore limited to the available photosynthate after these primary processes have achieved energy equilibrium (Gonzalez-Benecke et al. 2014; Ryan et al. 2006; Ryan and Yoder, 1997, West et al. 1999). Therefore, both temperature and water availability can limit tree growth (Hacker and Bilan 1992; Koirala et al. 2021; Niklas and Spatz 2004; Ryan 2010; Silva et al. 2022; White et al. 2022). Climate is known to impact tree growth and therefore introduced species in South Africa are typically planted in areas with similar conditions to their native habitat to ensure optimal survival and growth (Louw and Scholes 2006; Nedlo et al. 2009; Manson et al. 2013). Deployment of tree species to specific geo-climatic environments is known as site species matching (Peltier and Ogle 2020; Bernardo Pirovani et al. 2018; Schönau and Schulze, 1984). Therefore, forest growers have characterised the inherent growing conditions of plantation forests by means of climate modelling and soil surveys, to ensure the successful deployment of plantation forests and accurate prediction of future timber stock manufacturing (Koirala et al. 2021; Louw et al. 2011; Wang et al. 2007).

Climate change may however impact the local climate of the forestry regions. South Africa is deemed a “climate change hotspot” (Betts et al., 2018; Engelbrecht et al., 2015; Engelbrecht and Monteiro, 2021; Hoegh-Guldberg et al., 2018), with forecasts predicting a faster rate of heating than global averages. Together with expected increases in temperatures, regional downscaled models show generally drier conditions in the medium- and long-term future of South Africa (Engelbrecht et al. 2015; Engelbrecht and Monteiro 2021). In addition, it is predicted that rainfall variability will increase and extreme events such as droughts and floods will become more frequent and more intense (Engelbrecht et al. 2015). These changing climatic conditions could therefore have significant negative effects on plantation forestry production.

Draft

© The Author(s) or their Institution(s)

Plantation site productivity is broadly characterised at a compartment level by a site index (SI), which is defined as the dominant height at a predefined age (Louw and Scholes 2006; Tarmu et al. 2020; Wang et al. 2007). Plantation forests are enumerated regularly over the course of their growing cycle to predict future volumes of timber stock (Santiago-García et al. 2017; Santiago-García et al. 2020). This study aims to model dominant height for *P. elliottii*, *P. taeda*, *P. patula*, and *P. patula x P. tecunumanii* trees in South Africa, using recently developed climate models (Van der Merwe et al. 2023c) and historic enumeration data. Furthermore, site index at a plantation age of 20 years was projected for the Lowveld and Highveld forestry regions, respectively. The study investigated if climatic variables impact the growth of plantation forests in South Africa and how climate and height measurements can be used to improve site by species matching (Bauman et al. 2021; Leibing et al. 2009; Notivol et al. 2020).

Draft

Materials and methods

Study area, sampling strategy and measurement data

The study was conducted in the Mpumalanga forestry region of South Africa, which is grouped according to the Lowveld and Highveld sub-regions (Figure 1). The Lowveld is mountainous, has irregular terrain, and receives up to 1,500 mm of rainfall annually (Van der Merwe et al. 2023c; Van der Merwe et al. 2024). The Highveld has relatively regular terrain, is generally situated at a higher altitude, and receives less than 1,000 mm of rainfall annually (Van der Merwe et al. 2023c).

A total of 2,082 plantation compartments were enumerated, spanning ages from 3 to 25 years over the period from 2012 to 2023 (Figure 1, Table 1, and Supplemental Table S1). The enumerated plots were circular, covering an area of 500 m² per plot, and were replicated within each even-aged compartment until 6-7% of the total compartment surface area was

© The Author(s) or their Institution(s)

enumerated. Enumeration points were selected according to a grid system, dividing the plantation area into evenly sized polygons and determining the centroids of each polygon using ArcMap software (ESRI 2021).

Species and enumeration age of the corresponding trees were obtained from historic plantation records. The *P. patula x P tecunumanii* hybrids were established with a mixture of *P. tecunumanii* pollen that included both high and low altitude ecotypes, as was common practice during the time of commercial deployment of the hybrid pines. Enumeration measurements included dominant height, diameter at breast height (DBH), and tree count per hectare, however only dominant height measurements were assessed in this study. Centroid coordinates were determined for each enumerated compartment using QGIS software Version 3.24.1. Seasonal annual climate, including T-max (mean maximum temperature), T-min (mean minimum temperature), P-median (median rainfall) was obtained for each centroid position using existing climate models for the region (Van der Merwe et al. 2023c). The monthly demarcations of the respective seasons can be seen in Supplemental Table S2 (Van der Merwe et al. 2023c). Compartment stocking ranged between 1,111 and 1,333 trees per hectare. Over the course of the plantation rotation, compartments were systematically thinned to a mean stocking density of 800 trees per ha followed by a second thinning to a target planting of 500 trees per ha, while maintaining a mean target basal area of 24 m² per ha, as the trees matured. Pruning treatments included three stages of branch pruning; a first pruning to a height of 1.5 m, a second pruning to a height of 3.5 m, and a third pruning to a height of 5.5 m. Branch pruning was typically initiated according to tree size with the aim to maximize clear wood formation by restricting the knotty core of the pruned tree sections to less than 12 cm. Hence, pruning was initiated according to the growth rate of trees.

Draft

© The Author(s) or their Institution(s)

Data processing

All statistical analyses and data processing were completed using R version 3.4.1: A language and environment for statistical computing. Identification of significant climate variables for developing multiple linear regression models to predict dominant height was accomplished via stepwise regression, which also scrutinised the respective models by using the Bayesian information criterion (BIC) ['leaps'] (Lumley 2020). Only variables with significant probabilities ($\alpha = 0.05$), while maintaining a significant model intercept, were included in the final models. Resultant multiple linear regression models were developed for the respective species ['lm'] (R Core Team 2023). Model residuals were tested for normality ['qqPlot'] (Almeida et al. 2018). Model accuracies was determined by means of root mean square error (RMSE) and the resultant R-squared values. Landscape projections were done for dominant height at age 20 years, by harnessing existing climate models and the newly developed dominant height models from this study ['raster'] (Hijmans 2023; Van der Merwe et al. 2023c). The newly developed models were used to project dominant height of the respective sample plots to a normalised age of 20 years. The amount of theoretical annual growing days and growth temperature in a calendar year for the respective enumerated sites were derived by using the predicted climate and existing models (Louw et al. 2011). Growing days (GD) can be defined as the number of days with sufficient water available to facilitate evapotranspiration, while growth temperature (GT) refers to the mean monthly temperature, weighted according to the monthly growing days (Louw et al. 2011; Van der Merwe et al. 2023a).

Four principle component analyses (PCA) were completed to determine the key variables and relationships between variables impacting dominant height growth of the species at a normalised tree age of 20 years. Only climate variables that were found to

Draft

© The Author(s) or their Institution(s)

155 significantly impact height growth were included in the model [‘lfda’, ‘ggfortify’,
156 ‘rcompanion’] (Mangiafico 2024; Tang et al. 2016; Tang and Li 2019; Wei and Simko 2021).

157 A Pearson’s correlation coefficient matrix was calculated to establish the correlation
158 between annual climatic variables [‘corrplot’](Wei and Simko 2021). In addition, threshold
159 models were developed using Cate-Nelson analysis, incorporating annual maximum
160 temperature (T-max) and the projected dominant height at age 20 years [‘cateNelson’] (Cate
161 and Nelson 1971; Mangiafico 2013). Cate-Nelson analysis determined the threshold annual
162 maximum temperature and rainfall at which the respective species achieve optimal growth.
163 This analysis determined critical-x (climate variable) and critical-y (dominant height) values
164 by repetitively dividing the data into two groups and comparing the sum of squares of the
165 respective iterations. The critical x and y values were determined by maximising the amount
166 of data points within the second and fourth quadrants of an x-y plot in the case of a positive
167 trend, or first and third quadrants in the case of a negative trend (Cate and Nelson 1971; Lee
168 2016). Model accuracy was quantified using the chi-square test, also known as the Cramer’s
169 V statistic (Lee 2016).

170 Results and Discussion

171 Climate is widely recognised to be one of the most significant variables impacting the growth
172 of plantation-grown pine trees (Bauman et al. 2021; Nava-Nava et al. 2022; van der Merwe et
173 al. 2023a). Typically, tree height growth is significantly impacted by the most limiting
174 resource affecting photosynthesis, as tree growth is restricted when the atmospheric demand
175 for moisture or potential evaporation exceeds water availability (Belay 2016; Downes and
176 Drew 2008; Dye, 2001; Louw and Scholes 2006; Urban et al. 2017).

Draft

© The Author(s) or their Institution(s)

Seasonal climate impacts on dominant height growth

The study aimed to develop dominant height models for plantation grown *P. elliottii*, *P. taeda*, *P. patula*, and *P. patula x P. tecunumanii*, while including tree age and climate variables as independent variables. Multiple linear regression models were developed by identifying significant climatic variables impacting tree height growth by means of stepwise regression. Independent variables investigated included tree age at enumeration, together with annual maximum temperature (T-max), annual minimum temperature (T-min), seasonal maximum, and minimum temperatures for summer, autumn, winter and spring, respectively. Furthermore, seasonal precipitation and mean annual precipitation (MAP) were included for the respective sample sites. However, no precipitation values were included for winter, as the study area is located in a summer rainfall region, with little or no precipitation during the winter months (Van der Merwe et al. 2023c). The final models with variable probabilities and model R-square values for the respective species are summarised in Table 2. Furthermore, actual versus predicted plots are presented in Figure 2. Tree age was found to be the most important variable predicting tree height, with varying importance of climatic conditions per species, which further highlights the importance of genotype by environment (G x E) interaction on tree growth (Brawner et al. 2014; Zhai et al. 2015). The specific outcomes for each species examined are describe independently. Historic height measurements were archived per compartment *i.e.,* the number of sample plots and sample area was specified with a single dominant height value and not for the respective sample plots per compartment. Therefore, random effects were not included in the final multiple linear regression models.

Pinus elliottii

Dominant height (Ht_d) growth of *P. elliottii* was significantly impacted by annual maximum temperature and seasonal rainfall during the summer and spring months (Table 2, Figure 2A, and Supplemental Figure S1A). Trees were taller on sites with higher annual maximum

Draft

© The Author(s) or their Institution(s)

temperatures (Table 2). In addition, Ht_d was also larger on sites that received more summer and spring rainfall, allowing for more available water during the active growing season (Foster and Brooks 2001; Wang et al. 2007).

Previous international studies found height growth of southern pines, which included *P. elliottii* trees, to be positively correlated with annual temperature and seasonal rainfall (Harley et al. 2012; Huiqing et al. 2008; Loehle 1998; Teskey et al. 1994). In fact, *P. elliottii* has been found to be physiologically active throughout the year at sites with sufficient solar radiation and temperature (Martin 2000). Height growth of *P. elliottii* has been found to be negatively impacted by dry growing conditions, due to conservative water use efficiency (Ford and Brooks 2003; Foster and Brooks 2001; Samuelson et al. 2012). Of the species investigated, *P. elliottii* was one of the species significantly impacted by seasonal rainfall, which illustrates the importance of water availability to ensure optimal growth for the species.

Pinus taeda

Dominant height growth of *P. taeda* was significantly impacted by winter maximum temperatures, mean annual precipitation (MAP), and autumn rainfall (Table 2, Figure 2B, and Supplemental Figure S1B). Higher winter maximum temperatures and a higher annual rainfall were found to have a positive impact on dominant height growth, potentially due to an extended growing season, facilitated by higher temperatures (Martin 1999; Urban et al. 2017). Overall, annual precipitation was found to be positively correlated with height growth (Table 2). However, autumn rainfall was found to have a negative impact on height growth, possibly due to more cloud cover days and limited solar radiation on sites receiving more rainfall in autumn, during the seasonal transition period (Alvarado-Barrientos et al. 2014; Tang et al. 2003).

Draft

© The Author(s) or their Institution(s)

Similar to previous studies, height growth of *P. taeda* was significantly impacted by climate (Teodoro Zamin et al. 2013). Temperature and more specifically sufficient temperature during the winter season have been found to significantly impact height growth, potentially due to a longer growing season (Downs and Borthwick 1956; Loehle 1998; Martin 1999; Teodoro Zamin et al. 2013; Will et al. 2001). Dominant height growth of *P. taeda* was also significantly impacted and had a positive relationship with mean annual rainfall (MAP). These findings could be associated with lower water use efficiency as previously seen in southern pine species (Sameulson et al. 2012; Teodoro Zamin et al. 2013; Tang et al. 2003). The importance of sufficient soil water to maintain a positive stomatal water deficit in *P. taeda* has been reported (Ewers et al. 2001; Teskey et al. 2015). In fact, growth of *P. taeda* has been reported to stop, when grown in soils with moisture content below 40% (Dougherty et al. 1994).

Pinus patula

Winter maximum temperatures, and spring and autumn rainfall were found to be the most influential climatic variables, when predicting Ht_d for *P. patula* (Table 2, Figure 2C, and Supplemental Figure S1C). In particular, sites with higher winter maximum temperatures and more spring rainfall were found to have a positive impact on height growth, potentially due to an early onset of the growing season, while higher winter maximum temperatures may allow for an extended late growing season (Downs and Borthwick 1956) (Table 2). However, higher autumn rainfall had a negative impact on height growth, potentially due to radiation limitations associated with cloud cover during the autumn season (Alvarado-Barrientos et al. 2014) (Table 2). Thus, seasonal timing, rather than overall rainfall seems to have a more significant impact on the height growth of *P. patula*. Spring rainfall could lead to the onset of an early growing season, while autumn rainfall reduces radiation due to more frequent cloud cover (Van der Merwe et al. 2023a).

Draft

© The Author(s) or their Institution(s)

Previous studies have determined temperature to be the most influential climatic variable on height growth of *P. patula*, with larger trees being found on warmer sites displaying a negative relationship between temperature and rainfall (Nava-Nava et al. 2022; van der Merwe et al. 2023a). Similar to the current study, spring and summer rainfall had a positive relationship with the height growth, while autumn rainfall had a negative relationship with height growth of *P. patula* (Van der Merwe et al. 2023a). For example, higher altitude sites found on the Highveld received less rainfall and were found to be warmer than Lowveld sites that received more rainfall, located at similar altitude (Van der Merwe et al. 2023a).

Pinus patula x P. tecunumanii

Winter maximum temperature was the single most important variable impacting height growth in the hybrid, and sites with higher winter maximum temperatures had larger trees; none of the rainfall variables were observed to impact height growth (Table 2, Figure 2D, and Supplemental Figure S1D). These results indicate that *P. patula x P. tecunumanii* grown in the Mpumalanga forestry region is not restricted by water availability. In addition, the findings inherently imply a better water use efficiency, as no regional water limitations impact the Ht_d model. However, enumerations for the hybrid were mostly limited to the Lowveld forestry region, with only two sites enumerated in the drier Highveld region (Van der Merwe et al. 2023c). Plantings of *P. patula x P. tecunumanii* have been limited in the Highveld region, due to frost related mortality in the region.

Research on the environmental factors impacting height growth of the newly developed *P. patula x P. tecunumanii* hybrid has been limited. Previous studies have demonstrated that hybrid species, such as *P. patula x P. tecunumanii*, outperform pure pine species due to hybrid vigour (Nilsson et al. 2020). Additionally, studies conducted in Brazil, Columbia, Mexico, and South Africa have shown a positive correlation between height growth and temperature (Kanzler et al. 2012; Leibing et al. 2009; Nava-Nava et al. 2020; Van

Draft

© The Author(s) or their Institution(s)

der Merwe et al. 2023a). The results from the current study provide further insights into the growth dynamics of this hybrid, revealing that it is not limited by water availability, but rather by temperature, especially during winter months.

Annual climate impacts on dominant height growth

Dominant height measurement projections were carried out by employing local terrain features to determine climate conditions for the respective enumerated compartments, using recently developed climate models (Van der Merwe et al. 2023c)(Supplemental Figure S2).

Dominant height at age 20 was modelled for *P. elliottii*, *P. taeda*, *P. patula,* and *P. patula x P. tecunumanii* and projected for the Lowveld and Highveld landscape, according to the underlying climate conditions (Table 2, Supplemental Figure S4 and S5). A summary of the height distribution of the respective species according to the respective Lowveld and Highveld forestry regions are shown in Supplemental Table S3.

Draft

Furthermore, dominant height was predicted at age 20 years for all of enumerated sample plots using the newly developed dominant height models (Supplemental Figure S4 and S5). Dominant heights of the enumerated trees at a normalised tree age, allowed for the development of climate-dominant height threshold models. Three climate-independent variables were investigated, including annual maximum temperature (T-max), annual minimum temperature (T-min), and mean annual precipitation (MAP). Model significance was determined according to the chi-square-based Cramer's V statistic (Table 3). Model thresholds are visualised in Supplemental Figures S6, S7, and S8.

Pinus elliottii

T-max had a relatively strong positive relationship with height growth of *P. elliottii,* and 75 % of the observations were accounted for within the threshold quadrants (Cramer's V = 0.48, P-model = 0.75) (Table 3). Critical threshold T-max for dominant height growth was

© The Author(s) or their Institution(s)

300 determined to be 22.3°C and a threshold Ht_d of 24.4 meters was observed. The results
301 indicate that *P. elliottii* achieved superior growth above temperatures of 22.3°C and the
302 threshold height at this T-max is 24.4 meters. The P-model value indicates that 75% of the
303 measurements are explained between quadrants two and four (positive trend; Figure 3).

304 T-min was found to have strong positive relationship with the height growth of *P.*
305 *elliottii* and 89 % of the observations were accounted for within the threshold quadrants
306 (Cramer's V = 0.61, P-model = 0.89) (Table 3). Critical threshold T-min for dominant height
307 growth was determined to be 11.1°C and a threshold Ht_d of 23.2 meters was observed.
308 However, MAP was found to be the most influential climatic variable impacting Ht_d of *P.*
309 *elliottii*. A very strong positive relationship with height growth of *P. elliottii* was observed
310 and 96% of the observations were accounted for within the threshold quadrants (Cramer's V
311 = 0.87, P-model = 0.96) (Table 3). Critical threshold MAP for dominant height growth was
312 determined to be 828.8 mm and a threshold Ht_d of 23.5 meters was observed.

313 Recent studies have indicated that *P. elliottii* possesses the necessary physiological
314 and metabolic mechanisms to cope with warmer temperatures and drought conditions
315 (Celeste Dias et al. 2022). However, findings from this study suggest that *P. elliottii* grown in
316 Mpumalanga is most sensitive to MAP among the species investigated in this study (Table 3,
317 Supplemental Figure S8). Furthermore, the relatively high temperature thresholds and
318 positive relationship between Ht_d and temperature suggest that the species will benefit from
319 future warming climates, provided sufficient water is available for tree growth (Table 3,
320 Supplemental Figures S6, S7, and S8). The results suggest that *P. elliottii* height growth is
321 more sensitive and positively related to growth days (GD) than growth temperature (GT), as
322 precipitation and T-min were generally found to be positively correlated with GD and weakly
323 related to GT (Supplemental Figure S3) (Van der Merwe et al. 2023a).

Draft

© The Author(s) or their Institution(s)

Pinus taeda

T-max also had a relatively strong positive relationship with height growth of *P. taeda* and 88 % of the observations were accounted for within the threshold quadrants (Cramer's V = 0.46, P-model = 0.88) (Table 3). Critical threshold T-max for dominant height growth was determined to be 23.3°C and a threshold Ht_d of 24.2 meters was observed. T-min was found to have strong positive relationship with the height growth of *P. taeda* and 76% of the observations were accounted for within the threshold quadrants (Cramer's V = 0.57, P-model = 0.76) (Table 3). Critical threshold T-min for dominant height growth was determined to be 11.2°C and a threshold Ht_d of 23.1 meters was observed. Similar to the results for *P. elliottii*, MAP was found to be the most influential climatic variable impacting Ht_d of *P. taeda*. A strong positive relationship between MAP and height growth of *P. taeda* was observed, and 84% of the observations were accounted for within the threshold quadrants (Cramer's V = 0.87, P-model = 0.96) (Table 3). The critical threshold for MAP influencing dominant height growth was identified as 916.7 mm, marking the highest rainfall threshold value among the species investigated. A height threshold (Ht_d) of 23.2 meters was observed.

The results indicate that the growth of *P. taeda* plantations in the region, is fundamentally limited by rainfall and overall water availability (Table 3). The findings are supported by previous studies that reported stunted or halted growth of *P. taeda* under water-limiting conditions (Dougherty et al. 1994; Ewers et al. 2001; Teskey et al. 2015). However, a positive relationship between both T-max and T-min was also observed (Table 3). In fact, positive temperature responses for both T-min and T-max were found to be at the highest temperature thresholds of all the species investigated (T-max: 23.3 °C; T-min: 11.2 °C). The findings suggest that *P. taeda*, has high temperature-growth thresholds, while a positive soil water deficit is maintained. Therefore, water availability is essential for the species to maintain a positive growth trend, at higher temperatures. The findings indicated that

Draft

© The Author(s) or their Institution(s)

dominant height growth in southern pine species, including *P. taeda*, is more constrained and positively correlated with rainfall and associated growth days (GD) than with growth temperature (GT). This is attributed to the positive relationships observed between rainfall, T-min, and GD (Supplemental Figure S3).

Pinus patula

Temperature, specifically T-max, was identified to be the most influential climate variable affecting height growth of *P. patula*. The T-max threshold model showed a strong positive relationship between T-max and height growth of *P. patula*, with 92% of observations falling within the threshold quadrants (Cramer's V = 0.73, P-model = 0.92) (Table 3). The critical threshold for T-max was determined to be 20.4°C at a threshold (Ht_d) of 26.9 meters. Conversely, T-min exhibited a strong negative relationship with height growth of *P. patula*, with 84 % of observations within the threshold quadrants (Cramer's V = 0.67, P-model = 0.84), and a critical threshold of 11.6°C and Ht_d of 27.6 meters. Similarly, MAP displayed a strong negative relationship with 84% of observations within the threshold quadrants (Cramer's V = 0.69, P-model = 0.84), with a critical threshold of 868.0 mm and Ht_d of 27.7 meters observed for dominant height growth of *P. patula* (Table 3). It however remains unclear if the negative relationship between Ht_d, T-min and MAP was more related to temperature or rainfall, due to a strong colinear relationship between these two independent variables.

The results indicated that *P. patula* is most limited by T-max. *P. patula* grew tallest on sites with higher T-max and the results are substantiated by previous studies (Nava-Nava et al. 2022; Van der Merwe et al. 2023a; Van der Merwe et al. 2023b; Van der Merwe et al. 2024). In addition, *P. patula* was the most responsive to T-max and had the lowest T-max threshold value of the investigated species. A negative relationship was observed between

Draft

© The Author(s) or their Institution(s)

373 Ht_d and T-min. The Pearson's correlation coefficient matrix revealed a negative relationship
374 between T-min and sites with higher rainfall, a finding supported by previous studies (Van
375 der Merwe et al. 2023a) (Supplemental Figure S3). Similar to previous studies, *P. patula* was
376 found to be moderately drought tolerant and in fact more drought tolerant than *P. elliottii* and
377 *P. taeda*, which further supported the observations in this study (Sudhakara Reddy and
378 Natarajan 1996). The results indicate that *P. patula*, grown in the Mpumalanga forestry
379 region, currently faces limitations not in water supply, but in maximum temperature. The
380 findings imply that *P. patula*'s height growth is more responsive to growth temperature (GT)
381 than growth days (GD), since T-max was found to be positively correlated with GT and
382 weakly correlation with GD (Supplemental Figure S3).

383 ***Pinus patula x Pinus tecunumanii***

384 T-max was found to be the most significant climatic variable impacting height growth of *P.*
385 *patula x P. tecunumanii*. A very strong positive relationship was observed between T-max
386 and height growth of *P. patula x P. tecunumanii*, and 100% of the observations were
387 accounted for within the threshold quadrants (Cramer's V = 1.00, P-model = 1.00) (Table 3).
388 Critical threshold T-max for dominant height growth was determined to be 21.6°C and a
389 threshold Ht_d of 31.0 meters was observed. T-min had a relatively strong negative
390 relationship with height growth of *P. patula x P. tecunumanii* and 81% of the observations
391 were accounted for within the threshold quadrants (Cramer's V = 0.58, P-model = 0.81)
392 (Table 3). Critical threshold T-min for dominant height growth was determined to be 12.0°C
393 and a threshold Ht_d of 31.4 meters was observed. MAP had a negligible influence on height
394 growth of *P. patula x P. tecunumanii*, although 71% of the observations were accounted for
395 within the threshold quadrants (Cramer's V = 0.04, P-model = 0.71) (Table 3).

Draft

© The Author(s) or their Institution(s)

The results indicate that *P. patula x P. tecunumanii* significantly benefits from warm-dry conditions, as growth was superior on warmer sites with greater T-max. In addition, sites with lower T-min had larger trees and it is not clear if this observation is related to a colinear relationship between rainfall and T-min. The findings are supported by previous studies, which found *P. tecunumannii* to be more drought tolerant compared to *P. elliottii*, *P. taeda*, and *P. patula* which could explain why the *P. patula x P. tecunumanii* performed better on warm-dry sites (Dvorak et al. 2000; Hodge and Dvorak 1999; Kanzler et al. 2012; Kanzler et al. 2014). Furthermore, no significant threshold models were developed for the relationship between MAP and Ht_d, which further substantiates the drought tolerance of this species. The results indicate that *P. patula x P. tecunumanii* plantations in the Mpumalanga province of South Africa have not encountered growth limitations, due to water availability. Of the species investigated, *P. patula x P. tecunumanii* was the fastest growing, least impacted by rainfall, and grew best in warm conditions, which makes this hybrid the most promising for future deployment considering the anticipated impacts of global warming (Kanzler et al. 2012; Kanzler et al. 2014) (Supplemental Figure S4 and S5). Although the *P. tecunumanii* pollen used for the development and deployment of this hybrid was not segregated according to low and high ecotypes, no trend deviation in climatic suitability was observed during analysis of the commercially deployed populations investigated in this study.

Conclusion

The study emphasises the importance of understanding the impact of climate on the growth of pine plantation forests. In addition, the study identified species-specific climate limitations, and provides valuable insights into the underlying climate drivers impacting successful deployment and growth of commercially important introduced pine plantation species in South Africa. Future studies should focus on modelling the future climate of the Mpumalanga

Draft

© The Author(s) or their Institution(s)

forestry region and assessing how climate changes may influence decision-making in the selecting of introduced pine species for site-species matching. In addition, geo-climatic limitations for tree growth including the interaction between soil water availability and climate should be further investigated and scrutinised.

Acknowledgements

We would like to thank all the York Timbers staff who assisted in the measurement of trees in the plantation forests over time. We would also like to thank the current Chief Executive Officer of York Timbers, Mr. Gerald Stoltz for allocating funding to this project and believing in the power of science.

Author Statement

Jaco-Pierre van der Merwe collected the data, completed the modelling, and wrote the first draft of the manuscript. Yolandi Ernst assisted in the climate literature review. Gerard Lindner, Lizette De Waal, Ilaria Germishuizen, Yolandi Ernst, Nanette Christie, Charlie Clarke, and Shawn Mansfield aided with project conceptualisation, and reviewed and edited the manuscript. Elane van Dijk extracted compartment centroids for the respective plantation compartments and developed the maps. Thabani Ndlovu, Leonard Mabaso, and Niel Zondach was involved in the tree breeding and commercial deployment and site species matching of the specific pine species and hybrids included in this study and gave valuable insights into the geo-climatic impacts on tree growth.

Data Availability

The data used in the manuscript is readily available upon request from the corresponding author.

Draft

© The Author(s) or their Institution(s)

Research Funding

Funding for this project was provided by York Timbers, South Africa, as well as sampling and the allocation of company resources.

Conflict of Interest Statement

The authors declare that they have no conflicts of interest regarding this article.

References

Almeida, A., Loy, A. and Hoffman, H. 2018. ggplot2: Compatible quantile-quantile plots in R. The R Journal. 10(2): 248-261. doi:10.32614/RJ-2018-051.

Alvarado-Barrientos, M.S., Holwerda, F., Asbjornsen, H., Dawson, T.E. and Bruijnzeel, L.A. 2014. Suppression of transpiration due to cloud immersion in a seasonally dry Mexican weeping pine plantation. Agric. For. Meteorol. 186: 12-25. doi:10.1016/j.agrformet.2013.11.002.

Bauman, D., Fortunel, C., Cernusak, L.A., Bentley, L.P., McMahon, S.M., Rifai, S.W., Aguirre-Gutiérrez, J., Oliveras, I., Bradford, M., Laurance, S.G.W., Delhaye, G., Hutchinson, M.F., Dempsey, R., McNellis, B.E., Santos-Andrade, P.E., Ninantay-Rivera, H.R., Paucar, J.R.C., Phillips, O.L. and Malhi, Y. 2021. Tropical tree growth sensitivity to climate is driven by species intrinsic growth rate and leaf traits. Glob. Chang. Biol. 28(1): 1-19. doi:10.1111/gcb.15982.

Belay, T.T. 2016. Climate-growth relationship of *Pinus patula Schldl. et Cham.* in Wondo Genet, South Central Ethiopia. J. Climatol. Wx. Fcst. 4(3): 1-8. doi:10.4172/23322594.1000181.

Bernardo Pirovani, D., Macedo Pezzopane, J.E., Cândido Xavier, A., Macedo Pezzopane, J.R., de Jesus Júnior, W.C., Herrera Machuca, M.A., Alves dos Santos, G.M.A.D., Ferreira da Silva, S., Hatum de Almeida, S.L., De Oliveira Peluzio, T.M., Coelho Eugenio, F., Rizzo Moreira, T., Sobreira Alexandre, R. and Rosa dos Santos, A. 2018. Climate change impacts on the aptitude area of forest species. Ecol. Indic. 95(1): 405-416. doi:10.1016/j.ecolind.2018.08.002.

Draft

© The Author(s) or their Institution(s)

Brawner, J.T., Hodge, G.R., Meder, R. and Dvorak, W.S. 2014. Visualising the environmental preferences of *Pinus tecunumanii* populations. Tree Genet. Genomes. 10: 1123-1133. doi: 10.1007/s11295-014-0747-8.

Betts, R. A., Alfieri, L., Bradshaw, C., Caesar, J., Feyen, L., Friedlingstein, P., Gohar, L., Koutroulis, A., Lewis, K., Morfopoulos, C., Papadimitriou, L., Richardson, K. J., Tsanis, I. and Wyser, K. 2018. Changes in climate extremes, fresh water availability and vulnerability to food insecurity projected at 1.5°C and 2°C global warming with a higher-resolution global climate model. Phil. Trans. R. Soc. A., 376, 20160452.20160452. doi:10.1098/rsta.2016.0452.

Cate, R.B. and Nelson, L.A. 1971. A Simple Statistical Procedure for Partitioning Soil Test Correlation Data Into Two Classes. Soil. Sci. Soc. Am. J. 35: 658-660.

Celeste Dias, M., Ferreira de Oliveira, J.M.P., Marum, L., Pereira, V., Almeida, T., Nunes, S., Araújo, M., Moutinho-Pereira, J., Correia, C.M. and Santos, C. 2022. *Pinus elliottii* and *P. elliottii x P. caribaea* hybrid differently cope with combined drought and heat episodes. Ind. Crops. Prod. 176: 114428. doi:10.1016/j.indcrop.2021.114428.

Dougherty, P.M., Whitehead, D. and Vose, J.M. 1994. Environmental influences on the phenology of pine. Ecol. Bull. 43: 64-75.

Downes, G.M. and Drew, D.M. 2008. Climate and growth influences on wood formation and utilisation. South. For. 70(2): 155–167. doi:10.2989/SOUTH.FOR.2008.70.2.11.539.

Downs, R.J. and Borthwick, H.A. 1956. Effects of photoperiod on growth of trees. Bot. Gaz. 117(4): 310-326. doi:10.1086/335918.

Dvorak, W.S., Hodge, G.R., Gutiérrez, E.A., Osorio, L.F., Malan, F.S. and Stanger, T.K. 2000. *Pinus tecunumanii*. In: Conservation and Testing of Tropical and Subtropical Forest Species by the CAMCORE Cooperative. College of Natural Resources, NCSU. Raleigh, NC. USA. pp: 188-209.

Dye, P.J. 2001. Modelling growth and water use in four *Pinus patula* stands with the 3-PG model. South. Afr. For. J. 191(1): 53-63. doi:10.1080/20702620.2001.10434151.

Engelbrecht, F. A., Adegoke, J., Bopape, M.-J., Naidoo, M., Garland, R., Thatcher, M., McGregor, J., Katzfey, J., Werner, M., Ichoku, C. and Gatebe, C. 2015. Projections of rapidly

Draft

© The Author(s) or their Institution(s)

498 rising surface temperatures over Africa under low mitigation. Environ. Res. Lett. 10(8),
499 085004. doi:10.1088/1748-9326/10/8/085004.

500 Engelbrecht, F. A. and Monteiro, P. 2021. The IPCC assessment report six working group 1
501 report and southern Africa: Reasons to take action. S. Afr. J. Sci. 117(11–12), 1–7.
502 doi:10.17159/sajs.2021/1267.

503 ESRI. 2021. ArcGIS Desktop: Release 10.8.2 Redlands, CA: Environmental Systems
504 Research Institute.

505 Ewers, B.E., Oren, R., Johnsen, K.H. and Landsberg, J.J. 2001. Estimating maximum mean
506 canopy stomatal conductance for use in models. Can. J. For. Res. 31(2): 198-207.
507 doi:10.1139/x00-159.

508 Ford, C.R. and Brooks, J.R. 2003. Hydrological and climatic responses of *Pinus elliottii* var.
509 densa in mesic pine flatwoods Florida, USA. Ann. For. Sci. 60: 385-392.
510 doi:10.1051/forest:2003030.

511 Fortin, M., Van Gouwenberghe, R., Perez, V. and Piedallu, C. 2019. Evidence of climate
512 effects on the height-diameter relationships of tree species. Ann. For. Sci. 76, 1.
513 doi:10.1007/s13595-018-0784-9.

514 Foster, T.E. and Brooks, J.R. 2001. Long-term trends in the growth of *Pinus palustris* and
515 *Pinus elliottii* along a hydrological gradient in central Florida. Can. J. For. Res. 31: 1661-
516 1670. doi:10.1139/x01-100.

517 Gauli, A., Neupane, P.R., Mundhenk, P. and Köhl, M. 2022. Effect of climate change on the
518 growth of tree species: Dendroclimatological analysis. Forests. 13(4): 496.
519 doi:10.3390/f13040496.

520 Gonzalez-Benecke, C.A., Jokela, E.J., Cropper Jr., W.P., Bracho, R. and Leduc, D.J. 2014.
521 Parameterization of the 3-PG model for *Pinus elliottii* stands using alternative methods to
522 estimate fertility rating, biomass partitioning and canopy closure. For. Ecol. Manag. 327: 55-
523 75. doi: 10.1016/j.foreco.2014.04.030.

524 Hacker, W.D. and Bilan, M.V. 1992. Site factors affecting growth of Loblolly pine in the post
525 oak belt. SJAF. 16: 197-200. doi:10.1093/sjaf/16.4.197.

Draft

© The Author(s) or their Institution(s)

Harley, G.L., Grissino-Mayer, H.D., Franklin, J.A., Anderson, C. and Köse, N. 2012. Cambial activity of *Pinus elliottii* var. densa reveals influence of seasonal insolation on growth dynamics in the Florida Keys. Trees. 26: 1449-1459. doi:10.1007/s00468-012-07192.

Hijmans, R. 2023. raster: Geographic data analysis and modelling, R package version 3.6-23, https://CRAN.R-project.org/package=raster.

Hodge G.R. and Dvorak W.S. 1999. Genetic parameters and provenance variation of *Pinus tecunumanii* in 78 international trials. For. Genet. 6: 157–180. doi:10.1139/x00-189.

Hodge, G.R. and Dvorak, W.S. 2007. Variation in pitch canker resistance among provenances of *Pinus patula* and *Pinus tecunumanii* from Mexico and Central America. New For. 33: 193-206. doi:10.1007/s11056-006-9023-6.

Hoegh-Guldberg, O., Jacob, D., Taylor, M., Bindi, M., Brown, S., Camilloni, I., Diedhiou, A., Djalante, R., Ebi, K., Engelbrecht, F. A., Guiot, J., Hijioka, Y., Mehrotra, S., Payne, A., Seneviratne, S. I., Thomas, A., Warren, R. and Zhou, G. 2018. Impacts of 1.5oC Global Warming on Natural and Human Systems. In V. Masson-Delmotte, P. Zhai, H.-O. Pörtner, D. Roberts, J. Skea, P. R. Shukla, A. Pirani, W. Moufouma-Okia, C. Péan, R. Pidcock, S. Connors, J. B. R. Matthews, Y. Chen, X. Zhou, M. I. Gomis, E. Lonnoy, T. Maycock, M. Tignor, & T. Waterfield (Eds.), (Global warming of 1·5°C. An IPCC Special Report on the impacts of global warming of 1·5°C above pre-industrial levels and related global greenhouse gas emission pathways, in the context of strengthening the global response to the threat of climate change. World Meteorological Organization.

Huiqing, Z., Qijing, L., Zongwei, F., Xiaoke, W. and Zeqing, M. 2008. Modeling the interannual variation and response to climate change scenarios in gross and net primary productivity of *Pinus elliottii* forest in subtropical China. Acta Ecol. Sin. 28(11): 5314-5321. doi:10.1016/S1872-2032(09)60008-1.

Kanzler, A., Nel, A. And Ford, C. 2014. Development and commercialisation of the *Pinus patula* × *P. tecunumanii* hybrid in response to the threat of *Fusarium circinatum*. New For. 45: 417-437. doi:10.1007/s11056-014-9412-1.

Kanzler, A., Payn, K. and Nel. A. 2012. Performance of two *Pinus patula* hybrids in southern Africa. South. For. 74(1): 19-25. doi:10.2989/20702620.2012.683639.

Draft

© The Author(s) or their Institution(s)

555 Koirala, A., Montes, C.R. and Bullock, B.P. 2021. Modeling dominant height using stand and
556 water balance variables for loblolly pine in the Western Gulf, US. For. Ecol. Manag. 479:
557 118610. doi:10.1016/j.foreco.2020.118610.

558 Lee, D.K. 2016. Alternatives to P value: confidence interval and effect size. Korean J
559 Anesthesiol. 69(6): 555-562. doi:10.4097/kjae.2016.69.6.555.

560 Leibing, C., Van Zonneveld, M., Jarvis, A. And Dvorak, W. 2009. Adaptation of tropical and
561 subtropical pine plantation forestry to climate change: Realignment of *Pinus patula* and *Pinus*
562 *tecunumanii* genotypes to 2020 planting site climates. Scand. J. For. Res. 24(6): 483-493.
563 doi:10.1080/02827580903378642.

564 Loehle, C. 1998. Height growth rate tradeoffs determine northern and southern range limits
565 for trees. J. Biogeogr. 25(4): 735-742. doi:10.1046/j.1365-2699.1998.2540735.x.

566 Louw, J.H., Germishuizen, I. and Smith, C.W. 2011. A stratification of the South African
567 forestry landscape based on climatic parameters. South. For. 73(1): 51-62.
568 doi:10.2989/20702620.2011.574825.

569 Louw, J.H. and Scholes, M. 2002. Forest site classification and evaluation: a South African
570 perspective. For. Ecol. Manag. 171(1-2): 153-168. doi:10.1016/S0378-1127(02)00469-3.

571 Louw, J.H. and Scholes, M.C. 2006. Site index functions using site descriptors for *Pinus*
572 *patula* plantations in South Africa. For. Ecol. Manag. 225(2006): 94-103.
573 doi:10.1016/j.foreco.2005.12.048.

574 Lumley, T. 2020. leaps: Regression subset selection. R package version 3.1, https://CRAN.R-
575 project.org/package=leaps. doi:10.2989/20702620.2012.741792.

576 Mangiafico, S.S. 2013. Cate-Nelson Analysis for Bivariate Data Using R-project. J. Ext.
577 51:5, 5TOT1.

578 Mangiafico, S.S. 2024. rcompanion: Functions to support extension education program
579 evaluation. version 2.4.35. Rutgers Cooperative Extension. New Brunswick, New Jersey.
580 https://CRAN.R-project.org/package=rcompanion.Manson, D.G., Schmidt, S., Bristow, M.,
581 Erskine, P.D. and Vanclay, J.K. 2013. Species-site matching in mixed species plantations of
582 native trees in tropical Australia. Agrofor. Syst. 87: 233-250. doi:10.1007/s10457-012-9538-
583 0.

Draft

© The Author(s) or their Institution(s)

584 Martin, T.A.M. 1999. Winter season tree sapflow and stand transpiration in an intensively-
585 managed loblolly and slash pine plantation. Frontier of Sustainable Forestry. 10(1-2): 155-
586 163. doi:10.1300/J091v10n01_18.

587 Mitchell, R.G., Coutinho, T.A., Steenkamp, E., Herbert, M. and Wingfield, M.J. 2012. Future
588 outlook for *Pinus patula* in South Africa in the presence of the pitch canker fungus (Fusarium
589 circinatum). South. For. 74(4): 203-210. doi:10.2989/20702620.2012.741792.

590 Muller, B.G., Louw, J.H. and Malan, F.S. 2017. Variation in selected solid wood properties
591 of young *Pinus patula* from diverse sites in the Mpumalanga escarpment area in South
592 Africa. South. For. 79(4), 317-327. doi:10.2989/00306525.2016.1255376.

593 Nava-Nava, A., Santiago-García, W., Quinonez-Barraza, G., De los Santos-Posadas, H.M.,
594 Valdez-Lazalde, J.R. and Ángeles-Pérez. 2022. Climatic and topographic variables improve
595 estimation accuracy of Patula pine forest site productivity in Southern Mexico. Forests. 13(8):
596 1277. doi:10.3390/f13081277.

597 Nedlo, J.E., Martin, T.A., Vose, J.M and Teskey, R.O. 2009. Growing season temperatures
598 limit growth of loblolly pine (*Pinus taeda* L.) seedlings across a wide geographic transect.
599 Trees. 23: 751-759. doi:10.1007/s00468-009-0317-0.

600 Niklas, K.J. and Spatz, H-C. 2004. Growth and hydraulic (not mechanical) constraints govern
601 the scaling of tree height and mass. PNAS. 101(44): 15661-15663.
602 doi:10.1073/pnas.0405857101.

603 Nilsson, O., Hodge, G.R., Frampton, L.J., Dvorak, W.S. and Bergh, J. 2020. Growth and
604 modulus of elasticity of pine species and hybrids three years after planting in South Africa.
605 South. For. 82(4): 367-376. doi:10.2989/20702620.2020.1733769.

606 Notivol, E., Santos-del-Blanco, L., Chambel, R., Climent, J. and Alía, R. 2020. Seed sourcing
607 strategies considering climate change forecasts: A practical test in Scots pine. Forests. 11(11):
608 1222. doi:10.3390/f11111222.

609 Peltier, D.M.P. and Ogle, K. 2020. Tree growth sensitivity to climate is temporally variable.
610 Ecol. Lett. 23(11): 1561-1572.

611 R Core Team. 2023. R: A language and environment for statistical computing. R Foundation
612 for Statistical Computing, Vienna, Austria. URL: https://www.R-project.org/.

Draft

© The Author(s) or their Institution(s)

Ryan, M.G. 2010. Temperature and tree growth. Tree Physiol. 30(6): 667-668. doi:10.1093/treephys/tpq033.

Ryan, M.G., Phillips, N. and Bond, B.J. 2006. The hydraulic limitation hypothesis revisited. Plant Cell Environ. 29(3): 367-381. doi:10.1111/j.1365-3040.2005.01478.x.

Ryan, M.G. and Yoder, B.J. 1997. Hydraulic limits to tree height and tree growth. BioScience. 47(4): 235-242. doi:10.2307/1313077.

Samuelson, L.J., Stokes, T.A. and Johnsen, K.H. 2012. Ecophysiological comparison of 50-year-old longleaf pine, slash pine and loblolly pine. For. Ecol. Manag. 274: 108-115. doi:10.1016/j.foreco.2012.02.017.

Santiago-García, W., Jacinto-Salinas, A.H., Rodríguez-Ortiz, G., Nava-Nava, A., Santiago-García, E., Ángeles-Pérez, G. and Enríquez-del Valle, J.R. 2020. Generalized height-diameter models for five pine species at Southern Mexico. For. Sci. Technol. 16(2): 49-55. doi:10.1080/21580103.2020.1746696.

Santiago-García, W., Pérez-López, E., Quiñonez-Barraza, G., Rodríguez-Ortiz, G., Santiago-García, E., Ruiz-Aquino, F. and Tamarit-Urias, J.C. 2017. A Dynamic System of Growth and Yield Equations for *Pinus patula*. Forests. 8(12): 465. doi:10.3390/f8120465.

Schönau, A.P.G. and Schulze, R.E. 1984. Climatic and altitudinal criteria for commercial afforestation with special reference to Natal. South. Afr. For. J. 130(1): 10-18. doi:10.1080/00382167.1984.9628947.

Silva, M.R.F., McHugh, I., Neto, A.M.L.P., Pauwels, V.R.N., Cartwright, I. and Daly, E. 2022. Trading a little water for substantial carbon gains during the first years of a Eucalyptus globulus plantation. Agric. For. Meteorol. 318(2022): 108910. doi:10.1016/j.agrformet.2022.108910.

Sudhakara Reddy, M. and Natarajan, K. 1996. In vitro ectomycorrhizal formation of *Pinus patula*, *P. pseudostrobus*, *P. oocarpa* and *P. elliotii* grown in southern India. New For. 11: 149-153. doi:10.1007/BF00033410.

Tang, Z., Chambers, J.L., Sword, M.A. and Barnett, J.P. 2003. Seasonal photosynthesis and water relations of juvenile loblolly pine relative to stand density and canopy position. Trees. 17: 424-430. doi:10.1007/s00468-003-0256-0.

Draft

© The Author(s) or their Institution(s)

642 Tang, Y., Horikoshi, M., and Li, W. 2016. ggfortify: Unified Interface to Visualize Statistical
643 Result of Popular R Packages." The R Journal 8(2): 478-489. doi:10.32614/RJ-2016-060

644 Tang, Y. and Li, W. 2019. lfda: Local Fisher Discriminant Analysis in R. J. open source
645 softw. 4(39): 1572. doi:10.21105/joss.01572.Tarmu, T., Laarmann, D. and Kiviste, A. 2020.
646 Mean height or dominant height – what to prefer for modelling the site index of Estonian
647 forests? For. Stud. 72(1): 121-138. doi:10.2478/fsmu-2020-0010.

648 Teodoro Zamin, N., Do Amaral Machado, S., Filho, A.F. and Soares Koehler, H. 2013. Effect
649 of Climate variables on monthly growth in 26odelling biological yield of Araucaria
650 angustifolia and *Pinus taeda* in the juvenile phase. Int. J. For. Res. 1: 1-8.
651 doi:10.1155/2013/646759.

652 Teskey, R.O., Gholz, H.L. and Cropper Jr, W.P. 1994. Influence of climate and fertilization
653 on net photosynthesis of mature slash pine. Tree Physiol. 14(11): 1215-1227.
654 doi:10.1093/treephys/14.11.1215.

655 Teskey, R., Wertin, T., Bauweraerts, I., Ameye, M., McGuire, M.A. and Steppe, K. 2015.
656 Responses of tree species to heat waves and extreme heat events. Plant Cell Environ. 38:
657 1699-1712. doi: 10.1111/ce.12417.

658 Urban, J., Ingwers, M., McGuire, M.A. and Teskey, R.O. 2017. Stomatal conductance
659 increases with rising temperature. Plant Signal. Behav. 12(8): e1356534.
660 doi:10.1080/15592324.2017.1356534.

661 Van der Merwe, J-P., Germishuizen, I., Clarke, C. and Mansfield, S.D. 2023a. The impact of
662 soil, altitude, and climate on tree form and wood properties of plantation grown *Pinus patula*
663 in Mpumalanga, South Africa. Holzforschung. 77(1): 1-15. doi:10.1515/hf-2022-0126.

664 Van der Merwe, J-P., Madiope, S., Spogter, O., Kuisis, H., Potgieter, J., Tait, O., Clarke, C.
665 and Mansfield, S.D. 2023b. The impact of site on tree form, heartwood content and veneer
666 production of plantation-grown *P. patula*. Holzforschung. 77(9): 670-687. doi:10.1515/hf-
667 2023-0031.

668 Van der Merwe, J-P., Wang, Clarke, C. And Mansfield, S.D. 2023c. Predicting temperature
669 and rainfall for plantation forestry in Mpumalanga, South Africa, using locally developed
670 climate models. Agric. For. Meteorol. 329: 109275. doi:10.1016/j.agrformet.2022.109275.

Draft

© The Author(s) or their Institution(s)

671 Van der Merwe, J-P., Bacher, M., Madiope, S., Ncongwane, T., Ngomane, R., Spogter, O.,
672 Kuisis H., Potgieter, J., Tait, O., Clarke, C., Cool, J., Avramidis, S., Ellis, S. and Mansfield.
673 S.D. 2024. The impact of site on tree form, wood properties, and lumber quality of
674 plantation-grown *Pinus patula*. Holzforschung. 78(1): 1-15. doi:10.1515/hf-2023-0075.

675 Van Wilgen, B.W. and Richardson, D.M. 2012. Three centuries of managing introduced
676 conifers in South Africa: Benefits, impacts, changing perceptions and conflict resolution. J.
677 Environ. Manage. 106: 56-68. doi:10.1016/j.jenvman.2012.03.052.

678 Wang, Q., Preda, M., Cox, M. and Bubb, K. 2007. Spatial model of site index based on γ-ray
679 spectrometry and a digital elevation model for two Pinus species in Tuan Toolara State
680 Forest, Queensland, Australia. Can. J. For. Res. 37(11): 2299-2312. doi:10.1139/X07-088.

681 Wei, T. and Simko, V. 2021. R package 'corrplot': Visualization of a correlation matrix
682 (Version 0.92).

683 West, G.B., Brown, J. and Enquist, B.J. 1999. A general model for the structure and
684 allometry of plant vascular systems. Nature. 400(6745): 664-667. doi:10.1038/23251.

685 White, D.A., Ren, S., Mendham, D.S., Balocchi-Contreras, F., Silberstein, R.P., Meason, D.,
686 Iroumé, A. and de Arellano, P.R. 2022. Is the reputation of Eucalyptus plantations for using
687 more water than Pinus plantations justified? Hydrol. Earth Syst. Sci. 26: 5357-5371.
688 doi:10.5194/hess-26-5357-2022.

689 Will, R.E., Barron, G.A., Burkes, E.C., Shiver, B. and Teskey, R.O. 2001. Relationship
690 between intercepted radiation, net photosynthesis, respiration, and rate of stem volume
691 growth of *Pinus taeda* and *Pinus elliottii* stands of different densities. For. Ecol. Manag. 154:
692 155-163. doi:10.1016/S0378-1127(00)00625-3.

693 Zhai, L., Jokela, E.J., Gezan, S.A. and Vogel, J.G. 2015. Family, environment and
694 silviculture effects in pure- and mixed-family stands of loblolly (*Pinus taeda* L.) and slash (*P.*
695 *elliottii* Engelm. Var. elliotttii) pine. For. Ecol. Manag. 337: 28-40.
696 doi:10.1016/j.foreco.2014.10.030.

Draft

© The Author(s) or their Institution(s)





Figure 1. Enumerated plantation area according to species, including *P. elliottii* (red), *P. taeda* (yellow), *P. patula* (green), *P. patula x P. tecunumanii* (blue) and the not sampled area (grey), within the Lowveld and Highveld forestry regions, respectively. Figure was created using ESRI ArcGIS version 10.8.2 and assembled from the following data sources: compartment species from compartment lists (Microforest), compartment boundaries layer (York Timbers). Base maps from ArcMap, World Terrain Base, courtesy of ESRI ArcGIS (ESRI 2021).

© The Author(s) or their Institution(s)




711

712 **Figure 2.** Comparison of tree dominant height growth from multiple linear regression model
713 predictions to the actual measured dominant height values for A) *P. elliottii* (red), B) *P. taeda*
714 (yellow), C) *P. patula* (green), and D) *P. patula x P. tecunumanii* (blue) trees.

715

© The Author(s) or their Institution(s)



716

717 **Figure 3.** Cate-Nelson analysis visualisation including the T-max threshold at 22.3 °C, at a
718 tree height threshold of 24.4 meters. All the visualisation of the computed thresholds for the
719 respective species can be seen in Supplemental Figures S6, S7, and S8.

720

Draft

© The Author(s) or their Institution(s)

721 **Table 1.** Summary of the enumeration compartment features for the different species of pine.

Species	Plantation count	Plantation area (ha)	Plot count	Measured area (ha)	Sample %
P. elliottii	619	7 290	9 840	487	6.7%
P. taeda	234	2 755	3 901	194	7.0%
P. patula	1 013	13 267	17 285	857	6.5%
P. patula x P. tecunumanii	217	2 949	3 714	184	6.2%

722

Draft

© The Author(s) or their Institution(s)

723 **Table 2.** Summary statistics for dominant height (Ht_d) regression independent variables per species. Significant = * (0.01<p<0.05), highly
724 significant = ** (0.001<p<0.01) and very highly significant *** (p<0.001). Rows containing "NS" indicate variables that were not significant
725 (p>0.05). Variables investigated include age, annual maximum temperature (T-max), annual minimum temperature (T-min), seasonal maximum
726 and minimum temperatures, annual rainfall (MAP) and seasonal rainfall.

Variable	*Pinus elliottii*						*Pinus taeda*						*Pinus patula*						*Pinus patula x Pinus tecunumanii*					
	Est.	**Std.Err**	**P**	**Sign.**	**RMSE**	**R²**	**Est.**	**Std.Err**	**P**	**Sign.**	**RMSE**	**R²**	**Est.**	**Std.Err**	**P**	**Sign.**	**RMSE**	**R²**	**Est.**	**Std.Err**	**P**	**Sign.**	**RMSE**	**R²**
(Intercept)	-17,28	2,39	0,000	***			-4,66	1,79	0,010	**			-6,17	1,52	0,000	***			-5,95	1,22	0,000	***		
Age	1,07	0,02	0,000	***			1,00	0,03	0,000	***			1,24	0,02	0,000	***			1,49	0,04	0,000	***		
T-max	0,19	0,04	0,000	***			NS	NS	NS	NS			NS	NS	NS	NS			NS	NS	NS	NS		
T-min	NS	NS	NS	NS			NS	NS	NS	NS			NS	NS	NS	NS			NS	NS	NS	NS		
T-max (summer)	NS	NS	NS	NS			NS	NS	NS	NS			NS	NS	NS	NS			NS	NS	NS	NS		
T-max (autumn)	NS	NS	NS	NS			NS	NS	NS	NS			NS	NS	NS	NS			NS	NS	NS	NS		
T-max (winter)	NS	NS	NS	NS			0,30	0,07	0,000	***			0,33	0,05	0,000	***			0,38	0,07	0,000	***		
T-max (spring)	NS	NS	NS	NS			NS	NS	NS	NS			NS	NS	NS	NS			NS	NS	NS	NS		
T-min (summer)	NS	NS	NS	NS	1,40	0,82	NS	NS	NS	NS	1,41	0,84	NS	NS	NS	NS	1,76	0,88	NS	NS	NS	NS	1,39	0,90
T-min (autumn)	NS	NS	NS	NS			NS	NS	NS	NS			NS	NS	NS	NS			NS	NS	NS	NS		
T-min (winter)	NS	NS	NS	NS			NS	NS	NS	NS			NS	NS	NS	NS			NS	NS	NS	NS		
T-min (spring)	NS	NS	NS	NS			NS	NS	NS	NS			NS	NS	NS	NS			NS	NS	NS	NS		
MAP	NS	NS	NS	NS			0,01	0,00	0,000	***			NS	NS	NS	NS			NS	NS	NS	NS		
Summer rainfall	0,01	0,00	0,000	***			NS	NS	NS	NS			NS	NS	NS	NS			NS	NS	NS	NS		
Autumn rainfall	NS	NS	NS	NS			-0,05	0,01	0,000	***			-0,02	0,00	0,000	***			NS	NS	NS	NS		
Spring rainfall	0,04	0,01	0,000	***			NS	NS	NS	NS			0,03	0,00	0,000	***			NS	NS	NS	NS		

727

Draft

© The Author(s) or their Institution(s)

728 **Table 3.** Climate and dominant height threshold values for annual maximum temperature
729 (C-T-max), annual minimum temperature (C-T-min), mean annual precipitation (C-MAP)
730 and the resultant dominant height (C-Ht_d) at age 20 years, categorised by species. Summary
731 statistics are provided, including P-model values, indicating the number of observations
732 accounted for within the respective opposing quadrants, and the chi-square-based Cramer's V
733 statistic, measuring the strength of association between the variables. Cramer's V statistic
734 values of 0.00-0.10 are considered negligible (no association), 0.10-0.20 weak, 0.20-0.40
735 moderate, 0.40-0.60 relatively strong, 0.60-0.80 strong, and 0.80-1.00 very strong.
736 Additionally, the relationship trend (positive or negative) between the respective climate
737 variables and dominant height growth was indicated (Lee 2016).

Variable	Species	C-T-max (°C)	C-Ht_d (m)	P-model	Cramer.V	Trend
T-max	*P. elliottii*	22.3	24.4	0.75	0.48	Positive
	P. taeda	23.3	24.2	0.88	0.46	Positive
	P. patula	20.4	26.9	0.92	0.73	Positive
	P. patula x P. tecunumanii	21.6	31.0	1.00	1.00	Positive
Variable	**Species**	**C-T-min (°C)**	**C-Ht_d (m)**	**P-model**	**Cramer.V**	**Trend**
T-min	*P. elliottii*	11.1	23.2	0.89	0.61	Positive
	P. taeda	11.2	23.1	0.76	0.57	Positive
	P. patula	11.6	27.6	0.84	0.67	Negative
	P. patula x P. tecunumanii	12.0	31.4	0.81	0.58	Negative
Variable	**Species**	**C-MAP (mm)**	**C-Ht_d (m)**	**P-model**	**Cramer.V**	**Trend**
MAP	*P. elliottii*	828.8	23.5	0.96	0.87	Positive
	P. taeda	916.7	23.2	0.84	0.70	Positive
	P. patula	868.0	27.7	0.84	0.69	Negative
	P. patula x P. tecunumanii	990.0	30.0	0.71	0.04	Negative

738

Draft

© The Author(s) or their Institution(s)




Figure 1

147x210mm (300 x 300 DPI)

© The Author(s) or their Institution(s)




Figure 2

317x223mm (96 x 96 DPI)

© The Author(s) or their Institution(s)




Figure 3

203x143mm (96 x 96 DPI)

© The Author(s) or their Institution(s)