



Molecular characterization and susceptibility of foodborne *Escherichia coli* to conventional antibiotics and natural extracts of *Solanum palinacanthum* and *Siparuna guianensis*

Larissa Alves Oliveira¹ · Mayara Bocchi² · Liliane Nebo³ · Bianca Ferreira Gonçalves³ · Flávio Barbosa da Silva⁴ · Ariel Eurides Stella¹ · Cecília Nunes Moreira¹

Received: 12 August 2025 / Accepted: 14 February 2026
© The Author(s) 2026

Abstract

The present study aimed to evaluate the antimicrobial susceptibility profile and molecular characteristics of *Escherichia coli* isolated from various foods, including ground beef, pork, Minas fresh cheese, lettuce, and chicken. A total of 150 *E. coli* strains were recovered and tested against conventional antimicrobials and crude extracts of *Solanum palinacanthum* and *Siparuna guianensis*. Susceptibility testing was conducted using disk diffusion and broth microdilution methods. Additionally, phylogenetic characterization and pathotype identification were performed via polymerase chain reaction (PCR). Results indicated that all *E. coli* isolates were resistant to at least two antimicrobial classes, with 55.3% ($n=83$) classified as multidrug-resistant (MDR). Most strains belonged to phylogenetic groups B1, A, or C. Regarding the natural extracts, the isolates were not sensitive to either *S. palinacanthum* or *S. guianensis*. In conclusion, the high prevalence of MDR *E. coli* across diverse food groups represents a critical concern for food safety and public health. Furthermore, the lack of susceptibility to the tested plant extracts highlights the challenge of identifying alternative natural antimicrobials, underscoring the urgent need for stricter control measures to mitigate the spread of resistant foodborne pathogens.

Keywords Food contamination · Gastrointestinal diseases · Multidrug-resistant *Escherichia coli*

Introduction

The World Health Organization (WHO) estimates that each year, around 600 million people worldwide become ill from eating contaminated food, resulting in 420,000 deaths. In Brazil, between 2014 and 2023, *Escherichia coli* was the primary etiological agent identified in foodborne disease

outbreaks, accounting for 34.8% of cases [1]. Beyond its role as a primary pathogen, *E. coli* is a major opportunistic agent and a critical indicator of fecal contamination and antimicrobial pressure in the environment. Its genomic plasticity allows it to act as a central reservoir for the transmission of antimicrobial resistance (AMR) genes, particularly through horizontal gene transfer, which poses a severe threat to public health by complicating the treatment of common infections [2].

E. coli is a commensal bacterium of the human and animal intestinal microbiota. However, some strains acquire virulence factors, becoming major causes of intestinal infections [3]. The rising global threat of AMR has turned foodborne *E. coli* into a global concern, as the food chain serves as a bridge between animal reservoirs and human populations. Consequently, continuous monitoring of food isolates is essential to track the evolution of resistance patterns and to inform surveillance programs under the “One Health” approach [4].

Communicated by Maria Aparecida Scatamburlo Moreira

✉ Mayara Bocchi
mayara.fernandes@ufjf.edu.br

¹ Postgraduate Program in Bioscience and One Health, Federal University of Jataí, Jataí, Goiás, Brazil

² Department of Medicine, Federal University of Jataí, BR 364 km 195, Industrial Park Sector, Jataí 75801-615, GO, Brazil

³ Postgraduate Program in Chemistry, Federal University of Jataí, Jataí, Goiás, Brazil

⁴ University of Rio Verde, Rio Verde, Goiás, Brazil

The indiscriminate use of antibiotics in the agricultural sector contributes to selection pressure on *E. coli*, allowing them to acquire resistance genes that infiltrate the food chain [5]. Strains recovered from food show high resistance rates to critically important antimicrobials, such as quinolones, cephalosporins, and tetracyclines [6]. Given this escalating resistance crisis, there is an urgent need to explore alternative therapeutic and sanitizing strategies. Plant-derived bio-cides have emerged as promising alternatives due to their complex chemical compositions and ability to exert antibacterial effects through multiple mechanisms, which may reduce the likelihood of developing further resistance [7].

Some plants, such as *Siparuna guianensis* and *Solanum palinacanthum*, contain diverse secondary metabolites, including tannins, alkaloids, and flavonoids, whose synergistic presence is associated with potent antibacterial activity [8–10]. These botanical compounds represent a sustainable frontier for the development of new antimicrobials capable of targeting resistant strains where conventional drugs fail. In this sense, considering the importance of monitoring *E. coli* in food and the search for control alternatives, this study aimed to evaluate the molecular characteristics of *E. coli* isolated from food and its susceptibility to conventional antimicrobials and crude extracts of *S. guianensis* and *S. palinacanthum*.

Materials and methods

Experimental design

The current study used 150 bacterial strains of *E. coli* isolated from various foods as part of research carried out between 2013 and 2019, belonging to the bacteriothèque of the Federal University of Jataí (UFJ). Among them, 41 were isolated from chicken meat, 41 from ground beef, and 21 from pork, 26 from lettuce, and 21 from fresh Minas cheese.

Microbiological recovery of strains

From the different food samples, 25-gram aliquots were taken from an initial 200-gram sample and added to 225 mL of sterile buffered peptone water. The mixtures were homogenized in a stomacher for 60 s and subsequently incubated at 36 °C for 18 h in a shaker at 60 RPM. The presence of the bacteria was confirmed by streak-plating onto Harlequin® Chromogenic Agar *E. coli*/Coliform, where purple colonies were considered positive for *E. coli* after incubation at 36 °C for 24 h in a bacteriological incubator. Typical *E. coli* colonies were further confirmed by phenotypic biochemical tests, including indole, methyl red, citrate utilization, and Voges-Proskauer (IMViC tests). Quality control

for the antimicrobial susceptibility tests was ensured by using the reference strain *Escherichia coli* ATCC 25,922, in accordance with the CLSI M100 35th edition (2025) guidelines [11].

Antimicrobial susceptibility testing by disk diffusion

Previously recovered *E. coli* strains were tested for susceptibility to 17 antimicrobials using the disk diffusion method on Mueller-Hinton agar (Oxoid Ltd., Hampshire, United Kingdom). Bacterial suspensions were adjusted to a 0.5 McFarland standard (1.5×10^8 CFU/mL) and inoculated onto Mueller-Hinton agar plates. After incubation at 35 °C for 18–24 h, inhibition zones were measured with a digital caliper. The results for clinical antibiotics were interpreted as Susceptible (S), Intermediate (I), or Resistant (R) according to CLSI M100 35th ed. guidelines [11]. For veterinary-specific drugs or those without established CLSI breakpoints for Enterobacteriaceae (Penicillin, Tilmicosin, and Lincospectin), susceptibility was evaluated based on manufacturer instructions or epidemiological cut-offs to maintain consistency with the previously generated profiles. The following antimicrobials were evaluated: Aminoglycosides [Gentamicin (10 µg), Neomycin (30 µg)], β-lactams [Amoxicillin-clavulanic acid (20/10 µg), Ampicillin (10 µg), Ceftiofur (30 µg), Penicillin (30 µg)], Quinolones [Norfloxacin (10 µg), Ciprofloxacin (5 µg), Enrofloxacin (5 µg)], Macrolides [Tilmicosin (15 µg)], Sulfonamides [Trimethoprim-Sulfamethoxazole (25 µg)], Phosphonics [Fosfomycin (200 µg)], Tetracyclines [Oxytetracycline (30 µg), Doxycycline (30 µg), Tetracycline (30 µg)], Lincosamides [Lincospectin (109 µg)], and Polymyxin [Colistin (10 µg)]. Isolates were classified as multidrug-resistant (MDR) when they showed resistance to three or more antimicrobial classes [12].

Obtaining botanical material and identifying species

The botanical material of the native Cerrado plants *S. palinacanthum* and *S. guianensis* were obtained in the municipality of Jataí, Goiás, Brazil, located in the southwest region of the state of Goiás (GO). Branches containing green leaves and free of pathogens were used. Botanical identification of the species was performed visually, accompanied by literary material from the Department of Botanical Biology of UFJ.

Obtaining the ethanolic extracts

The extracts were obtained from the Organic and Inorganic Chemistry Laboratory of UFJ, following the methodology described by Carvalho et al. [13]. First, healthy leaves were

separated, washed in running water, and placed in plastic trays for drying at room temperature and in a ventilated place for seven days.

After drying, the leaves were crushed in a blender to obtain a powder. This material was then subjected to the extraction process using ethanol as a diluent. The mixture was left at room temperature, with stirring, for at least 12 h. Three consecutive extractions of the same powder were performed to remove as many active biological compounds as possible.

After extraction, the material was filtered using a funnel and filter paper, in addition to a vacuum pump to remove the solid part, leaving only the crude ethanol extract. The extract was then placed in a rotary evaporator to evaporate the solvent at a temperature of 55 °C, leaving only the crude extract. The material was weighed and stored in properly identified and packaged glass containers. The material was left in the exhaust hood to completely eliminate the ethanol. After being weighed three times consecutively, on alternate days, the process was completed and the extract was stored in a freezer at −80 °C for later analysis.

Identification of secondary metabolites present in crude extracts of *S. palinacanthum* and *S. guianensis*

The phytochemical screening of the ethanolic extracts was performed according to the methodology described by Barbosa et al. [14], with modifications. For the tests of alkaloids, phenols, and tannins, general flavonoids, leucoanthocyanidins, catechins, steroids and triterpenoids and flavones, and saponins, the extracts were diluted in ethanol 92.8° NIWM.

Alkaloids

A total of 100 µL of the extract diluted in ethanol (10 mg/mL) were transferred to a contact plate and the medium was acidified with 3 drops of 1% HCl solution (v/v). Then, 4 drops of Dragendorff reagent were added and the color change for orange or reddish-orange in samples positive for the presence of alkaloids was observed.

Steroids and triterpenoids

In total, 10 mg of the extracts were diluted in 1000 µL of chloroform. They were then filtered and transferred to a test tube to which 100 µL of acetic anhydride and 3 drops of concentrated H₂SO₄ were added. After the procedure, a change in color was observed in the positive samples. The presence of steroids is confirmed by a blue-green coloration, while triterpenoids are indicated by a brown to red coloration.

Phenols and tannins

On a contact plate, 2 drops of 1% (m/v) FeCl₃ solution were added to 100 µL of extract diluted in ethanol (10 mg/mL). As an indicator, a blank containing ethanol and 1% (m/v) FeCl₃ was used. The presence of phenols or tannins results in a coloration that varies between blue-green, dark blue or black. Any change in color or formation of a precipitate is considered positive when compared with the blank test.

Flavonoids

A total of 100 µL of extract diluted in ethanol (10 mg/mL) were pipetted into three wells on the contact plate, acidifying one medium with pH 3, while the remaining two were alkalized to pH 8.5 and 11. The results were then observed. Initially, the extract is treated with an alkaline solution, such as sodium hydroxide, which can intensify the characteristic yellow color of flavonoids. Then, the addition of an acid, such as hydrochloric acid, can cause this color to disappear or change, indicating the presence of flavonoids.

Leucoanthocyanidins, catechins, and flavones

A total of 500 µL of the extract diluted in ethanol (10 mg/mL) were transferred to two numbered test tubes. Tube 1 was acidified with 4 drops of 1% HCl (v/v). In tube 2, 4 drops of 1% NaOH solution (v/v) were added. The solutions were heated in a water bath. Changes in color indicate a positive result for the test. Leucoanthocyanidins may present a reddish or purple coloration after acidification and heating, due to the formation of anthocyanidins. Catechins may exhibit a greenish or brown coloration after alkalization and heating, due to oxidation. Flavones may change color in an alkaline medium, exhibiting yellowish or orange tones.

Saponins

A total of 20 mg of the extracts were diluted in 2 mL of ethanol and transferred to a test tube, to which 5 mL of distilled water and 5 mL of hot water were added. The solution was stirred manually for about two minutes and then the result was observed. Saponins are substances with surface-active properties, forming stable foam when agitated in water.

Sensitivity test of crude extracts of *S. palinacanthum* and *S. guianensis* by disk diffusion

The sensitivity test for the crude extracts of *S. palinacanthum* and *S. guianensis* was performed in triplicate, using the disk diffusion method on filter paper, as adapted from the Kirby-Bauer method [15] using the bacterial strains that

showed resistance or multiresistance to the antimicrobials tested.

Concentrations were prepared from the crude extract at 50% (500 µL/mL) and diluted to 10% (100 µL/mL), 5% (50 µL/mL), 2.5% (25 µL/mL), 1% (10 µL/mL), and 0.1% (1 µL/mL) [16]. Discs of 6 mm in diameter were used, soaked with 10 µL of each extract under study.

Bacterial suspensions were prepared in 0.9% sterile saline to reach a turbidity equivalent to a 0.5 McFarland standard (approximately 1.5×10^8 CFU/mL). This standardized inoculum was then evenly spread across the surface of Mueller-Hinton agar plates (4 mm depth). Subsequently, disks impregnated with the plant extracts at different concentrations were added, along with negative control disks (solvents and distilled water) and positive control disks (fosfomycin 30 µg). Following a 24-hour incubation at $36^\circ \pm 1^\circ \text{C}$, the diameters of the bacterial growth inhibition zones were measured in millimeters (mm) [17, 18].

Broth microdilution test of crude *S. palinacanthum* and *S. guianensis*

The strains that showed resistance and multiresistance to the antimicrobials used in the study were tested using the microdilution technique in 96-well plates containing BHI broth and ethanol extract of *S. palinacanthum* and *S. guianensis*, which were diluted in distilled water, previously autoclaved in an amber bottle, obtaining a concentration of 10 mg/mL. In total, 200 µL of the extract (10 mg/mL) and 100 µL of the BHI broth were used, following the serial dilution principle ($\frac{1}{2}$), obtaining the following extract concentrations: 10 mg/mL, 5 mg/mL, 2.5 mg/mL, 1.25 mg/mL, 0.625 mg/mL, and 0.313 mg/mL. After the dilution procedure, 10 µL of the bacterial solution adjusted to the 0.5 MacFarland scale were added in triplicate. To validate the microdilution test, a positive control containing BHI broth and the bacterial inoculum of the tested strain was considered, as well as a negative control, containing only the BHI broth.

Molecular characterization of *E. coli* strains isolated from food

The DNA of all 150 *E. coli* strains was extracted by the boiling method, according to Olsvik and Strockbine [19]. The DNA was transferred to a 0.5 mL Eppendorf tube, free of DNase and RNase, previously sterilized and kept at -20°C until the time of analysis.

Phylogenetic classification

The determination of the phylogenetic group was performed by means of simultaneous amplification assays

(PCR-quadruplex), aiming at the sequential detection of DNA from certain genes, as described by Clermont et al. [20] (Supplementary Table 1). The reactions were performed with a final volume of 20 µL, containing PCR Buffer 1x, 3 mM MgCl₂, 0.2 mM dNTP, 2 units of Taq DNA polymerase, 3 µL of DNA, and the appropriate primers. In total, 0.2 µM of each primer were used, except for AceK.f, which had a concentration in the reaction of 0.4 µM. All reactions were performed in the Veriti thermocycler (Applied Biosystems) programmed for initial denaturation at 94°C (4 min), followed by 30 cycles of 94°C (5 s), 57°C (group E) or 59°C (quadruplex and group C) (20 s) and 72°C (10 s). After the cycles, the final extension was performed at 72°C (5 min).

The PCR amplified products were detected by 2.0% agarose gel electrophoresis (BioAmerica Biotech®) using Tris/Borate/EDTA - TBE X buffer (0.89 M Tris; 0.02 M EDTA; 0.89 M boric acid) and the 100 bp Ladder plus molecular weight standard (Sinapse Inc.). Electrophoretic separation was performed at 4 Volts/cm for a period of approximately 1 h. After electrophoresis, the gels were stained in 0.5 mg/mL ethidium bromide solution and visualized under an ultraviolet transilluminator.

For the classification of phylogroups, the combination of the presence and absence of primers is as follows: A: *arpA* or *arpA* and *yjaA*; B1: *arpA* and TspE4.C2; C: *arpA* and *yjaA*; E: *arpA* and *chuA* or *arpA* and *chuA* and *yjaA* or *arpA* and *chuA* and TspE4.C2; D: *arpA* and *chuA* or *arpA* and *chuA* and TspE4.C2; F: *chuA*; B2: *chuA* and TspE4.C2 or *chuA* and *yjaA* and TspE4.C2 or *chuA* and TspE4.C2. E. clades: *yjaA* or *arpA* and *chuA* and *yjaA* [20].

Characterization of diarrheagenic *E. coli* pathotypes

DNA samples obtained from the 150 *E. coli* isolates were grouped into pools of a maximum of 5 samples for the characterization of Shiga toxin-producing *E. coli* (STEC), enteropathogenic *E. coli* (EPEC), and enterohemorrhagic *E. coli* (EHEC), as described by OH et al. [21], with modifications. Two PCRs were performed: a triplex PCR, allowing the identification of the virulence factors *stx1* and *stx2*, in addition to the *E16S* gene (used as an internal control for *E. coli*), and the second PCR, performed to identify the virulence factor *eaeA*. The sequences of the primers used are described in supplementary Table 2.

Triplex PCR was performed in a final volume of 25 µL, containing 10.7 µL of nuclease-free ultrapure water, 1x PCR buffer, 3 mM MgCl₂, 0.2 mM dNTPs, 0.4 µM each of the primers *stx1.F*, *stx1.R*, *stx2.F*, and *stx2.R*, 0.1 µM of primers *E16S.F* and *E16S.R*, 1.5 units of Taq DNA Polymerase (Ludwig Biotec), and 3 µL of DNA. Amplification was performed in a Veriti thermocycler (Applied Biosystems) with initial denaturation at 95°C for 1 min, followed by 35 cycles

at 95°C for 1 min, 60°C for 1.5 min, and 72°C for 1.5 min, and a final extension at 72°C for 7 min.

For PCR of the *eaeA* gene, a final volume of 20 µL was used, containing 10.8 µL of ultrapure water, 1x PCR buffer, 2 mM MgCl₂, 0.2 mM dNTPs, 0.3 µM of each of the primers *eaeA200.F* and *eaeA200.R*, 1 unit of Taq DNA Polymerase (Ludwig Biotec), and 3 µL of DNA. DNA amplification was performed in a Veriti thermocycler (Applied Biosystems) with initial denaturation at 95°C for 5 min, followed by 35 cycles at 95°C for 30 s, 55°C for 30 s, and 72°C for 30 s, and a final extension at 72°C for 7 min.

The PCR amplified products were detected by 2% agarose gel electrophoresis (BioAmerica Biotech®) using Tris/Borate/EDTA - TBE 1X buffer (0.89 M Tris; 0.02 M EDTA; 0.89 M boric acid) and 100 bp Ladder plus molecular weight standard (Sinapse Inc.). Electrophoretic separation was performed at 4 V/cm for approximately 40 min. The gels were stained in 0.5 mg/mL ethidium bromide solution and visualized under an ultraviolet transilluminator.

Results

Susceptibility profile of *E. coli* strains to antimicrobials

The 150 *E. coli* strains were recovered and tested for susceptibility to 17 antimicrobials, as shown in Table 1. These antimicrobials were grouped according to their mechanism of action, resulting in 10 distinct groups: β-lactams, quinolones, tetracyclines, aminoglycosides, amphenicols, sulfonamides, lincosamides, polymyxins, macrolides, and fosfomycins. In the current study, all *E. coli* strains presented resistance to at least 2 groups of antimicrobials tested and 55.3% ($n=83$) of the samples were classified as multi-resistant. Lettuce was the food with the highest number of multiresistant strains, followed by Minas fresh cheese, pork, ground beef, and chicken (Fig. 1).

The antimicrobial classes with the highest resistance frequencies were β-lactams, followed by macrolides. Regarding the susceptibility profile, all *E. coli* strains were susceptible to fosfomycin (100%; $n=150$). Figure 2 shows the percentages of *E. coli* strains resistant to antimicrobials present in food.

Phytochemical characterization and susceptibility profile of *E. coli* strains to ethanolic extracts of *S. palinacanthum* and *S. guianensis*

The present study observed the presence of secondary metabolites in the ethanolic extracts of *S. guianensis* and

Table 1 Antimicrobial susceptibility profile for 150 *E. coli* isolates from food

Pharmaco-logical group	Antimicrobial	Susceptible	Intermediate	Resistant
		<i>n</i> (%)	<i>n</i> (%)	<i>n</i> (%)
β-lactams	Amoxicillin-clavulanic acid (AMC; 20 µg)	102 (68.0)	14 (9.3)	34 (22.7)
	Ampicillin (AMP; 10 µg)	101 (67.3)	1 (0.7)	48 (32.0)
	Penicillin (P; 30 µg)	0 (0)	0 (0)	150 (100)
	Ceftiofur (EFX; 30 µg)	77 (51.3)	26 (17.3)	47 (31.3)
	Lincospectin (LS; 109 µg)	143 (95.3)	0 (0)	7 (4.7)
Aminoglycosides	Gentamicin (CN; 10 µg)	143 (95.3)	4 (2.7)	3 (2.0)
	Neomycin (N; 30 µg)	130 (86.7)	12 (8.0)	8 (5.3)
Quinolones	Norfloxacin (NOR; 10 µg)	143 (95.3)	6 (4.0)	1 (0.7)
	Ciprofloxacin (CIP; 5 µg)	131 (87.3)	15 (10.0)	4 (2.7)
	Enrofloxacin (ENR; 05 µg)	100 (67.0)	40 (26.0)	10 (7.0)
	Oxytetracycline (OX; 30 µg), Doxycycline (DOX; 30 µg), Tetracycline (TRA; 30 µg)	111 (74.0)	3 (2.0)	36 (24.0)
Tetracyclines		112 (74.7)	7 (4.7)	31 (20.6)
		83 (55.3)	4 (2.7)	63 (42.0)
Polymyxins	Colistin (CL; 10 µg)	147 (98.0)	0 (0)	3 (2.0)
	Tilmicosin (TIL; 15 µg)	6 (4.0)	0 (0)	144 (96.0)
Macrolides		133 (88.7)	11 (7.3)	6 (4.0)
Sulfonamides	Trimethoprim-Sulfamethoxazole (SXT; 25 µg)			
Phosphonics	Fosfomycin (FOS; 200 µg)	149 (99.3)	1 (0.7)	0 (0)

S. palinacanthum. The extract of *S. guianensis* was positive for the presence of phenols and tannins, flavonoids, leucoanthocyanidins, catechins, flavones, and saponins. The extract of *S. palinacanthum* presented the following compounds as secondary metabolites: steroids and triterpenoids, phenols and tannins, flavonoids, leucoanthocyanidins, catechins, and flavones (Table 2).

Although both extracts present secondary metabolites with antimicrobial action, such as tannins and flavonoids, the present study did not observe such an effect in *E. coli* isolates by the disk diffusion and microdilution methods.




Fig. 1 Percentage of multidrug-resistant *E. coli* strains present in food

Molecular characterization of *E. coli* strains isolated from food

From the determination of phylogenetic groups performed by quadruplex PCR, the present study observed that the majority of the analyzed samples (62.6%; $n=94$) belonged to phylogenetic group B1, followed by group A or C (18.0%; $n=27$). The phylogenetic groups with the lowest frequency were: D or E (6.0%; $n=9$), F (4.0%; $n=6$), Clade I or II (2.7%; $n=4$), and B2 (0.7%; $n=1$). A total of 6.0% ($n=9$) of the analyzed samples were classified as undetermined (U) (Table 3).

Regarding the presence of STEC, EHEC, and EPEC pathotypes in the analyzed samples, no isolate presented positive amplification results for the virulence genes *stx1*, *stx2*, and *eaeA*. However, the gene for 16 S rRNA (E16S) was amplified in all *E. coli* pools used in the study, demonstrating the presence, physical quality, and purity of the DNA.

Discussion

Although most strains of *E. coli* present in the food chain are harmless, some strains have the ability to acquire virulence genes and are considered one of the main causes of foodborne illnesses associated with intestinal diseases [22]. An additional concern is the increasing levels of antibiotic-resistant *E. coli*. The misuse and overuse of antimicrobial agents in food production puts selection pressure on bacteria, allowing them to acquire resistance genes [23]. This phenomenon represents a global threat to public health, as the food chain acts as a primary bridge for the dissemination of MDR bacteria from animal reservoirs to human populations.

E. coli strains have been found in different foods, such as pasteurized milk, ground beef, cheese, vegetables (carrots, spinach, and lettuce), chicken, turkey, and pork, among others. These products can serve as a source of contamination and transmission of resistant strains that pose a risk to human health [24–26]. The present study observed that all *E. coli* strains recovered from food showed resistance to at least 2 classes of antimicrobials tested and that 55.3% were classified as multidrug-resistant. This high prevalence of MDR strains in diverse food groups underscores a critical risk; these findings suggest that foodborne *E. coli* is not merely a contaminant but a significant reservoir for antimicrobial resistance genes that can be horizontally transferred to human pathogens during consumption or through environmental contact.

High rates of antimicrobial-resistant *E. coli* present in food are described in the literature. According to Baloyi et al. [27], 85.71% of *E. coli* isolates from spinach, apples, carrots, cabbage, and tomatoes were multidrug-resistant, and the antibiotic classes with the highest resistance rate were: aminoglycosides (94.81%), cephalosporins (93.51%), β -lactams (93.51%), and chloramphenicol (87.01%). Igbinosa et al. [28] identified that 85.9% of the isolates were resistant to ≥ 3 and ≤ 7 antimicrobial classes. Even in broader screenings, such as the study by Guragain et al. [29] with 3,367 samples, 17.3% were resistant to at least three antibiotics.

This scenario highlights the necessity of the “One Health” approach, recognizing that the resistance found in animal products and vegetables, such as lettuce, is part of an interconnected cycle of environmental and human exposure. According to Popli et al. [30], approximately 73% of antimicrobials sold globally are used in livestock, which explains the high selection pressure observed in our animal-origin samples. Furthermore, the detection of resistance in lettuce suggests a bidirectional flow: contamination via irrigation water containing animal waste or urban effluents allows ARGs to migrate from the agricultural environment directly to the consumer’s Table [31].

Despite presenting high resistance rates, the *E. coli* samples recovered in the study belonged mostly to phylogenetic group B1, followed by group A or C. Jalil et al. [32] observed similar results, where the most prevalent phylogenetic group was B1 (57.0%), followed by A (33.0%), D (6.0%), and B2 (4.0%). While groups B1 and A are typically classified as commensals, their high rate of multiresistance in this study is a significant public health concern. It indicates that even non-pathogenic *E. coli* lineages are successfully adapting to antimicrobial pressure, serving as “silent” environmental reservoirs capable of transferring resistance traits to pathogenic strains within the human gut [4].




Fig. 2 Percentage of *E. coli* strains resistant to antimicrobial groups present in food. β-lac - β-lactams; Qui - quinolones; Tetra - tetracyclines; Ami - Aminoglycosides; Amp - amphenicols; Sulf - sulfonamides; Lin - lincosamides; Pol - polymyxins; Mac - macrolides

Table 2 Secondary metabolites found in ethanol extracts of *S. guianensis* and *S. palinacanthum*

Compounds analyzed	Ethanol extract of leaves	
	<i>S. guianensis</i>	<i>S. palinacanthum</i>
Alkaloids	-	-
Steroids and triterpenoids	-	+
Phenols and tannins	+	+
Flavonoids*	+	+
Leucoanthocyanidins, catechins and flavones	+	+
Saponins	+	-

Legend: + positive; - negative; *yellow coloration at pH 11, indicative of flavones, flavonols and xanthones

Regarding the presence of STEC, EHEC, and EPEC pathotypes in the analyzed samples, no isolate presented positive amplification results for the virulence genes *stx1*, *stx2*, and *eaeA*. This result corroborates the phylogenetic data found, since 80.0% of the samples in the present study belong to groups B1, A, or C, that is, they are commensal bacteria. Amézquita-Montes et al. [24] also observed a low prevalence of pathogenic strains. Isolates carrying virulence genes exclusive to EPEC, STEC, and ETEC were identified in 8 (2.0%) of the 380 food samples analyzed. Although the rates found were relatively low, the presence of diarrheagenic pathotypes such as STEC, EHEC, and EPEC are responsible for large outbreaks of foodborne illnesses.

The presence of virulence factors, combined with the high prevalence of multiresistant *E. coli* strains found in food and the ease of transmission of antimicrobial resistance genes, have sparked researchers' interest in developing new methods of controlling microorganisms, such as the use of plant-based products [33, 34]. *S. palinacanthum* and *S. guianensis* have been highlighted for their inhibitory actions on different microorganisms due to the presence of alkaloids, tannins, and flavonoids [35–37].

The literature highlights the inhibitory actions of *S. palinacanthum* extract on different microorganisms, such as *Staphylococcus aureus*, *Pseudomonas aeruginosa*, and *Salmonella typhimurium* [35]. Lobô et al. [36] also demonstrated the antimicrobial activity of the plant against bacteria such as *E. coli*, *Staphylococcus aureus*, and *Pseudomonas*

aeruginosa. This activity was related to the presence of large amounts of alkaloids and tannins.

The phytochemistry of the crude leaf extract of *S. guianensis* suggests antimicrobial potential [37]. Studies carried out by Alves et al. [38] and Pereira et al. [39] demonstrated that *S. palinacanthum* extract can inhibit the growth of several bacteria in vitro, including *E. coli*. Oliveira et al. [40] detected bactericidal activity of *S. guianensis* essential oil in vitro against *Streptococcus mutans*, *Enterococcus faecalis* and *E. coli* bacteria, as well as antifungal action against *Candida albicans*. Moura et al. [41] also detected bactericidal activity of *S. guianensis* against *E. coli*, *Pseudomonas aeruginosa*, *Staphylococcus aureus* and *Streptococcus pyogenes*.

According to literature data, both plants present large groups of secondary metabolites that give them antimicrobial action [42], which were identified in the present study by phytochemical screening. However, it was not possible to determine the potential for bacteriostatic or bactericidal action by the proposed methods, since they were not able to inhibit bacterial growth. Other studies also did not observe an antimicrobial effect of *S. guianensis* and *S. palinacanthum* extracts against strains of *E. coli* [35, 43].

The lack of susceptibility of *E. coli* to the tested extracts can be largely attributed to the intrinsic structural resistance of Gram-negative bacteria. Unlike Gram-positive organisms, *E. coli* possesses a complex outer membrane containing lipopolysaccharides (LPS), which creates a robust hydrophobic barrier that limits the diffusion of many plant-derived bioactive compounds [44, 45]. This architecture effectively restricts the access of larger or more polar molecules, such as certain tannins and flavonoids identified in our phytochemical screening, to their internal cellular targets [46]. This phenomenon is clearly reflected in our results, where 100% resistance was observed for Penicillin, Tilmicosin, and Lincospectin—drugs to which *E. coli* is naturally resistant due to membrane impermeability. Furthermore, the presence of specialized porins and active efflux pumps in these wild MDR isolates may further diminish the intracellular concentration of any internalized phytochemicals, rendering the crude extracts ineffective at the concentrations tested [47]. This structural complexity explains why botanical extracts often show more potent activity against Gram-positive than Gram-negative pathogens.

Table 3 Distribution of phylogenetic groups of *E. coli* isolated from food

Phylogenetic Group	Ground beef	Pork	Chicken	Cheese	Lettuce	Total by Group	% by Group
A or C	7	6	9	2	3	27	18.0%
B1	30	12	20	17	15	94	62.6%
B2	-	-	1	-	-	1	0.7%
Clade I or II	-	2	-	2	-	4	2.6%
D or E	2	-	1	-	6	9	6.0%
F	-	-	6	-	-	6	4.0%
U	2	1	4	-	2	9	6.0%
ALL	41	21	41	21	26	150	100%

This structural defense is likely amplified in our study by the extreme resistance profile of the foodborne isolates. Most previous studies reporting success, such as Alves et al. [38] and Oliveira et al. [40], often utilize standard laboratory strains (ATCC), which are significantly more susceptible than the wild, MDR isolates characterized here. This highlights a ‘resistance gap’, suggesting that the robust resistance mechanisms of these environmental strains exceeded the inhibitory capacity of the crude extracts. Consequently, while our findings provide a crucial baseline, they emphasize that future research must move beyond crude extracts toward purified fractions or synergistic combinations to overcome the high biological fitness and membrane complexity of MDR foodborne pathogens [48, 49].

Conclusion

The study demonstrates a high prevalence of MDR *E. coli* across diverse food groups, including animal-derived products and fresh produce. Although the isolates predominantly belong to commensal phylogenetic groups (B1 and A) and lack the tested virulence genes, their robust resistance profile poses a significant threat to public health. These strains act as silent reservoirs of resistance genes that can be disseminated through the food chain, emphasizing the urgent need for integrated surveillance under the “One Health” framework.

Furthermore, while the crude extracts of *S. palinacanthum* and *S. guianensis* did not show inhibitory activity against the MDR foodborne isolates, the phytochemical screening confirms the presence of bioactive compounds. This lack of susceptibility highlights the high biological fitness of wild MDR strains compared to laboratory lineages. Therefore, future studies should explore purified fractions or synergistic applications of these botanical extracts with conventional antibiotics to overcome bacterial resistance mechanisms and ensure food safety.

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s42770-026-01898-9>.

Author contributions Larissa Alves Oliveira: Methodology, Investigation, Data Curation, Writing - Original Draft. Mayara Bocchi: Writing - Review & Editing, Visualization. Liliane Nebo: Methodology, Investigation. Bianca Ferreira Gonçalves: Methodology, Investigation. Flávio Barbosa da Silva: Validation, Formal analysis. Ariel Eurides Stella: Conceptualization, Writing - Review & Editing. Cecília Nunes Moreira: Conceptualization, Resources, Supervision, Project administration.

Funding statement The Article Processing Charge (APC) for the publication of this research was funded by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior - Brasil (CAPES) (ROR identifier: 00x0ma614). Not applicable.

Data availability All data generated or analyzed during this study are included in this published article. Furthermore, the data supporting this study’s findings are available on request from the corresponding author.

Declarations

Ethical approval Not applicable.

Consent to publish Not applicable.

Consent to participate Not applicable.

Competing interests We confirm that there are no known conflicts of interest associated with this work, and there has been no significant financial support for this work that could have influenced its outcome. Furthermore, we ensure that the manuscript has been read and approved by all named authors and that no other persons have satisfied the criteria for authorship but are not listed. We further confirm that everyone has agreed on the order of authors listed in the manuscript of us. Due care has been taken to ensure the integrity of the work.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article’s Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article’s Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Brazil (2024) Report Ministry of Health. Outbreaks of Waterborne and Foodborne Diseases – Retrieved from <https://www.gov.br/saude/pt-br/assuntos/saude-de-a-a-z/d/dtha/situacao-epidemiologica>. Accessed June 6, 2025
2. Poirel L, Madec JY, Lupo A, Schink AK, Kieffer N, Nordmann P, Schwarz S (2018) Antimicrobial resistance in *Escherichia coli*. Microbiol Spectr 6(6):10.1128/microbiolspec.ARBA-0026-2017. <https://doi.org/10.1128/microbiolspec.ARBA-0026-2017>
3. O’Neill L, Manzanilla EG, Ekhlās D, Leonard FC (2023) Antimicrobial resistance in commensal *Escherichia coli* of the porcine gastrointestinal tract. Antibiotics (Basel, Switzerland) 12(11):1616. <https://doi.org/10.3390/antibiotics12111616>
4. McEwen SA, Collignon PJ (2018) Antimicrobial resistance: a one health perspective. Microbiol Spectr 6(2):10.1128/microbiolspec.ARBA-0009-2017. <https://doi.org/10.1128/microbiolspec.ARBA-0009-2017>
5. Pradhan SS, Mahanty A, Pattanaik KP, Adak T, Mohapatra PK (2025) Entry, fate and impact of antibiotics in rice agroecosystem: a comprehensive review. Environ Sci Pollut Res Int 32(3):1120–1138. <https://doi.org/10.1007/s11356-024-35765-7>
6. Babines-Orozco L, Balbuena-Alonso MG, Barrios-Villa E, Lozano-Zarain P, Martínez-Laguna Y, Del Carmen Rocha-Gracia R, Cortés-Cortés G (2024) Antimicrobial resistance in food-associated *Escherichia coli* in Mexico and Latin America. Biosci

- Microbiota Food Health 43(1):4–12. <https://doi.org/10.12938/bm.fh.2023-022>
7. Khameneh B, Iranshahy M, Soheili V, Bazzaz BSF (2019) Review on plant antimicrobials: a mechanistic viewpoint. *Antimicrob Resist Infect Control* 8:118. <https://doi.org/10.1186/s13756-019-0559-6>
 8. Chung PY, Navaratnam P, Chung LY (2011) Synergistic antimicrobial activity between pentacyclic triterpenoids and antibiotics against *Staphylococcus aureus* strains. *Ann Clin Microbiol Antimicrob* 10:25. <https://doi.org/10.1186/1476-0711-10-25>
 9. Rajeh MA, Zuraini Z, Sasidharan S, Latha LY, Amutha S (2010) Assessment of *Euphorbia hirta* L. leaf, flower, stem and root extracts for their antibacterial and antifungal activity and brine shrimp lethality. *Molecules* 15(9):6008–6018. <https://doi.org/10.3390/molecules15096008>
 10. Matias LJ, Rocha JA, Menezes EV, De Melo Júnior AF, De Oliveira DA (2019) Phytochemistry in medicinal species of *Solanum* L. (Solanaceae). *Pharmacogn Res* 11(1):47–50
 11. Clinical and Laboratory Standards Institute (2025) Performance Standards for Antimicrobial Susceptibility Testing. 35th ed. CLSI supplement M100. Wayne, PA. <https://clsi.org/standards/product/s/microbiology/documents/m100/>
 12. Magiorakos AP, Srinivasan A, Carey RB, Carmeli Y, Falagas ME, Giske CG, Harbarth S, Hindler JF, Kahlmeter G, Olsson-Liljequist B, Paterson DL, Rice LB, Stelling J, Struelens MJ, Vatopoulos A, Weber JT, Monnet DL (2012) Multidrug-resistant, extensively drug-resistant and pandrug-resistant bacteria: an international expert proposal for interim standard definitions for acquired resistance. *Clin Microbiol Infect* 18(3):268–281. <https://doi.org/10.1111/j.1469-0691.2011.03570.x>
 13. Carvalho VF, Ramos LDA, da Silva CA, Nebo L, Moraes D, da Silva FFA, da Costa NCA, Rodrigues Junior RO, de Souza LF, Rodrigues RM (2019) In vitro anthelmintic activity of *Siparuna guianensis* extract and essential oil against *Strongyloides venezuelensis*. *J Helminthol* 94:e50. <https://doi.org/10.1017/S0022149X19000282>
 14. Barbosa WLR, Quignard E, Tavares ICC, Pinto LN, Oliveira FQ, Oliveira RM (2004) Manual para análise fitoquímica e cromatográfica de extratos vegetais. *Revista científica da UFPA* 4(9):1–19
 15. Nardeli JV, De Carvalho CT, De Arruda EJ, Vila Nova PCC, De Oliveira LCS, Arakaki AH (2014) Síntese, caracterização e atividade biológica do acetato de Cu (II) para larvas de *Aedes aegypti* (Diptera: culicidae) e bactérias *Escherichia coli*, *Staphylococcus aureus*, *Salmonella typhimurium* e *Lysteria monocytogenes*. *Orbital Electron J Chem* 6(3):122–129
 16. Haida KS, Parzianello L, Werner S, Garcia DR, Inácio CV (2007) In vitro assessment of the antimicrobial activity from eight medicinal plant species. *Arquivos de Ciências da Saúde da UNIPAR* 11(3)
 17. Azevedo SMM (2014) Farmacologia dos antibióticos Beta-lactâmicos, Master's thesis, Universidade Fernando Pessoa (Portugal)
 18. Guimarães DO, Momesso LDS, Pupo MT (2010) Antibiotics: therapeutic importance and perspectives for the discovery and development of new agents. *Quim Nova* 33:667–679. <https://doi.org/10.1590/S0100-40422010000300035>
 19. Olsvik O, Strockbine NA (1993) Polymerase chain reaction detection of heat-stable, heat-labile, and Shiga-like toxin genes in *Escherichia coli*. *Diagn Mol Microbiol* 271–276
 20. Clermont O, Christenson JK, Denamur E, Gordon DM (2013) The Clermont *Escherichia coli* phylo-typing method revisited: improvement of specificity and detection of new phylo-groups. *Environ Microbiol Rep* 5(1):58–65. <https://doi.org/10.1111/1758-2229.12019>
 21. Oh KH, Kim SB, Park MS, Cho SH (2014) Development of a one-step PCR assay with nine primer pairs for the detection of five diarrheagenic *Escherichia coli* types. *J Microbiol Biotechnol* 24(6):862–868. <https://doi.org/10.4014/jmb.1312.12031>
 22. Furlan JPR, Ramos MS, Dos Santos LDR, da Silva Rosa R, Stehling EG (2023) Multidrug-resistant Shiga toxin-producing *Escherichia coli* and hybrid pathogenic strains of bovine origin. *Vet Res Commun* 47(4):1907–1913. <https://doi.org/10.1007/s11259-023-10141-x>
 23. Ratshilingano MT, du Plessis EM, Duvenage S, Korsten L (2022) Characterization of multidrug-resistant *Escherichia coli* isolated from two commercial lettuce and spinach supply chains. *J Food Prot* 85(1):122–132. <https://doi.org/10.4315/JFP-21-125>
 24. Amézquita-Montes Z, Tamborski M, Kopsombut UG, Zhang C, Arzuza OS, Gómez-Duarte OG (2015) Genetic relatedness among *Escherichia coli* pathotypes isolated from food products for human consumption in Cartagena, Colombia. *Foodborne Pathog Dis* 12(5):454–461. <https://doi.org/10.1089/fpd.2014.1881>
 25. Davis GS, Waits K, Nordstrom L, Grande H, Weaver B, Papp K, Horwinski J, Koch B, Hungate BA, Liu CM, Price LB (2018) Antibiotic-resistant *Escherichia coli* from retail poultry meat with different antibiotic use claims. *BMC Microbiol* 18(1):174. <https://doi.org/10.1186/s12866-018-1322-5>
 26. Guibert F, Espinoza K, Taboada-Blanco C, Alonso CA, Oporto R, Castillo AK, Rojo-Bezares B, López M, Sáenz Y, Pons MJ, Ruiz J (2025) Traditional marketed meats as a reservoir of multidrug-resistant *Escherichia coli*. *International microbiology: the official journal of the Spanish Society for Microbiology* 28:27–43. <https://doi.org/10.1007/s10123-023-00445-y>
 27. Baloyi T, Duvenage S, Du Plessis E, Villamizar-Rodríguez G, Korsten L (2022) Multidrug resistant *Escherichia coli* from fresh produce sold by street vendors in South African informal settlements. *Int J Environ Res Public Health* 32(7):1513–1528. <https://doi.org/10.1080/09603123.2021.1896681>
 28. Igbinosa EO, Beshiru A, Igbinosa IH, Cho GS, Franz CMAP (2023) Multidrug-resistant extended spectrum β -lactamase (ESBL)-producing *Escherichia coli* from farm produce and agricultural environments in Edo State, Nigeria. *PLoS One* 18(3):e0282835. <https://doi.org/10.1371/journal.pone.0282835>
 29. Guragain M, Schmidt JW, Bagi LK, Paoli GC, Kalchayanand N, Bosilevac JM (2024) Antibiotic resistance and disinfectant resistance among *Escherichia coli* isolated during red meat production. *J Food Prot* 87(6):100288. <https://doi.org/10.1016/j.jfp.2024.100288>
 30. Popli S, Girdhar M, Praba GS, Anand S (2024) A one-health perspective of antimicrobial resistance (AMR): human, animals and environmental health. *Antibiotics (Basel, Switzerland)* 13(10):972. <https://doi.org/10.3390/antibiotics13100972>
 31. Goyal SP, Shome R, Shome BR, Sreekala S, Hiremath J, Chandel SS, Pathak A, Chauhan S, Shinde VV, Venugopal N (2025) One health surveillance of multidrug-resistant diarrheagenic *Escherichia coli* in Northeast India. *Sci Rep* 15:372. <https://doi.org/10.1038/s41598-024-84279-8>
 32. Jalil A, Gul S, Bhatti MF, Siddiqui MF, Adnan F (2022) High occurrence of multidrug-resistant *Escherichia coli* strains in bovine fecal samples from healthy cows serves as rich reservoir for AMR transmission. *Antibiotics* 12(1):37. <https://doi.org/10.3390/antibiotics12010037>
 33. Yang SC, Lin CH, Aljuffali IA, Fang JY (2017) Current pathogenic *Escherichia coli* foodborne outbreak cases and therapy development. *Arch Microbiol* 199(6):811–825. <https://doi.org/10.1007/s00203-017-1393-y>
 34. Urban-Chmiel R, Marek A, Stępień-Pyśniak D, Wiecek K, Dec M, Nowaczek A, Osek J (2022) Antibiotic resistance in bacteria—a review. *Antibiotics (Basel, Switzerland)* 11(8):1079. <https://doi.org/10.3390/antibiotics11081079>

35. Bouzada ML, Fabri RL, Nogueira M, Konno TU, Duarte GG, Scio E (2009) Triagem antibacteriana, citotóxica e fitoquímica de algumas plantas medicinais tradicionais no Brasil. *Biol Farmacêutica* 47(1):44–52. <https://doi.org/10.1080/13880200802411771>
36. Lôbo KMS, Athayde ACR, Silva AMA, Rodrigues FFG, Lôbo IS, Bezerra DAC, Costa JGM (2010) Evaluation of antibacterial activity and phytochemical prospection of *Solanum paniculatum* Lam. and *Operculina hamiltonii* (G. Don) DF Austin & Staples from the semi-arid region of Paraíba State, Brazil. *Revista Brasileira de Plantas Medicinais* 12:227–235. <https://doi.org/10.1590/S1516-05722010000200016>
37. de Bessa NGF, Borges JCM, Beserra FP, Carvalho RHA, Pereira MAB, Fagundes R et al (2013) Preliminary phytochemical screening of native Cerrado plants of medicinal popular use by the rural community of the Vale Verde settlement - Tocantins. *Revista Brasileira de Plantas Medicinais* 15:692–707. <https://doi.org/10.1590/S1516-05722013000500010>
38. Alves A, Pires A, Linardi V, Reina L, Galvão C (2005) Atividade Antibacteriana e Antifúngica dos extratos brutos etanólicos de *Solanum palinacanthum*. Encontro de pesquisa de IES do sistema estadual de Minas
39. Pereira AC, Oliveira DF, Silva GH, Figueiredo HC, Cavalheiro AJ, Carvalho DA, Chalfoun SM (2008) Identification of the antimicrobial substances produced by *Solanum palinacanthum* (Solanaceae). *An Acad Bras Cienc* 80:427–432
40. Oliveira MS, da Cruz JN, da Almei Costa W, Silva SG, Brito MDP, de Menezes SAF, de Jesus Chaves Neto AM, de Aguiar Andra EH, de Carvalho Junior RN (2020) Chemical composition, antimicrobial properties of *Siparuna guianensis* essential oil and a molecular docking and dynamics molecular study of its major chemical constituent. *Molecules* (Basel, Switzerland) 25(17):3852. <https://doi.org/10.3390/molecules25173852>
41. de Souza Moura W, de Souza SR, Campos FS, Cangussu ASR, Santos EMS, Andrade BS et al (2020) Atividade antibacteriana do óleo essencial de *Siparuna guianensis* mediada pelo comprometimento da permeabilidade da membrana e da replicação de bactérias patogênicas. *Culturas e produtos industriais* 146:112142. <https://doi.org/10.1016/j.indcrop.2020.112142>
42. de Melo DC, Miranda MLD, Júnior WGF, de Andrade PM, Alcoba AET, de Souza Silva T et al (2017) Anticariogenic and antimycobacterial activities of the essential oil of *Siparuna guianensis* Aublet (Siparunaceae). *Orbital: Electron J Chem* 55–60. <https://doi.org/10.17807/orbital.v0i0.930>
43. Araruna MK, Brito SA, Morais-Braga MF, Santos KK, Souza TM, Leite TR, Costa JG, Coutinho HD (2012) Evaluation of antibiotic & antibiotic modifying activity of pilocarpine & rutin. *Indian J Med Res* 135(2):252–254
44. Breijyeh Z, Jubeh B, Karaman R (2020) Resistance of Gram-Negative Bacteria to current antibacterial agents and approaches to resolve it. *Molecules* 25(6):1340. <https://doi.org/10.3390/molecules25061340>
45. Masi M, Rifai S, Pagès JM (2022) The outer membrane of Gram-Negative Bacteria: a key player in resistance and a target for antibiotic discovery. *Antibiotics* 11(10):1351. <https://doi.org/10.3390/antibiotics11101351>
46. Khameneh B, Iranshahy M, Soheili V, Bazzaz BSV (2021) Review on plant-derived antimicrobial compounds and their mechanisms of action. *J Infect Public Health* 14(1):116–123. <https://doi.org/10.1016/j.jiph.2020.11.015>
47. Uddin TM, Chakraborty AJ, Khusro A, Zidan BRM, Mitra S, Emran TB, Dhama K, Ripon MKH, Gajdacs M, Ullah MA, Islam MA, Kloc M, Ghoneim MM, Morsy MA (2021) Antibiotic resistance in microbes: history, mechanisms, therapeutic strategies and future prospects. *J Infect Public Health* 14(12):1750–1766. <https://doi.org/10.1016/j.jiph.2021.10.020>
48. Bessa L, Ferreira Z, Soares B, Cruz-Martins N, Brandão B, Brás S, Alves A, Gomes C (2024) Plants with antimicrobial activity against *Escherichia coli*: a meta-analysis for green veterinary pharmacology applications. *Microorganisms* 12(4):718. <https://doi.org/10.3390/microorganisms12040718>
49. Montanari L, Santos-Filho PC, Sales TA, Rezende GAS, Vieira JDG, Silva FG (2023) Exploring the antimicrobial potential and medicinal properties of *Siparuna guianensis* in the Brazilian Cerrado. *J Med Plants Res* 17(8):213–224. <https://doi.org/10.5897/JMPR2023.7317>

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.