IntegrateMind: A Multidisciplinary Journal

**To Cite:**
Alagbe, J. O. Prosopis africana Extract as Potential Natural Alternatives to Synthetic Antibiotics and a Key for Sustainable Broiler Production: A Review. *IntegrateMind: A Multidisciplinary Journal* 2023; Vol 1, Issue 1, pp. 45-50.

**Author Affiliation:**
[1]Department of Animal Nutrition and Biochemistry, Sumitra Research Institute, Gujarat, India. Orcid: 0000-0003-0853-6144.

[*]**Corresponding Author**
[1]Department of Animal Nutrition and Biochemistry, Sumitra Research Institute, Gujarat, India.
Email:dralagbe@outlook.com

**Peer-Review History**
Received:6th February, 2024
Reviewed & Revised:07/February /2024 to 16th February/2024
Accepted:17thFebruary 2024
Published: 18th February 2024

**Peer-Review Model**
External peer-review was done through double-blind method.

IntegrateMind: A Multidisciplinary Journal

**ISSNxxxx–xxxx**

URL:https://sites.google.com/view/sherwanjournals/integratemind-journall

**Ethics**
Animal studies are presented in this manuscript.
No human studies are presented in this manuscript.
No potentially identifiable human images or data is presented in this study.

License: **Creative Commons Attribution 4.0 International**

**SHERWAN**
**Publishers**

# Prosopisafricana Extract as Potential Natural Alternatives to Synthetic Antibiotics and a Key for Sustainable Broiler Production: A Review

**John Olujimi Alagbe[1*]**

## ABSTRACT

The chicken sector is more vulnerable to antimicrobial resistance, the buildup of toxic or dangerous residues in meat and eggs, and environmental contamination as a result of the widespread and careless use of antibiotics in most nations. Natural solutions, such as medicinal plants, are required to address these issues. These plants have been shown to contain a variety of phytochemicals, including phenols, flavonoids, terpenoids, tannins, alkaloids, and saponins. These chemicals endow plants with a multitude of medicinal qualities. One of the many alternatives to antibiotics is Prosopis africana extract, which includes stem bark, leaves, roots, and their essential oils. Antioxidant, antibacterial, antifungal, anti-helminthic, antiviral, hepatoprotective, immune-stimulatory, and antimicrobial properties are among the many pharmacological activity of P. africana preparations. They are also an abundant supplier of vital minerals and amino acids that support enzyme function and provide defense against the effects of oxidative stress. Using extracts from Prosopis africana is the way forward for effective chicken production, environmental sustainability and food hygiene.

**Keywords:** Prosopisafricana, medicinal plants, phytochemicals, food safety, antibiotics.

## 1. INTRODUCTION

Public concern over potential antibiotic resistance risks related to human health has driven interest in poultry nutrition and adoption of antibiotic-free feeding systems (Ulrich and Arne, 2022). This has led to the development of feed additives such as Prosopis africana extracts that can be used as in-feed antibiotic alternatives in livestock feeding strategies (Lius, 2022; Jan, 2021). The extracts have a wide range of potential benefits, all targeting the enhancement of livestock performance.

The medium-sized, annual savannah tree Prosopis africana is a member of the fabaceae family. The majority of Asia and Africa (both East and West Africa) are home to the plant (Kolapo et al., 2009). There are roughly forty-five species of it, and it can reach heights of 4 to 20 meters. It has a thick tap root that expands quickly and deeply into the earth (Orwa et al. 2009). The plant's leaves range in size from 12 mm to 30 mm and are bipinnate, with 9–16 oblong leaf pairs (Aremu et al., 2007). When dried, the seeds of Prosopis africana can open freely inside a yellow-intermeshed pod (Weber et al. 2008).According to Kolapo et al. (2009), the plants' seeds, leaves, roots, and stem bark have lots in phytochemicals and are typically used to treat bronchitis, fever, gonorrhea, toothache, stomachache, dysentery, and body aches.

The variances in plant extract effectiveness can be attributed to a variety of factors, including climate, location, harvest stage, and storage conditions, as well as variations in the chemical content of the extracts made from Prosopis africana (Jan, 2020).Numerous phytochemicals found in P. africana extracts have been shown to have a variety of pharmacological effects, including anti-inflammatory, anti-helminthic, antioxidant, antimicrobial, immune-stimulatory, anti-fungal, and hepatoprotective qualities (Kalemba and Kunicka, 2003).

Essential minerals like potassium, phosphorus, calcium, manganese, magnesium, zinc, and copper are present in Prosopis africana (Ajiboye et al., 2013). According to Yanwuyi et al. (2010), these minerals are critical for the activation of essential enzymes and for protecting against a state of oxidative stress. Additionally, P. africana contains the amino acids methionine, lysine, threonine, leucine, alanine, phenyl alanine, cysteine, and proline, as well as vitamins A, B2, B5, B6, B9, and B12. These nutrients are essential for reducing cellular damage, scavenging free radicals, promoting collagen formation, and further enhancing antioxidant activities (Alagbe, 2022). Considering the wealth of potential found in Prosopis africana and the growing awareness of the need to offer natural antibiotic substitutes instead of manufactured ones. This review summarizes earlier research and emphasizes the need of advancing food safety and sustainable livestock practices.

## 2. THE PHYTOCHEMICAL MAKEUP OF EXTRACTS FROM PROSOPIS AFRICANA

### I. Phenols

Due to their antioxidant and free-radical scavenging qualities, phenolic compounds help keep animals healthy (Dai and Mumper, 2010). Phenols work by depriving bacteria of their substrate, which causes fragmentation of the bacterial cytoplasm (Cowan 1999). In their investigations, Cushnie and Lamb (2005) and Davies et al. (1998) found that the inhibition of enzymes by oxidized substrate was the cause of phenols' potency against bacteria.

### II. Alkaloids

According to Nath et al. (2018) and Singh (2008), alkaloids are naturally occurring organic molecules that have a nitrogen heteroatom in their framework. According to Ayushi et al. (2021) and Wink (2016), alkaloids may have a variety of medicinal effects on birds, including antibacterial, analgesic, and antioxidant qualities. It has been found that alkaloids prevent the synthesis of proteins and RNA (Nath et al., 2018).

### III. Flavonoids

According to Cowan (1999), the carbonyl group of flavonoids has hydroxyl groups at positions 3 and 7. Moreover, it has been proposed that they possess antioxidant and antibacterial qualities. Additionally, they may attach to bacterial cell proteins, inhibiting and deactivating enzymes (Cushnie and Lamb 2005).

### IV. Tannins

Every portion of a plant, including the stem, bark, roots, leaves, fruits, and flowers, contains tannins (Lim et al. 2006). The capacity of tannins to form complexes with proteins and polysaccharides via covalent and hydrogen bonding is one of their primary characteristics (Szallasi, 2005). Enzyme inactivation results from tannins' capacity to bind to surface proteins and form complexes that do so. Enzyme inactivation results from tannins' capacity to bind to surface proteins and form groups that do so. According to Lim et al. (2006), it also causes membrane rupture and substrate restriction as a result of complex formation.

### V. Saponins

A class of phytochemicals known as saponins is one of the main defense mechanisms plants have against microbial, fungal, and insect invasion. Whilst the quantity and concentration of saponins in plants differ from species to species, they are present in the majority of plant species (González-Lamothe et al. 2009). According to Morrissey and Osbourn (1999), saponins function by building compounds with sterols or polysaccharides inside the microbial cell membrane, which destroys the integrity of the cytoplasmic membrane.

Table 1 shows the active compound and its medicinal properties for ready reference.

**Table 1: Some bioactive compounds in *Prosopisafricana* oil and their properties (**

| Active compounds | Medicinal properties | References |
|---|---|---|
| Prosogerin A (6 Methoxy-7-hydroxyl dioxyflavone) | Antibacterial, antioxidant and immune-stimulaory | Guimarães (2010) |
| Prosogerin B (2,4 dihydroxyl -5 methyl dioxychacon) | Antimicrobial, hepato-protective and antioxidant | Jim´enez-Arellanes et al. (2003) |
| 2,4-bis (1-phenylethyl) phenol | Antioxidants, antifungal | Kokoska et al. (2002) |
| β-phenethylamine | Antifungal and immune-stimulatory | Lin et al. (2000) |
| 2,4,6-tris(1-phenylethyl) phenol | Antibacterial, antiviral | Loizou et al. (2010) |
| α-pinene | Antioxidant and digestive stimulants | Alagbe et al. (2022) |
| α-terpinene | Antioxidant, antibacterial, | Alagbe et al. (2023) |
| β-pinene | Antioxidant, antimicrobial | Egunyomi and Oladunjoye (2012) |
| β-myrcene | Antioxidant, digestive stimulants | Edeoga et al. (2005); Alagbe (2022) |
| α-phellandrene | Antioxidant, Antifungal and antimicrobial | Idowu et al. (2006) |

| Active compounds | Medicinal properties | References |
|---|---|---|
| α-terpinolene | Antioxidant, antibacterial | Guimarães (2010) |
| γ-terpinene | Antioxidant, immune-stimulatory | Jim´enez-Arellanes et al. (2003) |
| 1-terpineol | digestive stimulants | Lalitha et al. (2011) |
| 4-terpineol | digestive stimulants | Lin et al. (2000) |
| Humulene | Antibacterial, anti-helminthic | Loizou et al. (2010) |
| Caryophyllene | digestive stimulants | Alagbe et al. (2022) |
| Copaene | Antimicrobial | Wild (1994) |
| Cis-linaloxide | Antifungal | Xu and Chang (2007) |
| α-Selinene | Antioxidant, anti-inflammatory | Shai et al. (2008) |
| γ-Elemene | digestive stimulants | Shaheen et al. (2005) |
| α-Gurjunene | Immuno-stimulatory, antioxidant | Özçelik et al. (2011) |
| β-Elemene | Antimicrobial, antioxidant, digestive stimulants | Singh et al. (2023); Omokore and Alagbe (2019) |
| β-Cyclocitral | Anti-inflammatory, digestive stimulants | Kolapo et al. (2009) |
| 3-hexenyl-2-methylbutanoate | Anti-bacterial | Inngjerdingen et al. (2004) |
| Exo-methyl-camphenilol | Antimicrobial | Ezike et al. (2010) |
| Caryophyllene oxide | Antibacterial, antioxidant | Enright et al. (2002) |
| Benzene -1-methoxy-2-methyl | Antifungal | Fabricant and Farnsworth (2001) |
| Napthalene, 1,2 hydro-1,1, 6 –trimethyl | Antimicrobial | Inngjerdingen et al. (2004) |
| Cycloheptasiloxane, tetradecamethyl | Anti-inflammatory | Chua (2013); Daniel et al. (2023) |
| Nerolidyl acetate | Anti-inflammatory | Azmir et al. (2013) |
| 1-Cyclohexene-1- butanal, alpha, 2, 6, 6-tetramethyl | Anti-inflammatory | Atanasov et al., 2015; Alagbe (2024) |

Source: Adapted from (Alagbe, 2023)

## 3. PHARMACOLOGICAL PROPERTIES OF PROSOPISAFRICANA

### I. Antimicrobial Properties

A study carried out by Badri et al. (2016) showed that essential oils from Prosopis africana have the capacity to inhibit the activities of some bacteria such as Shigella flexneri, Salmonella typhi, Proteus mirabilis, Pseudomonas aeruginosa, Klebsiella pneumoniae, Enterococcus faecalis, Listeria monocytogenes, and Bacillus cereus due to the presence of phytochemicals like phenols, tannins, and flavonoids, which have been suggested to perform multiple biological activities against pathogenic organisms. Other studies have shown that methanolic and ethanolic extracts from the leaves, stem bark, and roots of Prosopis are effective against Salmonella typhi, Proteus mirabilis, Enterococcus faecalis, and Listeria monocytogenes (Tajbakhsh et al., 2015).

### II. The Properties of Antioxidants

Extracts that are aqueous, methanolic, and ethanolic have a number of medicinal uses and have the ability to scavenge free radicals that can lead to infections in an animal's body (Darogba et al., 2009). Additionally, according to Saad et al. (2017), prosopis oil demonstrated efficacy against Erwinia spp., C. albicans, P. vulgaris, E. coli, and Shigella spp. Additionally, research has demonstrated that adding 400 mg/kg of prosopis oil to broiler diets can have a major impact on the activity of enzymes such as glutathione peroxidase, catalase, reduced glutathione, superoxide dismutase, and malondialdehyde (Alagbe, 2023).

### III. Hypolipidemic Characteristics

According to earlier research by Rahman et al. (2011), Prosopis extracts administered to rats at doses of up to 600 mg/kg significantly affected their levels of triglycerides, total cholesterol, low density lipoprotein, very low density lipoprotein, and atherogenic index (meat safety index), indicating a potential preventive effect against cardiovascular diseases. Additionally, according to Alagbe et al. (2022), adding 800 mg/kg of Prosopis africana essential oil to the diet can lower the amount of saturated fat and raise the amount of polyunsaturated fat in broiler meat (Alagbe, 2022).

Table 2 shows the Effects of Prosopisafricana extracts on the general performance of birds.

**Table 2: Effects of *Prosopisafricana*extracts on the general performance of birds**

| Type of extracts used | Concentration used | Effect on broilers |
|---|---|---|
| *Prosopisafricana* essential oil | 400 mg/kg | Increased body weight gain |
| | | Increased pancreatic enzyme production |
| | | Improved nutrient utilization |
| | | Increased feed intake |
| | | Reduced mortality rate |

| Type of extracts used | Concentration used | Effect on broilers |
|---|---|---|
| | | Increased in carcass weight |
| | | ----- |
| *Prosopisafricana* essential oil | 800 mg/kg | Increased production of polyunsaturated fatty acid in meat sample |
| | | Improved sensory attributes of meat |
| | | Increased in pack cell volume, red blood cell, haemoglobin, white blood cell, total protein, amongst others |
| | | Scavenging free radicals |
| | | Increasing antibody titres in birds after vaccination |
| | | Decrease in population of pathogenic microorganisms |
| *Prosopisafricana* stem bark extracts (aqueous) | 8ml/litre of water | Increased final body weight gain, feed intake and *Lactobacillus sp* count in the gut of broilers |

## 4. CONCLUSIONS

Medicinal plants are infinitely capable of producing phytochemicals, which are chemical substances with a vast range of possible benefits, all aimed at improving bird performance. Prosopis africana extracts contain several substances, which have been demonstrated to be safe, efficacious, and environmentally benign. Additionally, studies have revealed that P. africana extract includes considerable amounts of phenols, tannins, alkaloids, terpenoids, and flavonoids, which provide them antibacterial, antioxidant, antifungal, antiviral, and other beneficial properties. Additionally, adding nutrition to broiler feed may improve the birds' performance. These will increase the output of chickens and lower the rising number of antibiotic-resistant illnesses.

**Ethical approval**

All international standards have been adopted and compliance.

**Informed consent**

The study was conducted with equal participation by all authors.

**Conflicts of interests**

The authors declare that there are no conflicts of interests.

**Funding**

The study has not received any external funding.

**Data and materials availability**

All data associated with this study are present in the paper.

## REFERENCESANDNOTES


1. Ajiboye, A., Agboola, D., Fadimu, O., & Afolabi, A. (2013). Antibacterial, Phytochemical and Proximate Analysis of Prosopisafricana (Linn) Seed and Pod Extract. FUTA Journal of Research in Sciences 9 (1), 101-109
2. Alagbe, J.O (2023a). Investigating the effects of dietary supplementation of Eucalyptus camaldulensis essential oil on the growth performance, nutrient digestibility and caecal fermentation of weaned rabbits. Research in: Agricultural and Veterinary Sciences, 7(3): 139 - 148.
3. Alagbe, J.O (2023b). Investigating the effects of dietary supplementation of Eucalyptus camaldulensis essential oil on haemato-biochemical indices, immune response and oxidative stress of weaned rabbits. International Journal of Agriculture and Animal Production, 4(1): 34 - 46.
4. Alagbe, J. O. (2023c). Sensory evaluation and fatty acid composition of broiler chickens fed diets containing Prosopisafricana oil. Journal of Healthcare and Biomedical Science, 1(2): 36-45.
5. Alagbe, J. O. (2023d). Impact of dietary supplementation of Carica papaya essential oil on the blood chemistry of broiler chickens. Food Science & Applied Microbiology Reports, 2(2), 58-67.
6. Alagbe, J. O., Anorue, D. N., Shittu, M. D., Ramalan, S. M., Faniyi, T. O., & Ajagbe, A. D. (2024). Growth performance and physiological response of weaned pigs fed diet supplemented with novel a phytogenics. Brazilian Journal of Science, 3(1), 43-57.
7. Alagbe, J.O (2024). Novel phytogenics impact on weaned pig's growth performance, haematology and serum biochemical indicators. Black Sea Journal of Agriculture, 7(2): 1-9.
8. Alagbe, J.O. (2021). Prosopisafricana stem bark as an alternative to antibiotic feed additives in broiler chick's diets: Performance and Carcass characteristics. Journal of Multidimensional Research and Reviews, 2(1): 64-77.
9. Alagbe, J.O. (2022a). Use of medicinal plants as a panacea to poultry production and food security: A review. Gospodarka I Innowacje 22(2022): 1-12.
10. Alagbe, J.O. (2022b). Prosopisafricana (African mesquite) oil as an alternative to antibiotic feed additives on broiler chicken's diets: haematology and serum biochemical indices. Central Asian Journal of Theoretical and Applied Sciences 3(2): 19-29.
11. Alagbe, J.O. (2022c). Prosopisafricana (African mesquite) oil as an alternative to antibiotic feed additives on broiler chickens diets: performance and nutrient retention. Discovery 58(314): 134 -142.
12. Anorue, D. N., Ubong, F., & John, A. O. (2023). Investigating the effects of pawpaw (Carica papaya) essential oil dietary supplementation on the growth performance and carcass characteristics of broilers. Research in: Agricultural and Veterinary Sciences, 7(3), 164-174.
13. Atanasov, A.G., Waltenberger, B., Pferschy-Wenzig, E.M., Linder, T., Wawrosch, C., Uhrin, P., Temml, V., Wang, L., Schwaiger, S., Heiss, E.H. & Rollinger, J.M., (2015). 'Discovery and resupply of pharmacologically active plant-derived natural products: A review'. Biotechnology advances, 33(8), 1582-1614.
14. Atawodi, S. E. & Ogunbusola, F. (2009). Evaluation of Anti-Trypanosomal Properties of Four Extracts of Leaves, Stem and Root Barks of Prosopis Africana in Laboratory Animals'. Biokemistri 21 (2), 101-108.
15. Ayanwuyi, L. O., Yaro, A. H., & Abodunde, O. M. (2010) 'Analgesic and Anti-inflammatory Effects of the Methanol Stem Bark Extract of Prosopis Africana'. Pharmaceutical Biology 48 (3), 296-299
16. Azmir, J., Zaidul, I., Rahman, M., Sharif, K., Mohamed, A., Sahena, F., Jahurul, M., Ghafoor, K., Norulaini, N., & Omar, A. (2013) 'Techniques for Extraction of Bioactive Compounds from Plant Materials: A Review'. Journal of Food Engineering 117 (4), 426-436
17. Chua, L. S. (2013). A Review on Plant-Based Rutin Extraction Methods and its Pharmacological Activities. Journal of Ethnopharmacology 150 (3), 805-817
18. Cowan, M. M. (1999). Plant Products as Antimicrobial Agents. Clinical Microbiology Reviews 12 (4), 564-582
19. Cushnie, T. T. & Lamb, A. J. (2005). Antimicrobial Activity of Flavonoids. International Journal of Antimicrobial Agents 26 (5), 343-356.
20. Dai, J. & Mumper, R. J. (2010). Plant Phenolics: Extraction, Analysis and their Antioxidant and Anticancer Properties. Molecules 15 (10), 7313-7352
21. Davies, D. G., Parsek, M. R., Pearson, J. P., Iglewski, B. H., Costerton, J. W., & Greenberg, E. P. (1998) 'The Involvement of Cell-to-Cell Signals in the Development of a Bacterial Biofilm'. Science (New York, N.Y.) 280 (5361), 295-298.
22. Edeoga, H. O., Okwu, D. E. & Mbaebie, B. O. (2005). Phytochemical constituents of some Nigerian medicinal plants. African Journal of Biotechnology 47, 685- 688.
23. Egunyomi, A. S. & Oladunjoye, S. (2012). Studies on the chemical composition and Nutritive value of the fruit of African star apple. African Journal of Agricultural Research 7 (31), 4256-42588.
24. Enright, M. C., Robinson, D. A., Randle, G., Feil, E. J., Grundmann, H., & Spratt, B. G. (2002) The Evolutionary History of Methicillin-Resistant Staphylococcus Aureus (MRSA). Proceedings of the National Academy of Sciences of the United States of America, 99 (11), 7687-7692
25. Ezike, A. C., Akah, P. A., Okoli, C. O., Udegbunam, S., Okwume, N., Okeke, C., & Iloani, O. (2010). Medicinal Plants used in Wound Care: A Study of Prosopis Africana (Fabaceae) Stem Bark. Indian Journal of Pharmaceutical Sciences 72 (3), 334-339
26. Fabricant, D. S. & Farnsworth, N. R. (2001). The Value of Plants used in Traditional Medicine for Drug Discovery'. Environmental Health Perspectives, 109 Suppl 1, 69- 75
27. Guimarães, L. G. L. (2010). ÓleosEssenciais de Lippiasidoides Cham., Alomiafastigiata (Gardner) Benth, Ocoteaodorifera (Vell.) Rohwer, Mikaniaglauca Mart. e Cordiaverbenacea D.C.: Identificação e QuantificaçãoQuímica, Caracterização das EstruturasSecretoras, AtividadesAntioxidanteAntibacteriana. Ph.D. Thesis, Universidade Federal de Lavras, Lavras, Brazil.
28. Idowu, T. O., Iwalewa, E. O., Aderogba, M. A., Akinpelu, B. A. & Ogundaini, A. O. (2006). Antinociceptive, Anti-inflammatory and Antioxidant activities of Eleagnine: An alkaloid isolated from seed cotyledon of Chrysophyllumalbidum. Journal of Biological Science 6, 1029–1034.
29. Inngjerdingen, K., Nergård, C. S., Diallo, D., Mounkoro, P. P., &d Paulsen, B. S. (2004) 'An Ethnopharmacological Survey of Plants used

for Wound Healing in Dogonland, Mali, West Africa'. Journal of Ethnopharmacology 92 (2-3), 233-244

30. Jim´enez-Arellanes, A. M., Meckes, M., Ram´ırez, R., Torres, J. & Luna-Herrera, J. (2003). Activity against multidrug-resistant Myocbacteriun tuberculosis in Mexican plants used to treat respiratory diseases. Phytotherapy Research, 17 (8), 903–908.
31. Kokoska, L., Polesny, Z., Rada, V., Nepovim, A. & Vanek, T. (2002). Screening of some Siberian medicinal plants for antimicrobial activity. Journal of Ethnopharmacology 8, 51–53.
32. Kolapo, A., Okunade, M., Adejumobi, J., & Ogundiya, M. (2009). Phytochemical Composition and Antimicrobial Activity of Prosopis Africana against some Selected Oral Pathogens. World Journal of Agricultural Sciences, 5 (1), 90-93
33. Lalitha, V., Kiran, B. & Raveesha, K. (2011). Antifungal and antibacterial potentiality of six essential oils extracted from plant source. International Journal of Engineering Science and Technology (IJEST) 3 (4), 3029-3038.
34. Lin, F. Y. C., Ho, V. A., Bay, P. V., Thuy, N. T. T., Bryla, D., Thanh, T. C. & Ha, B. K. (2000). The epidemiology of typhoid fever in the Dong Thap Province, Mekong Delta region of Vietnam. American Journal of Tropical Medical Hygiene, 62, 644-648.
35. Loizou, S., Lekakis, I., Chrousos, G. P. & Moutsatsou, P. (2010). Beta-sitosterol exhibits anti-inflammatory activity in human aortic endothelial cells. Molecule Nutrition and Food Research 54, 551-558.
36. Omokore, E.A &and Alagbe, J.O. (2019). Effect of replacing soya bean meal with Indigoferazollingweriana leaf meal on the performance and carcass characteristic of growing rabbits. International Journal of Multidisciplinary Research and Development, 6(5), 74-77.
37. Özçelik, B., Kartal, M., & Orhan, I. (2011). Cytotoxicity, Antiviral and Antimicrobial Activities of Alkaloids, Flavonoids, and Phenolic Acids. Pharmaceutical Biology, 49 (4), 396-402
38. Sandasi, M., Leonard, C., & Viljoen, A. (2005). The in Vitro Antibiofilm Activity of Selected Culinary Herbs and Medicinal Plants Against Listeria Monocytogenes. Letters in Applied Microbiology 50 (1), 30-35
39. Shaheen, A. Y., Sheikh, A. A., Rabbani, M., Aslam, A., Bibi, T., Liaqat, F., Muhammad, J., & Rehmani, S. F. (2015). Antibacterial Activity of Herbal Extracts Against Multi-Drug Resistant Escherichia Coli Recovered from Retail Chicken Meat. Pakistan Journal of Pharmaceutical Science 28 (4), 1295-1300
40. Shai, L., McGaw, L., Masoko, P., & Eloff, J. (2008). Antifungal and Antibacterial Activity of Seven Traditionally used South African Plant Species Active Against Candida Albicans. South African Journal of Botany 74 (4), 677-684
41. Sharma, S., John, A. O., Xing, L., Ram, S., & Amita, K. (2022). Comparative Analysis of Ethanolic Juniperus Thurifera Leaf, Stem Bark and Root Extract Using Gas Chromatography and Mass Spectrometry. International Journal of Agriculture and Animal Production (IJAAP) 2(06), 18-27.
42. Singh, J. (2008). Maceration, Percolation and Infusion Techniques for the Extraction of Medicinal and Aromatic Plants. Extraction Technologies for Medicinal and Aromatic Plants 67, 32-35.
43. Wild, R. (1994). The Complete Book of Natural and Medicinal Cures. Emmaus PA: Rodale Press
44. Xu, B. & Chang, S. (2007). A Comparative Study on Phenolic Profiles and Antioxidant Activities of Legumes as Affected by Extraction Solvents. Journal of Food Science 72 (2), S159-S166.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of Sherwan Journals and/or the editor(s). Sherwan Journals and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.