

Taxonomic novelty in Pleomonodictydaceae and new reports for *Ampelomyces quisqualis* (Phaeosphaeriaceae), *Melomastia maolanensis* and *M. oleae* (Pleurotremataceae)

Digvijayini Bundhun^{1,2,3}, E. B. Gareth Jones⁴, Ruvishika S. Jayawardena^{2,3}, Erio Camporesi^{5,6},
Dhanushka N. Wanasinghe^{7,8}, Indunil C. Senanayake⁹, Vinodhini Thiyagaraja¹, Kevin D. Hyde^{1,2,3}

¹ Key Laboratory of Phytochemistry and Natural Medicines, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming, Yunnan 650201, China

² School of Science, Mae Fah Luang University, Chiang Rai 57100, Thailand

³ Center of Excellence in Fungal Research, Mae Fah Luang University, Chiang Rai 57100, Thailand

⁴ Department of Botany and Microbiology, College of Science, King Saud University, P.O. Box 2455, Riyadh 11451, Saudi Arabia

⁵ Gruppo Micologico Forlivese "Antonio Cicognani", Via Roma 18, Forlì, Italy

⁶ A.M.B. Circolo Micologico "Giovanni carini", C.P. 314, Brescia, Italy

⁷ Center for Mountain Futures, Kunming Institute of Botany, Chinese Academy of Sciences, Honghe County 654400, China

⁸ Department of Soil Science, College of Food and Agriculture Sciences, King Saud University, P.O. Box 145111, Riyadh 11362, Saudi Arabia

⁹ Germplasm Bank of Wild Species & Yunnan Key Laboratory for Fungal Diversity and Green Development, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming, 650201, China

Corresponding author: Kevin D. Hyde (kdhyde3@gmail.com)

This article is part of:

**Exploring the Hidden Fungal Diversity:
Biodiversity, Taxonomy, and Phylogeny
of Saprobiic Fungi**

Edited by Samantha C. Karunarathna,
Danushka Sandaruwan Tennakoon,
Ajay Kumar Gautam



Academic editor:

Samantha C. Karunarathna

Received: 24 August 2024

Accepted: 26 October 2024

Published: 17 December 2024

Copyright: © Digvijayini Bundhun et al.

This is an open access article distributed under
terms of the Creative Commons Attribution
License (Attribution 4.0 International – CC BY 4.0).

Abstract

This study introduces a novel genus *Robiniigena*, with its type *R. hyalinospora*. The specimen was collected on dead aerial branches of *Robinia pseudoacacia* in Italy. Based on the examination of morphology and the results of phylogenetic analyses involving nuclear 18S rDNA (SSU), nuclear 28S rDNA (LSU), nuclear rDNA ITS1-5.8S-ITS2 (ITS), translation elongation factor 1- α (*tef1- α*) and RNA polymerase II second largest subunit (*rpb2*) sequences, *Robiniigena* is referred to the family Pleomonodictydaceae (Pleosporales). It is characterized by immersed to erumpent, ostiolate ascomata, filiform, septate and cellular pseudoparaphyses, bitunicate, clavate to cylindric-clavate asci and fusiform, hyaline ascospores surrounded by a mucilaginous sheath. This research also establishes the taxonomic placement of the previously unclassified *Inflatispora* (Pleosporales genus *incertae sedis*) within the Pleomonodictydaceae. The sexual morph of *Ampelomyces quisqualis* (Phaeosphaeriaceae) is described for the first time and it is characterized by immersed, perithecial ascomata, a peridium comprising two layers, branched, septate and filiform pseudoparaphyses, short-pedicellate, bitunicate asci with an ocular chamber and sub-hyaline, fusiform, septate ascospores. This species, previously known only in its asexual morph, has been found as a saprobe on *Sonchus* sp. in Italy. Our identification of the sexual morph was based on LSU rDNA and ITS rDNA sequence data. *Melomastia maolanensis* (Pleurotremataceae) is reported for the first time in Thailand, collected from *Chromolaena odorata*, while *M. oleae* is documented as a new record from *Duranta erecta* in Thailand.

Key words: 2 new taxa, Dothideomycetes, Dyfrolomycetales, fungal diversity, morphology

Citation: Bundhun D, Jones EBG, Jayawardena RS, Camporesi E, Wanasinghe DN, Senanayake IC, Thiyagaraja V, Hyde KD (2024) Taxonomic novelty in Pleomonodictydaceae and new reports for *Ampelomyces quisqualis* (Phaeosphaeriaceae), *Melomastia maolanensis* and *M. oleae* (Pleurotremataceae). MycoKeys 111: 147–180. <https://doi.org/10.3897/mycokeys.111.135456>

Introduction

The family Pleomonodictydaceae was established to accommodate *Pleomonodictys*, along with two monodictys-like taxa, which formed a distinct clade within the suborder Massarineae, Pleosporales (Hernández-Restrepo et al. 2017). *Pleomonodictys* is characterized by mononematous, micronematous to semi-macronematous conidiophores, which are sometimes reduced to intercalary and polyblastic conidiogenous cells. The conidia, which vary in shape, are blastic, pleurogenous, muriform, and verrucose to tuberculate, and they are often solitary or occur in short chains (Hernández-Restrepo et al. 2017; Bao et al. 2021). The second genus, *Pleohelicoon* was established in the family to accommodate *Helicoon* taxa, which belonged to Pleosporales (Jayasiri et al. 2019). *Pleohelicoon* taxa are characterized by macronematous, mononematous, unbranched and hyaline to brown conidiophores, monoblastic and terminal conidiogenous cells and pale brown to dark fuscous conidia with multi-septate conidial filaments coiled 7–9 times in 3 planes, giving the conidia different shapes (Jayasiri et al. 2019). It is worth noting that no sexual morph has been reported for these genera.

Inflatipora was introduced by Zhang et al. (2011) to include the only taxon and type species, *I. pseudostromatica*, characterized by solitary to aggregated, semi-immersed or erumpent ascomata which form under a pseudostroma and open through a broad, round pore, narrow cellular pseudoparaphyses, bitunicate, cylindric-clavate asci, and hyaline, 3-septate, fusiform to cylindrical ascospores with an enlarged upper central cell and surrounded by a mucilaginous sheath (Zhang et al. 2011). *Inflatipora* shares a morphological resemblance to *Nodulosphaeria*, in terms of the swollen upper central cell of the ascospores. However, ascospores of *Nodulosphaeria* have terminal appendages, while in *Inflatipora pseudostromatica*, ascospores are surrounded by a sheath (Zhang et al. 2011). Moreover, phylogenetic analyses of nuclear 28S rDNA (LSU), nuclear 18S rDNA (SSU) and RNA polymerase II second largest subunit (*rpb2*) sequence data of *I. pseudostromatica* showed it grouped with taxa in the Massarineae, forming a separate basal lineage (Zhang et al. 2011). It was however treated as genus *incertae sedis* in the suborder owing to its uncertain familial placement (Zhang et al. 2011). A second species, *Inflatipora caryotae*, was introduced in the genus based on morpho-phylogenetic analyses (Tibpromma et al. 2017).

Ampelomyces quisqualis belongs to Phaeosphaeriaceae (Pleosporales), and is a parasite and biocontrol agent of powdery mildew (Kiss et al. 2004; Liyanage et al. 2018; Manjunatha et al. 2020). This taxon parasitizes the powdery mildew taxa by penetrating hyphae and producing pycnidia in the hyphae, conidiophores, and ascomata of the powdery mildew host, which eventually interferes with the latter's development (Kiss et al. 2004; Liyanage et al. 2018). *Ampelomyces quisqualis* is characterized by pale to dark brown, pycnidial conidiomata, with walls which are angular textured, and hyaline, ovate to elliptical, aseptate and smooth-walled conidia (Manjunatha et al. 2020). The sexual morph of the species has not been reported so far.

Melomastia, typified by *M. mastoidea* (= *M. friesii*), has been subjected to several taxonomic revisions before being finally accommodated in Dothideomycetes (Pleurotremataceae, Dyfrolomycetales) (Barr 1989, 1994; Lumbsch and Huhndorf 2010; Maharachchikumbura et al. 2015, 2016; Norphanphoun et al. 2017; Wijayawardene et al. 2017). It was earlier accommodated in Clypeosphaeriaceae

(Sordariomycetes) and was later transferred to Pleurotremataceae, the latter family then assigned to Xylariales (Barr 1989, 1994). However, the genus was then excluded from the family and classified as Ascomycota, genera *incertae sedis* (Kirk et al. 2001; Lumbsch and Huhndorf 2010). While reviewing the backbone tree for Sordariomycetes, Maharachchikumbura et al. (2015) placed *Melomastia* in Sordariomycetes genera *incertae sedis*, while Pleurotremataceae (with the single genus *Pleurotrema*) was accommodated in Chaetosphaeriaceae, Chaetosphaeriales. However, Maharachchikumbura et al. (2016) later excluded Pleurotremataceae from Sordariomycetes, since it was regarded as an earlier name of the family Dyfrolomycetaceae (Dothideomycetes) based on morphology. Meanwhile, *Melomastia* was still considered as an *incertae sedis* genus in Sordariomycetes (Maharachchikumbura et al. 2016). Norphanphoun et al. (2017) finally introduced sequence data for the *Melomastia* taxon, *M. italica* for the first time and showed that *Melomastia* belongs to Dothideomycetes, in the family Pleurotremataceae, along with *Dyfrolomyces* and *Pleurotrema*. They also synonymized *Dyfrolomyces maolanensis*, a saprobe collected from China (Zhang et al. 2017), to *Melomastia maolanensis*, owing to its closer phylogenetic affinity with *M. italica* (Norphanphoun et al. 2017). Li et al. (2022) synonymized *Dyfrolomyces* under *Melomastia* based on morphological and phylogenetic analyses. Ascospore septation, which was considered as the principal differentiating character between *Dyfrolomyces* (multi-septate) and *Melomastia* (2-septate), was no longer considered a key for delimitation of the two genera (Li et al. 2022). Kularathnage et al. (2023), however, recently reinstated *Dyfrolomyces* in Pleurotremataceae.

Aims of study

In an ongoing survey on the fungal diversity in Italy and Thailand (Chaiwan et al. 2021; Manawasinghe et al. 2022), we recovered several taxa of Dothideomycetes. We herein introduce a new genus *Robiniigena* and accommodate the *Inflatipora* genus *incertae sedis* in Pleomonodictydaceae based on morpho-phylogenetic evidence. Furthermore, the sexual morph of *Ampelomyces quisqualis* is described for the first time, based on LSU–ITS sequence data. We also introduce two new collections of *Melomastia*. *Melomastia maolanensis* is reported as a new geographical record from Thailand, while *M. oleae*, which was initially introduced from *Olea europaea* in China (Li et al. 2022), is a new host and geographical record on *Duranta erecta* in Thailand based on morphology and phylogeny. Findings herein indicate that there is still much to be discovered vis-a-vis the diversity of Dothideomycetes. Future studies in this class will undoubtedly result in further important taxonomic discoveries and advances.

Materials and methods

Sample collection, specimen examination, and isolation

Dead stems and twigs bearing fungal fruiting bodies were collected from the terrestrial environment in Italy during the dry (June) and wet (November) seasons, and in Thailand during the wet (July) and dry (November) seasons. The samples were taken to the laboratory in a plastic Ziplock bag and stored inside paper envelopes. Specimens were externally examined with a Motic SMZ 168

stereomicroscope. Free-hand sections of the sporocarps were prepared and placed on water-mounted glass slides. They were then examined using a Nikon ECLIPSE 80i compound microscope with differential interference contrast (DIC) illumination. Microscopic photography was conducted for the ascomata, peridium, pseudoparaphyses, asci, and ascospores. The images were captured with a Canon EOS 750D digital camera fitted to the microscope. Measurements were made with the Tarosoft (R) Image Frame Work v. 0.9.7. using the necessary calibration values and images used for figures were processed with Adobe Photoshop CS6 Extended v. 13.0.1 software (Adobe Systems, San Jose, California).

Single spore isolation was carried out following the method of Senanayake et al. (2020). Ascospore suspensions were made and spread on potato dextrose agar (PDA; 39 g/L) or water agar (WA; 20 g/L). The germinated ascospores were examined after 24 h, after which they were transferred to PDA media. Cultures were incubated at 25 °C in the dark and colony color was determined according to Rayner (1970) after 2–3 weeks of growth on PDA. Herbarium specimens were deposited in Mae Fah Luang University Herbarium (MFLU) while, the living cultures at the Mae Fah Luang University Culture Collection (MFLUCC) in Thailand. Index Fungorum and Faces of Fungi numbers are provided as per Index Fungorum (<https://www.indexfungorum.org>) and Jayasiri et al. (2015). Taxa descriptions and notes are also uploaded on the GMS microfungi website (<https://gmsmicrofungi.org/>) (Chaiwan et al. 2021).

DNA extraction, PCR amplification, and sequencing

Genomic DNA was extracted from mycelium or directly from fungal fruiting bodies. Fresh mycelium was scraped from cultures grown on PDA at 25 °C for two weeks. Around 15–20 ascomata were removed from the sterilized plant materials using fine sterile needles, while being observed through the stereomicroscope, and placed in 1.5 ml micro-centrifuge tubes (Wanasinghe et al. 2018). DNA was extracted using the Omega Bio-tek kit according to the manufacturer's instructions. The nuclear 18S rDNA (SSU), nuclear 28S rDNA (LSU), nuclear rDNA ITS1-5.8S-ITS2 (ITS), translation elongation factor 1-alpha (*tef1-a*), and the RNA polymerase II second largest subunit (*rpb2*) were amplified (Table 1). The total volume of the PCR reaction was 25 µL, and it consisted of 12.5 µL of 2× Power Taq PCR MasterMix, 1 µL of each primer, 2 µL genomic DNA extract, and 8.5 µL double-distilled water (ddH₂O). The PCR protocols were adapted accordingly (Table 1). Thirty-five cycles were used for the denaturation, annealing, and extension steps for each locus (except for *tef1-a* and *rpb2*, where 40 cycles were used). All the PCR thermal cycles included a final extension of 72 °C for 10 mins and a final hold at 4 °C.

Table 1. Loci, primers, and PCR amplification conditions used in the present study.

Loci	Primers (forward/reverse)	Initial Denaturation	Denaturation	Annealing	Extension	References
SSU	NS1/NS4	94 °C, 3 mins	94 °C, 30s	54 °C, 50s	72 °C, 1.30 mins	White et al. (1990)
LSU	LR0R/LR5	95 °C, 5 mins	95 °C, 45s	53 °C, 45s	72 °C, 2 mins	Vilgalys and Hester (1990)
ITS	ITS5/ITS4	95 °C, 5 mins	95 °C, 45s	53 °C, 45s	72 °C, 2 mins	White et al. (1990)
<i>tef1-a</i>	983F/2218R	94 °C, 5 mins	94 °C, 30s	58 °C, 1.30 mins	72 °C, 1.20 mins	Carbone and Kohn (1999)
<i>rpb2</i>	fRPB2-5f /fRPB2-7cR	95 °C, 5 mins	95 °C, 1 min	54 °C, 2 mins	72 °C, 1.30 mins	Liu et al. (1999), Sung et al. (2007)

PCR products were verified by staining with FluoroDye™ DNA Fluorescent Loading Dye on 1% agarose electrophoresis gels. They were then purified and sequenced using the same primers at the Tsingke Biotech Co. Ltd., Kunming, China. The quality of the sequences obtained was confirmed by checking the chromatograms using BioEdit v. 7.0 (Hall 2004), after which the sequences were assembled into contigs using SeqMan v. 7.1.0. The sequence data derived in this study have been deposited in GenBank (Table 2). New species are established based on recommendations provided by Jeewon and Hyde (2016) and Maharachchikumbura et al. (2021).

Sequence alignment and phylogenetic analyses

Generated SSU, LSU, ITS, *tef1-a* and *rpb2* sequences were subjected to BLASTn searches (<https://blast.ncbi.nlm.nih.gov>) and similar sequences were downloaded from GenBank based on BLAST similarities and following relevant papers (Tanaka et al. 2015; Phukhamsakda et al. 2016; Hongsanan et al. 2020; Tennakoon et al. 2020; Li et al. 2022; Hyde et al. 2023; Kularathnage et al. 2023; Table 2). Individual gene matrices were aligned using the default setting in MAFFT v. 7.036 (<http://mafft.cbrc.jp/alignment/server/>) (Katoh et al. 2019) and manually adjusted when necessary in BioEdit v. 7.0 (Hall 2004).

Table 2. Taxa used for the phylogenetic analyses in the present study, and their corresponding GenBank accession numbers. Type strains are denoted by ^T. Sequences derived in this study are shown in bold black.

Taxa	Strain	GenBank accession number				
		SSU	LSU	ITS	<i>tef1-a</i>	<i>rpb2</i>
<i>Acrospermum adeanum</i>	M133	EU940031	EU940104	–	–	–
<i>Acrospermum compressum</i>	M151	EU940012	EU940084	–	–	–
<i>Acrospermum graminum</i>	M152	EU940013	EU940085	–	–	–
<i>Ampelomyces quisqualis</i>	AMP	–	–	OP740825	–	–
<i>Ampelomyces quisqualis</i>	CBS 129.79	EU754029	EU754128	KY090653	–	–
<i>Ampelomyces quisqualis</i>	BRIP 72107	–	–	MZ054399	–	–
<i>Ampelomyces quisqualis</i>	CBS 131.79	–	MH872956	MH861188	–	–
<i>Ampelomyces quisqualis</i>	CBS 128.79	–	MH872954	MH861185	–	–
<i>Ampelomyces quisqualis</i>	CBS 133.32	–	MH866692	MH866692	–	–
<i>Ampelomyces quisqualis</i>	CBS 131.31	–	MH866605	AF035781	–	–
<i>Ampelomyces quisqualis</i>	HMLAC05119	–	OL739255	OL739255	–	–
<i>Ampelomyces quisqualis</i>	Chillan	–	–	MH997723	–	–
<i>Ampelomyces quisqualis</i>	CBS 130.79	–	–	HQ108039	–	–
<i>Ampelomyces quisqualis</i>	SMKC 22437	–	–	GQ324146	–	–
<i>Ampelomyces quisqualis</i>	SMKC 22438	–	–	GQ324128	–	–
<i>Ampelomyces quisqualis</i>	MFLU 23-0142	–	PP751625	PP751509	PP782197	–
<i>Anisomeridium phaeospermum</i>	MPN539	JN887374	JN887394	–	JN887418	–
<i>Anisomeridium ubianum</i>	MPN94	JN887379	–	–	JN887421	–
<i>Anthosulcatispora subglobosa</i>	MFLUCC 17-2065 ^T	MT226705	MT214592	MT310636	MT394649	MT394706
<i>Aquadictyospora lignicola</i>	MFLUCC 17-1318 ^T	–	MF948629	MF948621	MF953164	–
<i>Aquastroma magniostiolata</i>	HHUF 30122 ^T	AB797220	AB807510	LC014540	AB808486	–
<i>Asteromassaria pulchra</i>	CBS 124082	GU296137	GU301800	–	GU349066	GU371772
<i>Bambusicola aquatica</i>	MFLUCC 18-1031 ^T	MT864293	MN913710	MT627729	MT954392	MT878462
<i>Brunneomurispora lonicerae</i>	KUMCC 18-0157 ^T	MK356360	MK356346	MK356373	MK359065	–

Taxa	Strain	GenBank accession number				
		SSU	LSU	ITS	<i>tef1-α</i>	<i>rpb2</i>
<i>Clypeolocus akitaensis</i>	HHUF 27557 ^T	AB797253	AB807543	AB809631	AB808519	–
<i>Didymosphaeria rubi-ulmifolii</i>	MFLUCC 16-1000	MT226672	MT214555	MT310602	MT394734	–
<i>Dyrolomyces chromolaenae</i>	MFLUCC 17-1434 ^T	MT214413	–	–	MT235800	–
<i>Dyrolomyces tiomanensis</i>	MFLUCC 13-0440 ^T	KC692155	KC692156	–	KC692157	–
<i>Falciformispora lignatilis</i>	BCC 21117	GU371834	GU371826	KF432942	GU371819	–
<i>Flavomyces fulophazii</i>	CBS 135761 ^T	KP184082	KP184040	KP184001	–	–
<i>Fuscostagonospora camporesii</i>	MFLU 16-1362 ^T	MN750605	MN750590	MN750611	–	–
<i>Fuscostagonospora cytisi</i>	MFLUCC 16-0622 ^T	KY770977	KY770978	–	KY770979	–
<i>Fuscostagonospora sasae</i>	HHUF 29106 ^T	AB797258	AB807548	AB809636	AB808524	–
<i>Halobyssothecium estuariae</i>	MFLUCC 19-0386 ^T	MN598868	MN598871	MN598890	MN597050	–
<i>Halomassarina thalassiae</i>	BCC 17055	GQ925843	GQ925850	–	–	–
<i>Helicoon</i> sp.	JAC9590	–	–	MK432689	–	–
<i>Helminthosporium microsorum</i>	CBS 136910 ^T	KY984427	KY984329	KY984329	KY984448	KY984390
<i>Inflatispora caryotae</i>	MFLUCC 13-0825 ^T	KY264751	KY264747	KY264743	–	–
<i>Inflatispora pseudostromatica</i>	CBS 123110 ^T	JN231132	JN231131	–	–	JN231133
<i>Kaseifertia cubense</i>	CBS 680.96	AB797218	AB807508	LC014541	AB808484	–
<i>Katumotoa bambusicola</i>	HHUF 28661 ^T	AB524454	AB524595	LC014560	AB539108	AB539095
<i>Kazuakitanaka yuxiensis</i>	HKAS 122924 ^T	ON009092	ON009108	ON009124	ON009267	ON009290
<i>Latorua caligans</i>	CBS 576.65 ^T	–	MH870362	MH858723	–	–
<i>Lentithecium fluviale</i>	CBS 122367	GU296158	GU301825	–	GU349074	–
<i>Leptosphaeria doliolum</i>	CBS 505.75 ^T	GU296159	GU301827	JF740205	GU349069	KY064035
<i>Leucaenicola aseptata</i>	MFLUCC 17-2423 ^T	MK347853	MK347963	MK347746	MK360059	MK434891
<i>Longipedicellata megafusiformis</i>	MFLU 21-0062 ^T	–	MZ538546	MZ538512	MZ567090	–
<i>Macrodiplodiopsis desmazieri</i>	CPC 24971 ^T	–	KR873272	KR873240	–	–
<i>Massarina cisti</i>	CBS 266.62 ^T	AB797249	AB807539	LC014568	AB808514	–
<i>Melomastia beihaiensis</i>	KUMCC 21-0084 ^T	MZ727002	MZ726990	–	OK043822	–
<i>Melomastia clematidis</i>	MFLUCC 17-2092 ^T	MT226718	MT214607	–	MT394663	–
<i>Melomastia distoseptata</i>	NFCCI: 4377 ^T	–	MH971236	–	–	–
<i>Melomastia distoseptata</i>	MFLUC 21-0102	–	MT860427	–	–	–
<i>Melomastia fulvicomae</i>	MFLUCC 17-2083 ^T	MT226719	MT214608	–	MT394664	–
<i>Melomastia fusispora</i>	CGMCC 3.20618	OK623494	OK623464	–	OL335189	–
<i>Melomastia fusispora</i>	UESTCC 21.0001	OK623495	OK623465	–	OL335190	–
<i>Melomastia italica</i>	MFLUCC 15-0160 ^T	MG029459	MG029458	–	–	–
<i>Melomastia loropetalicola</i>	ZHKUCC 22-0174 ^T	OQ379411	OQ379412	–	–	–
<i>Melomastia maolanensis</i>	GZCC 16-0102 ^T	KY111906	KY111905	–	KY814762	–
<i>Melomastia maolanensis</i>	MFLU 23-0143	PP751954	PP751616	–	PP782198	–
<i>Melomastia neothailandica</i>	MFLU 17-2589 ^T	–	MN017857	–	–	–
<i>Melomastia oleae</i>	CGMCC 3.20619 ^T	OK623496	OK623466	–	OL335191	–
<i>Melomastia oleae</i>	UESTCC 21.0003	OK623497	OK623467	–	OL335192	–
<i>Melomastia oleae</i>	UESTCC 21.0005	OK623498	OK623468	–	OL335193	–
<i>Melomastia oleae</i>	UESTCC 21.0006	OK623499	–	–	OL335194	–
<i>Melomastia oleae</i>	MFLUCC 23-0086	PP751953	PP751621	–	PP782199	–
<i>Melomastia phetchaburiensis</i>	MFLUCC 15-0951 ^T	MF615403	MF615402	–	–	–
<i>Melomastia puerensis</i>	ZHKUCC 23-0802 ^T	OR922340	OR922309	–	OR966284	–
<i>Melomastia puerensis</i>	ZHKUCC 23-0803	OR922341	OR922310	–	OR966285	–
<i>Melomastia pyriformis</i>	ZHKUCC 22-0175	OP739334	OP791870	–	OQ718392	–
<i>Melomastia rhizophorae</i>	BCC 15481	KF160009	–	–	–	–
<i>Melomastia rhizophorae</i>	JK 5456A	GU479766	GU479799	–	GU479860	–
<i>Melomastia septata</i>	MFLUCC 22-0112 ^T	–	OP749870	–	OP760198	–

Taxa	Strain	GenBank accession number				
		SSU	LSU	ITS	<i>tef1-α</i>	<i>rpb2</i>
<i>Melomastia sichuanensis</i>	CGMCC 3.20620	OK623500	OK623469	–	OL335195	–
<i>Melomastia sichuanensis</i>	UESTCC 21.0008	OK623501	OK623470	–	OL335196	–
<i>Melomastia sinensis</i>	MFLUCC 17-1344 ^T	MG836700	MG836699	–	–	–
<i>Melomastia sinensis</i>	MFLUCC 17-2606	–	OL782048	–	OL875098	–
<i>Melomastia thailandica</i>	MFLUCC 15-0945 ^T	KX611367	KX611366	–	–	–
<i>Melomastia thamplaensis</i>	AND12	OL700222	OL457709	–	–	–
<i>Melomastia thamplaensis</i>	MFLUCC 15-0635 ^T	KX925436	KX925435	–	KY814763	–
<i>Melomastia winteri</i>	CGMCC 3.20621	OK623502	OK623471	–	OL335197	–
<i>Montagnula camporesii</i>	MFLUCC 16-1369 ^T	MN401744	MN401742	MN401746	MN397908	MN397909
<i>Morosphaeria velatispora</i>	PUFD25	MK026765	MK026764	MK026766	MN532688	MN532683
<i>Muyocopron castanopsis</i>	MFLUCC 14-1108	KU726968	KU726965	–	MT136753	–
<i>Muyocopron dipterocharpi</i>	MFLU 17-2608	KU726969	KU726966	–	MT136754	–
<i>Muyocopron heveae</i>	MFLUCC 17-0066 ^T	MH986828	MH986832	–	–	–
<i>Muyocopron lithocarpi</i>	MFLUCC 14-1106 ^T	KU726970	KU726967	–	MT136755	–
<i>Neoaquastroma bauhiniae</i>	MFLU 17-0002 ^T	MH023315	MH023319	MH025952	MH028247	MH028251
<i>Neohelicascus aquaticus</i>	MFLUCC 10-0918	KC886638	KC886640	KC886639	–	–
<i>Neokalmusia jonahhulmei</i>	KUMCC 21-0818 ^T	ON007048	ON007039	ON007043	ON009133	ON009137
<i>Neosetophoma lonicerae</i>	KUMCC 18-0155 ^T	MK356363	MK356349	MK356375	MK359067	–
<i>Palawania thailandensis</i>	MFLICC 14-1121	–	KY086494	–	–	–
<i>Palawania thailandensis</i>	MFLU 16-1873	KY086495	KY086493	–	–	–
<i>Palmiascoma gregariascomum</i>	MFLUCC 11-0175	KP753958	KP744495	KP744452	–	KP998466
<i>Parabambusicola bambusina</i>	KT 2637	AB797248	AB807538	LC014580	AB808513	–
<i>Periconia cortaderiae</i>	MFLUCC 15-0457 ^T	KX986345	KX954401	KX965732	KY310703	–
<i>Periconia pseudodigitata</i>	KT1395 ^T	AB797274	AB807564	LC014591	AB808540	–
<i>Pleohelicon fagi</i>	MFLUCC 17-2538	MK347925	MK348036	MK347816	–	MK434851
<i>Pleohelicon fagi</i>	MFLUCC 15-0182 ^T	MK347926	MK348037	MK347817	–	MK434853
<i>Pleohelicon richonis</i>	CBS 282.54	AY856952	–	MH857332	–	–
<i>Pleomonodictys capensis</i>	DLUCC:1323	–	MZ420757	MZ420742	–	MZ442696
<i>Pleomonodictys capensis</i>	HR 1	AB797261	AB807551	LC014570	AB808527	–
<i>Pleomonodictys capensis</i>	CBS 968.97 ^T	–	KY853521	KY853460	–	–
<i>Pleomonodictys descalsii</i>	FMR_12716 ^T	–	KY853522	KY853461	–	–
<i>Pseudoasteromassaria aquatica</i>	MFLUCC 18-1397 ^T	MT864322	MN913721	MT627674	MT954378	–
<i>Pseudochaetosphaeronema kunmingense</i>	KUMCC 19-0215 ^T	MN792814	MN792815	MN792812	MN794017	–
<i>Pseudochaetosphaeronema larense</i>	CBS 640.73 ^T	KF015652	KF015611	KF015656	KF015684	KF015706
<i>Pseudocoleophoma calamagrostidis</i>	KT3284 ^T	LC014604	LC014609	LC014592	LC014614	–
<i>Pseudosplanchnonema phorcioides</i>	CBS 122935	KY984434	KY984360	KY984360	KY984467	KY984418
<i>Pseudoxylomyces elegans</i>	KT 2887	AB797308	AB807598	LC014593	AB808576	–
<i>Robiniigena hyalinospora</i>	MFLU 23-0141^T	PP759387	PP759385	PP759383	PP782200	PP476210
<i>Robiniigena hyalinospora</i>	MFLUCC 23-0074^T	PP759386	PP759384	PP751493	PP782201	PP476211
<i>Stemphylium vesicarium</i>	CBS 191.86 ^T	DQ247812	DQ247804	MH861935	DQ471090	KC584471
<i>Stigmatodiscus labiatus</i>	CBS 144700 ^T	MH756065	MH756065	–	MH756083	–
<i>Stigmatodiscus oculatus</i>	CBS 144Z01 ^T	–	MH756069	–	MH756086	–
<i>Stigmatodiscus pruni</i>	CBS 142598	KX611110	KX611110	–	KX611111	–
<i>Submersispora variabilis</i>	MFLUCC 17-2360 ^T	MT864310	MN913682	–	–	–
<i>Sulcatispora acerina</i>	KT2982 ^T	LC014605	LC014610	LC014597	LC014615	–
<i>Trematosphaeria grisea</i>	CBS 332.50 ^T	KF015641	KF015614	KF015662	KF015698	KF015720
<i>Triseptata sexualis</i>	MFLUCC 11-0002 ^T	MN977850	MN977833	MN977832	–	–
<i>Vikalpa grandispora</i>	KUNCC 22-12425 ^T	OP526628	OP526648	OP526638	OP542240	–

Maximum likelihood (ML) and Bayesian posterior probability (BYPP) analyses were conducted using individual and combined loci datasets. The sequence alignments were converted from FASTA into PHYLIP format using the ALTER (alignment transformation environment, <http://www.singgroup.org/ALTER/>) bioinformatics web tool (Glez-Peña et al. 2010) before conducting the ML analyses. The latter were then performed in RAxML-HPC2 on XSEDE (v. 8.2.10) (Stamatakis 2014) with the GTRGAMMA substitution model and bootstrapping with 1000 replicates in the CIPRES Science Gateway (Miller et al. 2010) to generate ML trees.

The BYPP analyses were generated using Markov Chain Monte Carlo sampling in MrBayes v. 3.1.2 (Huelsenbeck and Ronquist 2001; Zhaxybayeva and Gogarten 2002). MrModeltest 2.3 (Nylander 2004) was initially used to estimate the best evolutionary models for each locus under the Akaike Information Criterion (AIC) implemented in PAUP v. 4.0b10 (Swofford 2002). Six simultaneous Markov chains were then run for the necessary number of generations, as mentioned under the phylogenetic trees. Trees were sampled every 1000th generation. Convergence was declared when the standard deviation of split frequencies was less than 0.01. The first 20% of generated trees, representing the burn-in phase, were discarded. The remaining 80% of trees were used to calculate the posterior probabilities in the majority rule consensus tree.

The resulting trees were viewed in FigTree v. 1.4.0 (Rambaut 2012) and modified in Microsoft PowerPoint. The ML bootstrap values (ML BS) equal to or greater than 60% and BYPP equal to or greater than 0.95 are given above or below the branches in each phylogenetic tree. The final alignments were deposited in TreeBASE, with the submission IDs mentioned under the respective phylogenetic trees.

Results

Phylogenetic analyses

Analysis 1

The combined SSU–LSU–ITS–*tef1-α*–*rpb2* dataset for the suborder Massariineae comprised 61 strains, including the strains MFLU 23-0141 and MFLUCC 23-0074 collected in the present study and the outgroup taxa *Leptosphaeria doliolum* (CBS 505.75) and *Stemphylium vesicarium* (CBS 191.86). The concatenated alignment consisted of 5220 characters. The ML and BYPP trees were almost similar in topology and did not differ significantly. The best RAxML tree with a final likelihood value of -50756.398321 was obtained and is illustrated in Fig. 1. The alignment had 2542 distinct alignment patterns, and 39.0% of gaps or undetermined characters were present. Estimated base frequencies were as follows: A = 0.241009, C = 0.249659, G = 0.270618, T = 0.238714 and the substitution rates were: AC = 1.287337, AG = 3.234605, AT = 1.329219, CG = 1.078417, CT = 6.074542, GT = 1.0. The gamma distribution rate parameter was 0.243991, while the Tree length was 6.343450.

Robiniigena hyalinospora MFLU 23-0141 and MFLUCC 23-0074 grouped in Pleomonodictydaceae with 99% ML BS, 1.00 BYPP statistical support (Fig. 1). They were basal in the family, shared by '*Helicoon*' sp. (JAC9590), *Inflatipora caryotae*




Figure 1. Phylogram generated from maximum likelihood analysis based on SSU–LSU–ITS– *tef1*–*rpb2* combined data-set for the families in Massarinaceae. The evolutionary model GTR+I+G was applied as the best-fit model for all the loci. The BYPP analysis was run for 3 million generations. The tree is rooted with *Leptosphaeria doliolum* (CBS 505.75) and *Stemphylium vesicarium* (CBS 191.86). Maximum likelihood bootstrap, ML BS ($\geq 60\%$), and BYPP (≥ 0.95) supports are respectively shown above or below the branches. Type strains are in bold black, and the novel taxon is in bold red. (-) indicate ML BS $< 60\%$ or BYPP < 0.95 . TreeBASE submission ID: 31443.




Figure 1. Continued.

(MFLUCC 13-0825), *I. pseudostromatica* (CBS 123110), *Pleohelicoon fagi* (MFLUCC 15-0182, MFLUCC 17-2538), *P. richonis* (CBS 282.54), *Pleomonodictys capensis* (CBS 968.97, DLUCC:1323, HR 1) and *Pl. descalsii* (FMR 12716). *Inflatipora caryotae* and *I. pseudostromatica* grouped in a subclade with 87% ML BS, and 1.00 BYPP support values. The two strains of *Inflatipora* are grouped with the three strains of *Pleohelicoon* and the strain '*Helicoon*' sp. (JAC9590) with 54% ML BS statistical support. The four strains of *Pleomonodictys* formed a subclade, sister to the '*Helicoon*' sp., *Inflatipora*, and *Pleohelicoon* taxa.

Analysis 2

The LSU–ITS matrix for *Ampelomyces* comprised 14 strains, including our strain MFLU 23-0142 and the outgroup taxon *Neosetophoma lonicerae* (KUMCC 18-0155). The concatenated alignment consisted of 1360 characters. The ML and BYPP trees were similar in topology and did not differ significantly. The best RAxML tree with a final optimization likelihood value of -3245.931378 was yielded and is presented below (Fig. 2). The alignment had 178 distinct patterns and 27.7% of gaps or undetermined characters. Estimated base frequencies were as follows: A = 0.247457, C = 0.216767, G = 0.254577, T = 0.281199 and the substitution rates were: AC = 4.277905, AG = 11.081952, AT = 4.397096, CG = 3.194723, CT = 27.614052, GT = 1.0. The gamma distribution rate parameter was 0.265280 while the Tree-length was 0.291945.

Strain MFLU 23-0142 nested with *Ampelomyces quisqualis* (AMP, Chillan, BRIP 72107 and CBS 133.32) in a subclade (A) with 100% ML BS, 1.00 BYPP support (Fig. 2).




Figure 2. Phylogenetic tree generated from RaxML analysis, based on LSU–ITS matrix for *Ampelomyces*. The evolutionary model GTR+I was applied for LSU, while GTR+I+G was the best-fit model for ITS. The BYPP analysis was run for 1 million generations. The tree is rooted with *Neosetophoma lonicerae* (KUMCC 18-0155). Maximum likelihood bootstrap, ML BS ($\geq 60\%$), and BYPP (≥ 0.95) supports are shown above the branches, respectively. The type strain is in bold, and the newly collected strain is in bold red. TreeBASE submission ID: 31444.

Analysis 3

An SSU–LSU–*tef1-α* dataset for Dyfrolomycetales and related orders and families, consisting of 48 strains, was used for analyzing the two strains MFLU 23-0143 and MFLUCC 23-0086 collected in the present study. The outgroup taxa used were *Anisomeridium phaeospermum* (MPN539) and *A. ubianum* (MPN94). The ML and BYPP trees were almost similar in topology and did not differ significantly. The best RaxML tree with a final likelihood value of -13964.812497 was yielded and is presented below (Fig. 3). The matrix had 976 distinct alignment patterns, with 23.70% of gaps or undetermined characters. Estimated base frequencies were as follows: A = 0.241398, C = 0.259224, G = 0.291514, T = 0.207864 and the substitution rates were: AC = 0.884199, AG = 2.067368, AT = 1.098051, CG = 0.895791, CT = 7.809461, GT = 1.0. The gamma distribution rate parameter was 0.210247 while the Tree-length was 1.259771.

The strain MFLU 23-0143 clustered with *Melomastia maolanensis* (GZCC 16-0102) with 100% ML BS, 1.00 BYPP statistical support. The isolate MFLUCC 23-0086 was sister to *M. oleae* (CGMCC3.20619, UESTCC 21.0003, UESTCC 21.0005, UESTCC 21.0006) with 55% ML BS and 0.67 BYPP support values (Fig. 3).

Taxonomy

In this section, we follow the classification of Wijayawardene et al. (2022) and revise it as needed. The amended description for Pleomonodictydaceae is provided, along with notes on *Inflatispora*, which is herein accommodated in the family. Descriptions, notes and illustrations are provided for: *Robiniigena hyalinosporea* gen. et sp. nov., *Ampelomyces quisqualis*, *Melomastia maolanensis* and *M. oleae*.

Pleomonodictydaceae Hern.-Restr., J. Mena & Gené, Stud. Mycol. 86: 76 (2017), amended

Index Fungorum: IF820279

Facesoffungi Number: FoF08344

Description. *Saprobic* on woody substrates in terrestrial and aquatic habitats. **Sexual morph:** *Ascomata* solitary or in groups, often growing under or in a pseudostroma, immersed, semi-immersed to erumpent, perithecial, brown to dark brown, ostiolate. **Ostiole** neck papillate. **Peridium** multi-layered, outer layer usually fusing with the stroma or host tissue. **Pseudoparaphyses** hyaline, filiform, cellular. **Asci** bitunicate, 8-spored, clavate to cylindric-clavate, pedicellate, with an ocular chamber. **Ascospores** 1–3-seriate, fusiform to cylindrical, hyaline, septate, surrounded by a mucilaginous sheath. **Asexual morph:** hyphomycetous. See Hernández-Restrepo et al. (2017), Jayasiri et al. (2019), Bao et al. (2021) for further details.

Type genus. *Pleomonodictys* Hern.-Restr., J. Mena & Gené

Notes. Pleomonodictydaceae has so far accommodated the two hyphomycetous genera, *Pleohelicoon* and *Pleomonodictys* (Hernández-Restrepo et al. 2017; Jayasiri et al. 2019), whose sexual morphs are still undetermined. In the present study, *Inflatispora* and the novel genus, *Robiniigena* are accepted in the family based




Figure 3. Phylogram generated from maximum likelihood RaxML analysis based on combined SSU, LSU, and *tef1-α* sequence data. The evolutionary model SYM+I+G was chosen as the best-fit model for SSU while GTR+I+G was applied for LSU and *tef1-α*. The BYPP analysis was run for 2 million generations. The tree is artificially rooted with *Anisomeridium phaeospermum* (MPN539) and *A. ubianum* (MPN94). Maximum likelihood bootstrap, ML BS (≥60%), and BYPP (≥0.95) are given respectively above or below the branches. Type strains are in bold. The newly generated sequences are indicated in bold red. (–) indicates ML BS < 60% or BYPP < 0.95. TreeBASE submission ID: 31445.

on the combined SSU–LSU–ITS–*tef1-α*–*rpb2* phylogenetic analyses (Fig. 1). Since both genera are known in their sexual morphs, the description of Pleomonodictydaceae is amended to include their sexual morphological characteristics.

***Inflatispora* Y. Zhang ter, J. Fourn. & K.D. Hyde, Sydowia 63(2): 290 (2011)**

Index Fungorum: IF561844

Facesoffungi Number: FoF11812

Description and illustrations. see in Zhang et al. (2011), Tibpromma et al. (2017)

Type species. *Inflatispora pseudostromatica* Y. Zhang ter, J. Fourn. & K.D. Hyde

Notes. When Tanaka et al. (2015) conducted a revision of the Massarineae, *Inflatispora pseudostromatica* clustered with *Monodictys capensis* (now synonymized to *Pleomonodictys capensis*, Hernández-Restrepo et al. 2017) in a clade which was considered as *incertae sedis*. When the second taxon, *I. caryotae* was introduced, it was sister to *I. pseudostromatica*, and together both species clustered with *M. capensis* and *Asteromassaria pulchra* (fig. 39 in Tibpromma et al. 2017). None of these taxa had any familial placement and thus, *Inflatispora* was referred to Massarineae *incertae sedis* (Tibpromma et al. 2017). At the same time, in the phylogenetic analyses conducted by Hernández-Restrepo et al. (2017), *Inflatispora* (*I. pseudostromatica*) grouped with taxa of the novel genus *Pleomonodictys* (*Pleomonodictys capensis* and *Pl. descalsii*) described in that study, and those of *Bactrodesmium* (*B. cubense*), *Clypeolocus* (*C. akitaensis*) and *Morosphaeria* (*M. velatispora*). However, *Pleomonodictys* was placed in the new family Pleomonodictydaceae, while the other genera were reported as *incertae sedis* (Hernández-Restrepo et al. 2017). Later, when Jayasiri et al. (2019) introduced *Pleohelicoon* in Pleomonodictydaceae, *Inflatispora* clustered with *Pleohelicoon* and *Pleomonodictys* in a clade, where it formed a distinct basal lineage to the two genera with low statistical support and was still considered as *incertae sedis* taxon. In the present study *Inflatispora* clusters in Pleomonodictydaceae with 99% ML BS, 1.00 BYPP statistical support (Fig. 1). *Inflatispora* resembles (in its sexual morph) *Robiniigena hyalinospora* gen. et sp. nov. in Pleomonodictydaceae and is accepted in the family based on morpho-phylogenetic evidence.

***Robiniigena* Bundhun, Camporesi & K.D. Hyde, gen. nov.**

Index Fungorum: IF902255

Facesoffungi Number: FoF14884

Etymology: The name is based on the host genus *Robinia*, from which the fungus was isolated.

Description. **Saprobic** on woody substrates. **Sexual morph:** **Ascomata** immersed, with black dots present on host surface or erumpent, scattered, solitary or growing in groups in a pseudostroma, perithecial, globose to subglobose, usually unilocular, brown to dark brown, ostiolate. **Ostiole neck** papillate, composed of hyaline pseudoparenchymatous cells. **Peridium** comprising two regions, outer region multi-layered, composed of brown to dark brown, thick-walled cells of **textura angularis**; inner layer made up of lightly pigmented to hyaline cells. **Hamathecium** composed of hyaline, filiform, cellular, branched, indistinctly septate pseudoparaphyses. **Asci** bitunicate, fissitunicate, 8-spored, hyaline, clavate to cylindric-clavate, thin-walled, short-pedicellate, apically rounded, with an ocular chamber. **Ascospores** overlapping 1–2-seriate, hyaline, narrow to broadly fusiform, straight to slightly curved, euseptate, constricted at the septum, both ends conically rounded, smooth-walled, usually guttulate,

surrounded by a thick or spreading mucilaginous sheath at maturity. **Asexual morph:** Undetermined.

Type species. *Robiniigena hyalinospora* Bundhun, Camporesi & K.D. Hyde

Notes. This monotypic genus forms a basal lineage in Pleomonodictydaceae with 99% ML BS and 1.00 BYPP statistical support (Fig. 1). *Robiniigena* shares several characteristics with *Inflatisporea*, in terms of cellular pseudoparaphyses, clavate to cylindric-clavate, short-pedicellate asci, and hyaline, septate ascospores surrounded by a mucilaginous sheath (Zhang et al. 2011; Tibpromma et al. 2017). The ascospores of *Robiniigena* are, however, narrow to broadly fusiform and sometimes contain few large guttules, while those of *Inflatisporea* are mostly narrowly fusiform to almost cylindrical and are ornamented with small guttules (Zhang et al. 2011; Tibpromma et al. 2017). The two genera are also phylogenetically distinct, with *Inflatisporea* grouping with *Pleohelicoon* and *Pleomonodictys* in a clade and *Robiniigena* forming a basal separate lineage (Fig. 1).

***Robiniigena hyalinospora* Bundhun, Camporesi & K.D. Hyde, sp. nov.**

Index Fungorum: IF902256

Facesoffungi Number: FoF14885

Fig. 4

Etymology. The epithet refers to the hyaline ascospores.

Holotype. MFLU 23-0141

Description. *Saprobic* on *Robinia pseudoacacia*. **Sexual morph:** **Ascomata** 320–470 µm high, 250–600 µm diam. ($\bar{x}$ = 396 × 471 µm, n = 5), immersed, with black dots present on host surface or erumpent, visible in bark fissures, scattered, solitary or aggregated in a pseudostroma, perithecial, globose to subglobose, usually unilocular, rarely bilocular, brown to dark brown, coriaceous, ostiolar **neck** 90–110 µm wide, papillate, comprising amorphous hyaline cells. **Peridium** 30–50 µm thick near the apex, 20–35 µm wide at the sides and base, comprising two regions; outer region multi-layered, composed of brown to dark brown, thick-walled cells of **textura angularis**, fusing and becoming indistinguishable from the pseudostroma or host cells towards the outermost side; inner layer made up of lightly pigmented to hyaline cells merging with the pseudoparaphyses. **Pseudoparaphyses** 1.5–2.5 µm wide, numerous, filiform, cellular, branched, indistinctly septate, usually guttulate, surrounding the asci and along the innermost layer of the peridium. **Asci** 80–145(–160) × 15–25 µm ($\bar{x}$ = 111.5 × 20.9 µm, n = 15), bitunicate, fissitunicate, 8-spored, hyaline, clavate to cylindric-clavate, straight to slightly curved, thin-walled, short-pedicellate, apically rounded, with an ocular chamber. **Ascospores** (25–)30–40(–45) × 6–10(–12) µm ($\bar{x}$ = 36.8 × 8.6 µm, n = 55), overlapping 1–2-seriate, hyaline, narrow to broadly fusiform, straight to slightly curved, 1-euseptate, constricted at the septum, symmetrical or upper cell slightly longer than lower cell, wider upper cell, broad to acute and conically rounded at both ends, smooth-walled, sometimes guttulate, surrounded by a thick or spreading mucilaginous sheath when mature. **Asexual morph:** Not observed.

Culture characteristics. Ascospores germinated on WA within 24 hr. Colony on PDA, reaching 2 cm diam. after 15 days at 25 °C; above view dark grey in the middle and pale grey edges, dense, circular, umbonate, surface rough, radially furrowed, fimbriate; reverse dark brown, radiating outwardly.




Figure 4. *Robiniigena hyalinospora* (MFLU 23-0141, holotype) **a** host substrate **b, c** ascomata on substrate **d** vertical section of an ascoma **e** section of an ostiolar neck **f** peridium **g** pseudoparaphyses **h–k** immature and mature asci **l–q** immature and mature ascospores **r** upper view of colony **s** reverse view of colony. Scale bars: 200 μm (**b**); 400 μm (**c**); 100 μm (**d**); 50 μm (**e, h–k**); 20 μm (**f, l–q**); 3 μm (**g**).

Material examined. ITALY • Padova, near Torreglia; on dead aerial branches of *Robinia pseudoacacia*, 18 Nov 2021, E. Camporesi IT 4807 (holotype MFLU 23-0141), ex-type culture MFLUCC 23-0074.

Notes. *Robiniigena hyalinospora* resembles *Inflatospira pseudostromatica* by its globose to sub-globose ascomata, short-pedicellate asci, and ascospores whose upper cells are comparatively broader than the lower part (Zhang et al. 2011). The ascomata of *R. hyalinospora* are however, coriaceous and occur in a pseudostroma mainly when they are aggregated, while the ascomata of *I. pseudostromatica* are hard and form under a black pseudostroma both when the ascomata are solitary or occur in groups (Zhang et al. 2011). Furthermore, *R. hyalinospora* has narrow to broadly fusiform, 1-euseptate ascospores with conically rounded ends, whereas *I. pseudostromatica* comprises narrowly fusiform to nearly cylindrical, 3-septate ascospores with broadly or narrowly rounded ends (Zhang et al. 2011).

Robiniigena hyalinospora is similar to *I. caryotae*, in terms of 1-(eu)septate ascospores with a constricted middle septum and surrounded by a mucilaginous sheath (Tibpromma et al. 2017). However, *I. caryotae* has immersed ascomata while *R. hyalinospora* has immersed or erumpent ascomata, and the ascospores of *I. caryotae* are narrowly fusiform with acute ends while those of *R. hyalinospora* are narrow to broadly fusiform, with comparatively broader and conically rounded ends. *Robiniigena hyalinospora*, *I. caryotae* and *I. pseudostromatica*, are the only taxa in Pleomonodictydaceae with known sexual morphs.

***Ampelomyces quisqualis* Ces., in Klotzsch, Bot. Ztg. 10: 301 (1852)**

Index Fungorum: IF121267

Facesoffungi Number: FoF11631

Fig. 5

Description. **Saprobic** on stem of *Sonchus* sp. **Sexual morph:** **Ascomata** 140–180 µm high, 200–260 µm diam. ($\bar{x}$ = 159 × 221 µm, n = 5), immersed, appearing as black dots on the host surface, solitary to aggregated, scattered, perithecial, unilocular, globose to subglobose, dark brown, ostiolate. **Ostiole** centric, comprising hyaline cells. **Peridium** 20–30 µm thick near the ostiole, 10–25 µm wide at the sides and 10–20 µm thick at the base, 3–4-layered, outer layer made up of thick-walled, brown cells of **textura angularis**; inner layer made up of thin-walled, pale brown to hyaline cells of **textura angularis**. **Pseudoparaphyses** 1–1.5 µm wide, numerous, hyaline, filiform, branched, septate, cellular, usually guttulate, surrounding the asci. **Asci** 45–65(–68) × 5–7.5 µm ($\bar{x}$ = 56.7 × 6.4 µm, n = 40), bitunicate, 8-spored, cylindrical, straight to slightly curved, thin-walled, short-pedicellate, bulbous, with an apical ocular chamber. **Ascospores** 12–17 × 2–4 µm ($\bar{x}$ = 15.3 × 3 µm, n = 50), overlapping uni- to bi-seriate, hyaline when immature, sub-hyaline on maturity, fusiform, straight to slightly curved, 2-celled with a median septum, symmetrical or upper cell slightly longer than lower cell, cell above septum slightly enlarged and with round or conical ends, lower cell mostly with round ends, minutely guttulate, sometimes both ends containing hyaline appendages which disappear with age. **Asexual morph:** see Manjunatha et al. (2020).




Figure 5. *Ampelomyces quisqualis* (MFLU 23-0142) **a** ascomata on host substrate **b** vertical section of an ascoma **c** section of an ostiole **d** peridium **e** pseudoparaphyses **f-h** asci **i-m** ascospores **n-p** hyaline appendages at both ends of ascospores (shown by arrows). Scale bars: 100 µm (**b**); 50 µm (**c**); 10 µm (**d, i-p**); 3 µm (**e**); 30 µm (**f-h**).

Material examined. ITALY • Forlì-Cesena, Valico Tre Faggi - Premilcuore; on dead aerial stems of *Sonchus* sp., 22 Jun 2021, E. Camporesi IT 4713, Herbarium material MFLU 23-0142.

Notes. Phylogenetic analyses based on the combined LSU–ITS dataset showed that strain MFLU 23-0142 grouped with *Ampelomyces quisqualis* (AMP, Chillan, BRIP 72107, and CBS 133.32) strains with 100% ML BS, 1.00 BYPP support (Fig. 2). While there was no base pair (bp) difference among the three strains of *A. quisqualis* (Chillan, BRIP 72107 and CBS 133.32) and MFLU 23-0142 vis-à-vis the ITS sequence (based on the aligned untrimmed dataset, including gaps), 1.2% (6/518 bp) difference between *A. quisqualis* (AMP) and strain MFLU 23-0142 was observed. Similarly, there was 0.1% bp (1/876 bp) difference between *A. quisqualis* (CBS 133.32) and MFLU 23-0142 with regards to the LSU sequence (no LSU sequence data are available for strains AMP, Chillan, and BRIP 72107 in GenBank). A morphological comparison could not be made since our strain was collected in its sexual morph while *Ampelomyces* has so far been reported in its asexual morph. Therefore, the strain MFLU 23-0142 is described as the sexual morph of *A. quisqualis* based on phylogenetic support.

Melomastia maolanensis (Jin F. Zhang, Jian K. Liu, K.D. Hyde & Zuo Y. Liu)
Norphoun, T.C. Wen & K.D. Hyde, Cryptog. Mycol. 38(4): 518 (2017)

Index Fungorum: IF554041

Facesoffungi Number: FoF02695

Fig. 6

Description. *Saprobic* on dead stems of *Chromolaena odorata*. **Sexual morph:** **Ascomata** 420–440 µm high (including ostiole), 330–365 µm diam. ($\bar{x}$ = 426 × 351.3 µm, n = 5), visible as dark, raised, black spots on host, scattered, solitary to gregarious, immersed, with erumpent ostiole, perithecial, subglobose to obpyriform, coriaceous, dark brown to black, papillate, with a black clypeus. **Ostiole** 170–190 × 160–190 µm, central, oblong, dark brown to black, periphysate. **Peridium** 18–25 µm wide, comprising two layers; outer layer made up of thick-walled, dark brown cells fusing with host tissue, inner layer composed of thin-walled, hyaline cells. **Pseudoparaphyses** 1.5–3 µm wide, numerous, filiform, generally aseptate and unbranched, tapering towards the apex. **Asci** 80–125 × 4–7 µm ($\bar{x}$ = 102.7 × 5.5 µm, n = 20), 8-spored, non fissitunicate, long cylindrical, slightly flexuous, short-pedicellate, truncate or rounded at the apex, with a small ocular chamber. **Ascospores** 14–18.5 × 3–5 µm ($\bar{x}$ = 17.2 × 4.2 µm, n = 20), uniseriate, partially overlapping, hyaline, fusiform, with rounded or acute ends, 3-septate, constricted at the septa, smooth-walled, guttulate, surrounded by a mucilaginous sheath which vanishes with age. **Asexual morph:** Not observed.

Material examined. THAILAND • Chiang Mai, on dead stems of *Chromolaena odorata*, 7 Jul 2021, N.S. Wijesinghe CMN1, Herbarium material MFLU 23-0143.

Notes. In this study, strain MFLU 23-0143 clustered with *M. maolanensis* (GZCC 16-0102) with 100% ML BS, 1.00 BYPP support in the multi-locus phylogeny (Fig. 3). Nucleotide comparison of LSU and *tef1-α* revealed 0.1% (1/911 bp) and 1.3% (12/921 bp) differences between the two strains (MFLU 23-0143 and GZCC 16-0102), indicating insufficient phylogenetic differences to separate them as two different taxa (Jeewon and Hyde 2016; Maharachchikumbura et al. 2021). The two strains are also morphologically similar with ostiolate and papillate, immersed ascomata, long cylindrical and short-pedicellate asci, and ascospores that are fusiform, guttulate, and 3-septate. Therefore, strain MFLU 23-0143 is




Figure 6. *Melomastia maolanensis* (MFLU 23-0143) **a** stem of *Chromolaena odorata* **b** ascomata on host **c** vertical section of an ascoma **d** ostiole **e** peridium **f** pseudoparaphyses **g, h** asci **i–m** ascospores. Scale bars: 200 μm (**b**); 100 μm (**c**); 50 μm (**d**); 20 μm (**e**); 5 μm (**f**); 30 μm (**g, h**); 10 μm (**i–m**).

identified as *M. maolanensis* based on morphological and phylogenetic evidence. The main morphological difference between the two strains is the presence of a mucilaginous sheath around the ascospores of *M. maolanensis* MFLU 23-0143, while no such report was made for the strain GZCC 16-0102 (Zhang et al. 2017). Variations in the size of the morphological characteristics between the two strains

may be accounted for by environmental differences. Zhang et al. (2017) reported *M. maolanensis* from an undetermined tree branch in China. The same taxon is reported here from dead stems of *Chromolaena odorata* in Thailand.

***Melomastia oleae* W.L. Li, Maharachch. & Jian K. Liu, Journal of Fungi 8(1, no. 76): 10 (2022)**

Index Fungorum: IF841500

Facesoffungi Number: FoF10534

Fig. 7

Description. *Saprobic* on dead branches of *Duranta erecta*. **Sexual morph:** **Ascomata** 415–420 µm high, 500–520 µm diam. ($\bar{x}$ = 418.9 × 511.1 µm, n = 5), visible as black, cone-shaped structures on host surface, usually solitary, scattered, semi-immersed to erumpent, globose to ampulliform, carbonaceous, dark brown to black, ostiolate. **Ostiole** central, dark brown to black, papillate, carbonaceous, periphysate. **Peridium** 18–35 µm diam., comprising two regions in vertical section; outer region carbonaceous, made up of 5–7-layered, thick-walled, brown cells of *textura angularis* to *textura epidermoidea*, innermost region composed of compressed, hyaline cells. **Pseudoparaphyses** 1.5–2.5 µm wide, numerous, filiform, unbranched, sometimes septate. **Asci** 100–180 × 5.5–8 µm ($\bar{x}$ = 142.7 × 6.7 µm, n = 22), 8-spored, non fissitunicate, cylindrical, straight to flexuous, rounded at the apex, with a small ocular chamber, short-pedicellate. **Ascospores** 20–25 × 5–6 µm ($\bar{x}$ = 22.2 × 5.5 µm, n = 30), uniseriate, hyaline, fusiform, with rounded or obtuse ends, 3-septate, slightly constricted at the septa, guttulate, smooth-walled. **Asexual morph:** Not observed.

Culture characteristics. Colonies on PDA reaching 20 mm diam. in 3 weeks at 25 °C. Culture from above circular, regular, entire margin, dense, white; reverse pale brown.

Material examined. THAILAND • Chiang Rai, Mae Fah Luang University, Mueang, Tha Sut; on dead stems and twigs of *Duranta erecta*, 30 Nov 2021, V. Thiyagraja DB 184, Herbarium material MFLU 23-0144, living culture MFLUCC 23-0086.

Notes. The isolate MFLUCC 23-0086 in the present study is basal to all the strains of *M. oleae* (Fig. 3). There was 0.2% (2/986 bp) nucleotide difference in SSU, while 0.9% (8/872 bp) in LSU and 0.1% (1/894 bp) in *tef1-a* between strain MFLUCC 23-0086 and the type strain (CGMCC3.20619) of *M. oleae*. Isolate MFLUCC 23-0086 is morphologically similar to *M. oleae* in terms of ascomata which appear as cone-shaped on the host surface, filiform and unbranched pseudoparaphyses, cylindrical, pedicellate asci and uniseriate, fusiform, 3-septate ascospores. However, the asci and ascospore sizes of the type of *M. oleae* are larger than our collection. Furthermore, the peridium of strain MFLUCC 23-0086 is observed as comprising thick-walled, brown cells of *textura angularis* to *textura epidermoidea* in the outer region and compressed, hyaline cells in the innermost region (Fig. 7). The type of *M. oleae* is reported to have a peridium with an outer thick, carbonaceous layer and the inner one made up of 5–6 layers of hyaline cells of *textura angularis* to *textura prismatica* (Li et al. 2022). Despite these morphological differences, MFLUCC 23-0086 is recognized as *M. oleae* as there is insufficient genetic variation to distinguish it as a different species (Maharachchikumbura et al. 2021; Pem et al. 2021). Furthermore,




Figure 7. *Melomastia oleae* (MFLU 23-0144) **a** ascomata on the stem of *Duranta erecta* **b** an ascoma eruptent through the host tissue **c** vertical section of an ascoma **d** peridium **e** pseudoparaphyses **f–h** asci **i–k** ascospores. Scale bars: 200 μ m (**c**); 20 μ m (**d**, **i–k**); 5 μ m (**e**); 30 μ m (**f–h**).

the morphological differences may be due to different hosts and environmental conditions. Therefore, *M. oleae* is herein introduced as a new record from *Duranta erecta* in Thailand.

Discussion

In the present study, the familial description of the sexual morph of taxa in Pleomonodictydaceae is emended to include *Robiniigena* gen. nov. and *Inflatispora*, as both share several common morphological characteristics. *Robiniigena* also shares similar features with taxa belonging to the sister clade Morosphaeriaceae (Hongsanant et al. 2020). *Robiniigena* has cellular pseudoparaphyses, similar to *Aquihelicascus* and *Neohelicascus* taxa (Dong et al. 2020) and short-pedicellate asci resembling those of *Aquilomyces* and *Clypeoloculus* (Tanaka et al. 2015). *Robiniigena* also has hyaline ascospores with a gelatinous sheath similar to species of *Aquilomyces*, *Clypeoloculus*, and *Morosphaeria* (Suetrong et al. 2009; Tanaka et al. 2015; Devadatha et al. 2018). *Robiniigena* differs from Morosphaeriaceae species in that its ascomata are not covered with brown hyphae, as in *Aquilomyces* and *Clypeoloculus* (Tanaka et al. 2015). It has short-pedicellate asci while taxa of *Aquihelicascus* comprise asci with long pedicels (Dong et al. 2020). Ascospores are hyaline in *Robiniigena* while those of several *Helicascus* and *Neohelicascus* taxa are brown (Preedanont et al. 2017; Zeng et al. 2018; Dong et al. 2020). The introduction of *Robiniigena* and the inclusion of the *incertae sedis* *Inflatispora* in Pleomonodictydaceae indicates that the family is under constant taxonomic review. Further taxon sampling will undoubtedly give a better insight into the family as well as its relationship with other families in Massariaceae.

Robiniigena hyalinosporea was isolated from dead branches of *Robinia pseudoacacia* in Italy. Similarly, other fungi have been reported from the same host in Italy. *Cladosporium nigrellum* and *Camarosporidiella mirabellensis*, for instance, were isolated from dead or decorticated branches of *R. pseudoacacia* (Bensch et al. 2012; Wanasinghe et al. 2017). *Dothidotthia robiniae* was also retrieved from the same host as a saprobe, but from the Russian Federation (Senwanna et al. 2019). *Camarosporidiella elongata* (syn. *Cucurbitaria elongata*) and *Massaria anomia* (syn. *Aglaospora profusa*) were pathogenic on *R. pseudoacacia* in Greece (Michalopoulos-Skarmoutsos and Skarmoutsos 1999). Other fungi from the class Sordariomycetes have also been recorded from this host. For instance, *Colletotrichum nymphaeae* was identified as a pathogen of *R. pseudoacacia* in Japan (Yamagishi et al. 2016), while *Diaporthe oncostoma* was isolated from dead twigs of *R. pseudoacacia* (even though the species was also reported as a weak parasite) in Bulgaria (Stoykov 2012). These findings indicate that *Robinia pseudoacacia* hosts a diverse range of fungi with different lifestyles, across a large geographical area.

In this study, we have also discovered and reported the previously unknown sexual morph of *Ampelomyces quisqualis* (Wijayawardene et al. 2021). *Ampelomyces* comprises 16 species (Species Fungorum 2024), but sequence data are available for only *Ampelomyces quisqualis*. The inclusion of the genus in Phaeosphaeriaceae has been accepted by Phookamsak et al. (2014) based on the type species *A. quisqualis* (CBS 129.79). However, the latter strain is not the type or a verified strain of *A. quisqualis* (de Gruyter et al. 2009; Phookamsak et al. 2014; Vu et al. 2019; Hongsanant et al. 2020). Herein, we have included several representative strains of *A. quisqualis* (from NCBI) in our phylogeny, and they all cluster in a clade, but with three subclades (Fig. 2). Our collection MFLU 23-0142 grouped with *A. quisqualis* strains (AMP, Chillan, BRIP 72107, and CBS 133.32) in subclade A. A recent study by Huth et al. (2021) compared the draft genome assemblies for *A. quisqualis* strains (BRIP 72107 and HMLAC 05119; NCBI names) and revealed

that the two strains are not conspecific. In the present study, strain HMLAC 05119 clustered with other *A. quisqualis* strains in subclade B, while *A. quisqualis* BRIP 72107 is nested in subclade A (Fig. 2). The presence of the subclades potentially indicates that all the strains may not actually be *A. quisqualis*; they probably belong to several different lineages in *Ampelomyces*. Several previous studies have also made such observations (Lee et al. 2007; Park et al. 2010; Angeli et al. 2011; Liyanage et al. 2018). Therefore, it is vital that *A. quisqualis* is recollected and the type sequenced to resolve these ambiguities. Until this taxonomic uncertainty is resolved, strain MFLU 23-0142, collected in the present study, is referred to *A. quisqualis*, as the sexual morph. In another scenario, *Ampelomyces* has been reported to cluster with *Neosetophoma* taxa in other phylogenetic studies (Phookamsak et al. 2014; Tibpromma et al. 2017; Hyde et al. 2018; Hongsanan et al. 2020). Until *Ampelomyces* is typified, its generic status, as well as its phylogenetic placement (including that of '*A. quisqualis*' strains), remain uncertain.

Melomastia maolanensis was originally described as a saprobe on dead branches of an undetermined host from Guizhou, China (Zhang et al. 2017), while *M. oleae* was collected from dead branches of *Olea europaea* at the foot of a mountain or mountainside in Sichuan, China (Li et al. 2022). As mentioned above, the strains collected in this study were similarly obtained from terrestrial habitats, although from different geographical locations and hosts. Some *Melomastia* taxa, such as *M. aquatica*, *M. neothailandica*, and *M. thailandica* have been collected from aquatic habitats as saprobes (Hyde 1992; Hyde et al. 2016; Dayarathne et al. 2020). *Melomastia septemseptata*, a recently introduced novel taxon based on morphological support, and characterized by dark green ascomata and multi-septate ascospores, has been found to occur on living tree bark (corticolous) in the terrestrial environment (Naziazeno and Aptroot 2023). *Melomastia distoseptata* has been reported from both terrestrial and freshwater habitats (Hongsanan et al. 2020; Boonmee et al. 2021). It is valuable to document new records of existing species, either from new geographical locations, habitats, hosts, or with different lifestyles; this enables a better understanding of the diversity and ecology of the fungal species, host jumping, and adaptations of fungi to various environmental conditions (Hyde et al. 2020; Chethana et al. 2021).

The phylogeny of the genera *Dyfrulomyces* and *Melomastia* has been confusing and not truly resolved. *Dyfrulomyces* was introduced by Pang et al. (2013) as part of a study on marine *Saccardoella* species, a genus variously classified in Clypeosphaeriaceae, Xylariales (Barr 1989), Pleurotremataceae, Xylariales (Barr 1994), and 'Unitunicate Ascomycota genera incertae sedis' (Jones et al. 2009). Suetrong et al. (2009) earlier established that *Saccardoella rhizophorae* did not show any affinities to members of the Sordariomycetes (Maharachchikumbura et al. 2015), but grouped within the Dothideomycetes, although it did not cluster with any known families or orders in the class. Pang et al. (2013) confirmed the monophyly of *S. rhizophorae* and the new marine fungus *S. tiomanensis*, in a sister clade to Acrospermaceae (Dothideomycetes) and separately from *S. montellica*, the type species of the genus referred to Sordariomycetes at that time. Pang et al. (2013), therefore, introduced a new genus *Dyfrulomyces* to accommodate *S. rhizophorae*, *S. tiomanensis* as well as *S. mangrovei* and *S. marinospora*, and also established the family Dyfrulomycetaceae in the order Dyfrulomycetales (Hyde et al. 2013). Later, Maharachchikumbura et al. (2016) synonymized Dyfrulomycetaceae under Pleurotremataceae. Norphanphoun et al. (2017) noted that *Melomastia italica* and

Dyfrulomyces maolanensis formed a sister clade to five *Dyfrulomyces* species, and referred *D. maolanensis* to *Melomastia*. They concluded that *Dyfrulomyces*, *Pleurotrema*, and *Melomastia* belonged in the Pleurotremataceae, and shared many features in common, but considered *Melomastia* and *Dyfrulomyces* “as distinct genera”. It is important to record that there were no sequences of *Melomastia* available when Pang et al. (2013) introduced the genus *Dyfrulomyces*.

In the study by Li et al. (2022), *Dyfrulomyces* and *Melomastia* species clustered together, which led to *Dyfrulomyces* and its taxa being synonymized under *Melomastia*. In the current phylogenetic study, *Melomastia* species form three clades (A, B, and C; Fig. 3). *Melomastia tiomanensis* (type species of *Dyfrulomyces*) and *M. chromolaenae* form a basal clade (clade C) with 100% ML BS, 1.00 BYPP support. Although they share common features with several *Melomastia* species, especially in their short-pedicellate, apically rounded asci, and multi-septate ascospores, they mainly differ from the other *Melomastia* taxa in their distinct spindle-shaped ascospores with tapering narrow ends (Pang et al. 2013; Mapook et al. 2020). The status of *Dyfrulomyces* is therefore justified for *M. tiomanensis* and *M. chromolaenae* based on morphological and phylogenetic evidence, as is equally reported by Kularathnage et al. (2023).

Meanwhile, the remaining *Melomastia* taxa (with molecular data available) form two clades (A and B; Fig. 3). Morphological delineation between the two clades is challenging since the taxa have overlapping features. The type species, *M. mastoidea* is characterized by 2-septate, ovoid ascospores with rounded ends, which are surrounded by a mucilaginous sheath, but it lacks sequence data. Kularathnage et al. (2023) have classified clade B as *Melomastia sensu stricto* since the taxa in that clade share most of the ascospore features with *M. mastoidea*. Clade A, which has been termed as *Melomastia sensu lato*, contains taxa whose ascospore morphology is less similar to that of *M. mastoidea*. However, this classification may not be totally accurate since *M. fusispora* in clade A also has a gelatinous sheath (Li et al. 2022). Taxa such as *M. septata*, *M. maolanensis*, and *M. sichuanensis*, which are accommodated in clade B, have 3-septate ascospores, similar to *M. oleae*, *M. distoseptata*, *M. fusispora* and *M. winteri* in clade A (Zhang et al. 2017; Hongsanan et al. 2020; Li et al. 2022; Hyde et al. 2023). Furthermore, the shape of the ascospores in both clades A and B are cylindrical, ellipsoidal to fusoid or fusiform, with rounded to acute ends. Clades A and B can indeed be indicative of two or more different genera. Moreover, *M. phetchaburiensis* and *M. sinensis*, which have been treated as uncertain species in *Melomastia* by Dong et al. (2023), may represent different genera. However, the definite classification of these taxa will be possible upon the availability of molecular data for the type species, *M. mastoidea*. Also, it could be that other characters are still underplayed. For instance, none of the taxa in either clade A or B have their asexual morphs reported so far. Therefore, with the collection of more taxa in both clades, a better resolution of their position in Pleurotremataceae will be possible. Moreover, employing more genetic markers in the phylogenetic analysis of Pleurotremataceae may provide a better and more accurate picture of the taxa in the family.

The phylogeny of *Melomastia* and *Dyfrulomyces* has long been challenging in that no sequences of *Melomastia* species were available when Pang et al. (2013) introduced *Dyfrulomyces*. Subsequent phylogenetic studies have failed to undertake an analysis of pairwise distances between the two genera or of

the family Pleurotremataceae. Likewise, base pair differences between *Melomastia* species might have clarified their taxonomy if the guidelines of Jeewon and Hyde (2016) and Maharachchikumbura et al. (2021) had been followed, especially when the *Dyfolomyces* species were synonymized by Li et al. (2022).

It is also to be noted that the phylogram in Fig. 1 involves the taxa *Pleomonodictys capensis* and *Pl. descalsii* which are clustered together but with no robust phylogenetic segregation. *Pleomonodictys descalsii* was introduced principally based on its smaller conidial size as compared to that of *Pl. capensis* (Hernández-Restrepo et al. 2017). However, Bao et al. (2021), while introducing a novel collection for *Pl. capensis*, noticed that the conidial size of their new collection closely matched that of *Pl. descalsii*, and there was also no remarkable difference vis-à-vis the other asexual features. This indicates that *Pl. capensis* could be conspecific with *Pl. descalsii* and this is equally being shown in the present phylogeny (Fig. 1). Additional morpho-phylogenetic studies are necessary to solve this taxonomic uncertainty.

Acknowledgements

Digvijayini Bundhun extends her heartfelt gratitude to Mae Fah Luang University for granting the tuition scholarship for PhD studies. She is also grateful to Professor Qi Zhao for providing experimental facilities and to Nuwanthika S. Wijesinghe for helping with specimen collection. Janith V.S. Aluthmuhandiram, Binu C. Samarakoon, and Chayanard Phukhamsakda are acknowledged for their valuable suggestions. E. B. Gareth Jones and Dhanushka N. Wanasinghe thank the King Saud University, Kingdom of Saudi Arabia for the award of a Distinguished Scientist Fellowship (DSFP). Dhanushka N. Wanasinghe thanks the Yunnan Department of Sciences and Technology of China (Grant No: 202101AS070045, 202205AM070007, 202302AE090023, 202303AP140001). The authors thank Shaun Pennycook for checking the nomenclature of the new taxa.

Additional information

Conflict of interest

The authors have declared that no competing interests exist.

Ethical statement

No ethical statement was reported.

Funding

This study was supported by the Chinese Research Fund, grant number E1644111K1, titled “Flexible introduction of the high-level expert program, Kunming Institute of Botany, Chinese Academy of Sciences”.

Author contributions

Conceptualization: DB, RSJ, KDH. Data curation: EC, DB. Formal analysis: DB, RSJ, DNW. Funding acquisition: VT, KDH. Investigation: DB, GEBJ, DNW. Methodology: DB, RSJ, DNW. Supervision: RSJ, KDH. Writing – original draft: DB, DNW. Review & editing: GEBJ, RSJ, ICS, DNW, VT, KDH.

Author ORCIDs

Digvijayini Bundhun  <https://orcid.org/0000-0002-0790-215X>
E. B. Gareth Jones  <https://orcid.org/0000-0002-7286-5471>
Ruvishika S. Jayawardena  <https://orcid.org/0000-0001-7702-4885>
Erio Camporesi  <https://orcid.org/0000-0002-2801-9462>
Dhanushka N. Wanasinghe  <https://orcid.org/0000-0003-1759-3933>
Indunil C. Senanayake  <https://orcid.org/0000-0002-7165-2394>
Vinodhini Thiyagaraja  <https://orcid.org/0000-0002-8091-4579>
Kevin D. Hyde  <https://orcid.org/0000-0002-2191-0762>

Data availability

All of the data that support the findings of this study are available in the main text.

References

- Angeli D, Maurhofer M, Gessler C, Pertot I (2011) Existence of different physiological forms within genetically diverse strains of *Ampelomyces quisqualis*. *Phytoparasitica* 40(1): 37–51. <https://doi.org/10.1007/s12600-011-0197-x>
- Bao DF, Hyde KD, McKenzie EH, Jeewon R, Su HY, Nalumpang S, Luo ZL (2021) Biodiversity of lignicolous freshwater hyphomycetes from China and Thailand and description of sixteen species. *Journal of Fungi* (Basel, Switzerland) 7(8): 669. <https://doi.org/10.3390/jof7080669>
- Barr ME (1989) The genus *Dothidotthia* (Botryosphaeriaceae) in North America. *Mycotaxon* 34: 517–526.
- Barr ME (1994) Notes on the Amphisphaeriaceae and related families. *Mycotaxon* 51: 191–224.
- Bensch K, Braun U, Groenewald JZ, Crous PW (2012) The genus *Cladosporium*. *Studies in Mycology* 72: 1–401. <https://doi.org/10.3114/sim0003>
- Boonmee S, Wanasinghe DN, Calabon MS, Huanraluek N, Chandrasiri SKU, Jones EBG, Rossi W, Leonardi M, Singh SK, Rana S, Singh PN, Maurya DK, Lagashetti AC, Choudhary D, Dai YC, Zhao CL, Mu YH, Yuan HS, He SH, Phookamsak R, Jiang HB, Martín MP, Dueñas M, Telleria MT, Kałucka IL, Jagodziński AM, Liimatainen K, Pereira DS, Phillips AJL, Suwannarach N, Kumla J, Khuna S, Lumyong S, Potter TB, Shivas RG, Sparks AH, Vaghefi N, Abdel-Wahab MA, Abdel-Aziz FA, Li GJ, Lin WF, Singh U, Bhatt RP, Lee HB, Nguyen TTT, Kirk PM, Dutta AK, Acharya K, Sarma VV, Niranjana M, Rajeshkumar KC, Ashtekar N, Lad S, Wijayawardene NN, Bhat DJ, Xu RJ, Wijesinghe SN, Shen HW, Luo ZL, Zhang JY, Sysouphanthong P, Thongklang N, Bao DF, Aluthmuhandiram JVS, Abdollahzadeh J, Javadi A, Dovana F, Usman M, Khalid AN, Dissanayake AJ, Telagathoti A, Probst M, Peintner U, Garrido-Benavent I, Bóna L, Merényi Z, Boros L, Zoltán B, Stielow JB, Jiang N, Tian CM, Shams E, Dehghanizadeh F, Pordel A, Javan-Nikkhah M, Denchev TT, Denchev CM, Kemler M, Begerow D, Deng CY, Harrower E, Bozorov T, Kholmuradova T, Gafforov Y, Abdurazakov A, Xu JC, Mortimer PE, Ren GC, Jeewon R, Maharachchikumbura SSN, Phukhamsakda C, Mapook A, Hyde KD (2021) Fungal diversity notes 1387–1511: Taxonomic and phylogenetic contributions on genera and species of fungal taxa. *Fungal Diversity* 111(1): 1–335. <https://doi.org/10.1007/s13225-021-00489-3>
- Carbone I, Kohn LM (1999) A method for designing primer sets for speciation studies in filamentous ascomycetes. *Mycologia* 91(3): 553–556. <https://doi.org/10.1080/00275514.1999.12061051>

- Chaiwan N, Gomdola D, Wang S, Monkai J, Tibpromma S, Doilom M, Wanasinghe DN, Mortimer PE, Lumyong S, Hyde KD (2021) an online database providing updated information of microfungi in the Greater Mekong Subregion. *Mycosphere* 12(1): 1513–1526. <https://doi.org/10.5943/mycosphere/12/1/19>
- Chethana KWT, Niranjana M, Dong W, Samarakoon MC, Bao DF, Calabon MS, Chaiwan N, Chuankid B, Dayarathne MC, de Silva NI, Devadatha B, Dissanayake AJ, Goonasekara ID, Huanraluek N, Jayawardena RS, Karunarathna A, Luo ZL, Marasinghe DS, Ma XY, Norphanphoun C, Pem D, Perera RH, Rathnayaka AR, Raspé O, Samarakoon BC, Senwanna C, Sun YR, Tang X, Thiagaraja V, Tennakoon DS, Zeng M, Zeng XY, Zhang JY, Zhang SN, Bulgakov TS, Camporesi E, Sarma VV, Wang Y, Bhat DJ, Hyde KD (2021) AJOM new records and collections of fungi: 101–150. *Asian Journal of Mycology* 4(1): 113–260. <https://doi.org/10.5943/ajom/4/1/8>
- Dayarathne MC, Jones EBG, Maharachchikumbura SSN, Devadatha B, Sarma VV, Khongphinitbunjong K, Chomnunti P, Hyde KD (2020) Morpho-molecular characterization of microfungi associated with marine based habitats. *Mycosphere* 11(1): 1–188. <https://doi.org/10.5943/mycosphere/11/1/1>
- de Gruyter J, Aveskamp MM, Woudenberg JHC, Verkley GJM, Groenewald JZ, Crous PW (2009) Molecular phylogeny of *Phoma* and allied anamorph genera: Towards a reclassification of the *Phoma* complex. *Mycological Research* 113(4): 508–519. <https://doi.org/10.1016/j.mycres.2009.01.002>
- Devadatha B, Sarma VV, Jeewon R, Jones EBG (2018) *Morosphaeria muthupetensis* sp. nov. (Morosphaeriaceae) from India: Morphological characterization and multigene phylogenetic inference. *Botanica Marina* 61(4): 395–405. <https://doi.org/10.1515/bot-2017-0124>
- Dong W, Wang B, Hyde KD, McKenzie EHC, Raja HA, Tanaka K, Abdel-Wahab MA, Abdel-Aziz FA, Doilom M, Phookamsak R, Hongsanan S, Wanasinghe DN, Yu XD, Wang GN, Yang H, Yang J, Thambugala KM, Tian Q, Luo ZL, Yang JB, Miller AN, Fournier J, Boonmee S, Hu DM, Nalumpang S, Zhang H (2020) Freshwater Dothideomycetes. *Fungal Diversity* 105(1): 319–575. <https://doi.org/10.1007/s13225-020-00463-5>
- Dong W, Hyde KD, Jeewon R, Liao CF, Zhao HJ, Kularathnaga ND, Li H, Yang YH, Pem D, Shu YX, Gafforov Y, Manawasinghe IS, Doilom M (2023) *Mycosphere* notes 449–468: Saprobic and endophytic fungi in China, Thailand, and Uzbekistan. *Mycosphere* 14(1): 2208–2262. <https://doi.org/10.5943/mycosphere/14/1/26>
- Glez-Peña D, Gómez-Blanco D, Reboiro-Jato M, Fdez-Riverola F, Posada D (2010) AL-TER: Program-oriented conversion of DNA and protein alignments. *Nucleic Acids Research* 38: W14–W18. <https://doi.org/10.1093/nar/gkq321>
- Hall TA (2004) BioEdit Sequence Alignment Editor 7.0. 1; Isis Pharmaceuticals, Carlsbad, CA, USA.
- Hernández-Restrepo M, Gené J, Castañeda-Ruiz RF, Mena-Portales J, Crous PW, Guarro J (2017) Phylogeny of saprobic microfungi from Southern Europe. *Studies in Mycology* 86: 53–97. <https://doi.org/10.1016/j.simyco.2017.05.002>
- Hongsanan S, Hyde KD, Phookamsak R, Wanasinghe DN, McKenzie EHC, Sarma VV, Boonmee S, Lücking R, Bhat DJ, Liu NG, Tennakoon DS, Pem D, Karunarathna A, Jiang SH, Jones EBG, Phillips AJL, Manawasinghe IS, Tibpromma S, Jayasiri SC, Sandamali DS, Jayawardena RS, Wijayawardene NN, Ekanayaka AH, Jeewon R, Lu YZ, Dissanayake AJ, Zeng XY, Luo ZL, Tian Q, Phukhamsakda C, Thambugala KM, Dai DQ, Chethana KWT, Samarakoon MC, Ertz D, Bao DF, Doilom M, Liu JK, Pérez-Ortega S, Suija A, Senwanna C, Wijesinghe SN, Konta S, Niranjana M, Zhang SN, Ariyawansa HA, Jiang HB, Zhang JF, Norphanphoun C, de Silva NI, Thiagaraja V, Zhang H, Bezerra JDP, Miranda-González R, Aptroot A, Kashiwadani H, Harishchandra D, Sérusiaux E, Aluthmuhandiram JVS,

- Abeywickrama PD, Devadatha B, Wu HX, Moon KH, Gueidan C, Schumm F, Bundhun D, Mapook A, Monkai J, Chomnunti P, Suetrong S, Chaiwan N, Dayarathne MC, Yang J, Rathnayaka AR, Bhunjun CS, Xu JC, Zheng JS, Liu G, Feng Y, Xie N (2020) Refined families of Dothideomycetes: Dothideomycetidae and Pleosporomycetidae. *Mycosphere* 11(1): 1553–2107. <https://doi.org/10.5943/mycosphere/11/1/13>
- Huelsenbeck JP, Ronquist F (2001) MRBAYES: Bayesian inference of phylogenetic trees. *Bioinformatics* (Oxford, England) 17(8): 754–755. <https://doi.org/10.1093/bioinformatics/17.8.754>
- Huth L, Ash GJ, Idnurm A, Kiss L, Vaghefi N (2021) The “bipartite” structure of the first genome of *Ampelomyces quisqualis*, a common hyperparasite and biocontrol agent of powdery mildews, may point to its evolutionary origin from plant pathogenic fungi. *Genome Biology and Evolution* 13(8): evab182. <https://doi.org/10.1093/gbe/evab182>
- Hyde KD (1992) The genus *Saccardoella* from intertidal mangrove wood. *Mycologia* 84(5): 803–810. <https://doi.org/10.1080/00275514.1992.12026209>
- Hyde KD, Jones EBG, Liu JK, Ariyawansa H, Boehm E, Boonmee S, Braun U, Chomnunti P, Crous P, Dai DQ, Diederich P, Dissanayake A, Doilom M, Doveri F, Hongsanan S, Jayawardena R, Lawrey JD, Li YM, Liu YX, Liucking R, Monkai J, Nelsen MP, Phookamsak R, Muggia L, Pang KL, Senanayake I, Shearer CA, Wijayawardene N, Wu HX, Thambugala M, Suetrong S, Tanaka K, Wikee S, Zhang Y, Hudson BA, Alias SA, Aptroot A, Bahkali AH, Bezerra LJ, Bhat JD, Camporesi E, Chukeatirote E, Hoog SD, Gueidan C, Hawksworth DL, Hirayama K, Kang JC, Knudsen K, Li WJ, Liu ZY, McKenzie EHC, Miller AN, Nadeeshan D, Phillip AJL, Mapook A, Raja HA, Tian Q, Zhang M, Scheuer C, Schumm F, Taylor J, Yacharoen S, Tibpromma S, Wang Y, Yan J, Li X (2013) Families of Dothideomycetes. *Fungal Diversity* 63(1): 1–313. <https://doi.org/10.1007/s13225-013-0263-4>
- Hyde KD, Hongsanan S, Jeewon R, Bhat DJ, McKenzie EHC, Jones EBG, Phookamsak R, Ariyawansa HA, Boonmee S, Zhao Q (2016) Fungal diversity notes 367–490: Taxonomic and phylogenetic contributions to fungal taxa. *Fungal Diversity* 80: 1–270. <https://doi.org/10.1007/s13225-016-0373-x>
- Hyde KD, Chaiwan N, Norphanphoun C, Boonmee S, Camporesi E, Chethana KWT, Dayarathne MC, de Silva NI, Dissanayake AJ, Ekanayaka AH, Hongsanan S, Huang SK, Jayasiri SC, Jayawardena RS, Jiang HB, Karunarathna A, Lin CG, Liu JK, Liu NG, Lu YZ, Luo ZL, Maharachchimbura SSN, Manawasinghe IS, Pem D, Perera RH, Phukhamsakda C, Samarakoon MC, Senwanna C, Shang QJ, Tennakoon DS, Thambugala KM, Tibpromma S, Wanasinghe DN, Xiao YP, Yang J, Zeng XY, Zhang JF, Zhang SN, Bulgakov TS, Bhat DJ, Cheewangkoon R, Goh TK, Jones EBG, Kang JC, Jeewon R, Liu ZY, Lumyong S, Kuo CH, McKenzie EHC, Wen TC, Yan JY, Zhao Q (2018) *Mycosphere* notes 169–224. *Mycosphere* 9(2): 271–430. <https://doi.org/10.5943/mycosphere/9/2/8>
- Hyde KD, de Silva N, Jeewon R, Bhat DJ, Phookamsak R, Doilom M, Boonmee S, Jayawardena RS, Maharachchikumbura SSN, Senanayake IC, Manawasinghe IS, Liu NG, Abeywickrama PD, Chaiwan N, Karunarathna A, Pem D, Lin CG, Sysouphanthong P, Luo ZL, Wei DP, Wanasinghe DN, Norphanphoun C, Tennakoon DS, Samarakoon MC, Jayasiri SC, Jiang HB, Zeng XY, Li JF, Wijesinghe SN, Devadatha B, Goonasekara ID, Brahmanage RS, Yang EF, Aluthmuhandiram JVS, Dayarathne MC, Marasinghe DS, Li WJ, Dissanayake LS, Dong W, Huanraluek N, Lumyong S, Liu JK, Karunarathna SC, Jones EBG, Al-Sadi AM, Xu JC, Harishchandra D, Sarma VV (2020) *AJOM* new records and collections of fungi: 1–100. *Asian Journal of Mycology* 3(1): 22–294. <https://doi.org/10.5943/ajom/3/1/3>
- Hyde KD, Norphanphoun C, Ma J, Yang HD, Zhang JY, Du TY, Gao Y, Gomes de Farias AR, He SC, He YK, Li CJY, Li JY, Liu XF, Lu L, Su HL, Tang X, Tian XG, Wang SY, Wei DP, Xu RF, Xu RJ, Yang YY, Zhang F, Zhang Q, Bahkali AH, Boonmee S, Chethana KWT,

- Jayawardena RS, Lu YZ, Karunarathna SC, Tibpromma S, Wang Y, Zhao Q (2023) Mycosphere notes 387–412 – novel species of fungal taxa from around the world. *Mycosphere* 14(1): 663–744. <https://doi.org/10.5943/mycosphere/14/1/8>
- Jayasiri SC, Hyde KD, Ariyawansa HA, Bhat DJ, Buyck B, Cai L, Dai YC, Abd-Elsalam KA, Ertz D, Hidayat I, Jeewon R, Jones EBG, Bahkali AH, Karunarathna SC, Liu JK, Luangsa-ard JJ, Lumbsch HT, Maharachchikumbura SSN, McKenzie EHC, Moncalvo JM, Ghobad-Nejhad M, Nilsson H, Pang KL, Pereira OL, Phillips AJL, Raspé O, Rollins AW, Romero AI, Etayo J, Selçuk F, Stephenson SL, Suetrong S, Taylor JE, Tsui CKM, Vizzini A, Abdel-Wahab MA, Wen TC, Boonmee S, Dai DQ, Daranagama DA, Dissanayake AJ, Ekanayaka AH, Fryar SC, Hongnan S, Jayawardena RS, Li WJ, Perera RH, Phookamsak R, Silva NI, Thambugala KM, Tian Q, Wijayawardene NN, Zhao RL, Zhao Q, Kang JC, Promputtha I (2015) The Faces of Fungi database: Fungal names linked with morphology, phylogeny and human impacts. *Fungal Diversity* 74(1): 3–18. <https://doi.org/10.1007/s13225-015-0351-8>
- Jayasiri SC, Hyde KD, Jones EBG, McKenzie EHC, Jeewon R, Phillips AJL, Bhat DJ, Wanasinghe DN, Liu JK, Lu YZ, Kang JC, Xu J, Karunarathna SC (2019) Diversity, morphology and molecular phylogeny of Dothideomycetes on decaying wild seed pods and fruits. *Mycosphere* 10(1): 1–186. <https://doi.org/10.5943/mycosphere/10/1/1>
- Jeewon R, Hyde KD (2016) Establishing species boundaries and new taxa among fungi: Recommendations to resolve taxonomic ambiguities. *Mycosphere* 7(11): 1669–1677. <https://doi.org/10.5943/mycosphere/7/11/4>
- Jones EBG, Sakayaroj J, Suetrong S, Somrithipol S, Pang KL (2009) Classification of marine Ascomycota, anamorphic taxa and Basidiomycota. *Fungal Diversity* 35: 1–187.
- Katoh K, Rozewicki J, Yamada KD (2019) MAFFT online service: Multiple sequence alignment, interactive sequence choice and visualization. *Briefings in Bioinformatics* 20(4): 1160–1166. <https://doi.org/10.1093/bib/bbx108>
- Kirk PM, Cannon PF, David JC, Stalpers JA (2001) *Ainsworth and Bisby's Dictionary of the Fungi*, 8th edn. CABI Publishing, London.
- Kiss L, Russell JC, Szentivanyi O, Xu X, Jeffries P (2004) Biology and biocontrol potential of *Ampelomyces mycoparasites*, natural antagonist of powdery mildew fungi. *Biocontrol Science and Technology* 14(7): 635–651. <https://doi.org/10.1080/09583150410001683600>
- Kularathnage N, Tennakoon D, Zhu X, Zhou J, Su B, Xie Y, Chen Q, Calabon M, Kirk P, Senanayake I, Doilom M, Xu B, Dong W, Song J (2023) Reinstating *Dyfratomyces* and introducing *Melomastia pyriformis* sp. nov. (Pleurotremataceae, Dyfratomycesales) from Guangdong Province, China. *Current Research in Environmental & Applied Mycology* 13(1): 13. <https://doi.org/10.5943/cream/13/1/16>
- Lee SY, Hong SK, Kim YK, Kim HG (2007) Selection and identification of a hyperparasite, *Ampelomyces quisqualis* 94013 for biocontrol of cucumber powdery mildew. *Han'guk Kyunhakhoe Chi* 35(2): 121–127. <https://doi.org/10.4489/KJM.2007.35.2.121>
- Li WL, Maharachchikumbura SS, Cheewangkoon R, Liu JK (2022) Reassessment of *Dyfratomyces* and four new species of *Melomastia* from Olive (*Olea europaea*) in Sichuan Province, China. *Journal of Fungi (Basel, Switzerland)* 8(1): 76. <https://doi.org/10.3390/jof8010076>
- Liu YJ, Whelen S, Hall BD (1999) Phylogenetic relationships among ascomycetes: Evidence from an RNA polymerase II subunit. *Molecular Biology and Evolution* 16(12): 1799–1808. <https://doi.org/10.1093/oxfordjournals.molbev.a026092>
- Liyanage KK, Khan S, Brooks S, Mortimer PE, Karunarathna SC, Xu J, Hyde KD (2018) Morpho-molecular characterization of two *Ampelomyces* spp. (Pleosporales) strains mycoparasites of powdery mildew of *Hevea brasiliensis*. *Frontiers in Microbiology* 9: 12. <https://doi.org/10.3389/fmicb.2018.00012>

- Lumbsch HT, Huhndorf SM (2010) Myconet Volume 14. Part One. Outline of Ascomycota - 2009. Fieldiana. Life and Earth Sciences 1: 1–42. <https://doi.org/10.3158/1557.1>
- Maharachchikumbura SSN, Hyde KD, Jones EBG, McKenzie EH, Huang SK, Abdel-Wahab MA, Daranagama DA, Dayarathne M, D'souza MJ, Goonasekara ID, Hongsanan S, Jayawardena RS, Kirk PM, Konta S, Liu J-K, Liu Z-Y, Norphanphoun C, Pang K-L, Perera RH, Senanayake IC, Shang Q, Shenoy BD, Xiao Y, Bahkali AH, Kang J, Somrothipol S, Suetrong S, Wen T, Xu J (2015) Towards a natural classification and backbone tree for Sordariomycetes. Fungal Diversity 72(1): 199–301. <https://doi.org/10.1007/s13225-015-0331-z>
- Maharachchikumbura SSN, Hyde KD, Jones EBG, McKenzie EH, Bhat JD, Dayarathne MC, Huang SK, Norphanphoun C, Senanayake IC, Perera RH, Shang QJ, Xiao Y, D'souza MJ, Hongsanan S, Jayawardena RS, Daranagama DA, Konta S, Goonasekara ID, Zhuang W-Y, Jeewon R, Phillips AJL, Abdel-Wahab MA, Al-Sadi AM, Bahkali AH, Boonmee S, Boonyuen N, Cheewangkoon R, Dissanayake AJ, Kang J, Li Q-R, Liu JK, Liu XZ, Liu Z-Y, Luangsa-ard JJ, Pang K-L, Phookamsak R, Promputtha I, Suetrong S, Stadler M, Wen T, Wijayawardene NN (2016) Families of Sordariomycetes. Fungal Diversity 79(1): 1–317. <https://doi.org/10.1007/s13225-016-0369-6>
- Maharachchikumbura SSN, Chen Y, Ariyawansa HA, Hyde KD, Haelewaters D, Perera RH, Samarakoon MC, Wanasinghe DN, Bustamante DE, Liu J, Lawrence DP, Cheewangkoon R, Stadler M (2021) Integrative approaches for species delimitation in Ascomycota. Fungal Diversity 109(1): 155–179. <https://doi.org/10.1007/s13225-021-00486-6>
- Manawasinghe IS, Zhang YX, Liao CF, Xiong YR, Kularathnage ND, Luo M, Chen JW (2022) Mycosphere notes 345–386. Mycosphere 13(1): 454–557. <https://doi.org/10.5943/mycosphere/13/1/3>
- Manjunatha L, Singh S, Ravikumara BM, Reddy GN, Senthilkumar M (2020) *Ampelomyces*. In: Amaresan N, SenthilKumar M, Annapurna K, Kumar K, Sankaranarayanan A (Eds) Beneficial Microbes in Agro-Ecology. Academic Press, 833–860. <https://doi.org/10.1016/B978-0-12-823414-3.00044-7>
- Mapook A, Hyde KD, McKenzie EHC, Jones EBG, Bhat DJ, Jeewon R, Stadler M, Samarakoon MC, Malaithong M, Tanunchai B, Buscot F, Wubet T, Purahong W (2020) Taxonomic and phylogenetic contributions to fungi associated with the invasive weed *Chromolaena odorata* (Siam weed). Fungal Diversity 101(1): 1–175. <https://doi.org/10.1007/s13225-020-00444-8>
- Michalopoulos-Skarmoutsos H, Skarmoutsos G (1999) Pathogenicity of fungi affecting black locust (*Robinia pseudoacacia*) in Greece. Phytoparasitica 27(3): 239–240. <https://doi.org/10.1007/BF02981464>
- Miller MA, Pfeiffer W, Schwartz T (2010) Creating the CIPRES Science Gateway for inference of large phylogenetic trees. Gateway Computing Environments Workshop (GCE). New Orleans, LA, USA, 1–8. <https://doi.org/10.1109/GCE.2010.5676129>
- Naziazeno LM, Aptroot A (2023) New species and records of mostly lignicolous dothideomycetous ascomycetes from Brazil. Cryptogamie. Mycologie 44(1): 1–9. <https://doi.org/10.5252/cryptogamie-mycologie2023v44a1>
- Norphanphoun C, Jeewon R, Mckenzie EHC, Wen TC, Camporesi E, Hyde KD (2017) Taxonomic position of *Melomastia italica* sp. nov. and phylogenetic reappraisal of Dyfrolomycetales. Cryptogamie. Mycologie 38(4): 507–525. <https://doi.org/10.7872/crym/v38.iss4.2017.507>
- Nylander JAA (2004) MrModeltest v2. Program distributed by the author. Evolutionary Biology Centre, Uppsala University.

- Pang KL, Hyde KD, Alias SA, Suetrong S, Guo SY, Idid R, Jones EBG (2013) Dyfrolomycetaceae, a new family in the Dothideomycetes, Ascomycota. *Cryptogamie. Mycologie* 34(3): 223–232. <https://doi.org/10.7872/crym.v34.iss3.2013.223>
- Park MJ, Choi YJ, Hong SB, Shin HD (2010) Genetic variability and mycohost association of *Ampelomyces quisqualis* isolates inferred from phylogenetic analyses of ITS rDNA and actin gene sequences. *Fungal Biology* 114(2–3): 235–247. <https://doi.org/10.1016/j.funbio.2010.01.003>
- Pem D, Jeewon R, Chethana KW, Hongsanan S, Doilom M, Suwannarach N, Hyde KD (2021) Species concepts of Dothideomycetes: Classification, phylogenetic inconsistencies and taxonomic standardization. *Fungal Diversity* 109(1): 283–319. <https://doi.org/10.1007/s13225-021-00485-7>
- Phookamsak R, Liu J-K, McKenzie EHC, Manamgoda DS, Ariyawansa H, Thambugala KM, Dai D-Q, Camporesi E, Chukeatirote E, Wijayawardene NN, Bahkali AH, Mortimer PE, Xu J-C, Hyde KD (2014) Revision of Phaeosphaeriaceae. *Fungal Diversity* 68(1): 159–238. <https://doi.org/10.1007/s13225-014-0308-3>
- Phukhamsakda C, Hongsanan S, Ryberg M, Ariyawansa HA, Chomnunti P, Bahkali AH, Hyde KD (2016) The evolution of Massarineae with Longipedicellataceae fam. nov. *Mycosphere* 7(11): 1713–1731. <https://doi.org/10.5943/mycosphere/7/11/7>
- Preedanon S, Klayuban A, Suetrong S, Promchoo W, Gundool W, Sangtiean T, Sakayaroj J (2017) *Helicascus mangrovei* sp. nov., a new intertidal mangrove fungus from Thailand. *Mycoscience* 58(3): 174–180. <https://doi.org/10.1016/j.myc.2017.01.001>
- Rambaut A (2012) FigTree v. 1.4. Molecular evolution, phylogenetics and epidemiology. University of Edinburgh, Institute of Evolutionary Biology, Edinburgh, UK.
- Rayner RW (1970) A mycological colour chart. Commonwealth Mycological Institute, Kew, Surrey.
- Senanayake IC, Rathnayaka AR, Marasinghe DS, Calabon MS, Gentekaki E, Lee HB, Hurdeal VG, Pem D, Dissanayake LS, Wijesinghe SN, Bundhun D, Nguyen TT, Goonasekara ID, Abeywickrama PD, Bhunjun CS, Jayawardena RS, Wanasinghe DN, Jeewon R, Bhat DJ, Xiang MM (2020) Morphological approaches in studying fungi: Collection, examination, isolation, sporulation and preservation. *Mycosphere* 11(1): 2678–2754. <https://doi.org/10.5943/mycosphere/11/1/20>
- Senwanna C, Wanasinghe DN, Bulgakov TS, Wang Y, Bhat DJ, Tang AM, Mortimer PE, Xu J, Hyde KD, Phookamsak R (2019) Towards a natural classification of *Dothidotthia* and *Thyrostroma* in Dothidotthiaceae (Pleosporineae, Pleosporales). *Mycosphere* 10(1): 701–738. <https://doi.org/10.5943/mycosphere/10/1/15>
- Species Fungorum (2024) [accessed 2 April 2024] <https://www.speciesfungorum.org/>
- Stamatakis A (2014) RAxML version 8: A tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* (Oxford, England) 30(9): 1312–1313. <https://doi.org/10.1093/bioinformatics/btu033>
- Stoykov DY (2012) Ecological interactions between invasive alien vascular plants, and essential saprophytic and parasitic fungi in Bulgaria. *Phytologia Balcanica* 18: 113–116.
- Suetrong S, Schoch CL, Spatafora JW, Kohlmeyer J, Volkmann-Kohlmeyer B, Sakayaroj J, Phongapaichit S, Tanaka K, Hirayama K, Jones EBG (2009) Molecular systematics of the marine Dothideomycetes. *Studies in Mycology* 64: 155–173. <https://doi.org/10.3114/sim.2009.64.09>
- Sung GH, Sung JM, Hywel-Jones NL, Spatafora JW (2007) A multi-gene phylogeny of Clavicipitaceae (Ascomycota, Fungi): Identification of localized incongruence using a combinational bootstrap approach. *Molecular Phylogenetics and Evolution* 44(3): 1204–1223. <https://doi.org/10.1016/j.ympev.2007.03.011>

- Swofford DL (2002) PAUP* Version 4.0 b10. Phylogenetic Analysis Using Parsimony (*and Other Methods). Sinauer Associates, Sunderland, Massachusetts.
- Tanaka K, Hirayama K, Yonezawa H, Sato G, Toriyabe A, Kudo H, Hashimoto A, Matsumura M, Harada Y, Kurihara Y, Shirouzu T, Hosoya T (2015) Revision of the Massarineae (Pleosporales, Dothideomycetes). *Studies in Mycology* 82(1): 75–136. <https://doi.org/10.1016/j.simyco.2015.10.002>
- Tennakoon DS, Thambugala KM, Wanasinghe DN, Gentekaki E, Promputtha I, Kuo CH, Hyde KD (2020) Additions to Phaeosphaeriaceae (Pleosporales): *Elongaticollum* gen. nov., *Ophiosphaerella taiwanensis* sp. nov., *Phaeosphaeriopsis beaucarnea* sp. nov. and a new host record of *Neosetophoma poaceicola* from Musaceae. *MycKeys* 70: 59–88. <https://doi.org/10.3897/mycokeys.70.53674>
- Tibpromma S, Hyde KD, Jeewon R, Maharachchikumbura SSN, Liu JK, Bhat DJ, Jones EBG, McKenzie EHC, Camporesi E, Bulgakov TS, Doilom M, de Azevedo Santiago ALCM, Das K, Manimohan P, Gibertoni TB, Lim YW, Ekanayaka AH, Thongbai B, Lee HB, Yang JB, Kirk PM, Sysouphanthong P, Singh SK, Boonmee S, Dong W, Raj KNA, Latha KPD, Phookamsak R, Phukhamsakda C, Konta S, Jayasiri SC, Norphanphoun C, Tennakoon DS, Li JF, Dayarathne MC, Perera RH, Xiao Y, Wanasinghe DN, Senanayake IC, Goonasekara ID, de Silva NI, Mapook A, Jayawardena RS, Dissanayake AJ, Manawasinghe IS, Chethana KWT, Luo ZL, Hapuarachchi KK, Baghela A, Soares AM, Vizzini A, Meiras-Ottoni A, Mešić A, Dutta AK, de Souza CAF, Richter C, Lin CG, Chakrabarty D, Daranagama DA, Lima DX, Chakraborty D, Ercole E, Wu F, Simonini G, Vasquez G, da Silva GA, Plautz Jr HL, Ariyawansa HA, Lee H, Kušan I, Song J, Sun J, Karmakar J, Hu K, Semwal KC, Thambugala KM, Voigt K, Acharya K, Rajeshkumar KC, Ryvarden L, Jadan M, Hosen MI, Mikšić M, Samarakoon MC, Wijayawardene NN, Kim NK, Matočec N, Singh PN, Tian Q, Bhatt RP, de Oliveira RJV, Tulloss RE, Aamir S, Kaewchai S, Svetasheva STY, Nguyen TTT, Antonín V, Li WJ, Wang Y, Indoliya Y, Tkalčec Z, Elgorban AM, Bahkali AH, Tang AMC, Su HY, Zhang H, Promputtha I, Luangsa-ard J, Xu JC, Yan J, Chuan KJ, Stadler M, Mortimer PE, Chomnunti P, Zhao Q, Phillips AJL, Nontachaiyapoom S, Wen TC, Karunarathna SC (2017) Fungal diversity notes 491–602: Taxonomic and phylogenetic contributions to fungal taxa. *Fungal Diversity* 83(1): 1–261. <https://doi.org/10.1007/s13225-017-0378-0>
- Vilgalys R, Hester M (1990) Rapid genetic identification and mapping of enzymatically amplified ribosomal DNA from several *Cryptococcus* species. *Journal of Bacteriology* 172(8): 4238–4246. <https://doi.org/10.1128/jb.172.8.4238-4246.1990>
- Vu D, Groenewald M, de Vries M, Gehrman T, Stielow B, Eberhardt U, Al-Hatmi A, Groenewald JZ, Cardinali G, Houbaken J, Boekhout T, Crous PW, Robert V, Verkley GJM (2019) Large-scale generation and analysis of filamentous fungal DNA barcodes boosts coverage for kingdom fungi and reveals thresholds for fungal species and higher taxon delimitation. *Studies in Mycology* 92(1): 135–154. <https://doi.org/10.1016/j.simyco.2018.05.001>
- Wanasinghe DN, Hyde KD, Jeewon R, Crous PW, Wijayawardene NN, Jones EBG, Bhat DJ, Phillips AJL, Groenewald JZ, Dayarathne MC, Phukhamsakda C, Thambugala KM, Bulgakov TS, Camporesi E, Gafforov YS, Mortimer PE, Karunarathna SC (2017) Phylogenetic revision of *Camarosporium* (Pleosporineae, Dothideomycetes) and allied genera. *Studies in Mycology* 87(1): 207–256. <https://doi.org/10.1016/j.simyco.2017.08.001>
- Wanasinghe DN, Phukhamsakda C, Hyde KD, Jeewon R, Lee HB, Jones EBG, Tibpromma S, Tennakoon DS, Dissanayake AJ, Jayasiri SC, Gafforov Y, Camporesi E, Bulgakov TS, Ekanayake AH, Perera RH, Samarakoon MC, Goonasekara ID, Mapook A, Li WJ, Senanayake IC, Li J, Norphanphoun C, Doilom M, Bahkali AH, Xu JC, Mortimer PE, Tibell L, Tibell S, Karunarathna SC (2018) Fungal diversity notes 709–839: Taxonomic and

- phylogenetic contributions to fungal taxa with an emphasis on fungi on Rosaceae. *Fungal Diversity* 89(1): 1–236. <https://doi.org/10.1007/s13225-018-0395-7>
- White TJ, Bruns T, Lee SJ, Taylor JW (1990) Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: Michael AI, David HG, John JS, Thomas JW (Eds) *PCR Protocols, A Guide to Methods and Applications*. Academic Press, Inc., New York, 315–322. <https://doi.org/10.1016/B978-0-12-372180-8.50042-1>
- Wijayawardene NN, Hyde KD, Rajeshkumar KC, Hawksworth DL, Madrid H, Kirk PM, Braun U, Singh RV, Crous PW, Kukwa M, Lucking R, Kurtzman CP, Yurkov A, Haelewaters D, Aptroot A, Lumbsch HT, Timdal E, Ertz D, Etayo J, Phillips AJL, Groenewald JZ, Papizadeh M, Selbmann L, Dayarathne MC, Weerakoon G, Jones EBG, Suetrong S, Tian Q, Castaneda-Ruiz RF, Bahkali AH, Pang KL, Tanaka K, Dai DQ, Sakayaroj J, Hujislova M, Lombard L, Shenoy BD, Suija A, Maharachchikumbura SSN, Thambugala KM, Wanasinghe DN, Sharma BO, Gaikwad S, Pandit G, Zucconi L, Onofri S, Egidi E, Raja HA, Kodsueb R, Caceres MES, Perez-Ortega S, Fiuza PO, Monteiro JS, Vasilyeva LN, Shivas RG, Prieto M, Wedin M, Olariaga I, Lateef AA, Agrawal Y, Fazeli SAS, Amooze-gar MA, Zhao GZ, Pfliegler WP, Sharma G, Oset M, Abdel-Wahab MA, Takamatsu S, Bensch K, de Silva NI, Kesel AD, Karunarathna A, Boonmee S, Pfister DH, Lu YZ, Luo ZL, Boonyuen N, Daranagama DA, Senanayake IC, Jayasiri SC, Samarakoon MC, Zeng XY, Doilom M, Quijada L, Rampadarath S, Heredia G, Dissanayake AJ, Jayawardana RS, Perera RH, Tang LZ, Phukhamsakda C, Hernandez-Restrepo M, Ma X, Tibpromma S, Gusmao LFP, Weerahewa D, Karunarathna SC (2017) Notes for genera: Ascomycota. *Fungal Diversity* 86(1): 1–594. <https://doi.org/10.1007/s13225-017-0386-0>
- Wijayawardene NN, Hyde KD, Anand G, Dissanayake LS, Tang LZ, Dai DQ (2021) Towards incorporating asexually reproducing fungi in the natural classification and notes for pleomorphic genera. *Mycosphere* 12(1): 238–405. <https://doi.org/10.5943/mycosphere/12/1/4>
- Wijayawardene NN, Hyde KD, Dai DQ, Sánchez-García M, Goto BT, Saxena RK, Erdođdu M, Selçuk F, Rajeshkumar KC, Aptroot A, Błaszowski J, Boonyuen N, da Silva GA, de Souza FA, Dong W, Ertz D, Haelewaters D, Jones EBG, Karunarathna SC, Kirk PM, Kukwa M, Kumla J, Leontyev DV, Lumbsch HT, Maharachchikumbura SSN, Marguno F, Martínez-Rodríguez P, Mešić A, Monteiro JS, Oehl F, Pawłowska J, Pem D, Pfliegler WP, Phillips AJL, Pošta A, He MQ, Li JX, Raza M, Sruthi OP, Suetrong S, Suwannarach N, Tedersoo L, Thiyagaraja V, Tibpromma S, Tkalc̃ec Z, Tokarev YS, Wanasinghe DN, Wijesundara DSA, Wimalaseana SDMK, Madrid H, Zhang GQ, Gao Y, Sánchez-Castro I, Tang LZ, Stadler M, Yurkov A, Thines M (2022) Outline of fungi and fungus-like taxa–2021. *Mycosphere* 13(1): 53–453. <https://doi.org/10.5943/mycosphere/13/1/2>
- Yamagishi N, Sato T, Chuma I, Ishiyama Y, Tosa Y (2016) Anthracnose of black locust caused by *Colletotrichum nymphaeae* (Passerini) Aa. *Journal of General Plant Pathology* 82(3): 174–176. <https://doi.org/10.1007/s10327-016-0649-8>
- Zeng M, Huang SK, Hyde KD, Zhao Q (2018) *Helicascus alatus* (Morosphaeriaceae), a new freshwater species from southwestern China. *Phytotaxa* 351(3): 210–218. <https://doi.org/10.11646/phytotaxa.351.3.2>
- Zhang Y, Fournier J, Bahkali AH, Hyde KD (2011) *Inflatipora*, a novel lignicolous genus of Pleosporales from France. *Sydowia* 63: 287–295.
- Zhang JF, Liu JK, Hyde KD, Chen YY, Liu XZ, Liu ZY (2017) Two new species of *Dyfrulomyces* (Dyfrulomycetaceae, Dothideomycetes) from karst landforms. *Phytotaxa* 313: 267–277. <https://doi.org/10.11646/phytotaxa.313.3.4>
- Zhaxybayeva O, Gogarten JP (2002) Bootstrap, Bayesian probability and maximum likelihood mapping: Exploring new tools for comparative genome analyses. *BMC Genomics* 3(1): 4. <https://doi.org/10.1186/1471-2164-3-4>