

Acute Toxicity Profile of Crude Methanolic Stem Bark Extract of *Parkia Biglobosa* in West African Dwarf (Wad) Goats

James Gana Josiah

josiah.james@miu.edu.ng

Mewar International University (MIU)

John Yisa Adama

Federal University of Technology

Okechukwu Anthony Obi

University of Agriculture

Hannah Edim Etta

Cross River University of Technology

Research Article

Keywords: Acute toxicity, biochemical Profile, Extract, Goats, Histopathology

Posted Date: October 20th, 2023

DOI: <https://doi.org/10.21203/rs.3.rs-3422231/v1>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Additional Declarations: No competing interests reported.

Version of Record: A version of this preprint was published at Journal of Bioscience and Biotechnology Discovery on February 28th, 2024. See the published version at <https://doi.org/10.31248/JBBD2024.203>.

Abstract

Despite the high demand for herbal medicines for treatment of health disorder, there are still concerns associated with not only their use, but their safety. This study aimed at providing information on the potential toxicity profile of the stem bark extracts of *P. biglobosa* in goats. For this purpose, the stem bark parts of *P. biglobosa* were collected and processed. The acute toxicity, biochemical and histopathological analysis in goats were determined. The results of phytochemical constituents revealed the presence of alkaloids, anthraquinones, cardiac glycosides, glycosides, oils, phlobatannins, reducing sugar, saponins and tannins (condensed and hydrolysable). The results of acute toxicity studies showed no mortality with up to 5000 mg/ kg of the body weight. The blood collected revealed that, CMSBE, have no significant ($P>0.05$) effect on biochemical parameters of WAD goats. However, there were slight changes in glucose concentration, cholesterol and triglyceride level, though they were not statistically significant ($P>0.05$). The extracts did not also have deleterious effect on morphology of the tissues observed during histopathology. The study has scientifically demonstrated that, stem bark extract of *P. biglobosa* have LD_{50} that is greater than 5,000 mg/kg and it can be inferred that the plant might have a wide margin of safety typical of treatment of some diseases. It is therefore recommended that despite the non-toxicity as demonstrated in this present study, detailed toxic and pharmacological effects of the extracts of *P. biglobosa* stem bark in other animals should be investigated.

1. Introduction

Medicinal plants are plants in which one or more of their parts contain substances that can be used for therapeutic purposes or which are precursors for the synthesis of therapeutic agents [1]. Globally, the focus on plant research as anthelmintic is on the increase and this evidence has been collected to show the immense potentials of medicinal plants used in various traditional systems [2].

The relevance of medicinal plants, as a normative basis for the maintenance of good health, has been widely observed in most developing countries and world at large [3, 4]. A survey carried out by the World Health Organization (WHO) showed that about 70–80% of the world populations rely on non-conventional medicine mainly of herbal sources in their primary healthcare [5].

Despite the high demand for herbal medicines for treatment of health disorder, there are still concerns associated with not only their use, but their safety. It is a common belief that the phytotherapeutic or nutraceutical options, as well as other “natural” products, are safer than synthetic drugs [6]. This statement is incorrect because, in fact, many plant secondary metabolites with anthelmintic effects may also have negative effects on the hosts. For example, tannins may affect diet digestibility [7].

Over the last two decades, different retrospective studies done indicated that the incidence of deaths occurring due to exposure to plants ranged from 1.5–7.2% in different countries of the world [8]. However, the total number of deaths due to exposure to plants throughout the world is very difficult to

establish and must certainly be underestimated since all cases of such deaths were from analytical and forensic points of view, not always well documented and thus, rarely published [9].

One of the basic standards set by WHO for the use of herbs as medicines is that they should be indicate to be non-toxic [10]. Toxicology testing of herbal drugs has also been found to have a lot of benefits [11]. Notably, it is easy to identify the toxic effects and thus determine the limit of exposure levels especially to sensitive population.

Plants such as *Parkia biglobosa* which grow naturally in West Africa are one of the most important economic trees in the Northern part of Nigeria. It has been used in West African rural communities to treat a variuos forms of diseases [12]. The efficacy of the various preparations of *P. biglobosa* is widely acclaimed by the Hausa communities of Northern Nigeria for the treatment of such diseases as malaria, diabetes-mellitus and pains [12, 13, 14, 15]

So far, most of the claims of anthelmintic efficacy of the extracts of stem bark of *P biglobosa* have been scientifically established, however, little or no information exist on the potential toxicity profile of the stem bark extract of *P. biglobosa* in goats. Therefore, the aim of this study is to provide information on adverse effects of extract of stem bark of *P biglobosa* on goats and to determine the extent at which such effects occur. This would then give us the information about the safety of the plant because of its widespread use in the management of various disorders.

2. Material and Methods

2.1 Collection, processing and extraction of plant materials

The Fresh stem bark of *P. biglobosa* was collected in North central, Nigeria. The sample was identified and authenticated by Mr Namadi Sanusi, a taxonomist in the Department of Botany, Ahmadu Bello University Zaria, Kaduna State, Nigeria. The Voucher number ABU/7064 was given to the sample which was deposited to the herbarium in the same Department for reference purposes. The methods of Soetan *et al.* and Meraiyebu *et al.* [16, 17] was used to processed plant sample. The extration was done with a Soxhlet's apparatus for 4 hours using the methods of Asuzu and Onu and Builder *et al.* [18, 19]. The determination of percentage yield of extract was calculated using the formula of Ezekwe *et al.* [20] as follows:

$$\% \text{Yield} = \frac{\text{Weight of the extract}}{\text{Weight of pulverized stem bark of } P. biglobosa} \times 100$$

2.2 Qualitative Phytochemical Screening of stem bark extract of *P. biglobosa*

The phytochemical screening of the extract was carried out to determined the constituents using standard phytochemical methods. The crude methanol extracts of stem bark was screened to determine

the possible presence of alkaloids, flavonoids, saponins, tannins, terpenoids, anthraquinones, glycosides, cardiac glycoside/cardenolides, phlobatannins, sterols and steroids, carbohydrates, starch, proteins and oils [21, 22].

2.3 Determination of maximum convenient concentrations (MCCs), maximum convenient doses (MCDs) and LD₅₀ of extract used in toxicity studies

The maximum convenient doses were calculated, due to lack of information on the actual doses of the plant extracts. The MCCs of the stem bark of *P. biglobosa* extract was prepared using the methods of Lorke and Ibrahim [23, 24]. One gramme (1g) of each extract was thoroughly mixed with 1 mL of distilled water (0.05 mL at a time) until the solution could be delivered through a needle at room temperature. The MCCs of extract in g/mL were determined from the volume of the solvent used. The maximum convenient volume (MCV) that could be administered to the goat by the oral route (gavage) is 5mL/kg [23, 24, 25]. The MCDs (g/kg) was calculated by multiplying the MCCs (g/ml) by the MCV (ml/kg) [26].

$$\text{Volume (ml)} = \frac{\text{Dose (mg/kg)} \times \text{Body weight (kg)}}{\text{Concentration (mg/mL)}}$$

The established MCD was then used as the basis for the administration of the plant extracts in acute toxicity studies

2.4 Acute toxicity studies

LD₅₀ experiment was done with modification [11, 27, 28, 29, 30]. The test extract was administered generally in a single dose by gavage to goats fasted 24 hours prior to dosing and was administered to five groups. Firstly, a lower dose of 1000mg/kg was administered to single goat at a time within interval of 24 hours. After 24 hours of the administration of the first dose, the next dose was determined by the outcome of the first dose administered. If the animal survives at the first dose, the dose is adjusted upward, but if mortality is recorded at first dose, it is adjusted downward. The adjustment of dose either upward or downward was by a constant factor. Each animal was observed for 1 or 2 days before dosing the next animal. Surviving animals monitored for delayed death for a total of 7 days. Testing was terminated when the upper limit (5000mg/kg) have been reached without mortality or when the LD₅₀ have been established from the test.

$$\text{LD}_{50} = (M_0 + M_1)/2$$

Where M₀ = Highest dose of test substance that gave no mortality.

M₁ = Lowest dose of test substance that gave mortality.

Immediately after oral administration of the CMSBE of *P. biglobosa*, the goats were critically examined for survival time, behavioral changes and clinical signs of toxicity.

2.5 Biochemical Analysis of Serum

Blood samples (2ml) for serum analysis were collected from individual goat per group during pre administration and post administration via jugular venipuncture using a 5 ml syringe into non heparinized tubes. This was allowed to coagulate at room temperature and then centrifuged for 10 minutes at 1500 x g. The serum was separated and analysed to evaluate the liver enzymes [Aspartate, aminotransferase (AST), alanine aminotransferase (ALT) and Alkaline phosphatase (ALP)] [31]. The glucose concentration, Serum urea, uric acid, creatinine, bilirubin and total protein as well as serum cholesterol and triglyceride were determined [32, 33, 34] as well as the serum analysis for electrolytes (sodium, chloride and potassium ions) levels [35]. The change in glucose /serum lipids was evaluated as follow.

$$\% \text{ Change in glucose/ serum lipid concentration} = \frac{\text{FinalConcentration}}{\text{Baselineconcentration}} \times 100$$

2.6 Post mortem examination of animals in the LD₅₀ experiment

Post mortem examination was performed on the goats that died within 24 hours after dosing with extracts. All goats with prolonged signs of toxicity and showing no signs of recovery at 48 hours were also euthanized and post mortem examination were performed. Histopathological investigation of the liver and kidney were carried out according to the method of Pieme *et al.*, (2006). This research work was conducted in accordance with the ethical rules on animals experimentation as approved by the Ethical Committee of Federal University of Technology (FUT), Minna.

2.7 Data Analysis

The data were computed in tables. The percentage yields of the extracts were calculated. Biochemical parameters were expressed as mean $\pm$ SEM. They were further subjected to t- test and one way analysis of variance (ANOVA). The Value of $P < 0.05$ was considered significant. GraphPad Instat version 3.05 Windows from Graphpad Software (2000) was used to analyze the data.

3. Results

The qualitative phytochemical screening results are shown in Table 1. The results revealed the presence of anthraquinones, alkaloids, cardiac glycosides, glycosides, oils, phlobatannins, reducing sugar, tannins (condensed and hydrolysable) and saponins. However, the extract tested negative for triterpenoid and protein. The level of availability of these phytochemical constituents varied. Glycosides, alkaloids, condensed tannins and saponins are present in higher amount .

Table 1
Qualitative Phytochemical Screening of Methanol Stem Bark Extracts
of *P. biglobosa*

Chemical constituents	Test methods	Methanol extract
Alkaloids	Mayer's test	+++
Anthraquinones	Bontrager's test	+++
Cardiac Glycosides	Keller-Kiliani test	+
Flavonoids	NaoH test	++
Glycosides	Benedict's test,	+++
Oil	Filter paper test	++
Protein	Millon reagent test,	
Phlobatannins	Hcl test	++
Reducing Sugar	Fehling test	++
Saponins	Frothing test	+++
Starch	Iodine test	
Sterols and Steroids	Conc H ₂ SO ₄ test	++
Tannin (Condensed)	Ferric chloride test	+++
Tannin (Hydrolysable)	Ferric chloride test	++
Terpenoid	Salkowski test	++
Triterpenoids	Salkowski test	
Key= - Absent, + Present, ++ Very present, +++ much present		

The percentage yield of the methanol extraction of pulverized stem bark of *P. biglobosa* are shown in Table 2.

The CMSBE of *P. biglobosa* was readily soluble in water. One (1 g) of extract was conveniently dissolved (pass through 18 G needle at room temperature) in 1.25 mL of distilled water to obtain its MCC (g/mL). The MCDs (g/kg) was obtained by multiplying MCCs (g/mL) by MCV (5 mL/kg). Therefore, the MCC (g/mL) and MCD (g/kg) of the extract were 0.8 and 4 respectively and is shown in Table 3

Table 2
Percentage Yield of Methanol Extracts from Pulverized stem bark of *Parkia biglobosa*

Components of <i>P. biglobosa</i>	Weight of Pulverized bark (g) at initial stage	Weight of the extracts(g) at final stage	Percentage yield (%)
Stem bark	3000	753.56	25.12

Table 3
Solubility, Maximum Convenient Concentrations (MCCs) (g/mL) and Maximum Convenient Doses (MCDs) (g/kg) of Stem Bark extract of *P. biglobosa*

<i>Parkia biglobosa</i>	MCC (g/mL)	MCD (g/kg)
Crude methanol stem bark extract (CMSBE)	0.8	4

The acute toxicity studies results is shown in Table 4. Varying doses of CMSBE of *P. biglobosa* were administered to five groups of goats at different time and there was no mortality recorded in all the animals administered with up to 5000 mg/ kg of their body weight. The absence of death at this higher dose showed that the LD₅₀ of the extracts (CMSBE) is greater than 5000mg/kg of body weight of the animal. The signs of toxicity recorded were in group 4 with dullness and group 5 showed dullness, enlargement of abdomen, dizziness and anorexia as signs of toxicity (Table 4). The signs of toxicity disappeared within 24 hours of extracts administration and all the groups were observed for seven days before postmortem.

Table 4
Result of Acute Toxicity Studies of crude methanol extract of stem bark of *P. biglobosa* Administered to WAD Goats

Group	Dosage(mg/Kg)	Survived	Dead	Clinical sign(s) or behavioural changes
1	1000	1	0	No sign of behavioural changes
2	2000	1	0	No sign of behavioural changes
3	3000	1	0	No sign of behavioural changes
4	4000	1	0	Dullness but mild
5	5000	1	0	Dullness, enlargement of abdomen, dizziness and anorexia

There was no significant ($P > 0.05$) changes in the serum, bilirubin, urea, Cl^- , Na^+ , K^+ and uric acid in all groups of WAD goats administered with CMSBE of *P. biglobosa* including the control group (Table 5). But glucose concentration differed significantly ($P < 0.05$) in group B and E with respect to pre-administration and post- administration. More also, group (A, D and E) of creatiine including control were not significantly different ($P > 0.05$) as shown in Table 5.

On the other hand, the CMSBE of *P. biglobosa* administered produced no significant change ($P > 0.05$) in the triglyceride, cholesterol, total protein levels as well as aspartate aminotransferase (AST), alkaline phosphate (ALP) and alanine aminotransferase (ALT) levels of WAD goats. The total protein for group B differs significantly ($P < 0.05$) (Table 6). The percentage change in glucose concentration of WAD goats administered with CMSBE of *P. biglobosa* are shown in Table 7. Elevated glucose concentration was observed in all the groups except in control group that showed reduced glucose concentration. The percentage increase in the cholesterol levels was evident compared to control group and group B that had reduced cholesterol. No change was observed in the cholesterol levels of WAD goat in group 4 (Table 8). However, reduction of triglycerides was observed among the experimental groups without significant ($P < 0.05$) (Table 8).

Table 5
Changes in the Biochemical Parameters of WAD Goats administered with CMSBE of *P. biglobosa*

Biochemical Parameters	Experimental Groups (mg/ml)	Pre administration-day	Post - administration day
Bilirubin (umol/l)	Control	0.33 ± 1.46^a	0.25 ± 0.05^a
	A	0.57 ± 0.03 ^a	0.50 ± 0.12 ^a
	B	0.60 ± 0.00 ^a	0.60 ± 0.00 ^a
	C	0.60 ± 0.00 ^a	0.40 ± 0.12 ^a
	D	0.60 ± 0.00 ^a	0.63 ± 0.03 ^a
	E	0.50 ± 0.10 ^a	0.63 ± 0.03 ^a
Glu Est. (mmol/l)	Control	2.63 ± 0.03^a	2.40 ± 0.30^a
	A	2.60 ± 0.06 ^a	2.67 ± 0.07 ^a
	B	2.50 ± 0.06 ^a	2.97 ± 0.18 ^b
	C	2.60 ± 0.12 ^a	2.70 ± 0.06 ^a
	D	2.60 ± 0.15 ^a	2.73 ± 0.09 ^a
	E	2.27 ± 0.33 ^a	2.73 ± 0.03 ^b
Urea (mmol/l)	Control	2.63 ± 0.48^a	3.10 ± 0.00^a
	A	6.23 ± 0.03 ^a	5.23 ± 1.02 ^a
	B	5.90 ± 0.35 ^a	4.87 ± 0.84 ^a
	C	5.23 ± 0.03 ^a	3.30 ± 0.06 ^a
	D	5.07 ± 0.53 ^a	3.77 ± 0.38 ^a
	E	4.50 ± 1.12 ^a	3.37 ± 0.12 ^a
Na⁺ (mmol/l)	Control	49.67 ± 6.81^a	49.50 ± 5.50^a
	A	64.67 ± 1.33 ^a	61.67 ± 0.33 ^a
	B	65.00 ± 1.53 ^a	59.00 ± 2.19 ^a
	C	62.33 ± 0.33 ^a	57.67 ± 2.96 ^a

Biochemical Parameters	Experimental Groups (mg/ml)	Pre administration-day	Post - administration day
Bilirubin (umol/l)	Control	0.33 ± 1.46^a	0.25 ± 0.05^a
	D	58.00 ± 2.00 ^a	59.67 ± 1.86 ^a
	E	58.00 ± 2.00 ^a	58.67 ± 1.76 ^a
K ⁺ (mmol/l)	Control	0.57 ± 0.13^a	0.55 ± 0.05^a
	A	0.83 ± 0.03 ^a	0.67 ± 0.03 ^a
	B	0.80 ± 0.06 ^a	0.60 ± 0.06 ^a
	C	0.83 ± 0.03 ^a	0.63 ± 0.03 ^a
	D	0.57 ± 0.03 ^a	0.67 ± 0.09 ^a
	E	0.70 ± 0.06 ^a	0.73 ± 0.07 ^a
Cl ⁻ (mmol/l)	Control	46.00 ± 0.0^a	45.50 ± 2.50^a
	A	43.67 ± 1.20 ^a	44.67 ± 1.33 ^a
	B	42.00 ± 0.58 ^a	43.33 ± 1.33 ^a
	C	43.33 ± 0.88 ^a	41.33 ± 0.33 ^a
	D	41.00 ± 0.00 ^a	43.33 ± 1.33 ^a
	E	42.00 ± 0.58 ^a	42.00 ± 0.00 ^a
Creatine (mmol/l)	Control	59.67 ± 7.13^a	55.50 ± 7.50^a
	A	68.33 ± 3.67 ^a	62.67 ± 5.49 ^a
	B	70.00 ± 2.00 ^a	55.00 ± 7.02 ^b
	C	71.33 ± 0.33 ^a	62.00 ± 0.58 ^b
	D	66.00 ± 0.33 ^a	65.67 ± 0.67 ^a
	E	63.67 ± 1.67 ^a	64.00 ± 0.58 ^a
Uric acid (u/l)	Control	3.33 ± 1.33^a	3.00 ± 0.00^a
	A	5.67 ± 0.33 ^a	6.00 ± 0.00 ^a
	B	6.33 ± 0.33 ^a	5.00 ± 0.00 ^a

Biochemical Parameters	Experimental Groups (mg/ml)	Pre administration-day	Post - administration day
Bilirubin (umol/l)	Control	0.33 ± 1.46 ^a	0.25 ± 0.05 ^a
	C	6.33 ± 0.33 ^a	4.67 ± 1.20 ^a
	D	5.33 ± 0.33 ^a	6.00 ± 0.00 ^a
	E	5.00 ± 1.00 ^a	5.00 ± 0.58 ^a

Each value is mean ± SEM for three goats in each group. There were no significant ($P > 0.05$) difference between the pre-administration and post-administration group

Table 6
Changes in the Biochemical Parameters (Coronary, Heart and Liver Determination) of WAD Goats
administered with CMSBE of *P. biglobosa*

Parameters	Group	Pre- treatment- day	Post treatment- day 15
Cholesterol mmol/l	Control	0.57 ± 0.13^a	0.50 ± 0.10^a
	A	0.57 ± 0.03 ^a	0.60 ± 0.00 ^a
	B	0.63 ± 0.03 ^a	0.50 ± 0.06 ^a
	C	0.67 ± 0.03 ^a	0.70 ± 0.06 ^a
	D	0.67 ± 0.07 ^a	0.67 ± 0.07 ^a
	E	0.50 ± 0.10 ^a	0.57 ± 0.03 ^a
Triglyceride mmol/l	Control	0.77 ± 0.09^a	0.75 ± 0.07^a
	A	0.83 ± 0.03 ^a	0.77 ± 0.12 ^a
	B	0.90 ± 3.83 ^a	0.77 ± 0.15 ^a
	C	0.87 ± 0.03 ^a	0.77 ± 0.15 ^a
	D	0.93 ± 0.13 ^a	0.83 ± 0.06 ^a
	E	0.87 ± 0.03 ^a	0.80 ± 0.00 ^a
Total protein g/ml	Control	2.77 ± 0.15^a	2.80 ± 0.00^a
	A	2.80 ± 0.10 ^a	2.60 ± 0.00 ^a
	B	2.93 ± 0.06 ^a	2.43 ± 0.21 ^b
	C	2.87 ± 0.06 ^a	2.80 ± 0.10 ^a
	D	2.67 ± 0.12 ^a	2.57 ± 0.06 ^a
	E	2.80 ± 0.10 ^a	2.70 ± 0.17 ^a
Aspartate aminotransferase(ASP)(u/l)	Control	14.00 ± 1.53	14.50 ± 1.50^a
	A	15.00 ± 0.00 ^a	16.00 ± 0.00 ^a
	B	16.00 ± 0.00 ^a	16.00 ± 0.00 ^a
	C	15.67 ± 0.58 ^a	16.33 ± 1.20 ^a

Parameters	Group	Pre- treatment- day	Post treatment- day 15
Cholesterol mmol/l	Control	0.57 ± 0.13^a	0.50 ± 0.10^a
	D	16.00 ± 0.00 ^a	16.00 ± 0.00 ^a
	E	16.33 ± 0.58 ^a	16.00 ± 0.00 ^a
Alanine aminotransferase ALT) (u/l)	Control	9.67 ± 1.86^a	11.00 ± 0.00^a
	A	11.33 ± 0.33 ^a	11.00 ± 0.00 ^a
	B	11.00 ± 0.58 ^a	11.33 ± 0.33 ^a
	C	11.67 ± 0.33 ^a	12.33 ± 0.88 ^a
	D	11.00 ± 0.00 ^a	11.33 ± 0.33 ^a
	E	10.67 ± 0.33 ^a	11.33 ± 0.33 ^a
Alkaline phosphate (ALP))(u/l	Control	21.67 ± 2.96^a	22.00 ± 1.00^a
	A	25.67 ± 0.33 ^a	25.00 ± 1.00 ^a
	B	25.67 ± 0.33 ^a	24.33 ± 1.67 ^a
	C	26.00 ± 0.00 ^a	26.33 ± 0.33 ^{aaa}
	D	26.00 ± 0.58 ^a	25.67 ± 0.33 ^a
	E	24.33 ± 1.20 ^a	25.67 ± 0.67 ^a

Each value is mean ± SEM for three goats in each group. The mean for each parameter with different superscript alphabet in the same row are statistically significant different (P < 0.05)

Table 7
Percentage Change in Glucose Concentration of WAD Goats administered with
CMSBE of *P. biglobosa*

Experimental Groups (mg/mL)	%change of glucose conc.	Mode of change
Contro1	-8.75	Decrease
A	+ 2.69	Increase
B	+ 18.8	Increase
C	+ 3.85	Increase
D	+ 5.0	Increase
E	+ 20.26	Increase
Key		
Control = distilled water (DW) 5ml/kg		
A = CMSBE of <i>P. biglobosa</i> (1000mg/kg)		
B = CMSBE of <i>P. biglobosa</i> (2000mg/kg)		
C = CMSBE of <i>P. biglobosa</i> (3000mg/kg)		
D = CMSBE of <i>P. biglobosa</i> (4000mg/kg)		
E = CMSBE of <i>P. biglobosa</i> (5000mg/kg)		

Table 8
Percentage Change in Serum Lipid Concentration of WAD Goats
administered with different dosage of CMSBE of *P. biglobosa*

Group	% change of cholesterol	Mode of change	% change of Triglyceride	Mode of change
Control	-12.28	Decrease	-2.6	Decrease
A	+ 5.26	Increase	-7.23	Decrease
B	-20.63	Decrease	-14.44	Decrease
C	+ 4.48	Increase	-12.5	Decrease
D	0	The same	-10.75	Decrease
E	+ 14	Increase	-8.05	Decrease

The histopathological studies of liver and kidney of goats administered with CMSBE of *P. biglobosa* are presented in Figures I and II. In all the treatment groups including controls, there were no structural

changes or abnormalities in the liver and kidney. The liver hepatocytes were preserved in all groups as well as prominent present of central vein. But group C, D and E showed passive congestion of central vein and infiltration of eosinophils around the portal vein, respectively (figure I). The architectural morphology of kidney showed no observable lesion in the group examined (Figure II).

4. Discussion

Medicinal plants offered great prospect for the development of novel chemotherapeutic agents that are essential for the management of various diseases in food animals and human [36]. The qualitative phytochemicals screening results in this study showed that tannin (condensed and hydrolysable), anthraquinones, cardiac glycosides, alkaloids, glycosides, flavonoides, oil, sterols/steroids, phlobatannins, reducing sugar, Proteins, saponins and terpenoids were present. These results were similar to that of Ezekwe *et al.*, Millogo Kone *et al.* and Banwo *et al.* [20, 37, 38] in methanol extracts of stem bark of *P. biglobosa*. They reported the presence of saponins, Glycosides, tannins and other phenolics with trace quantity of alkaloids. However, the report of Builder *et al.* [19] differed slightly from their results due to absence of alkaloids from the methanol stem bark extracts of *P. biglobosa*. The absence may not be a minus for the medicinal efficacy but could be the methods of processing and geographical location of this plant that might have led to differences in phytochemical constituents in the this work and the work of Builder *et al.* [19].

The separation of plant materials for pharmacological and medicinal uses. have different methods of extraction. In this study, the extract (CMSBE) yielded 25.12%, which is higher than that of Salit *et al.* [39] who reported 4.0% yield extracts of stem bark of *P. biglobosa*. This showed that, this would have been the best method of extracting phytochemical constituent of plant materials for toxicity studies. The CMSBE of *P. biglobosa* was readily soluble in water, with MCC and MCD of 0.8 g/ml and 4g/kg, respectively. These findings are quite similar to those obtained by Oshadu [40] in his MCC and MCD determination of different portions of *Acanthus montanus* leaves where he reported MCCs (g/ml) of 0.8 and MCDs (g/kg) of 4, for crude ethanol extract of *Acanthus montanus* leaves.

The acute toxicity study of CMSBE *P. biglobosa* showed no any obvious signs of toxicity or change of behaviour at doses ranging from 1000–4000 mg/kg. However, dullness, dizziness, enlargement of abdomen and anorexia were observed at 5000mg/kg but disappeared within 24 hours without mortality. This result suggested that CMSBE of *P. biglobosa* had an oral LD₅₀ greater than 5000 mg/kg. The high LD₅₀ value implies a remote risk of acute intoxication and high degree of relative safety when extract is administered orally. Though *P. biglobosa* has been used by Traditional Medical Practitioners (TMPs) without report of any mortality due to toxicity, this claim has been authenticated by the lack of death in oral treatment of WAD goats at the dose rate of 5000 mg/kg body weight. These finding agreed with the work of other researchers [15, 19, 40], they reported LD₅₀ greater than 5000 mg/kg body weight of rat administered with CMSBE of *P. biglobosa*. The low toxicity obtained may have been responsible for its widespread use in different ethnotherapeutic interventions and it is therefore considered practically non-toxic [42]. However, this result contradicted the report of Ezekwe *et al.* [20] who opined that LD₅₀ of

CMSBE of *P. biglobosa* was 400 mg/ kg body weight of albino rat. The differences in acute oral LD₅₀ values observed could be attributed to possible differences between species in drug metabolism and susceptibility to toxic effects [43].

The dose-toxicity relationships of toxic substances under experimental conditions have been graded by number of researchers. One attempts of such grading according to Loomis, Mutschler *et al.* and Wink and van Wyk [25, 44, 45], they defined < 1 mg /kg as extremely toxic, 1–50 mg/kg as highly toxic, 50–500 mg/kg as moderately toxic, 500–5000 mg/kg slightly toxic, 5000 -15,000 mg/kg as practically non-toxic and > 15,00 mg/kg as relatively harmless. Based on this grading, in this study, the extract tested in goat is considered as practically non-toxic, since the LD₅₀ fall within the range 5000 -15,000 mg/kg. Femi-Ola *et al.* [46] also reported that *P. biglobosa* plant is non-toxic to humans and that is why African countries use *P. biglobosa* as herbal medications. But the bark and the pods of the same plant were considered to contained parkine and equally have piscicidal property [47].

The CMSBE of *P. biglobosa* in this study has no effect on biochemical parameters of WAD goats. There was no significant ($P > 0.05$) increase in serum creatinine, urea, uric acid, sodium, potassium and Chloride ions. These results were in contrast with the results of Ezekwe *et al.* [20] who recorded increase in the level of creatinine and urea when methanol extract of *P. biglobosa* stem bark was administered to albino rats. Also increase in serum creatinine, urea, uric acid, sodium and potassium ions were recorded by Njidda *et al.* [48] in Kano Brown and Red Sokoto breed of goats fed on natural grazing range land of Northern Nigeria. However, this results coincides with the finding of Builder *et al.* [19] who administered extract of *P. biglobosa* stem bark to albino rats without increase in serum creatinine, urea, uric acid, sodium and potassium ions. The normal levels of serum creatinine, urea, uric acid, sodium, potassium and Chloride ions shows that the CMSBE of *P. biglobosa* did not affect the renal function and that the integrity of renal was preserved [49].

Research findings have shown that, flavonoid, terpenoid coumarin and a host of other secondary plant metabolites such as arginine and glutamic acids posses' hypoglycemic effects in various experimental animals model [50]. However, this hypothesis stipulated that plants which contain terpenoid and/or favonoid posses hypoglycemic activities in diabetic and normal mammals. One or more of the other chemical constituents of the plant especially flavonoid is also likely to have played a crucial role in the hypoglycemic action of the plant extract [51, 52].. In this study, despite the presence of terpenoid and flavonoid, there was no significant hypoglycemic activity exhibited by CMSBE of *P. biglobosa*. However, the flavonoid present in CMSBE would have been responsible for regulation of glucose level to some extent. Certain drugs and other substances are known to affect and influence circulating bilirubin. Increase in bilirubin suggests increase in hemolysis [53]. The CMSBE of *P. biglobosa* did not alter the bilirubin level of the treated goats when compared to the control.

The CMSBE of *P. biglobosa* did not interfere with total protein since there was not significant difference between the groups except group B. This result disagreed with the result of Sachin and Sahni [54] who reported increase in total protein when extract of *Nigella sativa*, *Swertia chirata* and *Piper longum* were

used for deworming goats. The reasons for these discrepancies are not known but may be due to immune response of animals during administration of CMSBE of *P. biglobosa*

The serum enzymes (AST, ALP and ALT) as well as bilirubin are conventionally used as indicator of hepatic damage [55, 56]. The serum concentration of these constituents was within the reference values of goats and there was no difference between the administered groups. Therefore, CMSBE of *P. biglobosa* show possibly no negative effect on liver function. This result agreed with that of Raghuvansi *et al.* [57]. The study also agreed with the finding of Builder *et al.* and Ezekwe *et al.* [19, 20], they reported decrease in serum alkaline phosphatase level with respect to control when they investigated the toxicity of the stem bark extract of *P. biglobosa* in albino rats. A rise in alkaline phosphatase is usually a characteristics found in cholestatic liver disease. As such, the significant reduction in ALP shows that no possible cholestasis occurred at the dose levels tested [49].

Glucose is relatively a good indicator of energy balance. The Reduction in glucose level in animals, result in gross reduction in weight gain and milk yield and in change in the fatty acid composition of the milk [58]. The slight change in the glucose concentration in the current study is an indication that infection was responsible for reduction despite the feeds that was provided. In this study, all groups showed slight increase in glucose level except the control group. This indicate that CMSBE of *P. biglobosa* can provide energy. The result agreed with that of Builder *et al.* [19] who reported increase in serum glucose level of rat treated with aqueous extract of *P. biglobosa*.

The finding in this research showed reduction in triglyceride and cholesterol. The reduction in triglyceride and cholesterol indicates that the CMSBE of *P. biglobosa* may have lipid some beneficial effects on the cardiovascular risk factors by lowering the lipid activity [59]. The reduction may be as a result of presence of flavonoid [60, 61, 62]. Several research findings indicated that many plant sterols reduce serum cholesterol absorption [63]. Therefore, the lowering property of these extracts may be the synergistic interaction of polyphenol that is present [64].

From the histopathological findings, the results revealed no effect of CMSBE on kidney and liver tissues since there was no structural changes or abnormalities of these organs when compared with controls. The normal hepatocytes displayed by the liver tissue without any enlargement in central vein, sinusoidal vein and portal triad in all administered groups and control agreed with the finding of Builder *et al.* [19], when 5000mg/kg of methanol extract of stem bark of *P. biglobosa* was administered to albino rats. Similar findings were reported by Roy *et al.* [65] without structural changes in liver tissues when extract of *Senna alata* leaf of varying dosages of 1000 to 3000 mg/kg were administered to Swiss Albino mice. Bello *et al.* [66] also reported the same findings when methanolic extract of *Alstonia scholaris* stem bark was administered to albino rats. However, this result contradicts that of Builder *et al.* [19], where they administered 5000mg/kg of aqueous extract of stem bark of *P. biglobosa* to albino rats; there was fatty degeneration of liver cells.

Kidney micrograph revealed normal architecture of glomerulus and Bowman's capsules with no tubular degeneration and dilation of glomerular capillaries. The normal architecture of kidney in this study

agreed with the finding of Builder *et al.* [19] who administered 5000mg/kg of methanol extract of stem bark of *P. biglobosa* to albino rats without abnormalities but 5000mg/kg of water extract of stem bark of *P. biglobosa* showed tubular degeneration and dilation of glomerular capillaries. Other researchers such as Ping *et al.* and Nabukenya *et al.* [67, 68] reported normal architectural of kidney when *Euphorbia hirta* L. methanol extract and aqueous extracts of *Tephrosia vogelii*, *Vernonia amygdalina* and *Senna occidentalis* extracts were administered to albino rats respectively.

Thus, the histopathological evaluation during the administration of CMSBE of *P. biglobosa* protocol indicated that the extracts did not have any deleterious effect on morphology of the tissues observed. These findings supported the biochemical results observed in this study since the extracts used did not produce any toxic effect in WAD goats.

5. Conclusion

- In conclusion, the CMSBE of *P. biglobosa* seemed to be blood, kidney and liver friendly as none of these parameters were altered during biochemical and histopathological analyses. This study has scientifically demonstrated that the LD₅₀ of CMSBE of *P. biglobosa* is greater than 5,000 mg/kg and it can be inferred that the plant might have a wide margin of safety typical of anthelmintics and treatment of other diseases. It is therefore recommended that Despite the non-toxicity as demonstrated in this present study, detailed toxic and pharmacological effects of the extracts of *P. biglobosa* stem bark should be investigated in other animals.

Declarations

Conflict of interest: No conflict of interest among the authors

Funding: This research did not receive any specific grant from funding agencies in the public, commercial, or not for a profit sectors. The research was funded by the authors

Author contributions: Conceptualisation: Josiah, JG., Adama JY., collecting data: Josiah, JG, Adama JY, data analysis: Josiah, JG., Adama, JY., Etta HH., Obi OA., obtaining funding: Josiah, JG., Adama, JY , Obi, OA., Etta HE., and writing manuscript: Josiah, JG. Etta HE., Adama, JY.,Obi OA.

Acknowledgement

The authors would like to acknowledged the students of Etsu Aliyu Senior Secondary School Tsaragi, Mr Shaaba A. Gana of Niger State Veterinary Hospital, Minna, laboratory technical staff of department of biochemistry, federal University of technology, Minna as well as laboratory technical staff of histopathological section of Faculty of veterinary Medicine, Ahmadu Bello University,Zaria for their assistance during this research work.

References

1. Sofowora, A., 2008. Medicinal Plants and Traditional Medicine in Africa. . Third edition. Pub Spectrum Books Limited, Ibadan.Pp 197-204.
2. Adamu, M., Nwosu, C.O., Agbede, R. I. S., 2009. Anti-trypanosoma effects of Aqueous extract of *Ocimum gratissimum* (Lamiaceae) leaf in rats infested with *Trypanosome brucei brucei*. *Afri J. of Traditional Complementary and Alternative Medicine*,**6**(3), 262-267.
3. Githiori, J. B., 2004. Evaluation of anthelmintic properties of ethnoveterinary plant preparations used as livestock dewormers by Pastoralists and small holder farmers in Kenya. *Doctoral thesis* ISSN 1401-6257.
4. Jordan, S.A., Cunningham, D.G., Marles R.J., 2010. Assessment of herbal medicinal products: Challenges and opportunities to increase the knowledge base for safety assessment. *Toxicol. and Applied Pharmacol.* 243: 198-216.
5. WHO, 2003. *Traditional Medicine*. Retrieved from <http://www.who.int/mediacentre/>
6. Athanasiadou, S., Githiori, J., Kyriazakis, I., 2007. Medicinal plants for helminth parasite control: facts and fiction. *Animal* 1, 1392–1400.
7. Méndez-Ortíz, F.A., Sandoval-Castro, C.A., Torres-Acosta, J.F.J., 2012. Short term consumption of *Havardia albicans* tannin rich fodder: Effect on feed intake, diet digestibility and excretion of *Haemonchus contortus* eggs in sheep. *Anim. Feed Sci. Technol.* 176, 185–191.
8. Gaillard, Y., Pepin, G., 1999. Poisoning by plant material: review of human cases and analytical determination of toxins by high performance liquid chromatography-mass spectrometry. *J. Chromat.* 733: 181-229.
9. Bent, S., Ko, R., 2007. Commonly used herbal medicines in the United States: a review. *The American J. Med.* 116(7): 478-485.
10. WHO, 2011. The World Medicines Situation. Traditional medicines: Global situation, issues and challenges. WHO/EMP/MIE/2011. Retrieved from <http://apps.int/medicinesdocs/documents/s/8063en.pdf>. Accessed on 13/05/2015.
11. Organization for Economic Co-operation and Development (OECD), 2008. Guidelines for the testing of chemicals, OECD/OCDE 425. *Acute oral toxicity – up-and-down-procedure (UDP) adopted*
12. Shao, M., 2002. M.Sc. thesis. Michigan Technological University, Michigan,USA.
13. Asase, A., Alfred, A., Yeboah, O., Odamtten, G. T., Simmonds, S. J., 2005. Ethnobotanical study of some Ghanaian antimalarial plant. *J. Ethnopharmacol.* 99, 273-279.
14. Gronhaug, T. E., Glaeserud, S., Skogsrud, M., Ballo, N., Bah, S., Diallo, D., Paulsen, B. S., 2008. Ethnopharmacological survey of six medicinal plants from Mali. *West African J. Ethnopharmacol.* 4, 4-26.
15. Tijani, A.Y., Okhale, S. E., Salawu, T. A., Onigbanjo, H. O., Obianodo, L. A., Akingbasote, J. & Emeje, M., 2009. Anti diarrheal and antibacterial properties of crude aqueous stem bark extract and fractions of *P. biglobosa* (Jacq) R.Br Ex G. Don. *Afri. J. Pharm. and Pharmacol.* 7, 347-353.

16. Soetan, K. O., Lasisi O. T., Agboluaje, A. K., 2011. Comparative assessment of *in-vitro* anthelmintic effects of the aqueous extracts of the seeds and leaves of the African locust
17. Meraiyebu, A., Olaniyan, O., Abutu, S., Dare, J., Atsukwei, D., 2013. Hepatoprotective Effect of *Parkia Biglobosa* Stem Bark Methanolic Extract on Paracetamol Induced Liver Damage in Wistar Rats. *American J. Biomed. and Life Sci.* 1(4), 75-78.
18. Asuzu, I. U., Onu, O. U., 1994. Anthelmintic activity of the ethanolic extract of *Piliostigma thonningii* bark in *Ascaridia galli* infected chickens. *Fitoterapia*, LXV, 291–297.
19. Builders, M. I., Isichie, C. O., Aguiyi, J. C., 2012. Toxicity Studies of the Extracts of *Parkia biglobosa* Stem Bark in Rats. *British J. Pharmaceut. Res.* 2(1), 1-16.
20. Ezekwe, C. I., Anaya, C. A., Okechukwu, P. C. U., 2013. Effects of Methanol Extract of *Parkia biglobosa* Stem Bark on the Liver and Kidney Functions of Albino Rats. *Global J. Biotechnol. and Biochem.*, 8(2), 40-50.
21. Sofowora, A., 1993. Medicinal Plants and Traditional Medicine in Africa. *Spectrum Books Limited, Ibadan, Nigeria*, pp. 191-289.
22. Evans, W.C. (ed), 2002. Trease and Evans' Pharmacology 15th Edition, W.B. Saunders, New York, pp. 221-393.
23. Lorke, D., 1983. A new approach to practical acute toxicity testing. *Arch. Toxicol.* 54, 275-287.
24. Ibrahim, M. A., 1984. Evaluation of the activities of some West African Traditional anthelmintic herbs against *Nippostrongylus brasiliensis* in rats. M.Sc. Thesis, Department of Veterinary Physiology and Pharmacology, Ahmadu Bello University, Zaria, pp.116.
25. Loomis, T. A., 1978. Essentials of Toxicology. Third edition. Lea and Fibiger, Philadelphia, pp 198.
26. Kimani, D., Kareru, P. G., Njonge, F. K., Kutima, H. L., Nyagah, G. C., Rechab, S. O., Wamburu, R. W & Karanja, J. M., 2013. Control of gastro-intestinal nematodes in ruminants using plant extracts. *Scientific conference proceeding* PP 193-198.
27. Bruce, R. D., 1985. An up- and- down procedure for acute toxicity testing. *Fundamental Applied Toxicol.* 5, 151- 7.
28. Organization for Economic Co-operation and Development (OECD), 2001. Guidelines for the testing of chemicals 423. *Documentation on acute oral toxicity and acute class method, 2*. Available from: <http://www.oecd.org>
29. Deora, P. S., Mishra, C. K., Mavani, P., Asha, R., Shrivastava, B., Rajesh, K. N., 2010. Effective alternative methods of LD₅₀ help to save number of experimental animals. *J. Chem. Pharmaceut. Res.*, 2(6), 450-453.
30. Chinedu, E., Arome, D., Ameh, F. S., 2013. A new method for determining acute toxicity in animal models. *Toxicol. Internat.* 20, 224-6.
31. Pieme, C. A., Penlap, V. N., Nkegoum, B., Taziebou, C. L., Tekwu, E. M., Etoa, F. X., Ngongang, J., 2006. Evaluation of acute and subacute toxicities of aqueous ethanolic extract of leaves of *Senna alata* (L.) Roxb (Ceasalpiniaceae). *Afri. J. Biotechnol.* 5, 283- 289.

32. Dou, X., Yamaguchi, Y., Yamamoto, H., Uenoyama, H., Ozaki, Y., 1996. Biological applications of anti-stokes Raman spectroscopy: quantitative analysis of glucose in plasma and serum by a highly sensitive multichannel Raman spectrometer. *Application of Spectroscopy*, 50, 1301-1306.
33. Aniagu, S.O., Nwinyi, F.C., Akumka, D.D., Ajoku, G.A., Dzarma, S., Izebe, K.S., Ditse, M., Patrick, E., Nwaneri, C., Wambebe, C., Gamaniel, K., 2006. Toxicity studies in rats fed nature cure bitters. *Afri J. of Biotechnol.* 4, 72-78.
34. Taga, I., Kamseu, P., Nganzie, O., Ack, F. X., Ngongang, J., 1998. Etude comparative des méthodes de dosage chimique et enzymatique de quelque parameters biochimiques:cholestérol,glucose acide urique et phosphatase alcaline. Biosciences proceeding. Annal publication of the Cameroon Biosci. Society, 2, 251- 258.
35. Henry, J.B., 1991. Clinical diagnosis and management by laboratory methods. 18th ed, W.B. sanders company, New York. pp 141- 142.
36. Sofowora, A., Eyitope Ogunbodede, E., Adedeji Onayade, A., 2013. The role and place of medicinal plants in the strategies for disease prevention. *Afr. J. Trad. Complement Alternative Medicine*, 10(5):210-229<http://dx.doi.org/10.4314/ajtcam.v10i5.2>
37. Millogo-Kone, H., Guissou, I. P., Nacoulma, O., Traore, A. S., 2006. Study of the antibacterial activity of the stem bark and leaf extract of *Parkia biglobosa* (Jacq) Benth on *Staphylococcus aureus*. *African Journal of Traditional Complementary and Alternative Medicine*, 3(2), 74-78.
38. Banwo, G. O., Abdullahi, I., Duguryil, M., 2004. The antimicrobial activity of the stem bark of *Parkia clappertoniana* keay family Leguminosae against selected microorganisms. *Nigerian J. Pharmaceut. Res.* 1, 16-22.
39. Salit, B. S., Emmanuel, U. O., Idongesit, J. M., Pau, I N. O., Labumi L., 2014. *Parka Biglolosa* Plants Parts: Phytochemical, Antimicrobial, Toxicity and Antioxidant Characteristics. *J. Natural Sci. Res.* 4(2), 130-133.
40. Oshadu, O. D., 2014. Evaluation of anthelmintic activities of *Acanthus montanus* (acanthaceae)leaf extracts against experimental *heligmosomoides bakeri* infection in mice. *Ahmadu Bello University, Zaria. M sc. thesis.*
41. Udobi, C. E. & Onaolapo, J. A., 2009. Phytochemical analysis and antibacterial evaluation of leaf, stem bark and root of the African locust bean (*Parkia biglobosa*). *J. Med. Plants Res.* 3, 338-344.
42. Schorderet, M., 1992. Pharmacologie. Des concepts fondamentaux aux applications theurapeutiques. Editions Slatkine Geneve, Edition Frison-Roche ,Paris. pp 33-34.
43. Salawu, O. A., Chindo, B. A., Tijani, A. Y., Adzu, B., 2008. Analgesic, anti- inflammatory, anti-pyretic and anti-plasmodial effects of the methanolic extract of *Crossopteryx febrifuga*. *J. Med. Plants*, 2(8), 213-218.
44. Mutschler, E., Geisslinger, H., Kroemer, H.K., Ruth, P., Schäfer-Korting, M., 2008. *MutschlerArzneimittelwirkungen. Lehrbuch derPharmakologie und Toxikologie*. 9th edition, WVG, Stuttgart,

45. Wink, M. & Van Wyk, B. E., 2008. Mindaltering and poisonous plants of the world. *BRIZA*, Pretoria (SA)
46. Femi-Ola T. O., Ajibade, V. A., Afolabi, A., 2008. Chemical composition and termicidal properties of *Parkia biglobosa* (Jacq) Benth. *J. Biol. Sci.* 8(2), 494-497
47. Abalaka, S. E., Auta, J., 2010. Toxic Effects of Aqueous and Ethanol Extracts of *Parkia biglobosa* Pods on *Clarias gariepinus* Adults. *World J. Biol. Res.*, 3(1), 9-17.
48. Njidda, A. A., Hassan, I. T. & E. A. Olatunji, E. A., 2013. Haematological and Biochemical Parameters of Goats of Semi Arid Environment Fed On Natural Grazing Rangeland of Northern Nigeria, *J. Agric. and Vet. Sci.* 3(2), 01-08.
49. Kaneko, J. J., 1989. Clinical biochemistry of domestic animals. 4th Edition. Academic Press. San Diego, New York, Berkely, Boston, London, Sydney, Tokyo, Toronto.
50. Akah, P. A., Okafor, C. L., 1992. Blood Sugar lowering effect of *Vernonia amygdalina* (Del) in an experimental rabbit model. *Phytotherapy Res.*, 6, 171-173.
51. Goji, A.D., Dikko, A.A., Bakari, A.G., Mohammed, A., Ezekiel, I., Tanko, I., 2009. Effect of aqueous ethanolic stem bark extract of *Commiphora africana* on blood glucose levels on normoglycemic wistar rats. *Inter. J. Animal Vet. Advan.* 1, 22-24.
52. Aragao, D. M., Guarize, L., Lanini, J., Da costa, J. C., Garcia, R. M., Scio, E., 2010. Hypoglycemic effects of *Cecropia pachystachya* in normal and alloxan-induced diabetic rats. *J. Ethnopharmacol.* 128, 629-633.
53. Orisakwe, O.R., Afonne, O.J., Chude, M.A., Obi, E., Dioka, C.E., 2003. Sub chronic toxicity studies of the aqueous extract of *Boerhavia diffusa* leaves. *Journal of Health Science*, 49, 444-447.
54. Sachin, J. & Sahni, Y. P., 2010. Biochemical Changes in Goats treated with anthelmintic indigenous herbs. *J. Vet. World*, 3(7), 315-317.
55. Silanikove, N. & Tiomkin, D., 1992. Toxicity induced by poultry litter consumption: effect on parameters reflecting liver function in beef cow. *Animal Prod.* 54, 203-209.
56. Adama, J. Y., Ajanusi, O. J., Chiezey, N. P., Lawal, A., 2011. Haematological responses of Yankasa sheep to experimental *Fasciola gigantica* infection in Zaria, Nigeria. *Agricultural and biology J. North America*, 2(8), 1232-1238
57. Raghuvansi, S. K. S., Tripathi, M. K., Mishra, A. S., Chaturvedi, O. H., Prasad, R., Saraswat, B.L. & Jakhmola, R.C., 2007. Feed digestion, rumen fermentation and blood biochemical constituents in Malpura rams fed a complete feed-block diet with inclusion of tree leaves. *Small Rum. Res.* 71, 21-30.
58. Zapta, B., Fuentes, V., Bonacic, C., Gonzalez, B., Villouta, G., Bas, F., 2003. Haematological and clinical biochemical findings in captive juvenile guanacos in Central Chile. *Small Ruminant Res.* 48, 15-21.
59. Dixit, V. P., Varma, M., Marthur, N.T., Marthur, R., Sharma, S., 1992. Hypocholesterolaemic and anti-arteriosclerotic effects of solasodine (C₂₇H₄₂O₂N) in cholesterol-fed rabbits. *Phytotherap. Res.* 6, 270-273.

60. Viana, S.B., Medeiros, C.C., Lacerda, A.C., Leal, A.M., Vale, T.G., Matos, F.A., 2004. Hypoglycemic and anti-lipemic effects of the aqueous extract from *Cissus sicyoides*. *BMC Pharmacol.* 4, 9.
61. Sharma, B., Balomajumder, C., Roy, P., 2008. Hypoglycemic and hypolipidemic effects of flavonoid rich extract from *Eugenia jambolana* seeds on streptozotocin induced diabetic rats. *Food chem. Toxicol.* 7, 237- 238
62. Zhou, T., Lou, D., Li, X., Luo, Y., 2009. Hypoglycemic and hypolipidemic effects of flavonoids from lotus (*Nelumbo nucifera* Gaertn) leaf in diabetic mice. *J. Med. Plant Res.* 3, 290-293.
63. Sushruta, K., Satyanarayana, S., Srinivas, N., Sekhar, S., Raja, J., 2006. Evaluation of the blood-glucose reducing effects of aqueous extracts of the selected umbelliferous fruits used in culinary practice. *Trop. J. Pharmaceut. Res.* 5, 613-617.
64. Uma, R. B., 2010. Acute and sub acute toxicity of Amalakyadi Churna. *Pharmacology online*, 1, 625-633.
65. Roy, S., Ukil, B., Lynden, L. M., 2016. Acute and sub-acute toxicity studies on the effect of *Senna alata* in Swiss Albino mice, *J. Cogent biol.* 2(1), 1-10 Agboluaje
66. Bello, I., Bakkouri, A. S., Tabana, Y. M., Hindi, B. A., Mansoub, M. A. A., Mahmud, R., Asmawi, M. Z., 2016. Acute and sub-acute toxicity evaluation of the methanolic extract of *Alstonia scholaris* stem bark. *Med. Sciences*, 4, 4.
67. Ping, K. Y., Darah, I., Chen, Y., Sreeramanan, S., Sasidharan, S., 2013. Acute and subchronic toxicity study of *Euphorbia hirta* L. methanol extract in rats. *BioMed Res. Internat.* 182064.
68. Nabukenya, I., Rubaire-Akiiki, C., Mugizi, D., Kateregga, J., Olila, D., Hoglund, J., 2014. Sub-acute toxicity of aqueous extracts of *Tephrosia vogelii*, *Vernonia amygdalina* and *Senna occidentalis* in rats. *Nat. Prod. Chem. & Res.* 2, 143.

Figures




Figure 1

Comparative Photomicrographs of Section of Goat Liver Exposed to Different dosage of *CMESB* of *Parkia biglobosa*. No Observable Microscopic Difference Between treatment groups and control group

Keys

Control = Liver of goat exposed to 5 ml/kg of distilled water . The liver showed central vein and normal hepatocytes (H & E, 400).


A= Liver of goats administered with 1000mg/kg of *CMESB* of *P. biglobosa*. There was passive congestion of the central vein, dilated sinusoids and normal hepatocytes (H & E, 400).

B= Liver of goats administered with 2000mg/kg of CMESB of *P. biglobosa*. The liver cells showed central vein and normal hepatocyte (H & E, 400).

C=Liver of goat administered with 3000mg/kg of CMESB of *P. biglobosa*. There was Infiltration of eosinophils around the central vein and slight dilation of sinusoid (H & E, × 400).

D=Liver of goat treated with 4000mg/kg of CMESB of *P. biglobosa*. There was infiltration of eosinophils around the central vein and slight dilation of sinusoid (H & E, × 400).

E=Liver of goat treated with 5000mg/kg of CMESB of *P. biglobosa*. There was infiltration of eosinophils around the central vein and slight dilation of sinusoid (H & E, × 400).



CONTROL




Figure 2

Comparative Photomicrographs of Section of Goats Kidney Exposed to Different dosage of CMESB of *P. biglobosa*. No Observable Microscopic Lesions in all groups when compared to Ccontrol group

Keys

Control= Kidney of goat administered with 5 ml/kg of distilled water (control). Bowman's capsule and the glomerulus are intact (H & E, ×100)

A= Kidney of goat administered 1000mg/kg of CMESB of *P. Biglobosa* Bowman's capsule and the glomerulus are intact (H & E, ×100)

B=Kidney of goat administered with 2000mg/kg of CMSBE of *P. biglobosa*. Bowman's capsule and the glomerulus are intact (H & E, ×100)

C= Kidney of goat administered 3000mg/kg of CMESB of *P. Biglobosa* Bowman's capsule and the glomerulus are intact (H & E, ×100)

D= Kidney of goat administered 4000mg/kg of CMESB of *P. Biglobosa* Bowman's capsule and the glomerulus are intact (H & E, ×100)

E= Kidney of goat administered 5000mg/kg of CMESB of *P. Biglobosa* Bowman's capsule and the glomerulus are intact (H & E, ×100)