

A new synonym of *Rhododendron leishanicum* (subg. *Hymenanthes*) from China (Ericaceae)

Sheng Chen¹, Lei Hao¹, Yi-Hui Deng², Cheng-Hua Yang³, Jun-Mei Yuan¹, Jia-Xin Lou¹, Yun-Song He¹, Xiang Chen¹

¹ Institute of Biology, Guizhou Academy of Sciences, Guiyang 550009, Guizhou, China

² Liuzhou Forestry Research Institute, Rongshui 545300, Guangxi, China

³ Guizhou Academy of Forestry, Guiyang 550005, Guizhou, China

Corresponding author: Xiang Chen (xiangchengz@163.com)

Abstract

Based on a critical examination of type specimens, images of living plants, and the literature has shown *Rhododendron oligocarpum* to be conspecific with *R. leishanicum*. Although slight variations in corolla colour exist amongst different populations of *R. oligocarpum*, it does not serve as a key distinguishing trait. Therefore, we reduced *R. oligocarpum* to a synonym of *R. leishanicum*, and recommend placing it in Subsection *Maculifera*.

Key words: Morphology, new synonym, *Rhododendron*, *Rhododendron leishanicum*



Academic editor: Bing Liu
Received: 20 February 2024
Accepted: 28 May 2024
Published: 14 June 2024

Citation: Chen S, Hao L, Deng Y-H, Yang C-H, Yuan J-M, Lou J-X, He Y-S, Chen X (2024) A new synonym of *Rhododendron leishanicum* (subg. *Hymenanthes*) from China (Ericaceae). *PhytoKeys* 243: 9–14. <https://doi.org/10.3897/phytokeys.243.121263>

Copyright: © Sheng Chen et al.
This is an open access article distributed under terms of the Creative Commons Attribution License (Attribution 4.0 International – CC BY 4.0).

Rhododendron L. which contains ca. 1200 species, is the largest genus of Ericaceae (Chamberlain et al. 1996; MacKay and Gardiner 2017). This genus is widely distributed in Asia, Europe and North America, of which the great majority (ca. 900) occurs in China and the Malaysian archipelago, the centres of diversity being in the Himalayas and South East Asia (Fang et al. 2005; Brown et al. 2006; Gibbs et al. 2011; Liu et al. 2020). When we examined the type specimens of *Rhododendron* from Guizhou Province, China, *R. leishanicum* W. P. Fang et S. S. Chang ex D. F. Chamb. and *R. oligocarpum* W. P. Fang et S. S. Chang were found to be morphologically very similar and identified as very confusing in terms of leaf, flower, and fruit characters.

Rhododendron leishanicum was originally described by Chamberlain (1982), based on a single collection, Austro-Guizhou Exped 909, from Leigong mountain in Lei Shan Xian, Guizhou Province, China (Fig. 1A). In the protologue, Chamberlain placed *R. leishanicum* in the Subsection *Williamsiana*. The next year, *R. oligocarpum* was described by Fang (1983), based on six collections (including Z. S. Zhang et al. 401557 (Types: HGAS0088928, IBSC0481928, PE01297915, PE01297916), Z. S. Zhang 58 (HGAS007915), T. H. Tu 31739 (SZ0036179), from Fanjing Mountain; Austro-Guizhou Exped 1411 (PE00312607, PE00313389, KUN540382), from Leigong Mountain and G. Z. Li 6211 & 11277 (IBK00187538, IBK00187539 & IBK00187541, IBK00187559), from Maoer Mountain) from three different origins in China (Fig. 1B–D). According to the protologue, Fang placed *R. oligocarpum* in the Subsection *Maculifera*. Meanwhile, *R. leishanicum*

was again described as a new species by Fang in the same protologue. It is worth noting that a paratype of *R. oligocarpum*, Austro-Guizhou Exped 1141, was collected from Leigong Mountain (Fig. 1C). However, the relationship between *R. oligocarpum* and *R. leishanicum* was not mentioned by Fang when he described *R. leishanicum* as a new species again in the publication in 1983.

According to critical examination of the type specimens and morphological comparisons of *R. leishanicum* and *R. oligocarpum*, we found that both species have setose young branches and petioles, with small apiculate leathery leaves, racemose-umbellate inflorescences, corolla campanulate, with deep purple basal spots, pedicel, ovary and filament with identical indumentum (Fig. 2). It is worth noting that during our field surveys of populations of these two species, we observed a certain difference in colour of the corolla between the Guangxi population of *R. oligocarpum* and those in Guizhou. However, variation in corolla colour is known to occur within species of the genus *Rhododendron* (Jin 2006; Zhou 2008); hence, the differences in corolla colour between them fall within a normal range of variation. Therefore, we place *R. oligocarpum* in synonymy with *R. leishanicum* according to ICN rules (Turland et al. 2018).

Based on the setose branchlets and petioles, the small apiculate leaves, the campanulate corolla and the glabrous style, we suggest placing *R. leishanicum* in the Subsection *Maculifera*.

Taxonomy treatment

***Rhododendron leishanicum* W. P. Fang & S. S. Chang ex D. F. Chamb.**

Type. CHINA. Guizhou province: Leishan Xian, Leishan, 1850 m elev, 29 April 1959, Austro-Guizhou Exped 909 (holotype: SZ; isotype: HGAS007912!; KUN0540381!).

***Rhododendron oligocarpum* W. P. Fang & S. S. Chang, syn. nov.**

Type. CHINA. Guizhou province: Yinjiang county, Sanduodian, 2070 m elev, 14 April 1964, Z. S. Zhang et al. 401557 (holotype: HGAS0088928!; isotypes: IBSC0481928!; PE01297915!, PE01297916!; paratype: HGAS007915!; IBK00187538!, IBK00187539!, IBK00187541!, IBK00187559!; PE00312607!, PE00313389!; KUN540382!; SZ0036179!).

Description. Shrubs or small trees, 3–6 m tall; bark grey, dehiscent on drying; **branchlets** terete, setose when young, gradually glabrescent. **Petiole** rounded abaxially, furrowed adaxially, 5–15 mm, densely setose; **leaf** blade leathery, elliptic or obovate, 4–8 × 2–4 cm; base rounded or subcordate; margin revolute, apex rounded, with small apiculate; adaxial surface green, abaxial surface pistachio; mid-rib impressed abaxially, grooved adaxially, glabrous or sometimes sparsely hairy; lateral veins 11–15-paired, slightly conspicuous on both surfaces. **Inflorescence** racemose-umbellate, 3–5-flowered; rachis 5–7 mm, tomentose. **Pedicel** 1–2 cm, densely setose-tomentose; **Calyx** small, discoid, lobes 5, triangular, tomentose, 1–2 mm. **Corolla** campanulate, purple-red to rose-red, with deep purple basal spots, 3–3.5 cm; lobes 5, suborbicular, ca. 1.5 × 2 cm, apex emarginate; **stamens** 10, unequal, 1–3 cm, filaments puberulent at base;



Figure 1. *Rhododendron leishanicum* and *R. oligocarpum* A isotype of *R. leishanicum* B holotype of *R. oligocarpum* C, D paratype from Leigong Mountain and Maoer Mountain of *R. oligocarpum*, respectively.




Figure 2. Comparison of *Rhododendron leishanicum* and *R. oligocarpum* **A, D, G, J, M** *R. leishanicum* from Ligong mountain **B, E, H, K, N** *R. oligocarpum* from Fanjing mountain **C, F, I, L, O** *R. oligocarpum* from Maoer mountain **A–C** branchlets with flowers **D–F** adaxial and abaxial surface of leaf **G–I** corolla with deep purple basal spots **J–L** pedicel, Calyx, ovary and style indumentum **M–O** stamens side view.

ovary cone-shaped, 4–5 mm, densely densely setose and tomentose; **style** 2.5–3 cm, glabrous, stigma slightly expanded. **Capsule** cylindrical, 20–25 × 7 mm, rough. Flowering from April to May and fruiting from September to October.

Distribution and habitat. *Rhododendron leishanicum* is distributed in Eastern Guizhou and North-eastern Guangxi. It grows in Thickets at 1800–2500 m a.s.l.

Specimens examined. CHINA. Guizhou: Qiandongnan Prefecture, Leigong Mountain, *P. L. Song 1016* (GYBG barcode 0002282, 0002283), *4407* (CCNU barcode 9018123), *4717* (CCNU barcode 9018122); Tongren City, Fanjing Mountain, *Z. S. Zhang et al. 401673* (IBSC barcode 0481934), *401317* (HGAS barcode 0088940, IBSC barcode 0481929), *400681* (HGAS barcode 0088933), *400610* (HGAS barcode 0088939), *Z. P. Jian 32039* (HGAS barcode 0088938), *Wuling Mountain Expedition 731* (GFS barcode 0007355, KUN barcode 0339467, 0339468, PE barcode 00258535), *1348* (GFS barcode 0007356, KUN barcode 0339469). **Guangxi:** Guilin City, Maoer Mountain, *G. Z. Li 11272* (IBK barcode 00187540), *12378* (IBK barcode 00187542), *12084* (IBK barcode 00187561), *F. X. Jin 1066* (HTC barcode 0010523, 0010524, 0010525), *J. X. Zhong 83311* (IBK barcode 00187551, 00187552, IBSC barcode 0481913), *83523* (IBK barcode 00187560), *81647* (IBK barcode 00187566, 00187567), *L. M. Gao 20077* (KUN barcode 0767388, 0767389).

Acknowledgements

We thank the curators of PE, IBSC, KUN, IBK, SZ and HGAS for allowing us to examine specimens or use images of specimens and the National Specimen Information Infrastructure (NSII, <https://www.nsii.org.cn/2017/>) for availability of digitised specimens.

Additional information

Conflict of interest

The authors have declared that no competing interests exist.

Ethical statement

No ethical statement was reported.

Funding

This study was supported by Young Scientists Fund of Guizhou Academy of Sciences (Qiankeyuan Key-J [2024] 15) and National Natural Science Foundation of China (31160037) and Guizhou Provincial Basic Research Program (Natural Science)-Study on the Type Specimens of Vascular Plants in Guizhou Based on DELTA System (Qiankehe Fundamentals-ZK [2023] General 234).

Author contributions

Conceptualisation: XC, LH, SC. Data curation: YHD, CHY, JMY, JXL, YSH. Writing – original draft: SC, LH, XC. Writing – review and editing: SC, XC.

Author ORCIDs

Sheng Chen  <https://orcid.org/0000-0002-2727-6504>

Lei Hao  <https://orcid.org/0009-0001-0276-6523>

Yi-Hui Deng  <https://orcid.org/0009-0006-2821-0247>
Chen-Hua Yang  <https://orcid.org/0000-0002-5941-676X>
Jun-Mei Yuan  <https://orcid.org/0009-0007-1861-5790>
Jia-Xin Lou  <https://orcid.org/0009-0001-0276-6523>
Yun-Song He  <https://orcid.org/0009-0002-8791-558X>
Xiang Chen  <https://orcid.org/0009-0008-0773-4978>

Data availability

All of the data that support the findings of this study are available in the main text.

References

- Brown GK, Craven LA, Udovicic F, Ladiges PY (2006) Phylogenetic relationships of *Rhododendron* section *Vireya* (Ericaceae) inferred from the ITS nrDNA region. *Australian Systematic Botany* 19(4): 329–342. <https://doi.org/10.1071/SB05019>
- Chamberlain DF (1982) A Revision of *Rhododendron* II. Subgenus *Hymenanthes*. Notes from the Royal Botanic Garden Edinburgh 39(2): 209–486.
- Chamberlain DF, Hyam R, Argent G, Fairweather G, Walter KS (1996) The Genus *Rhododendron*: Its Classification and Synonymy. Royal Botanic Garden Edinburgh, Edinburgh, 181 pp.
- Fang WP (1983) New taxa of the genus *Rhododendron* from China. *Zhiwu Fenlei Xuebao* 21(4): 457–469.
- Fang MY, Fang RZ, He MY, Hu LZ, Yang HB, Chamberlain DF (2005) *Rhododendron*. Flora of China. vol. 14. Science Press, Beijing & Missouri Botanical Garden Press, St. Louis, 260–373.
- Gibbs D, Chamberlain DF, Argent G (2011) The Red List of *Rhododendrons*. Botanic Gardens Conservation International, Richmond, 4–11.
- Jin XF (2006) Taxonomic study on *Rhododendron* subgen. *Tsutsusi* sensu Sleumer (Ericaceae). PhD Thesis, Zhejiang University, China.
- Liu DT, Chang YH, Ma YP (2020) Unclear resource background seriously restricts biodiversity conservation of *Rhododendron* in China. *Plant Science Journal* 38(4): 517–524. <https://doi.org/10.11913/PSJ.2095-0837.2020.40517>
- MacKay M, Gardiner SE (2017) A model for determining ex situ conservation priorities in big genera is provided by analysis of the subgenera of *Rhododendron* (Ericaceae). *Biodiversity and Conservation* 26(1): 189–208. <https://doi.org/10.1007/s10531-016-1237-0>
- Turland NJ, Wiersema JH, Barrie FR, Greuter W, Hawksworth DL, Herendeen PS, Knapp S, Kusber WH, Li DZ, Marhold K, May TW, McNeill J, Monro AM, Prado J, Price MJ, Smith GF (2018) International Code of Nomenclature for algae, fungi, and plants (Shenzhen Code) adopted by the Nineteenth International Botanical Congress Shenzhen, China, July 2017. *Regnum Vegetabile* 159. Koeltz Botanical Books, Glashütten. <https://doi.org/10.12705/Code.2018>
- Zhou LY (2008) The study on Relationship and Genetic Diversity of *Rhododendron*. PhD Thesis, Sichuan Agricultural University, China.