



**INTERNATIONAL JOURNAL OF
PHARMACEUTICAL SCIENCES**
[ISSN: 0975-4725; CODEN(USA): IJPS00]
Journal Homepage: <https://www.ijpsjournal.com>



Review Paper

A Review on the Hepatoprotective Activity of Bombax ceiba Root Extracts and Exudates

Dr. Santosh Kumar Mishra, Vivek Singh*, Tushar Gupta, Durgesh Yadav

Department of Quality Assurance Techniques . MES's College of Pharmacy, Sonai, India.

ARTICLE INFO

Published: 14 May 2026

Keywords:

Hepatoprotective Activity,
Bombax ceiba Root,
Exudates, Extracts,
Traditional medicines,
prevention and cure

DOI:

10.5281/zenodo.20180015

ABSTRACT

Traditional medicines are becoming crucial and decisive role in the prevention and cure of diseases and disorders. Traditional medicines have various pharmacological activities which important for human health. Bombax Ceiba widely used from past decades for the treatment of diseases and disorders. Bombax ceiba is usually referred to as kapok tree and moca, which belongs to Bombacaceae. The root bark contains the 3 naphthalene derivatives. Bombax ceiba L. shows many pharmacological properties. The stem bark of the plant is used for a wound. Thorn of a plant used in different formulation to treat acne problem. Androecium is used for food purposes. The leaf shows Antidiarrheal property. Root used for the treatment of Piles. Bark contains lupeol, saponins, tannins, gums and 4, 5, 7- trihydroxyflavone-3-O- β -D-glucopyranosyl (1-4)- α -L-rhamnopyranoside, n hexacosanol. Root bark contains lactone and ether. The seed contains palmitic acid and flower contains polysaccharide. Leaves contain Shamimin. This review paper provides an overview on Pharmacological, Phytomedicinal properties of the plant. Plants have played a signi cant role in traditional medicine for treating a wide range of human ailments. Among the many medicinal herbs used in Unani medicine, Bombax ceiba Linn. has been employed for centuries. This herbaceous plant is renowned for its impressive height, reaching approximately 150 feet. It can be found in temperate and tropical regions of Australia, Africa, and, Asia with occurrences in India at altitudes of up to 1500 meters. The indigenous communities and forest dwellers extensively utilize various components of this plant, including the root, ower, gum, leaf, prickles, stem bark, fruit, seed, and heartwood, to address diverse diseases. Ethnobotanical research reveals that Bombax ceiba Linn. is effectively employed in the treatment of ailments such as diarrhea, boils, wounds, leprosy, acne, and various other skin conditions. Furthermore, it has been used as an anthelmintic since ancient times.

***Corresponding Author:** Vivek Singh

Address: Assistant Professor, Sagar college of pharmacy Barabanki, Uttar Pradesh

Email ✉: tusharggj@gmail.com

Relevant conflicts of interest/financial disclosures: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.



INTRODUCTION

Throughout history, nature has been an invaluable source of therapeutic agents, leading to the discovery of groundbreaking drugs derived from natural origins [1]. India, often referred to as the "Botanical Garden of the world," is renowned for its abundant variety of therapeutic plants. These plants have played a crucial role in advancing material medica, contributing significantly to human well being and promoting a healthy, disease-free life [2]. The preservation of our health is of utmost importance, and India holds a significant position in this regard. With its rich traditions in medicine and culture, India has a profound legacy of utilizing medicinal plants that continues to be respected even in modern times. The old systems of medicine in India, like Ayurveda, Unani, and Siddha, are vital in maintaining the well-being of individuals [3]. Plants have long been recognized for their remarkable biochemical capabilities and have been integral to the field of phytomedicine for centuries [4]. Different parts of plants, including bark, leaves, flowers, roots, fruits, and seeds, serve as potential sources of natural compounds with medicinal properties. In the tropical regions of India Silk Cotton Tree, scientifically known as *Bombax ceiba*, holds significant importance as a medicinal plant. This tall deciduous tree exhibits a straight buttressed trunk and expansive branches. According to Ayurveda, *Bombax ceiba* is renowned for its proven medicinal properties and finds applications in numerous formulations. Virtually every part of this plant possesses medicinal value, with the roots and flowers being particularly noteworthy for their efficacy in treating various ailments [5]. Belonging to the Bombacaceae family, *Bombax ceiba* Linnaeus is a notable species among approximately 26 species and nearly 140 pantropical classes. It is usually referred to as Semal, Simbal, Simul, Indian kapok, Katsavar,

Indian bombax, or Red Silk cotton tree. In India, it can be found at altitudes of up to 1500 meters. This tree thrives in both dry and moist deciduous forests, as well as in the vicinity of rivers, particularly in peninsular India. *Bombax ceiba* prefers well drained soils, especially deep sandy loams, and is characterized as a fast-growing species that requires ample sunlight. It flourishes in regions with annual rainfall ranging from 50 to 460 centimeters, which is evenly distributed throughout the year [6]. *Bombax ceiba* is a deciduous tree that can reach heights of up to 45 meters. Its trunk is straight and is supported by buttresses measuring 1-2 meters in height. The bark has a gray color with white mottling and is approximately 20-30 mm thick. The branches are arranged horizontally in a whorled pattern. The leaves are alternately arranged and have a digitately-compound structure with small lateral stipules. The central axis of the leaf, known as the rachis, is smooth and measures 12-25 cm in length. The leaflets, usually 5-7 in number, are elliptic or elliptic-obovate in shape, with dimensions of 10-20 x 2-6 cm. The leaf margins are smooth, and the entire leaf surface is free from hairs. The flowers of *Bombax ceiba* are bisexual and have a dark crimson color. They are large, with a diameter of 6-7 cm, and can appear individually or in clusters of 2-5 flowers.[7] The calyx, which forms the outer part of the flower, has a bell shaped structure, and its lobes measure about 3-4 x 3 cm. The outer surface of the calyx is smooth, while the inner surface is covered in silky hairs. The petals, numbering 5, are fleshy and elliptic-obovate, with measurements of 8.5 x 1.8 x 3.5-5 cm. The tree produces a substantial number of stamens, ranging from 65 to 80, which can reach lengths of 3-7.5 cm. The ovary is conical and covered in fine hairs, and it contains 5 compartments with numerous ovules. The style, the elongated part of the female reproductive structure, is longer than the stamens [8]. The fruit of *Bombax ceiba* is a capsule that



measures 8-10 x 3 cm. It has a cylindrical shape and is covered in soft, tomentose hairs. As it matures, the capsule turns blackish and becomes glabrous. Inside the fruit, numerous seeds are present. These seeds are pyriform (pear-shaped), smooth, and have a dark brown color. They are surrounded by a distinctive white cotton-like material. Systems of medicine like Ayurveda, various parts of *Bombax*, including seeds, fruits, roots, leaves, bark, flowers, and gum, have been identified for their medicinal properties. The *Bombax* tree is a tall deciduous tree that can reach heights of up to 40 meters. It features horizontally spreading branches, and the young stems are adorned with hard prickles. The bark of the tree displays a grey-brown or silver-grey color and is adorned with sharp conical prickles. The leaves are large, spreading, and possess a smooth texture. They consist of lanceolate leaflets, typically numbering 3-7, with smooth margins.[9]

1.0. Pharmacological Activity:

• Antioxidant

Bombax ceiba, is a medicinal plant belonging to the Bombacaceae family [10]. The plant extracts of *B. ceiba* have been studied for their antioxidant, anti-inflammatory, and antibacterial properties. They have shown potential in managing diabetes and hyperglycemia. While certain parts of *B. ceiba* flowers are consumed in Thai culture, limited research has been conducted to explore their biological and pharmacological effects. Chemical analysis of *B. ceiba* leaf and flower extracts has revealed many flavonoids, saponins, tannins, terpenoids, and cardiac glycosides. Notably, the flower extracts exhibit significant concentrations of total phenolics and flavonoids. In vitro studies have demonstrated the antioxidant and anti-diabetic activities of the flower extracts, which contain phytochemicals such as anthocyanin, rutin, quercetin, and apigenin [11]. In recent years,

various chemical-based tests have been developed and modified to detect antioxidant activities. Primary antioxidants play a vital role in inhibiting oxidation by interrupting chain reactions and neutralizing free radicals. Secondary antioxidants act as metal chelating agents, helping to restore primary antioxidants and stabilize singlet oxygen. These tests have diverse applications in evaluating antioxidant properties [12]. Flavonoid compounds, known as secondary natural antioxidants, have been found to exhibit stronger inhibitory effects on glucosidase and amylase compared to acarbose in vitro. The flavonoids and phenolics present in *B. ceiba* flower extracts have the potential to suppress the activity of glucosidase and amylase, which could aid in the management of diabetes. Additionally, a 50/50 ethanolic extract of *B. ceiba* stem bark and flowers has shown hypoglycemic effects, with mangiferin being attributed to the reduction of fasting blood glucose levels. Nitrite, a highly toxic compound found in high levels in leafy and root vegetables, can have adverse effects such as methemoglobinemia through the oxidation of hemoglobin. In a specific study, *B. ceiba* extract exhibited excellent scavenging properties against nitric oxide radicals. Phenolic compounds, particularly flavonoids, are known not only as scavengers of free radicals but also for their ability to stabilize nitric oxide, peroxynitrite, and reactive oxygen species during lipid peroxidation.[13]

• Anti-Inflammatory

Chronic inflammation is associated with conditions like gastritis, rheumatoid arthritis, atherosclerosis, inflammatory bowel disease (IBD), and cancer. Excessive production of free reactive nitric oxide (NO) by inducible NO synthase (iNOS) can contribute to inflammatory disorders and autoimmune diseases like rheumatoid arthritis. Compounds that inhibit the overproduction of NO are often referred to as anti-inflammatory drugs. However, synthetic anti-inflammatory drugs

available in the market can have significant toxic side effects and may lead to a recurrence of inflammation upon discontinuation [14]. Plants provide a diverse range of secondary metabolites with potential anti-inflammatory activity, targeting various molecular pathways. Exploring the anti-inflammatory properties of natural products offers a promising approach to developing new drugs with reduced risks of side effects [15]. *Bombax malabaricum*, a plant species, has been found to exhibit significant antifungal activity against the fungus responsible for ringworm infections when extracted from its leaves.[16] Catechutannic acid is present in the bark gum of this plant, while mangiferin, an antioxidant and analgesic compound, has been isolated from its leaves. The roots of the plant contain several compounds, and the root bark is a source of lupeol, β -sitosterol, isohemigossypol-1-methyl ether, and 7-hydroxycadalene [17]

• Antibacterial

The antibacterial activity of methanol extracts from *Salmalia malabarica* was found to be potent against multi drug resistant strains such as *Salmonella typhi*, *Staphylococcus aureus*, *Micrococcus luteus* (Gram positive), *Escherichia coli*, and *Pseudomonas aeruginosa* (Gram-negative) bacteria [18]. The N-hexane and methanol extracts of *Bombax malabarica* also exhibited significant antimicrobial activity. Gram-positive bacteria including *S. aureus*, *Bacillus cereus*, *E. coli*, and *Vibrio cholerae* showed high susceptibility to the methanol extract of *B. malabarica* [19]. Furthermore, the extract demonstrated antifungal activity against *Cryptococcus neoformans* but did not exhibit effectiveness against *C. albicans* [20]. In the case of *Bombax ceiba* extracts, certain fractions showed greater susceptibility against bacterial strains such as *B. subtilis*, *S. aureus*, *E. coli*, *P. aeruginosa*, as well as fungal strains *A. niger* and

C. albicans. The bark extract in different solvents exhibited susceptibility against *A. niger* and *C. albicans*, with ethanol showing the highest activity followed by acetone and aqueous extracts. However, petroleum ether and chloroform extracts did not show any effect. *E. coli* showed the highest sensitivity among the tested microbes, with a larger zone of inhibition observed in the carbon tetrachloride fraction compared to the n-hexane and chloroform fractions [2]

• Hepatoprotective Activity

The aqueous extract of *B. malabarica* has shown protective effects against CCl₄-induced hepatotoxicity. Specifically, the xylem of the stem and root play a role in protecting the liver from histological changes associated with CCl₄ induced damage, such as fatty degeneration, cell necrosis, ballooning necrosis, lymphocytes, and Kupffer cell aggregation. Mangiferin, isolated from *B. malabarica*, has also been found to alleviate liver damage caused by CCl₄. Furthermore, mangiferin has been shown to reverse alterations in liver biochemical markers induced by antitubercular medicines (Isoniazid and Rifampicin) and paracetamol, although it does not have an impact on liver histology [22]. Excessive and chronic alcohol consumption can lead to severe liver injury. The therapeutic effects of aqueous methanol extract from *Bombax ceiba* (Bombacaceae) on liver steatosis were investigated in a study that lasted eight weeks and involved seven groups. One group served as the control, while the other six groups were subjected to various conditions. One group was fed a fatty diet, another received ethanol and a high-fat diet, and each group was administered the same dose of uvastatin (2 mg/kg/d). Another group received oral administration of BCE extract (200 mg/kg/d), and the nal group did not receive any treatment. The BCE extract contributed to weight loss and



enhanced hepatic function in alcohol induced liver injury.[23]

• **Anticancer Activity**

The antiproliferative activity of diethyl ether (DE) and light petroleum (PE) extracts from *Bombax ceiba* flowers was assessed against seven human cancer cell lines, including ACHN, COR-L23, A549, Caco-2, Huh-7D12, and C32 [24]. Both DE and PE extracts displayed strong inhibition of tumor cell viability, particularly against ACHN cells, in a dose-dependent manner. The IC₅₀ values for PE and DE were determined as 45.5 µg/mL and 53.2 µg/mL, respectively [25]. Flavonoid-rich extracts obtained from *B. ceiba* flowers were screened for their impact on fatty acid synthase (FAS) in various cancer cells [26]. FAS is known to be overexpressed and hyperactive in certain cancers. The *B. ceiba* extract exhibited significant inhibition of FAS activity across different cancer cells. Among the tested cells, N87 gastric cancer cells exhibited the lowest FAS activity, while A549 lung cancer cells showed the highest. The flavonoid-rich extract demonstrated inhibitory effects on FAS with a minimum inhibitory concentration of 247.98 µg/mL, using A549 cells [27]. The methanolic extract of *B. ceiba* demonstrated minimal cytotoxicity in the Vero cell line based on a mitochondrial activity assay [28]. The anticancer potential of the methanol extract from *B. ceiba* root was evaluated using a brine shrimp lethality bioassay, with vincristine sulfate as the standard cytotoxic agent. The LC₅₀ (50% mortality) and LC₉₀ values for the crude extract were determined as 3.90 µg/mL and 150.0 µg/mL, respectively [29]. Additionally, the methanolic extract of *B. ceiba* leaves exhibited antioxidant activity and showed significant enhancements in neutrophil adhesion, carbon clearance from blood, delayed-type hypersensitivity (DTH) response.[30]

• **Anti-Diabetic Activity**

Shamimin, obtained from the leaves of *Bombax ceiba* at a dosage of 500 mg/kg, has been identified as a hypoglycemic agent in rats [31]. The hydromethanolic (2:3) extract of *Salmalia malabarica* sepals demonstrated a significant reduction in Fasting Blood Sugar and Glycated Hb (HbA1C) levels in STZ-induced diabetic rats. This extract also restored the activity of specific carbohydrate metabolic enzymes and countered the hyperactivity of glucose-6-phosphatase in the liver and skeletal muscle, which were impaired due to STZ induction. Moreover, the extract alleviated elevated oxidative stress levels and restored SGOT and SGPT levels [32]. The n-hexane fraction of this hydromethanolic extract also exhibited significant hypoglycemic and hypolipidemic effects. Additionally, the n-hexane fraction increased serum insulin levels and hemoglobin concentration while decreasing glycated hemoglobin levels. This fraction was also found to be beneficial in preserving the islets of Langerhans in diabetic rats [33]. Certain compounds present in *Bombax ceiba*, such as quercetin and epicatechin, were identified as potent inhibitors of the α -glucosidase enzyme, with inhibitory rates of 50.5% and 48.3%, respectively [34]. Furthermore, glucosylxanthone derived from the plant has been investigated as a potential target for new anti-diabetic medication, particularly due to its inhibitory effect on DPP-IV. In silico binding studies revealed that glucosylxanthone and its analogues exhibited comparable binding activity to FDA-approved medicines and other compounds under research.[35] Inhibiting DPP-IV could lead to increased serum Glucagon-like Peptide-1 (GLP-1), resulting in a net hypoglycemic effect. Additionally, the aqueous (at a dose of 100 mg/kg) and ethanolic (at a dose of 200 mg/kg) extracts of *Bombax malabarica* bark demonstrated beneficial effects in alloxan-induced diabetic rats [36]. The anti-diabetic activity observed of the extract could be attributed to the antioxidant properties induced

by compounds such as isoorientin, vitexin, isomangiferin, quercetin, hexoside, mangiferinisovitexin, and nigricanside [37]. In a study conducted on the therapeutic potential of a standardized extract derived from *Bombax ceiba* leaves (BCE) was investigated in rats with type 2 diabetes mellitus (T2DM) [38]. The administration of BCE resulted in a significant decrease in fasting blood glucose levels and improved oral glucose tolerance in T2DM rats.[39]

• Anti-Obesity Activity

Bombax ceiba Linn. has a rich traditional history of being used to address various ailments, including diarrhea, dysentery, digestive disorders, diabetes, and imbalances of the three doshas (tridoshas). It is recognized for its positive effects on digestion and its ability to modulate insulin, leptin, and integrin signaling by stimulating PTP-1B.[40] This stimulation results in increased fatty acid synthase (FAS) activity, which can contribute to obesity. The powdered root of *B. ceiba* has demonstrated significant effects in modifying coronary risk factors, such as atherogenic lipids, brinogen, and oxidative stress, in individuals with ischemic heart disease. Its antioxidant activity is attributed to its high phenolics and tannins content [41]. In a study conducted on male Wistar albino rats, the anti-obesity efficacy of *Bombax ceiba* Linn. was investigated using a high-fat diet-induced obesity model[42]. After ten weeks of being fed a high-fat diet, the rats developed experimental obesity. From the 7th to the 10th week, the rats were orally administered a dose of 100 mg/kg of *B. ceiba* extract and 50 mg/kg of gembrozil. Significant result shows in this study. However, treatment with *B. ceiba* extract and gembrozil effectively reduced these obesity induced alterations.[43]

2.0. Ethnomedicinal activity of *bombax ceiba* L.

Fruit: Good for leprosy used as a cooling agent, digestible, shows the diuretic property, antioxidant, tonic, aphrodisiac, expectorant, shows good action in the genital, urinary organ. It is also used for bladder, kidney, chronic inflammation, and blood purifier.[44]

Flower: The flowers show bitter action and also shows acrid cooling, dry, astringent to the bowels, anti-inflammatory action. It removes bile and phlegm of the body and purifies the blood; it is beneficial to the spleen and shows a good response in leucorrhoea. It is also used topically to skin affections as cooling and astringent. It having phenolic compound.[45]

Seed: Seeds show good action in chickenpox, smallpox, catarrhal affections, chronic cystitis, and genitourinary diseases.[46]

Bark: Stem bark is used in the healing of the wound and as a paste in water to skin eruptions, boils, acne, pimples. The stem bark is also used as a demulcent, styptic, and used to removes phlegmatic. Aqueous extract with curd shows great action in treatment given for dysentery with blood.[47]

Leaves: It is used in the treatment of diarrhea used in treatment for inflammation, larvicidal activity.[48]

Root: Root bark is used for the treatment of piles. The root shows diuretic and astringent property. It is useful in biliousness, inflammations, and excessive heat of the body.[49]

Gum: The gum is an acrid, astringent, demulcent, tonic, aphrodisiac, and removes black bile. In powder form alone or with other herbs, it is used internally to treat hemoptysis, diarrhea, dysentery, bleeding piles and menorrhagia, leucorrhoea, and spermatorrhoea, and blood disorders. Topically it is applied as a styptic, astringent, demulcent in stomatitis, dermatological ailment, and burn wound.[50]

Other uses of the Plant *Bombax ceiba* L.



- Mangiferin isolated from the plant *Bombax malabaricum* shows antifungal properties.
- Stem bark of plant *Bombax ceiba* L. contains some natural dyes (Vadwala and Kola).
- Antibacterial activity of aqueous bark extract of *Bombax ceiba* L.[51-52]
- Phenolic compound from the flower of *Bombax malabaricum* shows antioxidant and antiviral properties.
- *Bombax ceiba* L leaf extract shows larvicidal activity against the filarial vector *Culex quinquefasciatus*
- The seed of *Bombax ceiba* L shows good nutritional property.[53-54]

CONCLUSION

The above literature survey revealed that *Bombax ceiba* L contains so much medicinal properties. It is used in many herbal formulations and traditional medicinal systems. It contains many phytoconstituent, which shows great medicinal properties. *Bombax ceiba* L shows antioxidant, larvicidal, diuretic, tonic, anti-leprosy, expectorant properties. It is also used in treatment in blood purifier, inflammation of the bladder, kidney calculi affection; it shows antimicrobial, analgesic, oxytocic, urolithiasis property. *Bombax ceiba*, commonly known as *B. ceiba*, has a diverse and extensive traditional history of medicinal use. This traditional knowledge has been validated by scientific investigations, further supporting the plant's therapeutic potential. Moreover, considering its ecological and economic significance, it is crucial to emphasize the importance of conserving *B. ceiba* from an ecological standpoint. Throughout history, *B. ceiba* has been utilized in the treatment of a wide range of ailments. It has been employed for conditions such as dysentery, menorrhagia (excessive menstrual bleeding), various skin problems, hemorrhoids, snakebites, scorpion stings, boils, leucorrhoea (vaginal discharge), internal bleeding, calculus affections

(stones in the body), chronic inflammation, ulcers in the bladder and kidneys, gonorrhea, hemoptysis (coughing up blood), in uenza, enteritis (intestinal inflammation), pulmonary tuberculosis, cystitis (bladder inflammation), and bleeding piles.

REFERENCES

1. Rameshwar V, Kishor D, Tushar G, Siddharth G, Sudarshan G. A Pharmacognostic and pharmacological overview on *Bombax ceiba*. *Scholars Academic Journal of Pharmacy*. 2014; 3(2): 100-7.
2. Achour Y, Bahsis L, Ablouh EH, Yazid H, Laamari MR, El Haddad M. Insight into adsorption mechanism of Congo red dye onto *Bombax buonopozense* bark Activated-carbon using Central composite design and DFT studies. *Surfaces and Interfaces*. 2021 Apr; 23: 100977. doi: 10.1016/j.sur.2021.100977.
3. Rathore D and Singh R. Phytochemistry and pharmacology of genus *Bombax*. *The Natural Products Journal*. 2019 Sep; 9(3): 184-96. doi: 10.2174/2210315508666180831095836.
4. Karole S, Gautam G, Gupta S. Pharmacognostic and pharmacological profile of *Bombax ceiba*. *Asian Journal of Pharmaceutical Education and Research*. 2017 Jul; 6(3): 16-27.
5. Wang G, Lin B, Tian M, Zhu K, Xie G, Qin M. Studies on chemical constituents from the roots of *Bombax ceiba*. *Journal of Tropical and Subtropical Botany*. 2017; 25(4): 387-93.
6. Joshi KR, Devkota HP, Yahara S. Chemical analysis of flowers of *Bombax ceiba* from Nepal. *Natural Product Communications*. 2013 May; 8(5): 1934578X1300800508. doi: 10.1177/1934578X1300800508.
7. Zhang YB, Wu P, Zhang XL, Xia C, Li GQ, Ye WC, et al. Phenolic compounds from the flowers of *Bombax malabaricum* and their



- antioxidant and antiviral activities. *Molecules*. 2015 Nov; 20(11): 19947-57. doi: 10.3390/molecules201119660.
8. Assogba GA, Fandohan AB, Gandji K, Salako VK, Assogbadjo AE. *Bombax costatum* (Malvaceae): state of knowns, unknowns, and prospects in West Africa. *BASE*. 2018 Mar; 22(4): 267-75. doi: 10.25518/17804507.16652. doi: 10.25518/1780-4507.16652.
9. Musah M, Muhammad HI, Mathew JT, Azeh Y, Umar MT. Proximate, Minerals and Functional Properties of *Bombax buonopozense* Celyx. *Communication in Physical Sciences*. 2021 Jun; 7(2): 126-33.
10. Anandarajagopal K, Sunilson JA, Ajaykumar TV, Ananth R, Kamal S. In-vitro anti-inflammatory evaluation of crude *Bombax ceiba* extracts. *European Journal of Medicinal Plants*. 2013 Jan; 3(1): 99. doi: 10.9734/EJMP/2013/2113.
11. Tilaoui M, Achibat H, Lébri M, Lagou S, Ait Mouse H, Zazouli S, et al. Phytochemical screening, antioxidant and in vitro anticancer activities of *Bombax buonopozense* stem bark extracts. *Biotechnology & Biotechnological Equipment*. 2021 Jan; 35(1): 1662-8. doi: 10.1080/13102818.2021.1997156.
12. Akhtar N and Mustafa R. Antibacterial and antioxidant potential of stem bark extract of *Bombax ceiba* collected locally from south Punjab area of Pakistan. *African Journal of Traditional, Complementary and Alternative Medicines*. 2017 Feb; 14(2): 9-15. doi: 10.21010/ajtcam.v14i2.2.
13. Guang-Kai XU, Xiao-Ying QI, Guo-Kai WA, Guo-Yong XI, Xu-Sen LI, Chen-Yu SU, et al. Antihyperglycemic, antihyperlipidemic and antioxidant effects of standard ethanol extract of *Bombax ceiba* leaves in high-fat-diet-and streptozotocin-induced Type 2 diabetic rats. *Chinese Journal of Natural Medicines*. 2017 Mar; 15(3): 168-77. doi: 10.1016/S1875-5364(17) 30033-
14. X. Kriintong N and Katisart T. In vitro antioxidant and antidiabetic activities of leaf and ower extracts from *Bombax ceiba*. *Pharmacognosy Research*. 2020 May; 12(2): 194-8. doi: 10.4103/pr.pr_116_19.
15. Hossain E, Sarkar D, Chatterjee M, Chakraborty S, Mandal SC, Gupta JK. Effect of methanol extract of *Bombax malabaricum* leaves on nitric oxide production during in ammation. *Acta Poloniae Pharmaceutica*. 2013 Mar; 70(2): 255-60.
16. Meenakshi SC, Beldal BS, Londonkar RL. Evaluation of methanolic extract of *Bombax ceiba* bark for in vitro antioxidant and Anti-inflammatory activities. *Journal of Pharmacognosy and Phytochemistry*. 2019 Jan; 8(1): 1504-7.
17. Mane RS and Vedamurthy AB. Protection of *Bombax Ceiba* by Revealing their Fungal Endophytic Diversity and Therapeutic Applications.(2020). *International Journal of Life Science and Pharma Research*. 2020 Apr; 10(2): 18-25.
18. Shah S, Ghimire K, Gupta AK, Pokhrel P, Banerjee J, Khanal H, et al. Evaluation of phytochemical parameters, antibacterial and antihelminthic activity of leaves and bark extracts of plant *Bombax ceiba*. *Journal of Applied Pharmaceutical Research*. 2017 Jul; 5(3): 38-44. doi: 10.18231/2348-0335.2017.0006.
19. Sada A, Abdur R, Sumbul R, Zakir SM. Evaluation of Antibacterial Potential of Extracts of Mocharas (gum of *Bombax malabaricum*) against certain bacterial strains. Available at: *Indian Drugs*. 2020; 57(9): 45-52.
20. doi: 10.53879/id.57.09.11878. Shukla RK, Nandan K, Shukla A, Kaur A, Rana D. Review on Traditional uses, Biological

- activities, Phytoconstituents of *Bombax ceiba* Linn. *Research Journal of Pharmacy and Technology*. 2020 Jan; 13(11): 5607-12. doi: 10.5958/0974-360X.2020.00978.
21. Arafa AF, Foda DS, Mahmoud AH, Metwally NS, Farrag AR. *Bombax ceiba* ower extract ameliorates hepatosteatosis induced by ethanol and relatively moderate fat diet in rats. *Toxicology Reports*. 2019 Jan; 6: 401-8. doi: 10.1016/j.toxrep.2019.04.008.
22. Tundis R, Rashed K, Said A, Menichini F, Loizzo MR. In vitro cancer cell growth inhibition and antioxidant activity of *Bombax ceiba* (Bombacaceae) ower extracts. *Natural Product Communications*. 2014 May; 9(5): 1934578X1400900527. doi: 10.1177/1934578 X1400900527.
23. Rehan M and Sha ullah S. The anticancer activity of oleanane-type saponin from *Bombax ceiba* (in vitro) and theoretical investigation of the signaling pathway. *Malaysian Journal of Chemistry*. 2021 Mar; 23(1): 33-47.
24. Jadhav V, Dhande S, Kadam V. Angiogenic effect of indigenous herbal extracts: *Bombax Ceiba* and *Erythrina variegata*. *Indian Journal of Natural Products and Resources*. 2018 Jun; 9(2): 126-31.
25. Hoque N, Rahman S, Jahan I, Afroze Shanta M, Sultana Tithi N, Nasrin N. A comparative phytochemical and biological study between different solvent extracts of *Bombax ceiba* roots available in Bangladesh. *Pharmacology & Pharmacy*. 2018 Feb; 9(02): 53-66. doi: 10.4236/pp.2018.92005. doi: 10.4236/pp.2018.92005.
26. Sharma N, Kispotta S, Mazumder PM. Immunomodulatory and anticancer activity of *Bombax ceiba* Linn. leaf extract. *Asian Pacific Journal of Tropical Biomedicine*. 2020 Sep; 10(9): 426. doi: 10.4103/2221 1691.290134.
27. Sharmin R, Joarder HH, Alamgir M, Mostofa G, Islam M. Antidiabetic and hepatoprotective activities of *Bombax ceiba* young roots in alloxan-induced diabetic mice. *Journal of Nutritional Health and Food Science*. 2018 Aug; 6: 1-7. doi: 10.15226/jnhfs.2018.001140.
28. Zulca i AS, Lim C, Ling AP, Chye S, Koh R. Focus: Plant-based Medicine and Pharmacology: Antidiabetic Potential of *Syzygium* sp.: An Overview. *The Yale journal of Biology and Medicine*. 2020 Jun; 93(2): 307.
29. Bhargava S and Shah MB. Evaluation of efficacy of *Bombax ceiba* extract and its major constituent, mangiferin in streptozotocin (STZ)-induced diabetic rats. *Journal of Complementary and Integrative Medicine*. 2020 Sep; 18(2): 311-8. doi: 10.1515/jcim 2020-0027.
30. Maurya SK, Verma NK, Verma DK. *Bombax ceiba* Linn. : A review of its phytochemistry and pharmacology. *Current Research Journal of Pharmaceutical and Allied Sciences*. 2018 Sep; 2(3): 14-23.
31. Zahan R, Nahar L, Haque M, Nesa ML, Alam Z. Antioxidant and antidiabetic activities of *Alangium salvifolium* and *Bombax ceiba*. *Dhaka University Journal of Pharmaceutical Sciences*. 2014 Jan; 12(2): 159-63. doi: 10.3329/dujps.v12i2.17617.
32. Shah MA, Reanmongkol W, Radenahmad N, Khalil R, Ul-Haq Z, Panichayupakaranant P. Anti hyperglycemic and anti-hyperlipidemic effects of rhinacanthins-rich extract from *Rhinacanthus nasutus* leaves in nicotinamide-streptozotocin induced diabetic rats. *Biomedicine & Pharmacotherapy*. 2019 May; 113: 108702. doi: 10.1016/j.biopha.2019.108702.
33. Aly O, Elias TR, Agaibyi MN, Rasheed WI, Yassen NN, Diab Y. Antidiabetic and

- Hepatoprotective Activities of Bombax Ceiba Extract in Obese Rats with Metabolic Syndrome. *Plant Archives*. 2021; 21(1): 748-56. doi: 10.51470/PLANTARCHIVES.2021.v21.S1.113.
34. Bhavsar C and Talele GS. Potential anti-diabetic activity of Bombax ceiba. *Bangladesh Journal of Pharmacology*. 2013 Feb; 8(2): 102-6. doi: 10.3329/bjp.v8i2.13701.
 35. Oraebosi MI, Good GM, Chia T, Oyeniran OI. Bombax Costatum extract abrogates piroxicam-mediated hepatic and gastric toxicities in rats. *Annales Pharmaceutiques Françaises*. 2020 Nov; 78(6): 507-14. doi: 10.1016/j.pharma.2020.06.002.
 36. Chauhan and Ekta Singh: Comparative studies on nutritional analysis and phytochemical screening of Bombax ceiba. *Bark and Seeds Powde* 2017; 5(2): 129-32.
 37. Depani and Parth: Ethnobotanical potential and phytochemical screening of Bombax ceiba L. *European Journal of Medicinal Plants* 2019; 1-8.
 38. Digge VG: Screening of Antibacterial Activity of Aqueous Bark Extract of Bombax Ceiba against Some Gram Positive and Gram Negative Bacteria. *Divya N: Academic Sciences* 2012;
 39. Faizi and Shaheen: Shamiminol: A new aromatic glycoside from the stem barks of Bombax ceiba. *Natural Product Communications* 1897; 6(12): 2011-900.
 40. Hossain and Emdad: Larvicidal activity of Dregea volubilis and Bombax malabaricum leaf extracts against the filarial vector Culex quinquefasciatus. *Asian Pacific Journal of Tropical Medicine Hainan Medical College* 2011; 4(6): 436-41.
 41. Jain, Vartika and Verma SK: Assessment of credibility of some folk medicinal claims on Bombax ceiba L. *Indian Journal of Traditional Knowledge* 2014; 13(1): <http://hdl.handle.net/123456789/26029>.
 42. Joshi and Raj K: Chemical analysis of flowers of Bombax ceiba from Nepal. *Natural Product Communications* 2013; 8(5): 583-84.
 43. Aguoru UC: Qualitative and quantitative phytochemical analysis of the leaf, stem bark and root of Bombax ceiba (Red silk cotton tree) in north central Nigeria.
 44. Yoges V and Kola N: Natural dyes extracted from bark of Bombax ceiba Linn. locally known as semal and its application on various fabrics pretreated with eco-friendly
 45. Karole and Sarita: Physicochemical, qualitative and quantitative phytochemical analysis of the leaf and bark of Bombax ceiba L. (Red silk cotton tree). *Journal of Drug Delivery and Therapeutics* 2018; 8(6): 105-10.
 46. Meena A: Review on ethnobotany and phytopharmacology of Bombax ceiba. *Research Journal of Pharmacognosy and Phytochemistry* 2011; 3(2): 57-61.
 47. Refaat and John: Bombacaceae: a phytochemical review. *International Journal of Sciences* 2015; 1(5): 37-41. 12.
 48. Shahat and Abdelaaty A: Isolation of mangiferin from Bombax malabaricum and structure revision of shamimin. *Planta Medica* 2003; 69(11): 1068-70. 13.
 49. Sint and Khin Maung: Wood anatomy and topochemistry of Bombax ceiba L. and Bombax insigne Wall. *Bio Resources* 2013; 8(1): 530-44.
 50. World Health Organization. WHO Traditional Medicine Strategy 2002-2005. Geneva: World Health Organization; 2002. p. 1-74. Available from: <https://doi.org/WHO/EDM/TRM/2002.1> [Last accessed on 13 Dec 2022].



51. Newman DJ, Cragg GM. Natural products as sources of new drugs from 1981 to 2014. *J Nat Prod.* 2016;79(3):629-61. doi: 10.1021/acs.jnatprod.5b01055. PMID 26852623.
52. Rameshwar V, Kishor D, Siddharth G, Sudarshan G. A pharmacognostic and pharmacological overview on *Bombax ceiba*. *Scholars Acad J Pharmonline Sch Acad J Pharm.* 2014;3(2):2320.
53. Pharmatutor.org, New York. Available www.pharmatutor.org/articles/aphrodisiac-activity-of-bombax-ceiba-linn-extract-in-male-mice.
54. Gupta P, Goyal R, Chauhan Y, Sharma PL. Possible modulation of FAS and PTP-1B signaling in ameliorative potential of *Bombax ceiba* against high fat diet-induced obesity. *BMC*

HOW TO CITE: Dr. Santosh Kumar Mishra, Vivek Singh, Tushar Gupta, Durgesh Yadav, A Review on the Hepatoprotective Activity of *Bombax ceiba* Root Extracts and Exudates, *Int. J. of Pharm. Sci.*, 2026, Vol 4, Issue 5, 3415-3425, <https://doi.org/10.5281/zenodo.20180015>

