

Section-4: Update on Mother Tinctures & Speciality products

This section revisits the classical role of mother tinctures and specialized homeopathic formulations in organ-specific therapeutics. While these remedies have long-standing roots in traditional practice, renewed interest in their pharmacological potential and clinical utility continues to reinforce their relevance in modern healthcare. Highlighted examples include well-established tinctures like *Aconitum napellus*, *Datura metel*, and *Erigeron canadense*, alongside proprietary products such as *Hepar sulphuris pentarkan*®, *Anekind*™, and *Alpha-RC*™, which exemplify the synergy between traditional principles and evidence-informed formulation strategies.

Praveen Saxena¹, Praveen Kumar Singh¹, Snigdha Suman Dalua¹

¹Dr. Willmar Schwabe India Pvt. Ltd.,

A 36, Sector 60, Noida, India. Pin – 201304.

DOI: doi.org/10.5281/zenodo.17349798

Aconitum napellus

Aconitum napellus, commonly known as monkshood or aconite, is a tall, perennial herb native to mountainous regions of the Himalayas and Central/Southern Europe, Siberia, and North America. It features tapering roots, smooth upright stems up to 2m, and palmately lobed leaves, with deep violet helmet-shaped flowers that bloom from May to July. This plant is recognized in the Homoeopathic Pharmacopoeia of India and was first proved by Dr. Hahnemann.

Its mother tincture is used in homeopathy for acute collapse, nervous restlessness, cardiac irregularities, rheumatic fevers, localized pain, burning and tingling of lips and throat, extremity paresthesias, hypersalivation, gastric disturbances, arrhythmias, myocardial shock, central nervous system depression, muscular weakness, and even seizures. A recent Indian study observed that combining *Aconitum napellus* with *Lachesis* at 10⁻² potency enhanced phagocytosis by approximately 390%, suggesting notable immunomodulatory effects.

Despite its therapeutic use in dilute form, *Aconitum napellus* is highly toxic in crude preparations. It contains potent diester diterpene alkaloids (e.g., aconitine, mesaconitine) that pose serious cardiotoxic and neurotoxic threats. Even minimal amounts, around 1 g of plant material or just a couple of milligrams of aconitine, can be fatal by causing respiratory paralysis, ventricular arrhythmias, or hypotension. Symptoms of poisoning typically begin within an hour, manifesting as nausea, burning tingling in the mouth, extremity numbness, dizziness, sweating, and progressing to severe cardiac complications. Percutaneous absorption can also lead to



life-threatening toxicity.

Given this narrow therapeutic index, homeopathic formulations undergo rigorous dilution to minimize risks. Therapeutic use requires strict medical supervision, with attention to patient condition and contraindications. In cases of overdose or exposure, treatment involves immediate cardiovascular and respiratory monitoring, administration of activated charcoal if within 1 hour, and use of atropine or antiarrhythmics as needed.

Despite the inherent risks, *Aconitum napellus* MT remains a valuable homeopathic remedy for acute conditions like shock and cardiac disturbances, with growing interest in its potential immune-enhancing effects highlighting the delicate balance between harnessing therapeutic benefit and ensuring safe use.

Recommended dose: 1 drop of Ø/1x. Repeat under medical supervision. Externally, the Mother tincture is used.

Caution: Poisonous. Avoid using Ø/1x.

Datura metel

Datura metel L. is a glabrous annual herb found globally, including regions across India like Kashmir and Sikkim. It features large, toothed leaves (17–20 cm) and striking solitary flowers that can be white, yellowish, or violet. The seeds, containing up to 0.5% tropane alkaloids hyoscyne, hyoscyamine, atropine, scopolamine, and nicotine, lend it potent narcotic, antispasmodic, and anodyne properties.

Officially recognized in the Homoeopathic Pharmacopoeia of the United States, its seed-based mother tincture was first proven by Dr. Wright. In homeopathic practice, *Datura metel* is indicated for neurologically intense conditions: convulsions, delirium, epilepsy, mania, timidity, and eye affections such as flickering before the eyes and light sensitivity. Clinical symptoms can also include extreme alternations in pulse and temperature, which mirror the effects of *Belladonna* and *Stramonium*. It induces soporose states culminating in delirium, further characterized by restless, talkative behavior, visual distortions, and dilated pupils.

From a phytochemical perspective, the plant is rich in tropane alkaloids and also contains sterols, withanolides, and flavonoids. Modern pharmacological studies have confirmed anticancer and immunomodulatory effects: methanolic extracts display cytotoxicity against human lung (A549), breast (MCF-7), and colorectal cell lines, inducing apoptosis and cell-cycle arrests. Furthermore, active fractions from *Datura stramonium* potentiate immune cell activity by enhancing IL-2 and IFN- secretion and improving PBMC-mediated cancer cell killing.

Despite its toxicity manifesting as anticholinergic delirium, tachycardia, hyperthermia, and mydriasis, no serious adverse effects have been observed at homeopathic potencies. This dual profile underscores *Datura metel*'s value in both conventional phytomedicine and homeopathic therapy. However, due to its high potency and potential risks, its use demands strict professional oversight.

Recommended dose: Ø up to 10 drops and diluted in water. drop doses and diluted in water up to the equivalent of 3rd attenuation and above before taking.



Caution: (1) Repeated doses of 1x induce gastric irritation and subsequent diarrhoea.

(2) A narcotic generally not to be prescribed below 3x.

Erigeron canadense

Erigeron canadensis (syn. *Conyza canadensis*), commonly known as horseweed, Canadian fleabane, coltstail, or butterweed is a fast-growing annual weed reaching up to 1.5 m tall. Its slender, alternate leaves (2–10 cm long, ~1 cm wide) bear coarse teeth, fading early on the lower stem. Tiny flower heads (~1 cm across) bloom in dense clusters, featuring white or pale purple rays around yellow centers and producing seeds equipped with downy tufts.

Native to North America, it now thrives across most temperate zones, including India's Western Himalayas, Punjab, Kashmir, Assam, Western Ghats, and Nilgiri Hills up to 2100 m. Phytochemical analyses have identified a rich profile of bioactive compounds like saponins, diterpenoids, flavonoids, tannins, glycosides, anthraquinones, steroids, volatile oils (notably limonene, terpineol, and linalool), and diverse sesquiterpenes.

Erigeron mother tincture (from the whole flowering plant) is recognized by both the Homoeopathic Pharmacopeia of India and the U.S. counterpart and was initially proved by Burt. It is primarily prescribed for bright-red haemorrhages such as urethral bleeding, haematuria, menorrhagia, haemorrhoids, epistaxis and genitourinary discharges like dysuria and chronic gonorrhoea. Ulceration-like symptoms, inflammatory bladder conditions, respiratory haemorrhages, and haemorrhages from wounds also respond well to *Erigeron*.

In traditional herbal medicine, *Erigeron* is used to treat diarrhea, dysentery, urinary tract infections, respiratory illnesses like bronchitis, rheumatism, skin wounds, and gastrointestinal issues. It even received recognition as a styptic and astringent tonic in homeopathic repertories.

Modern pharmacological studies reinforce its traditional use: extracts exhibit antimicrobial, antioxidant, anti-inflammatory, anticoagulant, anticancer, gastric-protective, and skin-depigmenting activities. One study showed significant gastric ulcer suppression in mice, while platelet aggregation and coagulation modulation were also observed.

Moreover, *Erigeron canadensis* offers a compelling synergy of traditional homeopathic applications and scientifically validated therapeutic actions particularly in bleeding disorders and mucosal inflammations making it a versatile medicinal resource.



Recommended dose: Ø 10-30 drops and diluted in water

Bibliography

1. Erhard, M., Kellner, J., Wild, J., Lösch, U. and Hatiboglu, F. S. (1994), Effect of Echinacea, Aconitum, Lachesis and Apis extracts, and their combinations on phagocytosis of human granulocytes. *Phytotherapy Research*, 8: 14–17.
2. Persson Hans., 2010., *Poisonous plants and fungi.*, Oxford Textbook of Medicine., United States by Oxford University Press., pg. 1363.
3. Amor Bellila, Cindy Tremblay, André Pichette, Brahim Marzouk, Vakhtang Mshvildadze, Serge Lavoie, Jean Legault, Cytotoxic activity of withanolides isolated from Tunisian *Datura metel* L., *Phytochemistry Volume 72, Issue 16*, November 2011, Pages 2031–2036.
4. The Encyclopedia of medicinal plant.
5. P. N. Varma, Indu Vaid, *Encyclopaedia of Homoeopathic Pharmacopoeia*, Updated edition 2007, B. Jain Publishers, New Delhi.
6. Launert. E. *Edible and Medicinal Plants*. Covers plants in Europe. a drawing of each plant, quite a bit of interesting information.
7. http://www.herbs2000.com/herbs/herbs_horseweed.htm
8. A. L. Blackwood, *A Manual of Materia Medica, Therapeutics and Pharmacology*, Reprint Edition 1995, B. Jain Publishers, New Delhi.