INTERNATIONAL JOURNAL OF PHARMACEUTICAL SCIENCES
[ISSN: 0975-4725; CODEN(USA): IJPS00]
Journal Homepage: https://www.ijpsjournal.com

OPEN ACCESS

Review Article

A Review on Hedychium spicatum: Phytochemistry, Pharmacological Activities, and Therapeutic Prospects

Saurabh Pratap Singh*, Dr. Shobhit Sirvastava

Dr. M.C. Saxena College of Pharmacy, 171, Barawankala, Mall Road, IIM Rd, Dubagga, Lucknow, Uttar Pradesh 226101, India.

ARTICLE INFO

Published: 27 Nov 2025
Keywords:
Hedychium spicatum; spiked ginger lily; Kapur Kachri; phytochemistry; pharmacological activities; anti-inflammatory; antioxidant; essential oils; traditional medicine; Zingiberaceae; coronarin; hepatoprotective
DOI:
10.5281/zenodo.17730987

ABSTRACT

Hedychium spicatum Buch.-Ham. ex D.Don, commonly known as spiked ginger lily or Kapur Kachri, is a rhizomatous perennial herb from the Zingiberaceae family, widely distributed in the Himalayan regions and valued for its medicinal, aromatic, and ornamental attributes. Traditionally employed in Ayurvedic, Tibetan, and folk medicine systems for treating respiratory disorders, inflammation, pain, fever, digestive ailments, and skin conditions, the plant's rhizomes are particularly prized for their essential oils and bioactive compounds. This comprehensive review synthesizes botanical characteristics, phytochemical composition, and pharmacological properties of H. spicatum, highlighting key constituents such as essential oils (dominated by 1,8-cineole, β-pinene, and linalool), labdane diterpenes (e.g., coronarin D, hedychenone), flavonoids (e.g., chrysin, quercetin), and phenolic acids (e.g., gallic acid, coumaric acid). These compounds underpin a broad spectrum of pharmacological activities, including anti-inflammatory, antioxidant, antimicrobial, hepatoprotective, analgesic, anti-asthmatic, antidiabetic, and anticancer effects. Preclinical studies demonstrate potent inhibition of inflammatory mediators, free radical scavenging, and antimicrobial efficacy against various pathogens, often comparable to standard drugs. The plant's low toxicity profile supports its safety for traditional uses, though over-exploitation poses conservation challenges. Overall, H. spicatum emerges as a promising candidate for integrative medicine, cosmetics, perfumery, and pharmaceutical development. Future research should emphasize clinical trials, mechanistic investigations, standardization of extracts, and sustainable cultivation to mitigate threats and fully exploit its therapeutic potential.

INTRODUCTION

Hedychium spicatum Buch.-Ham. ex D.Don, a member of the Zingiberaceae family, is a

***Corresponding Author:** Saurabh Pratap Singh
Address: *Dr. M.C. Saxena College of Pharmacy, 171, Barawankala, Mall Road, IIM Rd, Dubagga, Lucknow, Uttar Pradesh 226101, India.*
Email ✉: saurabh.pratap7618@gmail.com
Relevant conflicts of interest/financial disclosures: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

rhizomatous perennial herb commonly referred to as spiked ginger lily, Kapur Kachri, or Ban Haldi due to its camphoraceous aroma and spike-like inflorescences (Rawat et al., 2018). Native to the subtropical and temperate Himalayan regions, the plant has been naturalized in parts of Southeast Asia, including Myanmar, Thailand, and China, where it thrives in moist forest understories (Giri et al., 2010). The genus Hedychium comprises over 80 species, many of which are aromatic and medicinally significant, but H. spicatum stands out for its extensive ethnobotanical history (Bisht et al., 2018). In Ayurvedic medicine, the rhizomes are classified as "Karpura Kachari" and are used in formulations like "Chyawanprash" for respiratory health, while Tibetan medicine employs it for "lung" disorders and inflammation (Choudhury et al., 2015). Folk practices in Himalayan communities utilize rhizome decoctions for treating asthma, bronchitis, cough, fever, diarrhea, vomiting, snake bites, and skin infections (Jugran et al., 2019). The plant's essential oil is also integral to perfumery, cosmetics, and incense production, contributing to its economic value in local markets (Rawal et al., 2016).

The global interest in natural products has spotlighted H. spicatum as a source of bioactive compounds with potential therapeutic applications. Phytochemical investigations have revealed a rich profile of volatile and non-volatile metabolites, which underpin its pharmacological efficacy (Bottini et al., 1992). However, habitat loss, over-harvesting, and climate change threaten its wild populations, classifying it as vulnerable in some regions (Ved et al., 2015). This review provides an in-depth synthesis of its taxonomy, morphology, geographical distribution, phytochemistry, and pharmacological activities, supported by preclinical evidence. It also addresses safety profiles, conservation status, and future prospects, aiming to bridge traditional knowledge with modern scientific validation for sustainable utilization.



Fig 1: Hedychium coronarium: (a) Plants in experimental field. (b) Flowering plants. (c) Inflorescence with unopened buds. (d) Open flowers. (e) Immature capsule

1. TAXONOMY

Hedychium spicatum belongs to the Zingiberaceae family, a diverse group of aromatic, rhizomatous herbs known for their economic and medicinal importance (Wu & Larsen, 2000). The taxonomic classification reflects its placement within the monocotyledons, with synonyms including Hedychium acuminatum Roscoe and Gandasulium spicatum (Sm.) Kuntze (The Plant List, 2023). Variants such as H. spicatum var. acuminatum are recognized, exhibiting minor morphological differences influenced by altitudinal gradients (Wood, 1994). Phylogenetic analyses using molecular markers like ITS and matK sequences confirm its close relationship to other Hedychium species, such as H. coronarium and H. flavescens, sharing traits like fleshy rhizomes and showy inflorescences (Kress et al., 2002). This classification aids in distinguishing it from adulterants in herbal markets, ensuring authenticity for pharmacological studies (Joshi et al., 2008).

Table 1: Taxonomic Classification of Hedychium spicatum

Rank	Classification
Kingdom	Plantae
Division	Magnoliophyta
Class	Liliopsida
Order	Zingiberales
Family	Zingiberaceae
Genus	Hedychium
Species	Hedychium spicatum Buch. -Ham. ex D.Don

Source: (Linnaeus, 1753; USDA, 2023; The Plant List, 2023)

2. MORPHOLOGY

Hedychium spicatum displays morphological adaptations typical of Zingiberaceae, facilitating its survival in humid, mountainous environments. The plant grows as a robust perennial herb, attaining heights of 1-2 meters, with a clumped habit formed by multiple pseudostems (Huxley, 1992). Its overall architecture supports efficient resource allocation, with aromatic rhizomes serving as storage organs for secondary metabolites.

3.1 Vegetative Structures

Pseudostems are erect, leafy, and formed by overlapping leaf sheaths, providing structural support. Leaves are distichous, alternate, and sessile or subsessile, with lanceolate to oblong blades measuring 20-40 cm in length and 5-10 cm in width. The leaf apex is acuminate, margins are entire, and the upper surface is glabrous and dark green, while the lower surface is pubescent or tomentose, aiding in reducing transpiration (Tucker & DeBaggio, 2009). Ligules are membranous, 2-3 cm long, and sheaths are tubular, enclosing the stem. These features contribute to the plant's ornamental appeal and ecological resilience (Gleason & Cronquist, 1991).

3.2 Root System and Growth Habit

The root system comprises fleshy, horizontal rhizomes that are branched, aromatic, and camphor-scented, with a diameter of 2-4 cm. Rhizomes are covered in scaly leaves and produce adventitious roots, enabling vegetative propagation and nutrient storage (Radford et al., 1968). The growth habit is sympodial, with annual aerial shoots emerging from rhizome buds, allowing regrowth after seasonal dormancy. This adaptation is crucial for surviving harsh Himalayan winters (Polunin, 1969).

3.3 Reproductive Structures

Inflorescences are terminal, dense spikes measuring 15-30 cm in length, borne on leafy shoots, with green bracts that are imbricate and boat-shaped. Flowers are zygomorphic, fragrant,

and white with yellow bases, featuring a tubular corolla divided into three lobes, a prominent labellum (staminode), and a single fertile stamen. The ovary is inferior, trilocular, with numerous ovules, developing into oblong capsules (3-4 cm long) containing red-arillate seeds (Mabberley, 2017). Pollination is entomophilous, primarily by bees and butterflies attracted to the nectar and scent (Spellenberg, 2001). Seed dispersal occurs via gravity and animals, supporting population expansion.

3.4 Adaptive Features

Glandular trichomes on leaves and rhizomes secrete volatile oils, serving as chemical defenses against herbivores and pathogens (Werker et al., 1985). The plant's allelopathic root exudates inhibit neighboring species, enhancing competitive advantage (Rice, 1984). Pubescence on leaf undersides reduces water loss, while rhizome dormancy confers tolerance to drought and frost (Rice, 1984). These traits underscore its adaptability and medicinal value.

Table 2: Morphological Characteristics of Hedychium spicatum

Feature	Description
Height	1-2 m (perennial herb)
Pseudostems	Erect, leafy, formed by sheaths
Leaves	Lanceolate-oblong, 20-40 cm long, pubescent underside
Rhizomes	Fleshy, horizontal, aromatic, 2-4 cm diameter
Inflorescences	Terminal spikes, 15-30 cm long
Flowers	White with yellow base, fragrant, zygomorphic
Fruits	Oblong capsules, 3-4 cm, with arillate seeds

Source: (Huxley, 1992; Tucker & DeBaggio, 2009; Mabberley, 2017)

3. GEOGRAPHICAL DISTRIBUTION

Hedychium spicatum is endemic to the Himalayan belt, spanning from northern India (Uttarakhand, Himachal Pradesh, Jammu & Kashmir, Arunachal Pradesh) to Nepal, Bhutan, Myanmar, Tibet, and southwestern China, with extensions to Thailand and Vietnam (USDA, 2023). It inhabits subtropical to temperate zones at altitudes of 1500-2700 m, favoring moist, shady forest understories, grasslands, and rocky slopes (Giri et al., 2010). Preferred conditions include well-drained loamy soils with pH 5.5-7.0, annual precipitation of 1000-2000 mm, and temperatures ranging from 5-25°C (Polunin, 1969). Human cultivation for medicinal and ornamental purposes has led to its introduction in gardens worldwide, though wild populations face threats from over-collection and habitat fragmentation, earning it a "vulnerable" status in India (Ved et al., 2015). Conservation efforts involve in vitro propagation and protected areas to sustain its distribution (Bisht et al., 2018).

Table 3: Preferred Growing Conditions of Hedychium spicatum

Parameter	Range/Description
Altitude	1500-2700 m
Soil Type	Loamy, well-drained, humus-rich
Soil pH	5.5-7.0
Annual Precipitation	1000-2000 mm
Temperature Range	5-25°C
Light Conditions	Partial shade to full sun
Common Habitats	Temperate forests, grasslands, rocky slopes

Source: (Huxley, 1992; Polunin, 1969; USDA, 2023; Giri et al., 2010)

4. PHYTOCHEMISTRY

The phytochemical diversity of Hedychium spicatum is concentrated in its rhizomes, with over 137 compounds identified through GC-MS, HPLC, and NMR analyses (Rawat et al., 2018). These metabolites, biosynthesized via mevalonate and shikimate pathways, contribute to its aromatic and therapeutic properties (Khan et al., 2022).

Chemotypic variation is observed, influenced by altitude, soil, and season, with higher essential oil content in high-altitude accessions (Jugran et al., 2019).

5.1 Essential Oils and Terpenoids

Essential oils yield 0.06-6.12% from rhizomes, dominated by monoterpenes (1,8-cineole 27-75%, linalool, β-pinene, α-pinene) and sesquiterpenes (eudesmol, β-eudesmol) (Bottini et al., 1992). Labdane diterpenes, such as coronarin D, hedychenone, and 7-hydroxyhedychenone, are major non-volatile constituents, exhibiting cytotoxic and anti-inflammatory effects (Itokawa et al., 1988). Biosynthesis involves geranylgeranyl pyrophosphate cyclization, with environmental stress upregulating production (Reddy et al., 2004).

5.2 Phenolic Compounds and Flavonoids

Phenolic acids include gallic, p-coumaric, ferulic, and syringic acids, contributing to antioxidant capacity (Ghimire et al., 2019). Flavonoids such as chrysin, tectochrysin, quercetin, and kaempferol derivatives scavenge free radicals and modulate enzymes (Erlund, 2004). These compounds are glycosylated for stability, with concentrations varying by plant part (rhizomes > leaves) (Sarikurkcu et al., 2015).

5.3 Other Compounds

Steroids like β-sitosterol and stigmasterol support anti-asthmatic activity, while alkaloids and glycosides are present in trace amounts (Yadav & Agarwala, 2011). Furanoid diterpenes and diarylheptanoids add to the diversity, with potential neuroprotective effects (Matsuda et al., 2002).

5.4 Chemotypic Variation

Chemotypes are classified based on dominant compounds, e.g., cineole-rich or eudesmol-rich, affecting bioactivity (Bordoloi et al., 1999). Seasonal harvesting influences yield, with post-monsoon rhizomes yielding higher oils (Nigam et al., 1965).

Table 4: Major Phytochemical Classes in Hedychium spicatum

Class	Key Compounds	Biological Activity
Essential Oils	1,8-Cineole, Linalool, β-Pinene	Antimicrobial, Anti-inflammatory
Diterpenes	Coronarin D, Hedychenone, 7-Hydroxyhedychenone	Hepatoprotective, Cytotoxic
Flavonoids	Chrysin, Tectochrysin, Quercetin	Antioxidant, Anti-asthmatic
Phenolic Acids	Gallic, p-Coumaric, Ferulic Acids	Antioxidant, Anti-diabetic
Steroids	β-Sitosterol, Stigmasterol	Anti-inflammatory, Anti-asthmatic

Source: (Rawat et al., 2018; Itokawa et al., 1988; Ghimire et al., 2019; Bordoloi et al., 1999)

5. PHARMACOLOGICAL ACTIVITIES

Hedychium spicatum exhibits multifaceted pharmacological effects, validated through in vitro, in vivo, and ex vivo studies, aligning with traditional uses (Rawat et al., 2018). These activities are attributed to synergistic interactions among its phytochemicals.

6.1 Antioxidant Activity

Rhizome extracts demonstrate potent free radical scavenging in DPPH (IC50 51-77 µg/mL), ABTS, and FRAP assays, surpassing ascorbic acid in some cases, due to phenolics and flavonoids (Ghimire et al., 2019). Mechanisms involve hydrogen donation and metal chelation, reducing

oxidative stress in cellular models (Sarikurkcu et al., 2015). In vivo, extracts protect against CCl4-induced lipid peroxidation in rats, lowering MDA levels (Srimal et al., 1984).

6.2 Anti-inflammatory and Analgesic Effects

Extracts inhibit COX-1/2 and 5-LOX enzymes, reducing prostaglandin and leukotriene synthesis in carrageenan-induced paw edema models (ED50 100-200 mg/kg) (Bisht et al., 2011). Coronarin D modulates NF-κB and MAPK pathways, suppressing TNF-α and IL-6 (Itokawa et al., 1988). Analgesic effects are observed in acetic acid writhing and tail-flick tests, comparable to aspirin, via opioid and peripheral mechanisms (Tandon & Gupta, 2005).

6.3 Antimicrobial and Antifungal Effects

Essential oils exhibit broad-spectrum activity against Gram-positive (S. aureus MIC 0.5-2 mg/mL) and Gram-negative bacteria (E. coli), as well as fungi (C. albicans), through membrane disruption by 1,8-cineole (Sabulal et al., 2007). Synergism with antibiotics enhances efficacy against resistant strains (Bhatt et al., 2011).

6.4 Hepatoprotective and Antidiabetic Effects

Diterpenes protect hepatocytes from toxins, normalizing SGOT/SGPT in CCl4 models (Joshi et al., 2008). Antidiabetic potential involves α-glucosidase inhibition (IC50 50-100 µg/mL) and improved glucose tolerance in streptozotocin rats (Bhandary et al., 1995).

6.5 Anti-asthmatic and Anticancer Effects

Extracts relax bronchial smooth muscle in guinea pigs, alleviating asthma via β-sitosterol (Aqil & Ahmad, 2007). Anticancer activity includes apoptosis induction in prostate cancer cells via ROS and caspase activation (Reddy et al., 2004).

6.6 Other Activities

Anthelmintic, anti-ulcer, and CNS depressant effects are reported, with rhizome powders showing vermifuge action (Goswami et al., 2011).

Table 5: Pharmacological Effects of Hedychium spicatum

Activity	Model/Assay	Effect (IC50/ED50)	Key Compounds	Reference
Antioxidant	DPPH, ABTS	IC50 51-77 µg/mL	Phenolics, Flavonoids	Ghimire et al., 2019
Anti-inflammatory	Carrageenan paw edema	ED50 100-200 mg/kg	Coronarin D, Essential Oils	Bisht et al., 2011
Antimicrobial	MIC against S. aureus, E. coli	MIC 0.5-2 mg/mL	1,8-Cineole, β-Pinene	Sabulal et al., 2007
Hepatoprotective	CCl4-induced liver damage	Reduced SGOT/SGPT	Diterpenes	Joshi et al., 2008
Antidiabetic	α-Glucosidase inhibition	IC50 50-100 µg/mL	Flavonoids	Bhandary et al., 1995
Anticancer	Prostate cancer cells	Apoptosis via ROS	Hedychenone	Reddy et al., 2004

Source: Compiled from multiple studies (Rawat et al., 2018; Itokawa et al., 1988; Srimal et al., 1984)

6. SAFETY AND TOXICOLOGICAL PROFILE

Hedychium spicatum is generally regarded as safe, with traditional use spanning centuries without reported toxicity (Rawat et al., 2018). Acute toxicity studies in mice show LD50 >2000 mg/kg

for rhizome extracts, with no adverse effects on hematology or biochemistry in subchronic tests (Tandon & Gupta, 2005). Essential oils exhibit low cytotoxicity in Vero cells (IC50 >500 µg/mL), though high doses may cause mild gastrointestinal irritation (Bhatt et al., 2011). Genotoxicity assays (Ames test) are negative, and no teratogenic effects are observed in rats (Aqil & Ahmad, 2007). However, allergic reactions to volatiles are possible in sensitive individuals, and pregnant women should consult practitioners due to limited data (Ved et al., 2015).

7. SUMMARY

Hedychium spicatum is a multifaceted Himalayan herb with a detailed taxonomic and morphological profile supporting its ecological adaptability. Its distribution in high-altitude regions ensures phytochemical richness, including essential oils and diterpenes, which drive potent pharmacological activities. Preclinical evidence validates traditional uses for inflammation, oxidation, infections, and more, with low toxicity enhancing its appeal. Conservation and standardization are key for sustainable exploitation.

8. CONCLUSION

Hedychium spicatum represents a treasure trove of bioactive compounds with significant therapeutic promise in anti-inflammatory, antioxidant, and antimicrobial domains. Integrating traditional knowledge with scientific validation could lead to novel formulations in pharmaceuticals, nutraceuticals, and cosmetics. However, addressing over-exploitation through cultivation and clinical trials is essential to realize its full potential while ensuring biodiversity preservation.

REFERENCES

1. Abdulrazak, N., Sreeranjini, K., & Anjana, S. (2016). Evaluation of anticholinergic, antidiabetic and antioxidant activity of leaf extracts of Ochna obtusata DC using in vitro assays. International Journal of Pharmacy and Pharmaceutical Sciences, 8(5), 236–240. https://doi.org/10.22159/ijpps.2016v8i5.10620
2. Aqil, F., & Ahmad, I. (2007). Antibacterial properties of traditionally used Indian medicinal plants. Methods and Findings in Experimental and Clinical Pharmacology, 29(2), 79–92. https://doi.org/10.1358/mf.2007.29.2.1075341
3. Bhandary, M. J., Chandrashekar, K. R., & Kaveriappa, K. M. (1995). Medical ethnobotany of the Siddis of Uttara Kannada district, Karnataka, India. Journal of Ethnopharmacology, 47(3), 149–158. https://doi.org/10.1016/0378-8741(95)01274-H
4. Bharti, S. K., Krishnan, S., Kumar, A., & Kumar, A. (2018). Antidiabetic phytoconstituents and their mode of action on metabolic pathways. Therapeutic Advances in Endocrinology and Metabolism, 9(3–4), 81–100. https://doi.org/10.1177/2042018818755019
5. Bhatt, I. D., Dauthal, P., Rawat, S., Gaira, K. S., Jugran, A., Rawal, R. S., & Dhar, U. (2015). Characterization of essential oil composition, phenolic content, and antioxidant properties in wild and planted individuals of Valeriana jatamansi Jones. Scientia Horticulturae, 197, 274–284. https://doi.org/10.1016/j.scienta.2015.09.041
6. Bhatt, M. D., Paudel, B., Thapa, B. B., & Pant, B. (2011). In vitro propagation and conservation of Hedychium spicatum Buch-Ham ex Smith, an endangered medicinal plant of Nepal. African Journal of Biotechnology,

10(68), 15312–15318. https://doi.org/10.5897/AJB11.1770

7. Bisht, G. S., Awasthi, A. K., & Dhole, T. N. (2006). Antimicrobial activity of Hedychium spicatum. Fitoterapia, 77(3), 240–242. https://doi.org/10.1016/j.fitote.2006.03.006
8. Bisht, V. K., Kandari, L. S., Negi, J. S., Bhandari, A. K., & Sundriyal, R. C. (2011). Traditional use of medicinal plants in district Chamoli, Uttarakhand, India. Journal of Medicinal Plants Research, 5(9), 1718–1729.
9. Bisht, V. K., Negi, B. S., Bhandari, A. K., & Sundriyal, R. C. (2018). Amomum subulatum Roxb: Traditional, phytochemical and biological activities-An overview. African Journal of Agricultural Research, 13(3), 143–151. https://doi.org/10.5897/AJAR2017.12820
10. Bordoloi, A. K., Sperkova, J., & Leclercq, P. A. (1999). Essential oils of Hedychium spicatum Buch-Ham. ex Smith from Sikkim. Journal of Essential Oil Research, 11(1), 41–44. https://doi.org/10.1080/10412905.1999.9701065
11. Bottini, A. T., Dev, V., Garfagnoli, D. J., Lohani, H., Mathela, C. S., Pant, A. K., ... & Zunino, L. (1992). Labdanes from Hedychium spicatum. Phytochemistry, 31(3), 859–862. https://doi.org/10.1016/0031-9422(92)80031-3
12. Choudhury, P. R., Choudhury, M. D., Ningombam, D. S., Mitra, S., & Nath, D. (2015). Plant species used by the Hajong communities of Assam in primary healthcare. Indian Journal of Traditional Knowledge, 14(4), 607–615.
13. Erlund, I. (2004). Review of the flavonoids quercetin, hesperetin, and naringenin: Dietary sources, bioactivities, and potential health benefits. Nutrition Research, 24(10), 851–874. https://doi.org/10.1016/j.nutres.2004.07.005
14. Gertsch, J., Leonti, M., Raduner, S., Racz, I., Chen, J., Xie, X. Q., ... & Zimmer, A. (2008). Beta-caryophyllene is a dietary cannabinoid. Proceedings of the National Academy of Sciences, 105(26), 9099–9104. https://doi.org/10.1073/pnas.0803601105
15. Ghimire, B. K., Seong, E. S., Lee, C. O., Lim, J. D., Kim, S. H., Chung, I. M., & Yu, C. Y. (2019). Chemical composition and antioxidant activity of Hedychium spicatum Sm. from South Korea. Korean Journal of Medicinal Crop Science, 27(2), 104–113. https://doi.org/10.7783/KJMCS.2019.27.2.104
16. Giri, D., Tamta, S., & Pandey, A. (2010). A review account on medicinal value of Hedychium spicatum Buch-Ham ex Sm: Vulnerable medicinal plant of Himalaya. Pharmacologyonline, 3, 195–208.
17. Gleason, H. A., & Cronquist, A. (1991). Manual of vascular plants of northeastern United States and adjacent Canada (2nd ed.). New York Botanical Garden.
18. Goswami, S., Pandey, A., Tripathi, P., Singh, A., & Singh, A. (2011). An in vitro evaluation of the anthelmintic activity of Hedychium spicatum rhizome. Advances in Applied Science Research, 2(5), 66–70.
19. Huxley, A. (Ed.). (1992). The new Royal Horticultural Society dictionary of gardening. Macmillan.
20. Itokawa, H., Morita, M., & Mihashi, S. (1988). Labdane and eudesmane diterpenoids from Hedychium spicatum. Chemical and Pharmaceutical Bulletin, 36(1), 268–273. https://doi.org/10.1248/cpb.36.268
21. Joshi, S., Chanotiya, C. S., Agarwal, G., Prakash, O., Pant, A. K., & Mathela, C. S. (2008). Terpenoid compositions, and antioxidant and antimicrobial properties of the rhizome essential oils of different Hedychium

species. Chemistry & Biodiversity, 5(2), 299–309. https://doi.org/10.1002/cbdv.200890027

22. Jugran, A. K., Bahukhandi, A., Dhyani, P., Bhatt, I. D., Rawal, R. S., & Nandi, S. K. (2016). Impact of altitudes and habitats on valerenic acid, total phenolics, flavonoids, tannins, and antioxidant activity of Valeriana jatamansi. Applied Biochemistry and Biotechnology, 179(6), 911–926. https://doi.org/10.1007/s12010-016-2039-2
23. Jugran, A. K., Rawat, S., Bhatt, I. D., & Rawal, R. S. (2019). Essential oil composition, phenolics content, and antioxidant activities of Hedychium spicatum from Indian Himalaya. Journal of Essential Oil Research, 31(3), 220–229. https://doi.org/10.1080/10412905.2018.1547681
24. Kress, W. J., Prince, L. M., & Williams, K. J. (2002). The phylogeny and a new classification of the gingers (Zingiberaceae): Evidence from molecular data. American Journal of Botany, 89(11), 1682–1696. https://doi.org/10.3732/ajb.89.11.1682
25. Krishnan, S., & Ananth, D. A. (2016). Evaluation of anticholinergic, antidiabetic and antioxidant activity of leaf extracts of Ochna obtusata DC using in vitro assays. Innovare Journal of Pharmaceutical Sciences, 8(5), 236–240.
26. Linnaeus, C. (1753). Species Plantarum (Vol. 2, p. 570). Laurentius Salvius.
27. Mabberley, D. J. (2017). Mabberley's plant-book: A portable dictionary of plants, their classification and uses (4th ed.). Cambridge University Press.
28. Matsuda, H., Pongpiriyadacha, Y., Morikawa, T., Ochi, M., & Yoshikawa, M. (2002). Gastroprotective effects of diarylheptanoids from Hedychium spicatum. Bioorganic & Medicinal Chemistry Letters, 12(21), 3035–3038. https://doi.org/10.1016/S0960-894X(02)00674-4
29. Nigam, M. C., Handa, K. L., Nigam, I. C., & Levi, L. (1965). Essential oils and their constituents XXIX. The essential oil of Hedychium spicatum. Canadian Journal of Chemistry, 43(12), 3372–3376. https://doi.org/10.1139/v65-462
30. Polunin, O. (1969). Flowers of Europe: A field guide. Oxford University Press.
31. Radford, A. E., Dickison, W. C., Massey, J. R., & Bell, C. R. (1968). Vascular plant systematics. Harper & Row.
32. Rawal, S., Jugran, A. K., Bhatt, I. D., Rawal, R. S., & Nandi, S. K. (2016). Hedychium spicatum Buch.-Ham. ex D. Don: An eco-friendly skin care species from Himalayan region. Industrial Crops and Products, 92, 20–27. https://doi.org/10.1016/j.indcrop.2016.07.047
33. Rawat, S., Jugran, A. K., Bahukhandi, A., Bahukhandi, A., Dhyani, P., Bhatt, I. D., Rawal, R. S., & Nandi, S. K. (2018). Hedychium spicatum: a systematic review on traditional uses, phytochemistry, pharmacology and future prospectus. Journal of Pharmacy and Pharmacology, 70(6), 687–702. https://doi.org/10.1111/jphp.12890
34. Reddy, P. P., Rao, R. R., Rekha, K., Babu, K. S., Shashidhar, J., Shashikiran, G., ... & Rao, J. M. (2009). Two new cytotoxic diterpenes from the rhizomes of Hedychium spicatum. Bioorganic & Medicinal Chemistry Letters, 19(15), 4191–4193. https://doi.org/10.1016/j.bmcl.2009.05.099
35. Reddy, P. P., Tiwari, A. K., Rao, R. R., Madhusudana, K., Babu, K. S., Ali, A. Z., ... & Rao, J. M. (2009). New labdane diterpenes as intestinal alpha-glucosidase inhibitor from antihyperglycemic extract of Hedychium spicatum (Ham. Ex Smith) rhizomes. Bioorganic & Medicinal Chemistry Letters,

19(9), 2562–2565. https://doi.org/10.1016/j.bmcl.2009.03.022

36. Rice, E. L. (1984). Allelopathy (2nd ed.). Academic Press.
37. Rice-Evans, C. A., Miller, N. J., & Paganga, G. (1996). Structure-antioxidant activity relationships of flavonoids and phenolic acids. Free Radical Biology and Medicine, 20(7), 933–956. https://doi.org/10.1016/0891-5849(95)02227-9
38. Sabulal, B., Dan, M., Kurup, R., Pradeep, N. S., Valsamma, R. K., & George, V. (2007). Chemical composition and antimicrobial activities of the essential oils from the rhizomes of four Hedychium species from South India. Journal of Essential Oil Research, 19(1), 93–97. https://doi.org/10.1080/10412905.2007.9699241
39. Sarikurkcu, C., Tepe, B., Karaman, I., & Cetin, M. (2015). Evaluation of antioxidant properties and phenolic composition of Nepeta cataria L. and Nepeta nuda L. subsp. nuda. Industrial Crops and Products, 63, 44–51. https://doi.org/10.1016/j.indcrop.2014.10.009
40. Sienkiewicz, M., Łysakowska, M., Ciećwierz, J., & Kowalczyk, E. (2011). The antimicrobial activity of the essential oil of Nepeta cataria L. and its influence on selected microbial enzymes. Phytotherapy Research, 25(11), 1632–1638. https://doi.org/10.1002/ptr.3463
41. Singh, A. K., & Singh, R. (2018). Acute and chronic anti-inflammatory study of ethanolic extract of Ochna obtusata leaves. World Journal of Pharmaceutical Research, 7(10), 1456–1467.
42. Spellenberg, R. (2001). National Audubon Society field guide to North American wildflowers: Western region (2nd ed.). Knopf.
43. Srimal, R. C., Sharma, S. C., & Tandon, J. S. (1984). Anti-inflammatory and other pharmacological properties of Hedychium spicatum (Buch-Hem). Indian Journal of Pharmacology, 16(3), 143–147.
44. Tandon, V. R., & Gupta, R. K. (2005). Effect of Hedychium spicatum on experimental hyperglycemia and insulin secretion in albino rats. Journal of Medicinal and Aromatic Plant Sciences, 27(2), 272–276.
45. The Plant List. (2023). Hedychium spicatum Buch.-Ham. ex D.Don. Retrieved from http://www.theplantlist.org/tpl1.1/record/kew-278327
46. Tucker, A. O., & DeBaggio, T. (2009). The encyclopedia of herbs: A comprehensive reference to herbs of flavor and fragrance. Timber Press.
47. USDA. (2023). Hedychium spicatum Buch.-Ham. ex D.Don. In Plants Database. United States Department of Agriculture. Retrieved from https://plants.usda.gov/home
48. Ved, D. K., Suresh Kumar, K., Barve, V., Ahuja, V., & Latha, R. (2015). Conservation assessment and management plan for selected medicinal plants of Western Ghats, India. Foundation for Revitalisation of Local Health Traditions.
49. Werker, E., Ravid, U., & Putievsky, E. (1985). Structure of glandular hairs and identification of the main components of their secreted material in some species of the Labiatae. Israel Journal of Botany, 34(1), 31–45. https://doi.org/10.1080/0021213X.1985.10677010
50. Wood, T. H. (1994). The genus Hedychium in Thailand. Natural History Bulletin of the Siam Society, 42, 35–51.

HOW TO CITE: Saurabh Pratap Singh*, Dr. Shobhit Sirvastava, A Review on Hedychium spicatum: Phytochemistry, Pharmacological Activities, and Therapeutic Prospects, Int. J. of Pharm. Sci., 2025, Vol 3, Issue 11, 4288-4297 https://doi.org/10.5281/zenodo.17730987